

Supplement of Atmos. Meas. Tech., 11, 4509–4529, 2018
<https://doi.org/10.5194/amt-11-4509-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

The importance of surface reflectance anisotropy for cloud and NO₂ retrievals from GOME-2 and OMI

Alba Lorente et al.

Correspondence to: Alba Lorente (alba.lorentedelgado@wur.nl)


The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Figure S1: Google Earth image of the Amazon region (Lat.: 5N-10S and Lon.: 70W-60W) analysed in the manuscript.



30

Figure S2: Monthly averaged BRDF parameters (f_{iso} , f_{vol} , f_{geo}) from MODIS Band 3 (459-479 nm) averaged over the Amazon in a $0.25^\circ \times 0.25^\circ$ grid for March. BRDF parameters are from a 16-year climatology (2000-2016) created by the QA4ECV land group.



35 **Figure S3:** Same as Fig. S2 but for Band2 (841 - 876 nm)




Figure S4. Box-plot of simulated cloud fractions with a Lambertian surface (blue) and with surface BRDF (green) for GOME-2A East, Nadir and West measurements over Amazonia for March 2008. Triangles and circles show the mean values and the box represents 25th and 75th percentiles and the dashed lines the minimum and maximum values.




Figure S5. Relative differences between total tropospheric NO₂ AMF computed with surface BRDF and with a Lambertian surface as a function of cloud fraction for (a) backward scattering direction and (b) forward scattering direction for $(\theta, \theta_0) = (30^\circ, 45^\circ)$ for a moderately polluted (circles) and unpolluted (squares) troposphere. Surface BRDF parameters are $(f_{\text{iso}}, f_{\text{vol}}, f_{\text{geo}}) = (0.04, 0.03, 0.008)$ and $A_{\text{ws}} = 0.036$ for the Lambertian surface.

