



Corrigendum to

“Microwave Radar/radiometer for Arctic Clouds (MiRAC): first insights from the ACLOUD campaign” published in Atmos. Meas. Tech., 12, 5019–5037, 2019

Mario Mech¹, Leif-Leonard Kliesch¹, Andreas Anhäuser¹, Thomas Rose², Pavlos Kollias^{1,3}, and Susanne Crewell¹

¹Institute for Geophysics and Meteorology, University of Cologne, Cologne, Germany

²Radiometer-Physics GmbH, Meckenheim, Germany

³School of Marine and Atmospheric Sciences, Stony Brook University, NY, USA

Correspondence: Mario Mech (mario.mech@uni-koeln.de)

Published: 15 June 2021

We would like to inform you that there was a mistake in Fig. 7 of the published article. The x axes of the panel for 89 GHz were reversed by mistake in the originally submitted manuscript.

On the next page we show the correct figure.




Figure 7. Vertical cross section of Z_e measured by CloudSat CPR (top) and the MiRAC radar on Polar 5 (second row) for the satellite underflight on 27 May 2017 between 10:06 and 10:44 UTC along the black dashed line in Fig. 4. Grey shaded areas define the zone of reduced sensitivity. The third row gives the sea ice coverage based on AMSR2 observations along the flight track. Rows four to six show the passive radiometer measurements at 89 GHz from MiRAC-A and those channels of MiRAC-P, i.e., the six channels along the 183.31 GHz water vapor absorption line, and the two channels at 243 and 340 GHz.