

Supplement of Atmos. Meas. Tech., 12, 873–890, 2019
<https://doi.org/10.5194/amt-12-873-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Tropospheric water vapor profiles obtained with FTIR: comparison with balloon-borne frost point hygrometers and influence on trace gas retrievals

Ivan Ortega et al.

Correspondence to: Ivan Ortega (iortega@ucar.edu) and James W. Hannigan (jamesw@ucar.edu)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.




Figure S1. The HR-FTIR horizontal location (air mass sensitivity location) calculated with the water vapor center of gravity, solar zenith angle, and solar azimuth angle of all measurements (black dots) at BLD (left) and MLO (right). The HR-FTIR ground-based location is indicated with the red dots and the launch of the sonde in blue dots.




Figure S2. Distributions of the difference between WP retrieved and FPH at different altitude layers using ERA-d as a priori. A Gauss-type normal distribution is constructed for each panel using the bins, mean values (blue dotted vertical lines) and standard deviation (red dotted vertical lines). Dotted black vertical lines represent the percentile 95 and green dotted lines are the median.




Figure S3. Percentile 95th of the difference between retrieval and sonde in VMR as a function of a priori source for different layers.