



Supplement of

Intercomparison of upper tropospheric and lower stratospheric water vapor measurements over the Asian Summer Monsoon during the StratoClim campaign

Clare E. Singer et al.

Correspondence to: Clare E. Singer (csinger@caltech.edu) and Elisabeth J. Moyer (moyer@uchicago.edu)

The copyright of individual parts of the supplement might differ from the article licence.




Figure S1. Time series of water vapor for each of the six flights (F2-F4 and F6-F8). Altitude (green) and in situ H_2O measurements from ChiWIS (black/grey), FLASH (blue), and clear-sky FISH (pink). Periods where ChiWIS cell pressure is just out of regulation (20 – 30 mbar), are shown in grey. There is impressive agreement between the three hygrometers. ChiWIS shows much higher precision than the other two instruments, which is evident from the sharper temporal structures in the time series. Nevertheless, all three instruments are able to capture the same fine-scale variability of the atmosphere.




Figure S2. Vertical profiles of water vapor against potential temperature for each of the six flights (F2-F4 and F6-F8). Shown are in situ H₂O measurements from ChiWIS (black/grey), FLASH (blue), and clear-sky FISH (pink), with periods where ChiWIS cell pressure is just out of regulation (20 – 30 mbar) shown in grey.




Figure S3. Time series of relative humidity for each of the six flights (F2-F4 and F6-F8). Shown are altitude (green) and in situ RH_{ice} measurements from ChiWIS (black/grey), FLASH (blue), and clear-sky FISH (pink) with periods where ChiWIS cell pressure is just out of regulation (20 – 30 mbar) shown in grey.




Figure S4. Vertical profiles of relative humidity against potential temperature for each of the six flights (F2-F4 and F6-F8). Shown are in situ RH_{ice} measurements from ChiWIS (black/grey), FLASH (blue), and clear-sky FISH (pink) with periods where ChiWIS cell pressure is just out of regulation (20 – 30 mbar) shown in grey.




Figure S5. Estimated in-flight precision of the three hygrometers based on a segment from flight 4 with approximately constant H₂O. Shown in colored dots are the H₂O values measured by FLASH, FISH, and ChiWIS (blue, pink, black), the mean value for this period (transparent lines), and the standard deviation of the measurements (plotted error bars with labeled values).




Figure S6. Normalized PDF of percentage differences between ChiWIS (grey) and FISH (colors) from FLASH during each flight (rows). FISH did not report measurements on F3. The first column (leftmost) shows values from all conditions. The second column shows percentage difference in clear-sky periods with zero time lag, $\tau_c = 0$. Clear-sky is defined as regions where $BR < 1.2$ or $N_{ice} = 0 \text{ cm}^{-3}$. Each subsequent column shows the percentage difference in clear-sky periods with increasing $\tau_c = 5, 10, 20$. Note that the last column (rightmost) has a smaller x-axis range.




Figure S7. The left column (a,c) reproduces Fig. 3b,e showing a point-by-point comparison of clear-sky H_2O measured by ChiWIS and FLASH during periods when the ChiWIS optical cell pressure is ≥ 30 mbar (well regulated). The right column (b,d) shows the same comparison between clear-sky ChiWIS and FLASH measurements, but exclusively for periods when the ChiWIS optical cell pressure is slightly unregulated (between 20 and 30 mbar). Points in (a,b) are colored by flight number. Frequency of counts (c,d) in each 0.1 by 0.1 ppmv bin are shown in color. The number of hours of measurement for each category are shown in the bottom right corner of each panel. For clear-sky periods, 56% of measurements were made when the cell pressure was within regulation. For in-cloud periods, nearly 100% of measurements were made when the cell pressure was within regulation (not shown). The one-to-one line is plotted in solid black with dashed (dotted) lines indicated $\pm 10\%$ ($\pm 20\%$) differences. The percentage bias and r^2 coefficients are shown above each panel.




Figure S8. Normalized frequency distribution of ChiWIS measurements in clear-sky at different temperatures and relative humidities. (a) Reproduces Fig. 7a, showing measurements when the ChiWIS optical cell pressure is ≥ 30 mbar. (b) Shows measurements when the cell pressure is ≥ 20 mbar. The measurements are binned by 0.5 K in temperature and 0.05 in relative humidity. The total hours of measurement is shown in the bottom left of each panel. The solid black lines shows $RH_{ice} = 1$ and the dotted line shows the homogeneous ice nucleation threshold from Koop et al. (2000).




Figure S9. The left column (a,c) reproduces Fig. 8b,e showing a point-by-point comparison of clear-sky RH_{ice} measured by ChiWIS and FLASH during periods when the ChiWIS optical cell pressure is ≥ 30 mbar (well regulated). The right column (b,d) shows the same comparison between clear-sky ChiWIS and FLASH measurements, but exclusively for periods when the ChiWIS optical cell pressure is slightly unregulated (between 20 and 30 mbar). Points in (a,b) are colored by flight number. Frequency of counts (c,d) in each 0.03 by 0.03 bin are shown in color. The number of hours of measurement for each category are shown in the bottom right corner of each panel. For clear-sky periods, 56% of measurements were made when the cell pressure was within regulation. For in-cloud periods, nearly 100% of measurements were made when the cell pressure was within regulation (not shown). The one-to-one line is plotted in black. The percentage bias and r^2 coefficients are shown above each panel.

References

Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, <https://doi.org/10.1038/35020537>, 2000.