



Supplement of

**Towards harmonized measurements of condensable vapors:
insights from the intercomparison of six chemical ionization mass
spectrometers at a boreal forest site**

Cecilia Righi et al.

Correspondence to: Cecilia Righi (cecilia.righi@helsinki.fi) and Nina Sarnela (nina.sarnela@helsinki.fi)

The copyright of individual parts of the supplement might differ from the article licence.

S1 Abstract – description of the supporting material

This Supporting Information provides additional material to clarify and extend the analysis presented in the manuscript text, including supporting figures and tables. It is organized into Sections S1–S16: S1 Abstract – description of the supporting material, S2 ACTRIS CI-FI1 campaign timeline, S3 SmartSMEAR metadata, S4 Atmospheric Pressure interface Time-of-Flight mass spectrometer, S5 Peak lists, S6 Inlet tube losses, S7 Sulfuric acid calibration setup, S8 Equations used for the analysis of bromide mode data, S9 Background measurements, S10 Considerations on the application of the transmission correction factor, S11 Measurements in nitrate mode – additional species, S12 Measurements in nitrate mode – correlation analysis, S13 Measurements in bromide mode – calibration factor, S14 Measurements in bromide mode – correlation analysis, S15 Measurements in different reagent ion modes: nitrate, bromide, and iodide – additional species, S16 References.

S2 ACTRIS CI-FI1 campaign timeline

Figure S1 shows a timeline of the ACTRIS CI-FI1 campaign.




Figure S1 Schematic timeline of the ACTRIS CI-FI1 campaign. The horizontal bars indicate the time span during which each instrument was employed and are colored for better contrast. The figure highlights the timing of sulfuric acid calibrations (magenta), transmission calibrations (green), and background measurements (grey), performed during the campaign. Transmission calibration on ML2 was performed post-campaign in the ACTRIS Laboratory of the University of Helsinki.

S3 SmartSMEAR metadata

Table S1 provides metadata for the meteorological data retrieved from the SmartSMEAR database (<https://smear.avaa.csc.fi/>, last access: 13 January 2026) for the SMEAR II station during the measurement campaign period. These data were used to support the interpretation of the measurements from the instruments used in this study.

Table S1 Metadata of the meteorological variables retrieved from the SmartSMEAR database for the SMEAR II station during the campaign period.

title	Air temperature 16.8 m (2)	Relative humidity 16.8 m	Precipitation	UVB radiation
tableId	HYY_META	HYY_META	HYY_META	HYY_META
name	Tmm168	RH168	Precipacc	UV_B
description	Air temperature at 16.8 m height measured with Pt100 inside Rotronic RS12T/RS24T radiation shield, inconsistent with the old 4.2-67.2 m temperature profile	Relative humidity at 16.8 m height	Precipitation (liquid water equivalent) accumulated during previous 1 min	Ultraviolet B radiation in wavelength range 280-315 nm at 18 m height (radiation tower 1997-2/2017) or 35 m height (35 m tower 2/2017-), data until 2023 converted from erythemally weighted UVB to unweighted UVB with factor 7.55, high relative uncertainty at solar elevation angle of <20 degrees
unit	°C	%	mm	W m ⁻²
source	Pt100 inside Rotronic shield	Rotronic MP102H RH sensor, mastmet33	Vaisala FD12P weather sensor at 18 m height	Solar Light SL501A (-2023) or K&Z SUV-B (2024-)
periodStart	2013-01-22T00:00:00.0	2013-01-22T15:31:00.0	2005-04-01T00:00:00.0	1997-03-07T17:21:00.0
periodEnd	null	null	null	null
coverage	0	0	0	0
rights	public	public	public	public
mandatory	FALSE	FALSE	FALSE	FALSE
derivative	FALSE	FALSE	FALSE	FALSE
timestamp	2020-06-30T11:31:43	2020-12-22T13:33:07	2021-03-08T07:49:55	2024-10-04T12:41:42
category	Meteorology	Meteorology	Meteorology	Radiation

S4 Atmospheric Pressure interface Time-of-Flight mass spectrometer

The Atmospheric Pressure interface Time-of-Flight (APi-ToF) mass spectrometer is a high-resolution instrument capable of detecting ions sampled at atmospheric pressure (Junninen et al., 2010). Once the sampled ions enter the APi section through a 0.3 mm pinhole at a flow rate of $\sim 0.8 \text{ L min}^{-1}$, neutral molecules are gradually pumped away, while the ions are focused at the center of the flow and guided from atmospheric pressure to the high vacuum of the ToF chamber ($\sim 10^{-6}$ mbar) via three differentially pumped stages. This gradual pressure drop helps prevent the loss of ions that could result from abrupt decompression. In the ToF chamber, ions are accelerated by a pulsed electric field, and their m/z are determined by measuring the time required to traverse a field-free region and reach the detector. Detection is performed using two microchannel plates (MCPs), each consisting of a dense array of microscopic channels that act as electron multipliers: when ions collide with the MCP surface, they generate an initial pulse of electrons, which then cascade through successive collisions along the channel walls, amplifying the signal. The resulting electron current is collected at the anode. Finally, data acquisition is carried out using either a time-to-digital converter, which records the arrival time of each ion exceeding a certain threshold, or an analog-to-digital converter, which continuously measures the ion signal intensity.

S5 Peak lists

This section presents the peak lists used for the comparison of the CI-APi-ToFs during the measurement campaign. Separate tables are provided for instruments operating in nitrate (NO_3^-), bromide (Br^-), and iodide (I^-) mode. In addition, tables listing the additional peaks included to account for the enhanced sensitivity of bromide toward less oxidized compounds are provided for both ionization modes. For convenience, nitrophenol is included within the group of strong organic and inorganic acids (HA), although it should be noted that it is not strictly a strong acid, given its relatively high pK_a (~ 7.15). Nevertheless, nitrophenol can be efficiently detected in NO_3^- mode and is commonly regarded as a marker compound for biomass burning.

Table S2 Peak list selected for the instruments operating in nitrate (NO_3^-) mode. The monoisotopic mass (Th) corresponds to the exact mass of the detected ion. The last column indicates the species or chemical group assigned to each peak. In addition to highly oxygenated organic molecules (HOMs), semi-volatile organic compounds (SVOCs) were included to account for less oxidized organic compounds observed in all the reagent ion modes considered. The volatility of HOMs and SVOCs was estimated using the parametrization of Donahue et al. (2011), updated by Mohr et al. (2019), defined at reference temperature of 300 K.

Ion	Monoisotopic Mass (Th)	Volatility	Species / Group
NO_3^-	61.988		reagent ion
$(\text{H}_2\text{O}) \cdot \text{NO}_3^-$	79.999		reagent ion
CH_3SO_3^-	94.981		methanesulfonic acid / HA
HSO_4^-	96.960		sulfuric acid

$C_3H_3O_4^-$	103.004		malonic acid / HA
$(HNO_3) \cdot NO_3^-$	124.984		reagent ion
$C_6H_4NO_3^-$	138.020		nitrophenol / HA
$(CH_4SO_3) \cdot NO_3^-$	157.976		methanesulfonic acid / HA
$(HNO_3) \cdot HSO_4^-$	159.956		sulfuric acid
$(C_3H_4O_4) \cdot NO_3^-$	165.999		malonic acid / HA
IO_3^-	174.890		iodic acid / HA
$(HNO_3)_2 \cdot NO_3^-$	187.980		reagent ion
$(H_2SO_4) \cdot HSO_4^-$	194.927		sulfuric acid
$(C_6H_5NO_3) \cdot NO_3^-$	201.015		nitrophenol / HA
$(HIO_3) \cdot NO_3^-$	237.885		iodic acid / HA
$(C_{10}H_{14}O_3) \cdot NO_3^-$	244.083	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{16}O_3) \cdot NO_3^-$	246.098	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{14}O_4) \cdot NO_3^-$	260.078	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{15}O_4) \cdot NO_3^-$	261.085	SVOC	organic peroxy radicals / SVOC
$(C_{10}H_{16}O_4) \cdot NO_3^-$	262.093	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{16}O_5) \cdot NO_3^-$	278.088	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{14}O_6) \cdot NO_3^-$	292.067	SVOC	closed shell non-nitrate monomers / SVOC
$(H_2SO_4)_2 \cdot HSO_4^-$	292.895		sulfuric acid
$(C_{10}H_{15}O_6) \cdot NO_3^-$	293.075	SVOC	organic peroxy radicals / SVOC
$(C_{10}H_{16}O_6) \cdot NO_3^-$	294.083	SVOC	closed shell non-nitrate monomers / SVOC
$(C_{10}H_{14}O_7) \cdot NO_3^-$	308.062	LVOC	closed shell non-nitrate monomers / HOM
$(C_{10}H_{16}O_7) \cdot NO_3^-$	310.078	LVOC	closed shell non-nitrate monomers / HOM
$(C_{10}H_{14}O_8) \cdot NO_3^-$	324.057	LVOC	closed shell non-nitrate monomers / HOM
$(C_{10}H_{15}O_8) \cdot NO_3^-$	325.065	LVOC	organic peroxy radicals / HOM
$(C_{10}H_{16}O_8) \cdot NO_3^-$	326.073	LVOC	closed shell non-nitrate monomers / HOM
$(C_{10}H_{15}O_8N) \cdot NO_3^-$	339.068	LVOC	closed shell organonitrate monomers / HOM
$(C_{10}H_{14}O_9) \cdot NO_3^-$	340.052	LVOC	closed shell non-nitrate monomers / HOM
$(C_{10}H_{16}O_9) \cdot NO_3^-$	342.068	LVOC	closed shell non-nitrate monomers / HOM

$(\text{C}_{10}\text{H}_{15}\text{O}_9\text{N}) \cdot \text{NO}_3^-$	355.063	LVOC	closed shell organonitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_{10}) \cdot \text{NO}_3^-$	356.047	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_{10}) \cdot \text{NO}_3^-$	357.055	LVOC	organic peroxy radicals / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_{10}) \cdot \text{NO}_3^-$	358.063	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_{11}) \cdot \text{NO}_3^-$	372.042	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_{11}) \cdot \text{NO}_3^-$	374.058	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_{11}\text{N}) \cdot \text{NO}_3^-$	387.053	LVOC	closed shell organonitrate monomers / HOM
$(\text{C}_{19}\text{H}_{28}\text{O}_{11}) \cdot \text{NO}_3^-$	494.152	ULVOC	dimers / HOM
$(\text{C}_{19}\text{H}_{28}\text{O}_{12}) \cdot \text{NO}_3^-$	510.146	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{12}) \cdot \text{NO}_3^-$	524.162	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{32}\text{O}_{13}) \cdot \text{NO}_3^-$	542.173	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{31}\text{O}_{13}\text{N}) \cdot \text{NO}_3^-$	555.168	ULVOC	nitrate dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{14}) \cdot \text{NO}_3^-$	556.152	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{32}\text{O}_{15}) \cdot \text{NO}_3^-$	574.162	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{16}) \cdot \text{NO}_3^-$	588.142	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{18}) \cdot \text{NO}_3^-$	620.132	ULVOC	dimers / HOM

60 **Table S3** Peak list selected for the instruments operating in bromide (Br^-) mode. The monoisotopic mass (Th) corresponds to the exact mass of the detected ion. The last column indicates the species or chemical group assigned to each peak. In addition to highly oxygenated organic molecules (HOMs), semi-volatile organic compounds (SVOCs) were included to account for less oxidized organic compounds observed in all the reagent ion modes considered. The volatility of HOMs and SVOCs was estimated using the parametrization of Donahue et al. (2011), updated by Mohr et al. (2019), defined at reference temperature of 300 K.

Ion	Monoisotopic Mass (Th)	Volatility	Species / Group
Br^-	78.919		reagent ion
CH_3SO_3^-	94.981		methanesulfonic acid / HA
$(\text{H}_2\text{O}) \cdot \text{Br}^-$	96.929		reagent ion
HSO_4^-	96.960		sulfuric acid
$\text{C}_3\text{H}_3\text{O}_4^-$	103.004		malonic acid / HA
$\text{C}_6\text{H}_4\text{NO}_3^-$	138.020		nitrophenol / HA
IO_3^-	174.890		iodic acid / HA
$(\text{CH}_4\text{SO}_3) \cdot \text{Br}^-$	174.907		methanesulfonic acid / HA

$(\text{H}_2\text{SO}_4) \cdot \text{Br}^-$	176.886		sulfuric acid
$(\text{C}_3\text{H}_4\text{O}_4) \cdot \text{Br}^-$	182.930		malonic acid / HA
$(\text{C}_6\text{H}_5\text{NO}_3) \cdot \text{Br}^-$	217.946		nitrophenol / HA
$(\text{HIO}_3) \cdot \text{Br}^-$	254.816		iodic acid / HA
$(\text{C}_{10}\text{H}_{14}\text{O}_3) \cdot \text{Br}^-$	261.013	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_3) \cdot \text{Br}^-$	263.029	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{14}\text{O}_4) \cdot \text{Br}^-$	277.008	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{15}\text{O}_4) \cdot \text{Br}^-$	278.016	SVOC	organic peroxy radicals / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_4) \cdot \text{Br}^-$	279.024	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_5) \cdot \text{Br}^-$	295.019	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{14}\text{O}_6) \cdot \text{Br}^-$	308.998	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{15}\text{O}_6) \cdot \text{Br}^-$	310.006	SVOC	organic peroxy radicals / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_6) \cdot \text{Br}^-$	311.014	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{14}\text{O}_7) \cdot \text{Br}^-$	324.993	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_7) \cdot \text{Br}^-$	327.008	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_8) \cdot \text{Br}^-$	340.988	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_8) \cdot \text{Br}^-$	341.996	LVOC	organic peroxy radicals / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_8) \cdot \text{Br}^-$	343.003	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_8\text{N}) \cdot \text{Br}^-$	355.999	LVOC	closed shell organonitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_9) \cdot \text{Br}^-$	356.983	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_9) \cdot \text{Br}^-$	358.998	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_9\text{N}) \cdot \text{Br}^-$	371.994	LVOC	closed shell organonitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_{10}) \cdot \text{Br}^-$	372.978	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_{10}) \cdot \text{Br}^-$	373.985	LVOC	organic peroxy radicals / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_{10}) \cdot \text{Br}^-$	374.993	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{14}\text{O}_{11}) \cdot \text{Br}^-$	388.972	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{16}\text{O}_{11}) \cdot \text{Br}^-$	390.988	LVOC	closed shell non-nitrate monomers / HOM
$(\text{C}_{10}\text{H}_{15}\text{O}_{11}\text{N}) \cdot \text{Br}^-$	403.983	LVOC	closed shell organonitrate monomers / HOM
$(\text{C}_{19}\text{H}_{28}\text{O}_{11}) \cdot \text{Br}^-$	511.082	ULVOC	dimers / HOM

$(\text{C}_{19}\text{H}_{28}\text{O}_{12}) \cdot \text{Br}^-$	527.077	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{12}) \cdot \text{Br}^-$	541.093	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{32}\text{O}_{13}) \cdot \text{Br}^-$	559.103	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{31}\text{O}_{13}\text{N}) \cdot \text{Br}^-$	572.098	ULVOC	nitrate dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{14}) \cdot \text{Br}^-$	573.082	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{32}\text{O}_{15}) \cdot \text{Br}^-$	591.093	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{16}) \cdot \text{Br}^-$	605.072	ULVOC	dimers / HOM
$(\text{C}_{20}\text{H}_{30}\text{O}_{18}) \cdot \text{Br}^-$	637.062	ULVOC	dimers / HOM

65 **Table S4** Peak list selected for the instruments operating in iodide (I^-) mode. The monoisotopic mass (Th) corresponds to the exact mass of the detected ion. The last column indicates the species or chemical group assigned to each peak. The volatility of the organic compounds was estimated using the parametrization of Donahue et al. (2011), updated by Mohr et al. (2019), defined at reference temperature of 300 K.

Ion	Monoisotopic Mass (Th)	Volatility	Species / Group
$(\text{C}_{10}\text{H}_{14}\text{O}_3) \cdot \text{I}^-$	308.999	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_3) \cdot \text{I}^-$	311.015	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{14}\text{O}_4) \cdot \text{I}^-$	324.994	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_4) \cdot \text{I}^-$	327.010	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{16}\text{O}_5) \cdot \text{I}^-$	343.005	SVOC	closed shell non-nitrate monomers / SVOC
$(\text{C}_{10}\text{H}_{14}\text{O}_6) \cdot \text{I}^-$	356.984	SVOC	closed shell non-nitrate monomers / SVOC

S6 Inlet tube losses

70 Sulfuric acid wall losses are already accounted for in the model used to calculate sulfuric acid concentrations in the calibration experiment (Kürten et al., 2012); therefore, no additional correction factor is required when deriving concentrations from ion count rates. For species other than sulfuric acid, however, a factor C_{inlet} was introduced to account for losses in the inlet tube, calculated following the steps outlined below.

1. A gas diffusion coefficient, D_g ($\text{cm}^2 \text{s}^{-1}$), is calculated for each compound as:

$$D_{AB} = \frac{1 \times 10^{-3} \cdot T^{1.75} \cdot \sqrt{\frac{1}{M_A} + \frac{1}{M_B}}}{p \cdot \left(\sqrt[3]{\sum_A v_i} + \sqrt[3]{\sum_B v_i} \right)^2}, \quad \text{Eq. (S1)}$$

where T (K) is the temperature, p (atm) is the pressure, M_{AB} (g mol⁻¹) are the molecular mass of air (A) and the species of interest (B), v_i (cm⁻³) is the atomic diffusion volume to be summed over the atoms/groups of each diffusing species. v_i values have been derived from Fuller et al. (1969).

2. The frequency of wall collision, k (s⁻¹) is calculated for each species using the following equation assuming a first-order loss process:

$$k = 3.6 \cdot \frac{D_{AB}}{r_{in}^2}, \quad \text{Eq. (S2)}$$

where r_{in} (cm) is the radius of the inlet tube. The factor 3.6 is the geometric factor accounting for diffusion-limited transport to the wall of a cylindrical tube under laminar-flow conditions (Tang et al., 2014).

3. The residence time of the gas inside the inlet tube, Rt_{in} (s), is calculated as:

$$Rt_{in} = \frac{\pi \cdot r_{in}^2 \cdot l_{in}}{\frac{Q_{in}}{60} \cdot 10^3}, \quad \text{Eq. (S3)}$$

where l_{in} (cm) and Q_{in} (L min⁻¹) are the inlet tube length and flow rate, respectively.

4. The inlet tube loss is calculated for each species as:

$$L_{in} = 1 - e^{-k \cdot Rt_{in}}. \quad \text{Eq. (S4)}$$

5. Finally, the loss correction factor is calculated as:

$$C_{inlet} = \frac{1}{1 - L_{in}}. \quad \text{Eq. (S5)}$$

Table S5 shows the C_{inlet} values calculated for the CI-API-ToF mass spectrometers participating in the ACTRIS CI-FI1 campaign for the species included in the nitrate peak list. For sulfuric acid, the C_{inlet} factor is equal to unity, since the losses to the wall of the inlet tube are already included in the factor C_{SA} for this species.

Table S5 C_{inlet} values calculated for the species included in the nitrate peak list for each CI-API-ToF mass spectrometer participating in the ACTRIS CI-FI1 campaign.

Ion	Instrument				
	EH1	EH2	EL	ML1	ML2
CH ₃ SO ₃ ⁻	1.643	1.800	1.368	1.318	1.256
C ₃ H ₃ O ₄ ⁻	1.571	1.707	1.330	1.286	1.230
C ₆ H ₄ NO ₃ ⁻	1.432	1.530	1.254	1.221	1.179
(CH ₄ SO ₃) · NO ₃ ⁻	1.512	1.631	1.298	1.259	1.209
(C ₃ H ₄ O ₄) · NO ₃ ⁻	1.471	1.579	1.276	1.240	1.194

IO_3^-	1.690	1.861	1.393	1.339	1.272
$(\text{C}_6\text{H}_5\text{NO}_3) \cdot \text{NO}_3^-$	1.381	1.465	1.226	1.197	1.160
$(\text{HIO}_3) \cdot \text{NO}_3^-$	1.547	1.676	1.317	1.275	1.222
$(\text{C}_{10}\text{H}_{14}\text{O}_3) \cdot \text{NO}_3^-$	1.302	1.366	1.181	1.158	1.129
$(\text{C}_{10}\text{H}_{16}\text{O}_3) \cdot \text{NO}_3^-$	1.298	1.362	1.179	1.156	1.127
$(\text{C}_{10}\text{H}_{14}\text{O}_4) \cdot \text{NO}_3^-$	1.297	1.360	1.178	1.156	1.127
$(\text{C}_{10}\text{H}_{15}\text{O}_4) \cdot \text{NO}_3^-$	1.295	1.358	1.177	1.155	1.126
$(\text{C}_{10}\text{H}_{16}\text{O}_4) \cdot \text{NO}_3^-$	1.293	1.356	1.176	1.154	1.125
$(\text{C}_{10}\text{H}_{16}\text{O}_5) \cdot \text{NO}_3^-$	1.289	1.350	1.174	1.152	1.123
$(\text{C}_{10}\text{H}_{14}\text{O}_6) \cdot \text{NO}_3^-$	1.287	1.348	1.173	1.151	1.123
$(\text{C}_{10}\text{H}_{15}\text{O}_6) \cdot \text{NO}_3^-$	1.286	1.346	1.172	1.150	1.122
$(\text{C}_{10}\text{H}_{16}\text{O}_6) \cdot \text{NO}_3^-$	1.284	1.344	1.171	1.149	1.122
$(\text{C}_{10}\text{H}_{14}\text{O}_7) \cdot \text{NO}_3^-$	1.283	1.343	1.170	1.149	1.121
$(\text{C}_{10}\text{H}_{16}\text{O}_7) \cdot \text{NO}_3^-$	1.280	1.339	1.169	1.147	1.120
$(\text{C}_{10}\text{H}_{14}\text{O}_8) \cdot \text{NO}_3^-$	1.278	1.337	1.168	1.147	1.119
$(\text{C}_{10}\text{H}_{15}\text{O}_8) \cdot \text{NO}_3^-$	1.277	1.336	1.167	1.146	1.119
$(\text{C}_{10}\text{H}_{16}\text{O}_8) \cdot \text{NO}_3^-$	1.276	1.334	1.166	1.145	1.118
$(\text{C}_{10}\text{H}_{15}\text{O}_8\text{N}) \cdot \text{NO}_3^-$	1.274	1.332	1.165	1.144	1.118
$(\text{C}_{10}\text{H}_{14}\text{O}_9) \cdot \text{NO}_3^-$	1.275	1.332	1.165	1.145	1.118
$(\text{C}_{10}\text{H}_{16}\text{O}_9) \cdot \text{NO}_3^-$	1.272	1.329	1.164	1.143	1.117
$(\text{C}_{10}\text{H}_{15}\text{O}_9\text{N}) \cdot \text{NO}_3^-$	1.270	1.327	1.163	1.143	1.116
$(\text{C}_{10}\text{H}_{14}\text{O}_{10}) \cdot \text{NO}_3^-$	1.271	1.328	1.163	1.143	1.116
$(\text{C}_{10}\text{H}_{15}\text{O}_{10}) \cdot \text{NO}_3^-$	1.269	1.326	1.163	1.142	1.116
$(\text{C}_{10}\text{H}_{16}\text{O}_{10}) \cdot \text{NO}_3^-$	1.268	1.325	1.162	1.141	1.115
$(\text{C}_{10}\text{H}_{14}\text{O}_{11}) \cdot \text{NO}_3^-$	1.267	1.323	1.161	1.141	1.115
$(\text{C}_{10}\text{H}_{16}\text{O}_{11}) \cdot \text{NO}_3^-$	1.265	1.320	1.160	1.140	1.114
$(\text{C}_{10}\text{H}_{15}\text{O}_{11}\text{N}) \cdot \text{NO}_3^-$	1.263	1.319	1.159	1.139	1.113
$(\text{C}_{19}\text{H}_{28}\text{O}_{11}) \cdot \text{NO}_3^-$	1.204	1.246	1.124	1.109	1.089
$(\text{C}_{19}\text{H}_{28}\text{O}_{12}) \cdot \text{NO}_3^-$	1.202	1.244	1.123	1.108	1.088

$(\text{C}_{20}\text{H}_{30}\text{O}_{12}) \cdot \text{NO}_3^-$	1.198	1.238	1.121	1.106	1.086
$(\text{C}_{20}\text{H}_{32}\text{O}_{13}) \cdot \text{NO}_3^-$	1.195	1.235	1.119	1.104	1.085
$(\text{C}_{20}\text{H}_{31}\text{O}_{13}\text{N}) \cdot \text{NO}_3^-$	1.194	1.234	1.119	1.104	1.085
$(\text{C}_{20}\text{H}_{30}\text{O}_{14}) \cdot \text{NO}_3^-$	1.195	1.234	1.119	1.104	1.085
$(\text{C}_{20}\text{H}_{32}\text{O}_{15}) \cdot \text{NO}_3^-$	1.192	1.231	1.117	1.103	1.084
$(\text{C}_{20}\text{H}_{30}\text{O}_{16}) \cdot \text{NO}_3^-$	1.192	1.231	1.117	1.103	1.084
$(\text{C}_{20}\text{H}_{30}\text{O}_{18}) \cdot \text{NO}_3^-$	1.189	1.227	1.115	1.101	1.083

S7 Sulfuric acid calibration setup

100 The calibration setup (Figure S2) was built based on the one presented by Kürten et al. (2012), with a few key differences. First, the dimensions of the box in our setup are $14 \times 14 \times 8$ cm (L $\times$ W $\times$ H), with a 3/4"-diameter tube passing through the box, whereas Kürten et al. (2012) used a box of $30 \times 20 \times 15$ cm and a 1/2"-diameter tube. Second, our setup does not include a photodiode for monitoring lamp stability.

105 The aluminum box houses a Pen-Ray® Hg Lamp (Analytik Jena US LLC), which serves as the UV source responsible for the photolysis of H_2O vapor and the in-situ production of H_2SO_4 via the following reactions:



110 Four mass flow controllers (MFCs; Alicat Scientific MC-series) were used to regulate the inflow rates of sulfur dioxide, synthetic air, and nitrogen. The nitrogen flow was first split into three separate lines: (1) a carrier gas line, (2) a line for H_2O vapor production, and (3) a line for flushing the Hg lamp. The carrier gas and H_2O vapor lines were each regulated by a MFC, while the Hg-lamp flush was adjusted using a needle valve. To generate water vapor, the N_2 flow from the second MFC passed through a bottle containing Milli-Q H_2O , which served as a humidifier. All tubing was made of Teflon® with a diameter of 6 mm, except for the 3/4"-diameter stainless steel tube passing through the box.

115 When connecting the calibration setup to the CI-API-ToF, the nitrogen flush to the aluminum box was set to 1.5 L min^{-1} , and the lamp was switched on and allowed to warm up for approximately 90 min. After this period, the gas flow rates were adjusted via the MFCs so that their sum slightly exceeded the flow drawn into the inlet. Each calibration step lasted 10 minutes, after which the incoming wet nitrogen flow was modified while keeping the other parameters constant. The calibration experiment consisted of five steps, repeated twice. In the calibration experiments conducted during the ACTRIS
120 CI-FI1 campaign, modeled sulfuric acid concentrations ranged from $5 \cdot 10^7$ to $1 \cdot 10^9$ molecules cm^{-3} .

Following Kürten et al. (2012), chemical actinometry was used to estimate the I^+ -product needed for modeling H_2SO_4 concentrations. For this study, the actinometry experiment was conducted in January 2024 at the ACTRIS Laboratory of the University of Helsinki. Further instructions and best practices for sulfuric acid calibration and actinometry experiment are available on GitHub at https://github.com/ceciliarighi/ACTRIS_CiGas_condensable_vapors.

125




Figure S2 Schematic diagram of the calibration setup used for the sulfuric acid calibration during the measurement campaign.

S8 Equations used for the analysis of bromide mode data

Here we report the equations used to calculate the concentrations of sulfuric acid, acidic organic and inorganic compounds (HAs), and highly oxygenated organic molecules (HOMs) from bromide mode measurements. Equations (S6)–(S8) are analogous to Eqs. (3)–(5) presented in the main text for nitrate mode data. In addition, Eq. (S9) describes the calculation of concentrations for the semi-volatile organic compounds (SVOCs) group.

$$[\text{H}_2\text{SO}_4] = \frac{\text{HSO}_4^- + \text{H}_2\text{SO}_4 \cdot \text{Br}^-}{\text{Br}^- + \text{H}_2\text{O} \cdot \text{Br}^-} \times C'_{\text{SA}}, \quad \text{Eq. (S6)}$$

$$[\text{HA}] = \frac{\text{A}^- + \text{HA} \cdot \text{Br}^-}{\text{Br}^- + \text{H}_2\text{O} \cdot \text{Br}^-} \times C'_{\text{SA}} \times C_{\text{inlet}}, \quad \text{Eq. (S7)}$$

$$[\text{HOM}] = \frac{\text{HOM} \cdot \text{Br}^-}{\text{Br}^- + \text{H}_2\text{O} \cdot \text{Br}^-} \times C'_{\text{SA}} \times C_{\text{inlet}} \times C_{\text{trans}}, \quad \text{Eq. (S8)}$$

$$[\text{SVOC}] = \frac{\text{HOM} \cdot \text{Br}^-}{\text{Br}^- + \text{H}_2\text{O} \cdot \text{Br}^-} \times C'_{\text{SA}} \times C_{\text{inlet}} \times C_{\text{trans}}. \quad \text{Eq. (S9)}$$

As discussed in Section 2.2.5, Br^- ionization is not fully kinetically limited for HOMs, implying lower charging efficiency than for sulfuric acid and HAs compared to NO_3^- , which in turn exhibits lower sensitivity toward the less oxygenated SVOCs.

Equation (S10) is analogous to Eq. (1) in the manuscript text and was used to calculate the sulfuric acid signal normalized to the reagent ion signal during the sulfuric acid calibration experiments.

$$S_{H_2SO_4} = \frac{HSO_4^- + H_2SO_4 \cdot Br^-}{Br^- + H_2O \cdot Br^-} \tag{S10}$$

S9 Background measurements

To evaluate the effectiveness of the HEPA filter for suppressing the signals of the gas-phase compounds targeted in this study, measurements with and without the HEPA filter were compared, as shown in Figure S3 for instrument ML2.




Figure S3 Normalized ion counts for sulfuric acid, malonic acid, iodic acid, closed shell non-nitrate monomers, and closed shell organonitrate monomers before, during, and after a HEPA filter was connected to ML2 inlet tube. The shaded area indicates the period during which the HEPA filter was connected to the inlet.

S10 Considerations on the application of the transmission correction factor

Figure S4 visually supports the considerations discussed in Sect. 3.2 of the manuscript text, highlighting the effect of applying different correction factors to the data measured by the five CI-APi-ToFs operating in nitrate mode.




Figure S4 Concentrations of (a) closed shell organonitrate monomers, and (b) dimers measured in nitrate mode by five mass spectrometers during 1-2 August 2024. For each panel, the left plot shows concentration time series obtained by applying only the sulfuric acid calibration factor (C_{SA}); the central plot shows concentrations obtained by applying C_{SA} , inlet loss correction (C_{inlet}), and transmission correction (C_{trans}) to all instruments; right plot shows concentrations obtained by applying C_{SA} , C_{inlet} , and C_{trans} to all instruments except EH1. the abrupt shift in the EL time series occurring around 05:40 LT on 2 August (visible as an outlier peak in the sulfuric acid time series) was caused by an issue encountered while adjusting the HNO_3 flow mixed with the sheath flow.


Table S6, analogous to Table 6 for nitrate mode, show the results of the correlation analysis performed on concentrations of different HOM classes measured by ML1 and ML2 operating in Br^- mode on 27–31 July 2024.

Table S6 Coefficients of determination (R^2) and relative median absolute deviation (rMAD) calculated from concentrations of different HOM classes measured by ML1 and ML2 on 27–31 July 2024. Values are shown for cases (a) where only C_{SA} is applied, and (b) where C_{SA} , C_{inlet} , and C_{trans} are applied.

HOM class	R_a^2	R_b^2	$rMAD_a$ [%]	$rMAD_b$ [%]
organic peroxy radicals	0.826	0.828	56.5	30.1
closed shell non-nitrate monomers	0.785	0.782	25.4	14.3
closed shell organonitrate monomers	0.730	0.725	27.2	39.4
dimers	0.132	0.131	94.4	88.5
semi-volatile organic compounds	0.877	0.853	22.3	6.1

S11 Measurements in nitrate mode – additional species

Figure S5 shows time series and box plots for additional HOM classes measured in nitrate mode on 1–2 August 2024, complementing Figure 5 of the manuscript text.



170 **Figure S5** Concentrations of (top panel) nitrophenol, (central panel) organic peroxy radicals, and (bottom panel) closed-shell non-nitrate
 monomers measured in nitrate mode by five CI-API-ToFs during 1-2 August 2024. For each panel, the left figure ((a), (d), and (g)) shows
 concentration time series, with instrument-specific LOD_{CIMS} values indicated by dotted horizontal lines. The central figure ((b), (e), and (h))
 shows a box plot summarizing the distributions over the same period; boxes indicate the interquartile range, horizontal lines indicate the
 175 median, whiskers represent the non-outlier range, and empty circles denote outliers. Data points below LOD_{CIMS} are excluded from the box
 plot. The right figure ((c), (f), and (i)) shows the hourly median concentration measured by each instrument against the median of the
 hourly median concentrations across all five instruments. Dashed lines indicate log-log linear regressions for each instrument, with the
 corresponding coefficient of determination (R^2) reported in the legend. Points below the corresponding LOD_{CIMS} are shown as crosses (X),
 while outliers are plotted as smaller semi-transparent points; both are excluded from the linear fits.

S12 Measurements in nitrate mode – correlation analysis

180 Table S7-Table S9, analogous to Table 7 in the manuscript text, report the results of the correlation analysis for the species
 targeted in NO_3^- mode based on data recorded by the five CI-API-ToFs. In Table S7, concentrations were calculated by

applying only C_{SA} to all species. In Table S8 and Table S9, concentrations were calculated by applying C_{SA} , C_{inlet} and C_{trans} ; however, in the latter, data from EH1 were excluded from the correlation analysis to assess whether some of the low R^2 values could be attributed to the different behavior of this instrument in specific m/z ranges.

185 **Table S7** Statistical parameters of the correlation analysis for the species measured in nitrate mode based on data recorded by five CI-APi-ToFs during 1-2 August 2024. Points below the corresponding LOD_{CIMS} are excluded from the statistical analysis. R^2 is the coefficient of determination, p is the probability value, and parameters a and b are the slope and the offset of the linear fit, respectively. Concentrations were calculated by applying only C_{SA} to all species.

Species	R^2	p	a	b
sulfuric acid	0.577	<0.001	0.897	0.525
methanesulfonic acid	0.271	<0.001	0.677	1.579
malonic acid	0.561	<0.001	0.995	0.010
iodic acid	0.339	<0.001	0.959	0.140
nitrophenol	0.314	<0.001	0.666	2.068
organic peroxy radicals	0.037	<0.05	0.394	3.859
closed shell non-nitrate monomers	0.625	<0.001	0.951	0.343
closed shell organonitrate monomers	0.730	<0.001	0.963	0.248
dimers	0.288	<0.001	0.653	1.907

190 **Table S8** Statistical parameters of the correlation analysis for the species measured in nitrate mode based on data recorded by five CI-APi-ToFs during 1-2 August 2024. Points below the corresponding LOD_{CIMS} are excluded from the statistical analysis. R^2 is the coefficient of determination, p is the probability value, and parameters a and b are the slope and the offset of the linear fit, respectively. Concentrations were calculated by applying C_{SA} , C_{inlet} , and C_{trans} to all species.

Species	R^2	p	a	b
sulfuric acid	0.577	<0.001	0.897	0.525
methanesulfonic acid	0.110	<0.001	0.587	2.188
malonic acid	0.450	<0.001	0.923	0.549
iodic acid	0.088	<0.001	0.599	1.815
nitrophenol	0.433	<0.001	0.815	1.185
organic peroxy radicals	0.012	>0.05	0.303	4.349
closed shell non-nitrate monomers	0.415	<0.001	0.917	0.644
closed shell organonitrate monomers	0.555	<0.001	0.956	0.369
dimers	0.003	>0.05	-0.186	6.395

195 **Table S9** Statistical parameters of the correlation analysis for the species measured in nitrate mode based on data recorded by four CI-API-ToFs during 1-2 August 2024; EH1 data were excluded from the analysis. Points below the corresponding LOD_{CIMS} are excluded from the statistical analysis. R^2 is the coefficient of determination, p is the probability value, and parameters a and b are the slope and the offset of the linear fit, respectively. Concentrations were calculated by applying C_{SA} , C_{inlet} , and C_{trans} to all species.

Species	R^2	p	a	b
sulfuric acid	0.379	<0.001	0.711	1.514
methanesulfonic acid	0.063	<0.05	0.487	2.728
malonic acid	0.291	<0.001	0.755	1.723
iodic acid	0.022	>0.05	0.189	3.627
nitrophenol	0.681	<0.001	0.829	1.057
organic peroxy radicals	0.320	<0.001	0.637	2.205
closed shell non-nitrate monomers	0.831	<0.001	0.934	0.435
closed shell organonitrate monomers	0.891	<0.001	0.965	0.207
dimers	0.004	>0.05	-0.073	5.539

S13 Measurements in bromide mode – calibration factor

200 Figure S6 reproduces the analysis presented in Figure 6 in the manuscript text, but with ML1 concentrations calculated using the sulfuric acid calibration factor derived during the campaign, C_{SA} , instead of the revised C'_{SA} . We decided to disregard this factor (1) because this calibration did not agree with the others performed on the same instrument, and (2) due to the behavior observed in Figure S6a. Specifically, ML1 and ML2 show a nearly identical response to temporal variations in SVOC concentrations, while their absolute concentrations are offset by an approximately constant factor. This systematic

205 shift is particularly evident in the scatter plot, where the two datasets align closely but are displaced relative to one another.




Figure S6 Concentrations of (top panel) semi-volatile organic compounds, and (bottom panel) closed-shell non-nitrate monomers measured in bromide mode by ML1 and ML2 during 27-30 July 2024. For each panel, the left figure ((a) and (d)) shows concentration time series, while the central figure ((b) and (e)) shows box plots summarizing the distributions over the same period. Boxes indicate the interquartile range, horizontal lines indicate the median, whiskers represent the non-outlier range, and empty circles denote outliers. The right figure ((c) and (f)) shows the hourly median concentration measured by each instrument against the mean of the corresponding hourly median concentrations from the two instruments. Dashed lines indicate log-log linear regressions for each instrument, with the corresponding coefficient of determination (R^2) reported in the legend. Outliers are plotted as smaller semi-transparent points and are excluded from the linear fits. The concentrations have been calculated using the original C_{SA} derived for ML1 during the campaign, in place of the revised C'_{SA} , as it is for Figure 6 in the manuscript text.

S14 Measurements in bromide mode – correlation analysis

Table S10 and Table S11 report the results of the correlation analysis for the species targeted in Br^- mode based on data recorded by ML1 and ML2. In Table S10, concentrations were calculated by applying only C_{SA} (or C'_{SA}) to all species. In Table S11, concentrations were calculated by applying C_{SA} (or C'_{SA}), C_{inlet} , and C_{trans} .

Table S10 Statistical parameters of the correlation analysis for the species measured in bromide mode based on data recorded by ML1 and ML2 during 27-30 July 2024. R^2 is the coefficient of determination, p is the probability value, and parameters a and b are the slope and the offset of the linear fit, respectively. Concentrations were calculated by applying only C_{SA} (or C'_{SA}) to all species.

Species	R^2	p	a	b
sulfuric acid	0.557	<0.001	0.812	1.201
closed shell non-nitrate monomers	0.785	<0.001	0.495	3.807


closed shell organonitrate monomers	0.730	<0.001	0.277	4.615
semi-volatile organic compounds	0.853	<0.001	0.785	1.811

225 **Table S11** Statistical parameters of the correlation analysis for the species measured in bromide mode based on data recorded by ML1 and ML2 during 27-30 July 2024. R^2 is the coefficient of determination, p is the probability value, and parameters a and b are the slope and the offset of the linear fit, respectively. Concentrations were calculated by applying C_{SA} (or C'_{SA}), C_{inlet} , and C_{trans} to all species.

Species	R^2	p	a	b
sulfuric acid	0.557	<0.001	0.812	1.201
closed shell non-nitrate monomers	0.782	<0.001	0.501	3.282
closed shell organonitrate monomers	0.725	<0.001	0.273	4.023
semi-volatile organic compounds	0.853	<0.001	0.783	1.554


S15 Measurements in different reagent ion modes: nitrate, bromide, and iodide – additional species

Figure S7 and Figure S8 extend the analysis shown in Figure 8 in the manuscript text to additional compounds.



230 **Figure S7** (a) Box plots showing the diurnal variation of $C_{10}H_{16}O_4$ based on measurements conducted on 1-2 August 2024. Data are from five instruments operating in nitrate mode (EH1, EH2, EL, ML1, ML2), one instrument operating in bromide mode (ML2), and one instrument operating in iodide mode (AB). Boxes indicate the interquartile range, whiskers represent the non-outlier range, and full circles denote outliers. For the instruments operating in nitrate mode, instrument-specific LOD_{CIMS} values are indicated by dotted horizontal lines.

235 (b) Scatter plots of hourly median concentrations showing, for each instrument and ionization mode combination, the concentrations measured by the individual instrument-mode against the median concentration measured across all instruments-mode combinations over the same period. Dashed lines indicate linear regressions for each instrument, with the corresponding coefficient of determination (R^2) reported in the figure.



240 **Figure S8** (a) Box plots showing the diurnal variation of $C_{10}H_{16}O_3$ based on measurements conducted on 1-2 August 2024. Data are from
 five instruments operating in nitrate mode (EH1, EH2, EL, ML1, ML2), one instrument operating in bromide mode (ML2), and one
 instrument operating in iodide mode (AB). Boxes indicate the interquartile range, whiskers represent the non-outlier range, and full circles
 denote outliers. For the instruments operating in nitrate mode, instrument-specific LOD_{CIMS} values are indicated by dotted horizontal lines.
 245 (b) Scatter plots of hourly median concentrations showing, for each instrument and ionization mode combination, the concentrations
 measured by the individual instrument-mode against the median concentration measured across all instruments-mode combinations over
 the same period. Dashed lines indicate linear regressions for each instrument, with the corresponding coefficient of determination (R^2)
 reported in the figure.

S16 References

- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-
 250 aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11, 3303–3318, <https://doi.org/10.5194/acp-11-3303-2011>, 2011.
- Fuller, E. N., Ensley, K., and Giddings, C. J.: Diffusion of Halogenated Hydrocarbons in Helium. The Effect of Structure on
 Collision Cross Sections, *J. Phys. Chem.*, 73, 3679–3685, <https://doi.org/10.1021/j100845a020>, 1969.
- Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostiainen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala,
 M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas.*
 255 *Tech.*, 3, 1039–1053, <https://doi.org/10.5194/amt-3-1039-2010>, 2010.
- Kürten, A., Rondo, L., Ehrhart, S., and Curtius, J.: Calibration of a chemical ionization mass spectrometer for the
 measurement of gaseous sulfuric acid, *Journal of Physical Chemistry A*, 116, 6375–6386, <https://doi.org/10.1021/jp212123n>,
 2012.

Mohr, C., Thornton, J. A., Heitto, A., Lopez-Hilfiker, F. D., Lutz, A., Riipinen, I., Hong, J., Donahue, N. M., Hallquist, M.,
260 Petäjä, T., Kulmala, M., and Yli-Juuti, T.: Molecular identification of organic vapors driving atmospheric nanoparticle
growth, *Nat. Commun.*, 10, <https://doi.org/10.1038/s41467-019-12473-2>, 2019.
Tang, M. J., Cox, R. A., and Kalberer, M.: Compilation and evaluation of gas phase diffusion coefficients of reactive trace
gases in the atmosphere: Volume 1. Inorganic compounds, *Atmos. Chem. Phys.*, 14, 9233–9247, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-14-9233-2014)
14-9233-2014, 2014.

265