



This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

**The detection of
cloud free snow
covered areas**

L. G. Istomina et al.

The detection of cloud free snow covered areas using AATSR measurements

L. G. Istomina, W. von Hoyningen-Huene, A. A. Kokhanovsky, and J. P. Burrows

Institute of Environmental Physics, University of Bremen, Bremen, Germany

Received: 26 January 2010 – Accepted: 8 March 2010 – Published: 24 March 2010

Correspondence to: L. G. Istomina (lora@iup.physik.uni-bremen.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

A new method to detect cloud free snow covered areas is developed using the measurements by the Advanced Along Track Scanning Radiometer (AATSR) on board the ENVISAT satellite in order to discriminate clear snow fields for the retrieval of aerosol optical thickness or snow properties. The algorithm uses seven AATSR channels from VIS to TIR and analyzes the spectral behavior of each pixel in order to recognize the spectral signature of snow. The algorithm includes a set of relative thresholds and combines all seven channels into one flexible criterion, which allows us to filter out all the pixels with spectral behavior similar to that of snow. The algorithm does not use any kind of morphological criteria and does not require the studied surface to have any special structure. The snow spectral shape criterion was determined by a comprehensive theoretical study, which included radiative transfer simulations for various atmospheric conditions as well as studying existing models and measurements of snow optical and physical properties in different spectral bands. The method has been optimized to detect cloud free snow covered areas, and does not produce cloud/land/ocean/snow mask. However, the algorithm can be extended and be able to discriminate various kinds of surfaces.

The presented method has been validated against Micro Pulse Lidar data and compared to MODIS cloud mask over snow covered areas, showing quite good correspondence to each other.

1 Introduction

For remote sensing retrievals of any kind, one of the most important initial steps is to understand what we are looking at, in order to only choose those data needed for further processing. Cloud detection over snow is important for aerosol retrievals (Istomina et al., 2009, 2010) as well as for retrievals of cloud coverage, cloud height and cloud properties, and also for retrievals of snow properties such as grain size,

AMTD

3, 1099–1132, 2010

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



soot concentration, etc. (Kokhanovsky et al., 2010). In this paper, we present a cloud screening technique which was created especially for aerosol optical thickness retrieval in Arctic regions. This means that as a result of the cloud screening we like to get clean snow scenes only.

5 Among all available cloud-screening algorithms of satellite imagery one can distinguish three basic approaches (applicable to a radiometer data):

10 1. Analysis of time-sequences of data, with the assumption that any short-term changes in reflectance can only be introduced by clouds (e.g. Key and Barry, 1989; Diner et al., 1999; Lyapustin et al., 2008; Lyapustin and Wang, 2009; Gafurov and Bárdossy, 2009).

15 2. Applying an absolute threshold for reflectance or brightness temperature in one or several spectral channels, or associating some combination of a few reflectances to some kind of surface, e.g. ratio of reflectances in the form of NDVI, where only few channels are used (e.g. Minnis et al., 2001; Bréon and Colzy, 1999; Lotz et al., 2009; Allen et al., 1990; Spangenberg et al., 2001; Trepte et al., 2001).

3. Spatial variability analysis (e.g. Martins et al., 2002).

These approaches alone cannot yield the required quality of cloud screening, as each of them has disadvantages: the first approach requires some sequence of data, basically the longer the better, which is not always available. Even with a sequence of 20 two images, the reflectance change can be introduced either by surface BRDF shape due to different illumination-observation geometries of these two measurements, or by aerosol optical thickness, or by a temporal change of surface properties. The available information about all these factors is often not sufficient for proper correction. For time-sequence analysis some visible non-changing structure of the surface is needed 25 (Lyapustin et al., 2008; Lyapustin and Wang, 2009), therefore cloud screening has difficulties over uniform or changing surface. The second mentioned cloud screening approach only requires one image, and reflectance thresholds can be set very

The detection of
cloud free snow
covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



well for some small set of images. However, one can never be sure that these absolute thresholds (often derived experimentally or manually) will work properly for all the illumination-observation geometries and all the types of surface available. Relative indices combined from several reflectances in different channels are better, but their disadvantage is that they only use few spectral bands and therefore are highly dependent on the selection of the channels. Spatial pattern analysis, thirdly, also has limitations: the surface can introduce temporally variable spatial structure, which has to be somehow distinguished from that introduced by clouds. This makes it important to choose wavelengths where the surface has no structure, which is not always possible.

Several available cloud screening methods over snow use either absolute (experimentally derived) thresholds in VIS, NIR and IR channels (Allen et al., 1990), and still have problems with ice clouds over snow (which is often the case in Arctic region), or they assume some snow BRDF model to derive the threshold value for NIR spectral channel and use only one NIR channel (Spangenberg et al., 2001; Trepte et al., 2001). The second approach cannot always distinguish cloudy conditions from clean but highly polluted scenes, which can often be seen during Arctic haze and smoke events.

Of course, the chosen cloud/clear sky detection methods depend on the spectral and spatial resolution of the instrument, and if the instrument has a lot of spectral channels, some combination of cloud screening approaches mentioned above will give a reliable cloud mask. An example of such a comprehensive combination is the MODIS cloud/clear sky determination algorithm (MODIS ATBD MOD-06, 2002). This algorithm has been validated against micropulse lidar and millimeter-wavelength cloud radar, and proved to be reliable.

However, there is another way of cloud/clear sky detection. In the current work we present a cloud screening approach which is based on the analysis of the spectral shape of a scene reflectance and does not require time sequences of data or absolute thresholds for reflectances or brightness temperatures. This approach has been invented for a specific task – to pick out cloud free snow covered regions to serve as an

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



input for aerosol optical thickness (AOT) retrieval over snow. Therefore, we did not pursue the task of recognizing different types of surface and creating a cloud mask similar to the cloud mask of e.g. MODIS. The product of presented cloud screening method consists of two values: “applicable for AOT retrieval over snow or determination of snow reflection (clear snow)” and “not applicable for AOT retrieval or determination of snow reflection (everything else but snow – clouds, land, ocean, etc.)”.

2 Theoretical basis of spectral cloud screening over snow

Any cloud screening method always depends on incoming data and available features of the instrument. The spectral coverage of AATSR, which has been used in this work,

enables the spectral behavior of a scene in seven different channels from VIS to TIR to be investigated. Observations and models (e.g. Warren, 1982) show that snow has a significant and recognizable spectral signature in the range from VIS to NIR (AATSR channels 550 nm, 660 nm, 870 nm, 1.6 μm), whereas clouds behave differently there, e.g. they do not show such a significant drop of reflectance between channels 870 nm and 1.6 μm (Kokhanovsky, 2006).

This different spectral behavior of snow and clouds can be seen in the reflectance charts of different AATSR channels (see Fig. 1). Possible snow impurities, snow grain size differences, and liquid water can affect the shape of the snow surface spectrum (Warren, 1982). Different atmospheric conditions, e.g. different aerosol loads, can also affect the spectral shape of the scene and should be taken into account.

In the TIR channels of AATSR (3.7 μm , 10.8 μm , 12 μm), the spectral shapes of snow and clouds are also different, and can be affected by physical parameters of both snow and clouds.

The following two sections are dedicated to a detailed description of our spectral cloud screening criteria in these two spectral regions (VIS-NIR and TIR) and possible disturbance factors which have to be taken into account.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.1 Cloud screening in VIS and NIR: disturbance from aerosols, trace gases and physical parameters of snow

AMTD

3, 1099–1132, 2010

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



is equal 0°), rather low Sun (solar zenith angle equal to 60°), and the relative azimuth angle is equal to 0° . According to Mendonca et al. (1981), the single scattering albedo of Arctic haze appears to be rather high, larger than 0.9. Delene and Ogren (2002) reported the SSA of haze at Barrow, Alaska, to be 0.94. As we do not know the actual value of the SSA, we assume it to be 1, and compensate the possible lack of absorption with too high AOT. The aerosol is assumed to be uniformly distributed in a 3 km layer above the surface.

The result of such a simulation is the wavelength-dependence of TOA reflectance of the system “surface+atmosphere”. Here TOA reflectance is the amount of radiation, scattered at the surface and in the atmosphere (as seen from the top of atmosphere), divided by the incident amount of radiation.

Figure 3a shows the result of our simulation for the discussed surface and an aerosol load from 0 to 3.0 (step 0.2). Figure 3b shows the same channels, surface and aerosols, but with included Rayleigh scattering and O_3 absorption.

It can be seen that the shape of the spectral curve does not depend on AOT for channels 550 nm, 660 nm, 870 nm in the “ideal” case (Fig. 3a) as well as in the “real” case with Rayleigh scattering and O_3 included. The inclusion of O_3 absorption and Rayleigh scattering results in a slight decrease of TOA reflectance in the 550 nm channel.

As can also be seen from Fig. 3, over such a bright surface, the increasing of aerosol load in the atmosphere causes a darkening of the scene in channels 550 nm, 660 nm and 870 nm for the given geometry, due to scattering of light reflected from the surface on aerosol layer and thus decreasing the upward flux at the TOA. Channel $1.6\text{ }\mu\text{m}$ reacts to an increase of AOT with an increase of the TOA reflectance, due to the fact that even a thin cloud of aerosol particles reflects more than snow in this spectral region (snow albedo is around 0.1 here).

As a result, various aerosol loads (AOT from 0 to 3) affect TOA reflectance in 550 nm, 660 nm and 870 nm channels, causing an equal decrease in all these channels of about 10%. However, this does not affect the shape of the spectral curve. It is important to note that these effects are only valid for the discussed geometry and a very bright

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

surface (Kokhanovsky et al., 2010).

Rayleigh scattering and O₃ absorption affect the shape of the spectrum, bringing the TOA reflectance in 550 nm channel down by 10% in comparison to the “ideal” case without this disturbance. The shape of the spectral curve is also affected in the 1.6 μm channel, where the TOA reflectance grows by more than 50% with the increase of the aerosol load until AOT equals 3. All these factors disturb the shape of the snow spectrum and have to be taken into account during the analysis of the spectral behavior of the scene reflectance.

The other factors which can disturb the shape of the spectral curve are caused by the physical parameters of snow and not by the atmosphere above it. They are snow grain size, liquid water content and possible impurities of the snowpack, which can be soot, dust, or atmospheric aerosols fallen out on the snow surface. All these factors have been studied and are presented in papers aimed at both modeling and measurements of snow optical properties.

The effect of dust and soot impurities in snow has been studied by Warren and Wiscombe (1980). It is shown that the VIS spectral region is particularly sensitive to these impurities, whereas NIR and longwave emissivity of snow are not affected (which is important for Sect. 2.2). Of course, the distortion of snow albedo spectrum depends on the concentration of dust and soot. For example, 10 ppmw of soot can decrease the albedo of snow by more than 50% in VIS for an extremely large snow grain size of 1000 μm (Warren and Wiscombe, 1980). However, this effect is smaller for smaller grains and Arctic snow is believed to have a soot concentration of around 0.2 ppmw, which reduces the snow albedo by approximately 20% in the 550 nm and 660 nm channels. For 870 nm, this effect is already less than 10%. Similar effects take place in case of dust impurities: depending on snow grain size, extremely high dust concentrations can reduce the VIS albedo of snow drastically, but dust concentrations of 10 ppmw decrease snow albedo by less than 10% in VIS, for both small and large grain sizes. Aoki et al. (2000) show the modeled spectral albedo of snow for several reasonable combinations of physical parameters such as snow density, impurities concentration,

The detection of
cloud free snow
covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



5 snow grain size and snow depth. These spectra differ from the pure snow spectrum only in the shortwave region, and by no more than 10%, which proves that earlier discussed scatter values caused by soot and dust are sufficient for our task. The effect of snow age on the snow spectrum was studied by Domine et al. (2006). The comparison of wind crust, fresh snows and depth hoar spectra show that these spectra are quite similar to each other in the visible, but differ more in the IR. The 1.6 μm channel reacts on the snow aging with a decreasing albedo, sometimes almost to zero. However, this effect does not destroy the main feature of snow spectral signature – the spectacular drop of albedo from 0.87 μm to 1.6 μm . Liquid water content, however, decreases the 10 snow albedo also in the VIS and NIR (Gerland et al., 1999), but it does not change the spectral shape in the channels of AATSR and the relation between all the discussed VIS and NIR channels remains the same (Warren, 1982).

15 The effect of snow grain size on the snow spectrum is studied, for example, by Domine et al. (2008), Wiscombe and Warren (1980), and Tedesco and Kokhanovsky (2007). The snow spectrum is affected by the grain size mostly in NIR and only a bit in VIS. Snow albedo in the 550 nm and 660 nm channels is affected only by a few percent, at 870 nm it goes up by around 10% for smaller grains and 10% down for larger grains. The snow albedo in the 1.6 μm channel increases for smaller grains up to 0.2 and drops down to almost zero for large grains, which is similar to the behavior of the albedo in 20 the 870 nm channel. This means that the relation between these two channels remains the same and the shape of the snow spectrum is not affected.

25 All the discussed disturbance factors (aerosol load, Rayleigh scattering, ozone, soot and dust impurities, age and liquid water content, and snow grain size) affect the spectrum of the cloud free snow scene in different wavelength regions. The performed literature study and RT simulations made it possible to develop a criterion of snow spectral shape in VIS and NIR channels of AATSR. It is schematically shown in Fig. 4 for two example spectra. They are the fresh snow spectrum (from the ASTER spectral library, CalTech, 2008), represented with the blue curve and the dry long grass spectrum (from the USGC Spectroscopy Lab library), represented with the black curve. Filled areas

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(violet for fresh snow and turquoise for dry long grass) show allowed freedom in the shape of each spectrum. As the criterion is relative, the allowed scatter will change from spectrum to spectrum. We would like to underline that for our cloud screening routine, we only need to analyze the shape of a spectrum in four VIS and NIR AATSR channels. These channels are shown in Fig. 4 by vertical lines. However, to make the concept more obvious, we extended the graphic representation of the shape criteria outside of the four discussed channels. The numerical criteria only exist for the four discussed channels and should connect the TOA reflectances, $R_{TOA}(\lambda)$, of those channels in the following way:

10 – R_{TOA} (870 nm) and R_{TOA} (1.6 μ m) should differ not less than 80%;
 – R_{TOA} (660 nm) should not be smaller than R_{TOA} (870 nm) by more than 10%;
 – R_{TOA} (660 nm) and R_{TOA} (550 nm) should differ not more than 40%.

15 Here the percentage is given for relative reflectance difference with the relation to R_{TOA} (870 nm) for the first two criteria and to R_{TOA} (660 nm) for the third one. From Fig. 4 it is visible that the fresh snow spectrum (blue curve) fits into our snow spectral shape criteria, whereas the spectrum of dry long grass does not. The R_{TOA} (1.6 μ m) of dry grass is too high and does not represent the main feature of a snow spectrum – the reflectance drop from 870 nm to 1.6 μ m. Also, the feature of vegetation spectra, the so called “red edge” (fast increase of reflectance at 0.4–0.7 μ m), is visible in the figure and 20 makes the R_{TOA} (660 nm) be at the edge of the snow spectral shape criteria.

25 The discussed criteria are already enough to screen out optically thick warm clouds, but will have difficulties with cirrus and any optically thin clouds, because these do not significantly disturb the spectral signature of snow in these spectral regions. To screen out such clouds, we need to use TIR channels of AATSR. This problem is discussed in the next section.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.2 Cloud screening in TIR: snow emissivity and cloud reflectance

AMTD

3, 1099–1132, 2010

The 3.7, 10.8 and 12 μm channels provide additional information to distinguish surface and clouds. As was mentioned by Spangenberg et al. (2001), brightness temperature (BT) of the 3.7 μm channel contains not only emitted radiation, but also radiation scattered by the object, in our case by a cloud or snow.

Thermal properties of snow have been measured (Hori et al., 2006; English et al., 1995) and show that snow is very close to a black body and emits according its physical temperature. The reflection of snow is very low in the 3.7, 10.8 and 12 μm channels (Wald, 1994). This is not the case for clouds, because R (3.7 μm) of cloud droplets is much higher depending on the effective radii of the particles and on the optical thickness of the cloud. Therefore the spectral shape of a cloud pixel in the discussed 3 channels will differ from that of snow due to the different physical nature of these two objects.

To estimate the amplitude of reflectance contamination of BT (3.7 μm), we calculate the R_{TOA} (3.7 μm) according to Spangenberg et al. (2001):

$$R_{\text{TOA}}(3.7\mu\text{m}) = \frac{[\varepsilon_{3.7\mu\text{m}} \cdot B_{3.7\mu\text{m}}(T_3) - \varepsilon_{3.7\mu\text{m}} \cdot B_{3.7\mu\text{m}}(T_4)]}{[(\mu_0 \cdot S_{3.7\mu\text{m}}) - \varepsilon_{3.7\mu\text{m}} \cdot B_{3.7\mu\text{m}}(T_4)]} \quad (1)$$

where T_3 is the measured 3.7 μm brightness temperature, T_4 is the measured 11 μm brightness temperature, μ_0 is the cosine of solar zenith angle, $S_{3.7\mu\text{m}}$ is the solar constant at 3.7 μm ($3.47 \text{ W m}^{-2} \mu\text{m}^{-1}$), $\varepsilon_{3.7\mu\text{m}}$ is the clear snow emissance, $B_{3.7\mu\text{m}}$ (BT) is the Planck function at 3.7 μm for some temperature BT.

An example of a calculated R_{TOA} (3.7 μm) is presented in Fig. 1e. One can clearly see the high reflectance of clouds and very low reflectance of snow and ocean (see ocean-land mask contours). This reflectance pattern is included in the AATSR BT(3.7 μm) (Fig. 1f), causing the differences in the shapes of the spectral curves of snow and clouds for 3.7, 10.8 and 12 μm channels. It is visible that the brightness temperature of snow in all three channels is approximately the same, which corresponds

The detection of
cloud free snow
covered areas

L. G. Istomina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to, e.g. measurements of snow emissivity in MODIS USCB Emissivity Library. Measured emissivity is quite stable throughout the thermal region of the spectrum (variation around 2%). Hori et al. (2006) show that emissivity of snow for a given temperature depends on its physical parameters such as grain size and liquid water content. However, these dependencies cause a variation of snow emissivity of less than 5%. This scatter of the spectral shape can be taken into account and is still much smaller than that caused by the high reflectance of clouds in the 3.7 μm channel.

Independently of the physical temperature of snow, the relative BT differences of the 3.7, 10.8 and 12 μm channels are not to be larger than 3%. This criterion will remove both warm and cold clouds. Both are present in the discussed scene; in Fig. 1f, the blue color corresponds to cold, and orange to warm clouds. BT of snow and ocean is similar in all three discussed channels within the allowed scatter.

All the factors mentioned in Sect. 2.1 and 2.2 create a spectral shape criterion for the relative reflectances and brightness temperatures in seven AATSR channels. It is not an absolute threshold in each of the channel separately, but rather a flexible relative connection of all the seven spectral channels in order to recognize the spectral signature of snow, if it is present in the spectrum of the scene.

To summarize, here are all the parts of the spectral shape criterion:

Relative BT difference at 3.7, 10.8 and 12 μm channels should not be larger than 3%; relative reflectance difference at 1.6 μm and 870 nm channels should be larger than 80%; reflectance in the 670nm channel should not be smaller than that in the 870 nm channel by more than 10%; and, finally, the relative reflectance difference of the 550 nm and 660 nm channels should be not more than 40%.

Experiments show that this order of criteria is computationally the most effective, because most clouds are screened out during TIR check and the snow signature is mostly stored in the large drop between 870 nm and 1600 nm channels. None of these criteria are related to the absolute value of reflectance or BT; therefore, the algorithm does not rely on spatial contrast of the image in any channel.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The result of our cloud screening method is shown in Fig. 5 (unscreened scene in all the 7 AATSR channels is shown in Fig. 1).

AMTD

3, 1099–1132, 2010

3 Validation of the algorithm

3.1 MPL

5 The discussed cloud screening method was applied for a sequence of AATSR data for spring 2006, at Spitsbergen (approximately 100 scenes). The results were compared to the Micro-Pulse Lidar network data (Welton et al., 2001; Campbell et al., 2002). In 95% of the cases, the discussed method detects cloudy and clear sky correctly. The resulting 5% of wrong detections are connected to thin clouds (Fig. 6), which disturb
10 the spectral shape of the scene only slightly, or to a particular case of surface (Fig. 7).

In Fig. 6 we see the scene for Spitsbergen, 11 May 2006. Micro Pulse Lidar located in Ny Ålesund ($78^{\circ}.91667$ N, $11^{\circ}.93300$ E), in the western part of the island. It shows the presence of a very thin cloud at 1 km altitude, however, neither discussed cloud screening technique nor the MODIS cloud mask show it. At the same time, their correspondence to each other is rather good throughout the rest of the image. The reason
15 can be the size of the cloud: subpixel thin clouds over snow simply do not change the spectral shape of the scene significantly.

In Fig. 7 we see another scene for Spitsbergen, 8 May 2006. According to Micro Pulse Lidar, the Ny Ålesund area should be cloud free. However, our cloud screening
20 shows that there is no clear snow at Ny Ålesund on this scene. The MODIS cloud product shows this area as “probably clear”. AATSR reflectance shows that the surface is slightly darker at Ny Ålesund. This can be caused by shadowing by mountains or extremely contaminated snow surface.

25 So we see that thin clouds are, unfortunately, still a problem for the discussed cloud screening method. Of course, one could give more strict spectral shape criteria, and reduce the allowed scatter. However, this could lead to screening out of pixels with

The detection of
cloud free snow
covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



present aerosol loads or pixels with those various kinds of snow which differ from the fresh clean snow. In other words, the spectral shape criterion can be adjusted to fit the certain task. If the task is to study physical properties of snow (where atmospheric correction is usually a problem), one could set the allowed scatter to be very narrow along the desired spectrum. Such a criterion would only give the scenes with clean fresh snow, according to the reference spectrum (see e.g. Fig. 4). The influence of the atmosphere here will be very small and all the exotic types of snow will also be absent. On the other hand, if the objective is to study atmosphere and for example to retrieve AOT over snow, then the allowed range of scatter should be rather wide, to ensure that high aerosol loads are not screened. However, a broad criterion for scatter yields scenes with various types of snow, possibly contaminated. Thus the target product for the retrieval (in this case AOT over snow retrieval) needs to be considered.

3.2 The comparison to MODIS cloud mask for different kinds of snow surface

In order to check the performance of the presented cloud screening method over different kinds of snow surface, we processed a set of AATSR scenes from both Northern and Southern hemispheres at various locations, and compared the results to the MODIS cloud product for the same location. Unfortunately, exact temporal collocation of these two instruments was sometimes impossible, in which case the closest available observation was chosen. Pixel-to-pixel collocation of the two instruments appears to be not easy, but visual analysis can already help to estimate the quality of the presented method. The results of the presented cloud screening method plotted together with MODIS cloud mask for eight different cases of clouds over snow are shown in Figs. 8–15. They include scenes over Alaska (snow, ice covered with snow), Greenland (snow, ice covered with snow), the Scandinavian Peninsula (snow covered mountains, snow covered forest), Northern part of Russia (snow covered forest, snow on ice), Himalayan mountains (snow covered peaks), Antarctica (snow, snow covered ice). Figures 8–13 shown that the presented cloud screening method gives rather good quality, screening out all the clouds and not confusing them with snow (Fig. 10). Flat

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



snow, either on ice or on land, is the best surface for the cloud screening (Figs. 8–9, ice part of Fig. 12). Mountain peaks covered with snow can be recognized (Figs. 10, 11, 14). Snow covered forest (Fig. 11) is also recognized (however, not continuously; only single brightest snow pixels are detected). In general, the presented method appears 5 to be a little stricter in comparison to the MODIS cloud mask. For example, the case of flat snow, according to the MODIS cloud product, is not recognized very well (Fig. 15, Antarctica). The reason for a not very accurate detection might be the illumination-observation geometry (low solar zenith angle), which could affect the ranges of applicability of the presented spectral shape criteria. Additionally, the reason might be some 10 type of snow here overlooked, which is located outside the suggested spectral shape criterion, but still is a cloud free and not extremely contaminated snow. Extensive research is needed to understand if this is the feature of all the Antarctic snow or just the feature of one special day or location. In general, according to the comparison above, 15 the quality of the presented cloud/snow discrimination appears to be reasonably good for such a simple discrimination algorithm.

4 Conclusions

A simple and robust cloud screening method was developed for AOT retrieval over snow using AATSR observations. The method uses a spectrum-shape control in seven 20 VIS, NIR and TIR AATSR channels and does not require time-sequences of data, morphological analysis of the scene or manual tuning to set absolute thresholds. The method was compared to MODIS cloud mask for various dates and locations and showed good correspondence with it over flat snow, snow-covered mountain peaks, 25 snow-covered forest and snow-covered ice. The presented cloud screening was validated against MPL data. The reliability of the method is 95% (on approximately 100 scenes). The presented cloud screening over snow can be extended to other types of surface and could be a candidate for the upcoming ESA Sentinel 3 mission.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Acknowledgements. The authors are grateful to ESA for providing AATSR data, to NASA for providing MODIS cloud mask product, to R. Treffeisen and M. Stock for providing their aerosol phase function measurements, to V. Rozanov and A. Rozanov for the help with the forward RT model SCIATRAN, and also to M. Schreier for helpful discussions on cloud screening and handling of satellite data.

References

Allen, R. C., Durkee, P. A., and Wash, C. H.: Snow/cloud discrimination with multispectral satellite measurements, *J. Appl. Meteor.*, 29, 994–1004, 1990.

Aoki, Te., Aoki, Ta., Fukabori, M., Hachikubo, A., Tachibana, Y., and Nishio, F.: Effects of snow physical parameters on spectral albedo and bidirectional reflectance of snow surface, *J. Geophys. Res.*, 105(D8), 10219–10236, doi:1999JD901122, 2000.

Bréon, F.-M. and Colzy, S.: Cloud detection from the spaceborne POLDER instrument and validation against surface synoptic observations, *J. Appl. Meteor.*, 38 pp. 777–785, 1999.

Campbell, J. R., Hlavka, D. L., Welton, E. J., Flynn, C. J., Turner, D. D., Spinhirne, J. D., Scott, V. S., and Hwang, I. H.: Full-Time, Eye-Safe Cloud and Aerosol Lidar Observation at Atmospheric Radiation Measurement Program Sites: Instruments and Data Processing, *J. Atmos. Oceanic Technol.*, 19, 431–442, 2002.

Delene, D. J. and Ogren, J. A.: Variability of aerosol optical properties at four North American surface monitoring sites, *J. Atmos. Sci.* 59, 1135–1150, 2002.

Diner, D., Clothiaux, E., Di Girolamo, L.: MISR Multi-angle imaging spectro-radiometer algorithm theoretical basis. Level 1 Cloud detection, Jet Propulsion Laboratory, JPL D-13397, 1999.

Domine, F., Albert, M., Huthwelker, T., Jacobi, H.-W., Kokhanovsky, A. A., Lehning, M., Picard, G., and Simpson, W. R.: Snow physics as relevant to snow photochemistry, *Atmos. Chem. Phys.*, 8, 171–208, 2008,
<http://www.atmos-chem-phys.net/8/171/2008/>.

Domine, F., Salvatori, R., Legagneux, L., Salzano, R., Fily, M., and Casacchia, R.: Correlation between the specific surface area and the short wave infrared (SWIR) reflectance of snow, *Cold Reg. Sci. Technol.*, 46, 60–68, 2006.

English, S. J., Jones, D. C., Hewison, T. J., Saunders, R. W., Hallikainen, M.: Observations of

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the emissivity of snow and ice surfaces from the SAAMEX and MACSI airborne campaigns, Geoscience and Remote Sensing Symposium, 1995. IGARSS '95, "Quantitative Remote sensing for Science and Applications", International, Vol. 2, 1493–1495, 1995.

5 Gafurov, A. and Bárdossy, A.: Cloud removal methodology from MODIS snow cover product, *Hydrol. Earth Syst. Sci.*, 13, 1361–1373, 2009,
<http://www.hydrol-earth-syst-sci.net/13/1361/2009/>.

Gerland, S., Winther, J.-G., Ørbaek, J. B., Liston, G. E., Ørtsland, N. A., Blanco, A., and Ivanov, B.: Physical and optical properties of snow covering Arctic tundra on Svalbard, *Hydrological Processes* 13, 2331–2343, 1999.

10 Hori, M., Aoki, T., Tanikawa, T., Motoyoshi, H., Hachikubo, A., Sugiura, K., Yasunari, T. J., Eide, H., Storvold, R., Nakajima, Y., and Takahashi, F.: In-situ measured spectral directional emissivity of snow and ice in the 8–14 μm atmospheric window, *Remote Sens. Environ.*, 100, 486–502, 2006.

15 Istomina, L. G., von Hoyningen-Huene, W., Kokhanovsky, A. A., and Burrows, J. P.: Retrieval of aerosol optical thickness in Arctic region using dual-view ATSR observations. *Proceedings of ESA Atmospheric Science Conference*, Barcelona, Spain, 7–11 Sept. 2009, ESA SP-676, 2010.

20 Istomina, L. G., von Hoyningen-Huene, W., Kokhanovsky, A. A., Rozanov, V. V., Schreier, M., Dethloff, K., Stock, M., Treffeisen, R., Herber, A., Burrows, J. P.: Sensitivity study of the dual-view algorithm for aerosol optical thickness retrieval over snow and ice, *Proceedings of the 2nd MERIS/(A)ATSR user workshop*, ESRIN, Frascati, Italy, 22–26 Sept. 2008, ESA SP-666, 2009.

Key, J. and Barry, R.G.: Cloud cover analysis with Arctic AVHRR data. 1. Cloud detection, *J. Geophys. Res.*, 94(D15), 18521–18535, 1989.

25 Kokhanovsky, A. A.: *Cloud Optics*, edited by: Mysak, L. A. and Hamilton, K., Publ.: Springer, 2006.

Kokhanovsky, A. A., Rozanov, V. V., Aoki, T., Odermatt, D., Brockmann, B., Krüger, O., Bouvet, M., Drusch, M., Hori, M.: Sizing of snow grains using backscattered solar light, submitted to *Int. J. Remote Sens.*, 2010.

30 Lotz, W. A., Vountas, M., Dinter, T., and Burrows, J. P.: Cloud and surface classification using SCIAMACHY polarization measurement devices, *Atmos. Chem. Phys.*, 9, 1279–1288, 2009,
<http://www.atmos-chem-phys.net/9/1279/2009/>.

Lyapustin, A., Tedesco, M., Wang, Y., Aoki, T., Hori, M., and Kokhanovsky, A.: Retrieval of snow

The detection of
cloud free snow
covered areas

L. G. Istomina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



grain size over Greenland from MODIS, *Remote Sens. Environ.*, 113, 1976–1987, 2009.

Lyapustin, A. and Wang, Y.: The time series technique for aerosol retrievals over land from MODIS, in: *Satellite Aerosol Remote Sensing over Land*, edited by: Kokhanovsky A. A. and de Leeuw, G., Springer Praxis Publ. Chichester, 69–99, 2009.

5 Lyapustin, A., Wang, Y., and Frey, R.: An automated cloud mask algorithm based on time series of MODIS measurements, *J. Geophys. Res.*, 113, D16207, doi:10.1029/2007JD009641, 2008.

Martins, J. V., Tanré, D., Remer, L., Kaufman, Y., Mattoo, S., and Levy, R.: MODIS Cloud screening for remote sensing of aerosols over oceans using spatial variability, *Geophys. Res. Lett.*, 29(12), 8009, doi:10.1029/2001GL013252, 2002.

10 Mendonca, B. G., DeLisi, J. J., and Schroeder, J. A.: Arctic Haze and perturbation in the solar radiation fluxes at Barrow, Alaska, *Proceedings from the 4th Conference on Atmospheric Radiation*. Atm. Met. Soc., Toronto, Ontario, Canada, 95–96, 1981.

Minnis, P., Chakrapani, V., Doelling, D. R., Nguyen, L., Palikonda, R., Spangenberg, D. A., Uttal, T., Arduini, R. F., and Shupe, M.: Cloud coverage and height during FIRE ACE derived from AVHRR data, *J. Geophys. Res.*, 106(D14), 15215–15232, 2001.

15 MODIS ATBD MOD-06, Discriminating clear-sky from cloud with MODIS, modis.gsfc.nasa.gov/data/atbd/atbd_mod06.pdf, 2002.

Quinn, P. K., Miller, T. L., Bates, T. S., Ogren, J. A., Andrews, E. and co-authors: A three-20 year record of simultaneously measured aerosol chemical and optical properties at Barrow, Alaska, *J. Geophys. Res.*, 107(D11), 4130, doi:10.1029/2001JD001248, 2002.

Rozanov, A. V., Rozanov, V. V., Buchwitz, M., Kokhanovsky, A. A., and Burrows, J. P.: SCIA-25 TRAN 2.0 – new radiative transfer model for geophysical applications in the 175–2400 nm spectral range, *Adv. Space Res.* 36, 1015–1019, 2005.

Spangenberg, D. A., Chakrapani, V., Doelling, D. R., Minnis, P., and Arduini, R. F.: Development25 of an automated Arctic cloud mask using clear-sky satellite observations taken over the SHEBA and ARM NSA sites, *Proc. 6th Conf. on Polar Meteor. and Oceanography*, San Diego, CA, May 14–18, 2001, 246–249, 2001.

Tedesco, M. and Kokhanovsky, A. A.: The semi-analytical snow retrieval algorithm and its30 application to MODIS data, *Rem. Sens. Env.*, 110, 317–331, 2007.

Tomasi, C., Vitale, V., Lupi, A., Di Carmine, C., Campanelli, M., Herber, A., Treffeisen, R., Stone, R. S., Andrews, E., Sharma, S., Radionov, V., von Hoyningen-Huene, W., Stebel, K., Hansen, G. H., Myhre, C. L., Wehrli, C., Aaltonen, V., Lihavainen, H., Virkkula, A., Hillamo, R., Ström,

**The detection of
cloud free snow
covered areas**

L. G. Istomina et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

J., Toledano, C., Cachorro, V. E., Ortiz, P., de Frutos, A. M., Blindheim, S., Frioud, M., Gausa, M., Zielinski, T., Petelski, T., Yamanouchi, T.: Aerosols in polar regions: A historical overview based on optical depth and in situ observations, *J. Geophys. Res.*, 112, D16205, doi:10.1029/2007JD008432, 2007.

5 Trepte, Q., Arduini, R. F., Chen, Y., Sun-Mack, S., Minnis, P., Spangenberg, D. A., and Doelling, D. R.: Development of a daytime polar cloud mask using theoretical models of near-infrared bidirectional reflectance for ARM and CERES, *Proc. AMS 6th Conf. Polar Meteorology and Oceanography*, San Diego, CA; May 4-18, 242–245, 2001.

10 Wald, A. E.: Modeling thermal infrared (2–14 μ m) reflectance spectra of frost and snow, *J. Geophys. Res.*, 99(B12), 94JB01560, 24241–24250, 1994.

15 Warren, S. G.: Optical Properties of Snow, *Rev. Geophys.*, 20(1), 67–89, 1982.

Warren, S. G. and Wiscombe, W. J.: A model for the spectral albedo of snow. II: Snow containing atmospheric aerosols, *J. Atmos. Sci.*, 37, 2734–2733, 1980.

Welton, E. J., Campbell, J. R., Spinhirne, J. D., Scott, V. S.: Global monitoring of clouds and aerosols using a network of micro-pulse lidar systems. *Proc. Int. Opt. Eng.* 4153, 151–158, 2001.

20 Wiscombe, W. J. and Warren, S. G.: A model for the spectral albedo of snow, I. Pure snow, *J. Atmos. Sci.*, 37(12), 2712–2733, 1980.

AMTD

3, 1099–1132, 2010

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 1. AATSR scene for 3rd of May 2006, Spitsbergen, reflectance at 550 nm (a), 660 nm (b), 865 nm (c), 1600 nm (d), 3700 nm (e). Brightness temperature for 3.7 (f), 10.8 (g) and 12 μm (h).

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The detection of cloud free snow covered areas

L. G. Istomina et al.




Fig. 2. Phase function of Arctic haze aerosol used in the simulation of aerosol effect on the spectral shape of snow surface. See text for details.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**The detection of
cloud free snow
covered areas**

L. G. Istomina et al.




Fig. 3. The sensitivity of TOA reflectance over snow surface to aerosol load, simulated with RT calculations, for Lambertian snow and aerosol only (3a) and for snow, aerosol and O_3 absorption (3b). AOT changes from 0 to 3, with the step 0.2. AOT = 0 is shown with the thick red line. AOT = 3.0 is shown with the thick violet line.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The detection of cloud free snow covered areas

L. G. Istomina et al.




Fig. 4. The allowed amount of TOA reflectance scatter is shown for fresh dry snow (blue curve) and dry long grass (black curve). Violet and turquoise areas are the schematic view of the spectral shape criterion. They show where the corresponding spectral curve should go to be still recognized as snow. It is visible that dry long grass spectrum does not satisfy the spectral shape criterion, e.g. at channel 1.6 μm .

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

**The detection of
cloud free snow
covered areas**

L. G. Istomina et al.




Fig. 5. The example of discussed cloud screening method: **(a)** the initial AATSR scene, 3rd of May 2006, Spitsbergen, 550 nm; **(b)** the probability of clear atmosphere over snow – the result of the cloud screening over snow routine.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)




Fig. 6. Left panel shows the false color composite (red is $11\text{ }\mu\text{m}$ BT, green is 550 nm reflectance, blue is $1.6\mu\text{m}$ reflectance) of the initial AATSR scene, 11 May 2006, $16^{\text{h}}26'26''$, orbit number 21938; middle panel is the screened AATSR scene, probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear. MODIS scene is for 11 May 2006, $16^{\text{h}}40'$.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 7. Left panel shows the false color composite (red is $11\text{ }\mu\text{m}$ BT, green is 550 nm reflectance, blue is $1.6\text{ }\mu\text{m}$ reflectance) of the AATSR scene, 8 May 2006, $13^{\text{h}}02'12''$, orbit number 21 893; middle panel is the screened AATSR scene, probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear. MODIS scene is for 8 May 2006, $12^{\text{h}}55'$.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 8. Alaska, AATSR: 30 April 2006, $22^{\text{h}}17'44''$, orbit number 21 783. MODIS: 30 April 2006, $22^{\text{h}}05'$. Left panel is the false color composite (red is $11 \mu\text{m}$ BT, green is 550 nm reflectance, blue is $1.6 \mu\text{m}$ reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 9. Greenland. AATSR: 4 May 2005, 15^h07'47", orbit number 21 836, MODIS: 4 May 2005, 15^h00'. Left panel is the false color composite (red is 11 μm BT, green is 550 nm reflectance, blue is 1.6 μm reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 10. Scandinavian Peninsula, AATSR: 14 April 2006, $10^{\text{h}}38'17''$, orbit number 21 547. MODIS: 14 April 2006, $10^{\text{h}}35'$. Left panel is the false color composite (red is $11 \mu\text{m}$ BT, green is 550 nm reflectance, blue is $1.6 \mu\text{m}$ reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 11. Scandinavian Peninsula, AATSR: 22 April 2006, 09^h47'27", orbit number 21 661. MODIS: 22 April 2006, 09^h40'. Left panel is the false color composite (red is 11 μm BT, green is 550 nm reflectance, blue is 1.6 μm reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

The detection of cloud free snow covered areas

L. G. Istomina et al.




Fig. 12. Northern part of Russia. AATSR: 22 April 2006, 04^h43'20'', orbit number 21 658. MODIS: 22 April 2006, 04^h45'. Left panel is the false color composite (red is 11 μm BT, green is 550 nm reflectance, blue is 1.6 μm reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)




Fig. 13. Northern part of Russia. AATSR: 21 April 2006, 5^h13'21", orbit number 21 644. MODIS: 21 April 2006, 5^h40'. Left panel is the false color composite (red is 11 μm BT, green is 550 nm reflectance, blue is 1.6 μm reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 14. Himalayan mountains. AATSR: 18 April 2006, 5^h17'59", orbit number 21 601. MODIS: 18 April 2006, 5^h20'. Left panel is the false color composite (red is 11 μ m BT, green is 550 nm reflectance, blue is 1.6 μ m reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 15. Antarctica, AATSR: 20 April 2006, 5^h16'00'', orbit number 20341. MODIS: 20 January 2006, 5^h00'. Left panel is the false color composite (red is 11 μm BT, green is 550 nm reflectance, blue is 1.6 μm reflectance) of initial AATSR scene; middle panel is the screened AATSR scene, the probability of clean snow; right panel is MODIS cloud mask: 1 – cloudy, 2 – probably cloudy, 3 – probably clear, 4 – clear.

The detection of cloud free snow covered areas

L. G. Istomina et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

