

**MAX-DOAS aerosol  
corrected  
tropospheric NO<sub>2</sub>  
columns**

T. Vlemmix et al.

# Retrieval of tropospheric NO<sub>2</sub> using the MAX-DOAS method combined with relative intensity measurements for aerosol correction

T. Vlemmix<sup>1,2</sup>, A. J. M. Pitters<sup>1</sup>, P. Stammes<sup>1</sup>, P. Wang<sup>1</sup>, and P. F. Levelt<sup>1,2</sup>

<sup>1</sup>Royal Netherlands Meteorological Institute, KNMI, The Netherlands

<sup>2</sup>Eindhoven University of Technology, The Netherlands

Received: 9 April 2010 – Accepted: 7 May 2010 – Published: 19 May 2010

Correspondence to: T. Vlemmix (vlemmix@knmi.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## Abstract

Multi-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) is a technique to measure trace gas amounts in the lower troposphere from ground-based scattered sunlight observations. MAX-DOAS observations are especially suitable for validation of tropospheric trace gas observations from satellite, since they have a representative range of several kilometers, both in the horizontal and in the vertical dimension.

A two-step retrieval scheme is presented here, to derive aerosol corrected tropospheric NO<sub>2</sub> columns from MAX-DOAS observations. In a first step, boundary layer aerosols, characterized in terms of aerosol optical thickness (AOT), are estimated from relative intensity observations, which are defined as the ratio of the sky radiance at elevation  $\alpha$  and the sky radiance in the zenith. Relative intensity measurements have the advantage of a strong dependence on boundary layer AOT and almost no dependence on boundary layer height. In a second step, tropospheric NO<sub>2</sub> columns are derived from differential slant columns, based on AOT-dependent air mass factors.

This two-step retrieval scheme was applied to cloud free periods in a twelve month data set of observations in De Bilt, the Netherlands. In a comparison with AERONET (Cabauw site) a mean difference in AOT (AERONET minus MAX-DOAS) of  $-0.01 \pm 0.08$  was found, and a correlation of 0.85. Tropospheric-NO<sub>2</sub> columns were compared with OMI-satellite tropospheric NO<sub>2</sub>. For ground-based observations restricted to uncertainties below 10%, no significant difference was found, and a correlation of 0.88.

## 1 Introduction

### 1.1 Validation of satellite NO<sub>2</sub> data

Tropospheric nitrogen dioxide (NO<sub>2</sub>) plays an important role in atmospheric chemistry. It is involved in many chemical cycles such as in the formation of tropospheric ozone,

AMTD

3, 2317–2366, 2010

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which is toxic to humans. High concentrations of NO<sub>2</sub> often indicate high levels of air pollution in general.

The trace gas NO<sub>2</sub> is monitored in various ways. Primarily by surface in-situ monitoring stations all over the world, but since the last decade also from space. Space borne observations of NO<sub>2</sub> form the basis for studies of regional and global trends, global transport and chemical cycles (e.g. Richter et al., 2005; van der A et al., 2006; Blond et al., 2007, and Boersma et al., 2008). In addition, satellite observations of tropospheric NO<sub>2</sub> are essential for validation of atmospheric chemical transport models and top-down construction of emission data bases (e.g. Martin et al., 2003; Mijling et al., 2009).

Despite the many results of observations from space, there is still a great demand for independent, quantitative validation of the NO<sub>2</sub> retrievals (Brinksma et al., 2008; Irie et al., 2008b; Hains et al., 2010). Validation should be performed under various atmospheric conditions and in different parts of the world, since it is known that clouds, aerosols, surface albedo, surface altitude, trace gas profile and other parameters all have significant impact on the satellite retrievals (e.g. Boersma et al., 2004; Zhou et al., 2009).

Whereas satellite validation of ozone retrievals is often done by comparison with ozone-sondes, there is not yet an equivalent in-situ profiling measurement for NO<sub>2</sub>. Comparing to in-situ surface monitors is problematic due to the large difference in spatial representativeness of the two observation techniques: In-situ surface monitors have a local character (e.g. street level), whereas even the smallest pixels of current trace gas monitoring satellite instruments have sizes of several hundreds of square kilometers. Although rural stations are representative for larger regions (Blond et al., 2007), comparison with satellite observations requires strong assumptions on the vertical distribution of NO<sub>2</sub>.

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1.2 MAX-DOAS method

The Multi-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) method (e.g. Wagner et al., 2004; Hönninger et al., 2004; Wittrock et al., 2004; Sinreich et al., 2005; Friess et al., 2006; Leigh et al., 2007; Irie et al., 2008a) offers an alternative in this respect, since it has a much larger spatial representativeness than in-situ surface monitors. MAX-DOAS instruments observe scattered solar radiation from the surface – in the UV and/or Visible – at a spectral resolution of typically 0.5 nm in multiple viewing directions (see Fig. 1). Small elevations have a relatively high sensitivity to the lower troposphere, since detected photons at small elevations have longer paths through these layers than photons observed at larger elevations. Radiative transfer simulations show that the horizontal representative range is about 5 to 10 km, whereas the vertical range is about 1 to 4 km. Both the horizontal and vertical range have a strong dependence on elevation, see e.g. Wittrock et al. (2004) and Pikelnaya et al. (2007). The increased sensitivity to the lower troposphere also distinguishes the MAX-DOAS technique from other ground-based passive DOAS techniques, such as direct-sun DOAS (total column NO<sub>2</sub>, see e.g. Herman et al., 2009), and zenith sky DOAS (stratospheric NO<sub>2</sub>, see e.g. Melo et al., 2005).

The various DOAS techniques (see Platt and Stutz, 2008, for an extensive overview) have in common that the analysis of spectral measurements of atmospheric radiation is based on the DOAS equation:

$$\ln \left( \frac{I(\lambda)}{I_{\text{ref}}(\lambda)} \right) = - \sum_{i=1}^n \Delta\sigma_i(\lambda) \Delta N_i^S + P(\lambda). \quad (1)$$

In this equation  $n$  differential cross-section spectra  $\Delta\sigma_i(\lambda)_{i=1,n}$  and a low-order polynomial  $P(\lambda)$  are fitted to the natural logarithm of the ratio of two atmospheric spectra  $I(\lambda)$ . A differential cross-section uniquely characterizes a trace gas and is obtained by subtracting a low-order polynomial from the cross-section. The two atmospheric spectra in the DOAS equation correspond to two different viewing directions, or times of

### MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observation, or both, depending on the specific DOAS application.

The fitting procedure yields a so-called differential slant column (density)  $\Delta N^S$  in [molecules/cm<sup>2</sup>] for each trace gas. The differential slant column represents the difference in trace gas absorption along the two light paths corresponding to the atmospheric spectra. In the case of MAX-DOAS, it is custom to combine a zenith spectrum with a spectrum corresponding to another, preferably small, elevation  $\alpha$ , since photons detected at small elevation have the largest path length difference with photons detected in the zenith, and this combination thus gives a high sensitivity to the lower troposphere.

In this study, MAX-DOAS observations are used for the retrieval of tropospheric columns of NO<sub>2</sub>. To convert a differential slant column to a corresponding tropospheric vertical column  $N^{\text{Tr}}$ , a so called differential air mass factor  $\Delta M$  is required. This factor is a function of the elevation  $\alpha$  – and to a lesser extent of many other parameters – and is defined here as the ratio of the differential slant column density and the tropospheric column density of NO<sub>2</sub>:

$$\Delta M_{\alpha} = \frac{\Delta N_{\alpha}^S}{N^{\text{Tr}}}. \quad (2)$$

The simplest calculation of MAX-DOAS differential air mass factors has become known as the geometrical approximation (Hönninger et al., 2004). The geometrical approximation is not based on radiative transfer simulations, but assumes that the last scattering altitude of photons detected at the surface is below the stratosphere and above the tropospheric layer of a trace gas. This assumption leads to the following relation:

$$\Delta M_{\alpha} = \frac{1 - \sin(\alpha)}{\sin(\alpha)}. \quad (3)$$

Although the geometrical approximation is known to be inaccurate for small elevations (see e.g. Wittrock et al., 2004; Pinardi et al., 2008), this approximation is believed to give an acceptable first estimate of the tropospheric column if applied to a relatively

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



large elevation, such as 30°. The geometrical approximation is used in e.g. Brinksma et al. (2008), Hains et al. (2010) and Wagner et al. (2010).

To exploit the high sensitivity of the MAX-DOAS technique at small elevations, it is desirable to have appropriate differential air mass factors for these viewing directions.

This requires a more sophisticated description of the radiative transfer than the geometrical approximation. Since photon paths through the atmosphere are affected by aerosols, it is essential to take the effect of aerosols on the differential slant columns into account.

Recently developed algorithms to estimate aerosol extinction profiles from MAX-DOAS observations, often depend on MAX-DOAS measurements of O<sub>4</sub> absorption (see e.g. Wagner et al., 2004; Sinreich et al., 2005; Friess et al., 2006; Irie et al., 2008a; Clemer et al., 2010). Absorption measurements of the collision complex of oxygen molecules, O<sub>4</sub>, can be related via inverse modeling to the aerosol extinction profile, since O<sub>4</sub> absorption depends on the photon path length through the atmosphere on which aerosols have significant impact. The profile shape of O<sub>4</sub> is well-known, it has the shape of the squared pressure profile.

### 1.3 This paper

In this study, we propose a simple algorithm for a first order aerosol correction on the differential air mass factors as an alternative to the full combined retrieval of NO<sub>2</sub> and aerosol extinction profiles based on both O<sub>4</sub> and NO<sub>2</sub> differential slant column observations. In our algorithm, aerosol characterization, in terms of aerosol optical thickness (AOT), is based on relative intensity measurements. Relative intensity observations are defined as the ratio of the sky radiance at elevation  $\alpha$  and the sky radiance in the zenith, and they have the advantage of a strong dependence on boundary layer AOT and almost no dependence on boundary layer height (Sect. 3.2). This characteristic of relative intensity observations distinguishes it from O<sub>4</sub> differential slant column measurements.

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





temperature dependence at this wavelength, which is not far from the spectral fitting window for NO<sub>2</sub> (Sect. 2.3.2).

Furthermore, the field-of-view (FOV) of the instrument was characterized (see Figs. 3 and 4). Measurements with a distant light source (−2.0 mm at 5 m) indicated that the FWHM of the FOV was around 0.45°. The pointing offset was determined from the same experiment. This pointing offset is defined as the angle between the top plane of the instrument and the horizontal plane, when the instrument is rotated such that it has maximum signal from a distant light source that is at the same height as the lens of the instrument. The pointing offset was measured each time after the instrument box was opened for maintenance and showed substantial variations (e.g. one occasion with a change from −2.0° to +0.4°) due to small displacements of the fiber entrance. The experiment was repeated including a black tube in front of the lens (not drawn), which is used normally to block stray light. Adding the tube did not change the results. The pointing offset was taken into account each time the instrument was started. The pointing of the instrument was checked on a weekly basis.

## 2.2 Operations

Observations were performed from the roof of the KNMI building in De Bilt, The Netherlands (52.101° N, 5.178° E), from 14 November 2007 until 29 April 2008 and from 11 September 2008 until 21 April 2009 with some days missing due to technical circumstances, resulting in 362 days of observations in total. The azimuth position of the instrument was fixed towards the North-East (at 46° azimuth relative to North). This direction was chosen for practical purposes (constraints at small elevations by trees and buildings surrounding the measurement site), and to look away from the sun for most of the day throughout the year, which is advantageous with respect to the sensitivity of the retrievals to estimated fixed parameters. Spectra were recorded at 0°, 2°, 4°, 8°, 16°, 30° and 90° elevation angles. Automated operation of the instrument was done with the DOASIS software developed by IUP Heidelberg in cooperation with Hoffmann GmbH. The integration time for each elevation was set to 30 s divided into

### MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







comparable to applying no correction. Not applying the dark current correction in the DOAS NO<sub>2</sub> fit will in general lead to somewhat larger fit residuals but not to systematic biases. Tests on temperature-controlled measurements, under representative measurement conditions, have shown that differences between including and not including DC-correction was less than 0.1% in the fitted NO<sub>2</sub> differential slant column. For other trace gases with smaller tropospheric column amounts, the effect of not correcting for dark current will be larger.

### 2.3.2 DOAS fitting

DOAS-fitting was performed with the “Qdoas” software developed at the Belgian Institute for Space and Aeronomy (IASB/BIRA) (Fayt and Van Roozendaal, 2001). For each spectrum at elevation  $\alpha$ , the nearest (in time) zenith spectrum was selected as a reference, in order to minimize the effect of unknown instrument instabilities on the DOAS fit. Since spectra were measured within 30 s to 2 min from the zenith measurement, the change in stratospheric path length was expected to be of little influence, except around sunrise and sunset. The cross-sections of NO<sub>2</sub> (298 K, Vandaele et al., 1997) O<sub>3</sub> (243 K, Bogumil et al., 2003) were included in the DOAS fitting routine as well as the Ring cross-section based on a solar spectrum from Kurucz et al. (1984). Cross-sections were convoluted with the measured instrumental slit function (line shape) of the instrument. The fitting interval was 415 to 431 nm. This interval was chosen because NO<sub>2</sub> has relatively little interference with other absorbers in this window and because of the relatively pronounced structures in the NO<sub>2</sub> differential cross-section. It was not possible to exploit the even more pronounced structures of NO<sub>2</sub> between 430 and 450 nm, since those are just outside the detector range of the instrument. Wavelength calibration of the spectra was done in Qdoas by applying a non-linear least squares fit of the spectrum to a high resolution solar spectrum (Kurucz et al., 1984) that was convoluted with the instrumental slit function.

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2.4 Relative intensity observations

The observation of relative intensity of skylight is another method to derive information on atmospheric constituents from the MAX-DOAS instrument. Intensity of skylight,  $I$ , is measured here as the MAX-DOAS detector signal averaged over a certain spectral interval in one viewing direction. Only relative values of intensity can be compared to their simulated counterparts, since the instrument is not radiometrically calibrated. In this work relative intensity,  $I_{\alpha}^{\text{rel}}$ , refers to the ratio of the intensity in direction  $\alpha$  to the intensity in the zenith direction:

$$I_{\alpha}^{\text{rel}} = \frac{I_{\alpha}}{I_{90}}. \quad (4)$$

The wavelength interval used for  $I_{\alpha}^{\text{rel}}$  was 426–429 nm, which is within the NO<sub>2</sub> DOAS fitting window, and contains the wavelength chosen for the differential air mass factor calculation (Sect. 3.1.3). In the absence of clouds, relative intensities in the visible are mainly influenced by Rayleigh scattering and aerosol scattering and absorption. Since Rayleigh scattering is accurately known, relative intensity observations contain information on aerosols, as will be shown in Sect. 3.2.

## 3 Retrieval algorithm

A two step approach is used to derive tropospheric vertical columns of NO<sub>2</sub> from MAX-DOAS differential slant column observations, see Fig. 5. The first step is to estimate aerosol optical thickness from relative intensity observations. This is done by interpolation of the observed relative intensity for a particular elevation and solar position (zenith and azimuth) on look-up table values of simulated relative intensity as a function of boundary layer AOT. Here it is assumed that both aerosols and tropospheric NO<sub>2</sub> are homogeneously distributed in the boundary layer. The second step is to use this

estimated AOT in another look-up table containing the NO<sub>2</sub> differential air mass factors for each elevation and solar position as a function of AOT.

By this method, the AOT and the tropospheric vertical column of NO<sub>2</sub> ( $N_{\alpha}^{\text{Tr}}$ ) are derived for each elevation independent of observations in the other elevations. A final NO<sub>2</sub> tropospheric vertical column ( $N^{\text{Tr}}$ ) is found by averaging over the elevations 4°, 8° and 16°:

$$N^{\text{Tr}} = \frac{N_{4^{\circ}}^{\text{Tr}} + N_{8^{\circ}}^{\text{Tr}} + N_{16^{\circ}}^{\text{Tr}}}{3}. \quad (5)$$

Smaller elevations were not used because the effect of the curvature of the earth is not captured by the radiative transfer model (see also Sect. 3.1.1). Larger elevations were not used because the signal-to-noise ratio of the measured differential slant column densities for this elevation was too low, namely often well below 5. This was due to the relative short integration time of 30 s, and the relative low sensitivity of the larger elevations.

As an estimate of the uncertainty on the retrieved average NO<sub>2</sub> tropospheric column we use the difference between the maximum and minimum NO<sub>2</sub> tropospheric column that is retrieved for 4°, 8° and 16° elevation (see Sect. 3.3).

### 3.1 Radiative transfer modeling

A multiple scattering radiative transfer model was used in this study for two purposes. Firstly to derive the look-up tables that relate relative intensity and differential air mass factors to boundary layer AOT. Those look-up tables are used in the retrieval algorithm. Secondly, to study the sensitivity of relative intensities and differential air mass factors to several parameters that cannot be retrieved and for which fixed (climatological) values are assumed.

The radiative transfer model used in this study is DAK (Doubling-Adding KNMI). The DAK model is based on the doubling-adding algorithm for multiple scattering of sunlight in a vertically inhomogeneous atmosphere with polarization included (De Haan

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al., 1987; Stammes et al., 1989). In the doubling-adding method, one starts with an optically thin layer for which the analytical solution for single and double scattering suffices to describe the radiation field. Next, subsequent doubling of this optically thin layer to an optically thick homogeneous layer, and adding different homogeneous layers together leads to a multi-layered atmosphere in which the reflected, transmitted, and internal radiation fields are calculated. In each layer, the extinction optical thickness, single scattering albedo and phase matrix (or phase function for unpolarized light) have to be prescribed. These atmospheric input parameters are calculated from temperature, pressure, trace gas mixing ratio, and aerosol profiles (Stammes, 2001). The inclusion of polarization in computing the radiation fields is especially important in the UV and blue parts of the solar spectrum, where atmospheric Rayleigh scattering is dominating.

### 3.1.1 Comparison with other radiative transfer models

The DAK model was compared to the results of the model intercomparison of Wagner et al. (2007), where a comparison of nine radiative transfer models was reported with special focus on box-air mass factors for MAX-DOAS viewing geometries. In general a good agreement was found between DAK and the models included in the comparison. A deviation from the spherical models was found only for box air mass factors in the Rayleigh atmosphere (no aerosols) at 577 nm and 2 degrees elevation (Fig. 8 from Wagner et al., 2007). This deviation at 577 nm was consistent with the other plane parallel models. The same comparison at 360 nm showed no difference between spherical and plane-parallel models.

Based on this comparison it was concluded that the DAK simulations could be applied to the MAX-DOAS observations at around 425 nm for elevations of 4 degrees and above.

### 3.1.2 Parameter settings

In the algorithm and radiative transfer model, several parameters were assumed fixed. These are also the standard settings in the sensitivity study of Sect. 3.2. These parameters are:

(a) Block functions to describe the aerosol extinction profile and  $\text{NO}_2$  profile in the planetary boundary layer (layer height: 1.0 km); (b) US-standard mid-latitude summer profiles for temperature and pressure at all heights and for  $\text{NO}_2$  above 5 km; (c)  $\text{NO}_2$  tropospheric vertical column is  $2 \times 10^{16}$  molecules/cm<sup>2</sup>; (d) Aerosol characterization: single scattering albedo is 0.92, asymmetry parameter is 0.70, Henyey-Greenstein phase function; (e) Surface albedo is 0.06; (f) Effect of linear polarization is included. The choice for this single scattering albedo and asymmetry parameter was based on AERONET observations from the Cabauw site (22 km from instrument location), for fourteen blue sky days in the years 2007–2009 throughout the various seasons.

### 3.1.3 Calculation of differential AMF

The differential air mass factor was calculated for one effective wavelength (428.22 nm), and this differential AMF was assumed to be representative for the whole fitting window 415–431 nm. Assumptions with respect to the model atmosphere for which differential AMFs and relative intensities were calculated, are described in the previous section.

The differential air mass factor was calculated in a way that is somewhat different from the conventional method where a differential AMF ( $\Delta M_\alpha$ ) for an elevation is found by subtracting from its AMF ( $M_\alpha$ ) the zenith-AMF ( $M_{90^\circ}$ ). In the case of this conventional method, the AMF itself is derived from two radiative transfer simulations: one for an atmosphere excluding  $\text{NO}_2$ , and another one for an atmosphere including  $\text{NO}_2$ , assuming a specific vertical distribution of  $\text{NO}_2$ .

In our study an approach is followed which is numerically equivalent to the conventional method within one percent. It was chosen because it closely resembles the MAX-DOAS method, where differential slant columns are derived directly from spectra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in two viewing directions: one spectrum measured at elevation  $\alpha$  and one zenith reference spectrum. Radiative transfer calculations were performed for just one atmosphere, including an assumed NO<sub>2</sub> vertical profile. Three wavelengths (426.48, 428.22 and 429.86 nm) were used to be able to subtract a background absorption spectrum, which also includes the low-pass filtered (or broad band) absorption by NO<sub>2</sub>. In this way, a simulated differential slant column can be derived from the ratio of the differential absorption slant optical thickness to the differential cross-section of NO<sub>2</sub>. The three wavelengths were chosen such that they were close together, at local maximum and minima of the NO<sub>2</sub> cross-sections, and within the DOAS spectral fitting window in which the measurements are analyzed (see Fig. 6).

The derivation of the differential AMF is as follows. We assume that the light paths in the atmosphere are not changed when adding a relatively weak absorber like NO<sub>2</sub>. The sky radiance for elevation  $\alpha$  and wavelength  $\lambda$  with NO<sub>2</sub> in the atmosphere ( $I_\alpha(\lambda)$ ) is equal to the sky radiance without NO<sub>2</sub> ( $I_\alpha^0(\lambda)$ ) times an attenuation term depending on the NO<sub>2</sub> slant column for this elevation ( $N_\alpha^S$ ), and the NO<sub>2</sub> absorption cross-section ( $\sigma(\lambda)$ ) at this wavelength:

$$I_\alpha(\lambda) = I_\alpha^0(\lambda) e^{-N_\alpha^S \sigma(\lambda)}, \quad (6)$$

also known as the law of Bouguer-Lambert-Beer. Taking the natural logarithm of the ratio of the radiances at elevation  $\alpha$  and the zenith ( $\alpha=90^\circ$ ), and writing:

$$R_\alpha^{(0)}(\lambda) = \ln \left[ \frac{I_\alpha^{(0)}(\lambda)}{I_{90^\circ}^{(0)}(\lambda)} \right], \quad (7)$$

and

$$\Delta N_\alpha^S = N_\alpha^S - N_{90^\circ}^S, \quad (8)$$

leads to:

$$R_\alpha(\lambda) = R_\alpha^0(\lambda) - \Delta N_\alpha^S \sigma(\lambda). \quad (9)$$

This equation may be written for the three wavelengths  $\lambda_1=426.48$  nm,  $\lambda_2=428.22$  nm and  $\lambda_3=429.86$  nm.  $R_\alpha^{(0)}$  and  $\sigma$  can be interpolated to  $\lambda_2$ , using the values at  $\lambda_1$  and  $\lambda_3$ , which leads to a second equation defined at  $\lambda_2$ . The two equations at  $\lambda_2$  are:

$$R_\alpha(\lambda_2) = R_\alpha^0(\lambda_2) - \Delta N_\alpha^S \sigma(\lambda_2), \quad (10)$$

5 and

$$R_\alpha^*(\lambda_2) = R_\alpha^{0*}(\lambda_2) - \Delta N_\alpha^S \sigma^*(\lambda_2), \quad (11)$$

where the \* refers to the interpolated values at  $\lambda_2$ . Now we assume that in this short interval,  $R^0$  can be approximated by a linear function, since it is not affected – in the model atmosphere – by absorbers with a fine-scale structure. Consequently:

$$10 \quad R_\alpha^{0*}(\lambda_2) = R_\alpha^0(\lambda_2). \quad (12)$$

If we define

$$\Delta R_\alpha(\lambda_2) = R_\alpha(\lambda_2) - R_\alpha^*(\lambda_2), \quad (13)$$

and

$$\Delta \sigma(\lambda_2) = \sigma^*(\lambda_2) - \sigma(\lambda_2), \quad (14)$$

15 (see Fig. 6), take the difference of Eqs. (10) and (11), and make use of Eq. (2), we find the following expression for the differential air mass factor ( $\Delta M_\alpha$ ) at elevation  $\alpha$ :

$$\Delta M_\alpha(\lambda_2) = \frac{\Delta R_\alpha(\lambda_2)}{N^{\text{Tr}} \Delta \sigma(\lambda_2)}. \quad (15)$$

### 3.2 Sensitivity study

20 A sensitivity study was performed to quantify the effect of variations in several parameters on the differential air mass factors, relative intensities and the combined steps in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the algorithm that lead to the retrieved tropospheric NO<sub>2</sub> column. These parameters were: boundary layer height, tropospheric NO<sub>2</sub> vertical column, aerosol optical thickness, asymmetry parameter, single scattering albedo, surface albedo and polarization. In each DAK model run only one parameter was changed with respect to the standard settings described in Sect. 3.1.2. The range of the variation of each parameter is given by columns case 1 and case 2 in Table 1.

The first column of each elevation (4°, 8°, 16°) in this table gives the sensitivity of the relative intensity to a parameter change from case 1 to case 2 (where all other parameters do not change), and each second column gives the sensitivity of the differential air mass factor. The third column of each elevation shows the sensitivity of the retrieved tropospheric NO<sub>2</sub> column for that elevation.

Table 1 shows that both the relative intensity and the differential AMF are most sensitive to the amount of boundary layer aerosols, as seen by the effect of a change in the aerosol optical thickness. The retrieval of tropospheric NO<sub>2</sub> columns however, is insensitive to a change in AOT, since the algorithm is designed to correct for this change. Figure 7 shows the effect of the AOT on relative intensity and differential AMF in more detail.

The sensitivity to a simultaneous change of both the NO<sub>2</sub> and aerosol vertical block profiles is given in the second line of Table 1. This shows that relative intensity observations – and thus the AOT retrieval – are quite insensitive to a change in boundary layer height, when compared with the sensitivity to a change in AOT (first line). However, the combined effect of relative intensity and differential AMF leads to a relatively large change in the tropospheric NO<sub>2</sub> column, especially for 4° elevation. If only the NO<sub>2</sub> vertical block profile is changed with respect to the reference situation (third line), then there is a larger change of the differential AMF and the retrieved tropospheric NO<sub>2</sub> column. This underlines that knowledge of the NO<sub>2</sub> profile shape is crucial to have an accurate tropospheric NO<sub>2</sub> column retrieval.

Measured differential slant columns are influenced not only by the vertical profile shape of NO<sub>2</sub>, but also by the vertical sensitivity to NO<sub>2</sub>. A useful quantity to describe

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the sensitivity to NO<sub>2</sub> at various altitudes is the height-dependent air mass factor, a quantity that is also referred to as (differential) box-AMF (see e.g. Hönninger et al., 2004; Wagner et al., 2007). The height-dependent differential AMF ( $\Delta m_\alpha(z)$ ) at height  $z$  for elevation  $\alpha$ , was calculated by perturbing the background NO<sub>2</sub> profile at height  $z$ :

$$\Delta m_\alpha(z) = \frac{[\Delta M_\alpha]^{z+} - [\Delta M_\alpha]^{\text{ref}}}{[N^V]^{z+} - [N^V]^{\text{ref}}}, \quad (16)$$

where the superscript “ref” refers to a simulation for a background NO<sub>2</sub> profile, and the superscript  $z+$  refers to a simulation where NO<sub>2</sub> is added to the background profile at height  $z$ .

Figure 8 shows the differential box-AMF for different elevations and for two values of the AOT (see also Wittrock et al., 2004, and Pikelnaya et al., 2007, for similar results obtained with different radiative transfer models). The low elevations have a sensitivity to NO<sub>2</sub> that decreases rapidly with height, whereas the vertical sensitivity of the higher elevations is more constant. Increasing the amount of aerosols in the boundary layer leads, for the low elevations, to a pronounced decrease in sensitivity to NO<sub>2</sub> with increasing altitude.

Figure 9 shows the dependence of relative intensity on boundary layer height and aerosol optical thickness, for two elevations (4° and 30°). The relatively weak sensitivity of relative intensity to a change in boundary layer height, is the reason that relative intensity observations are more suitable for boundary layer aerosol optical thickness estimations than measurements of O<sub>4</sub>. A relative intensity observation of only a single elevation is needed to estimate the aerosol optical thickness. O<sub>4</sub> observations on the other hand, contain more information on the aerosol profile, which is the reason why they are used in MAX-DOAS aerosol extinction profile retrievals (Friess et al., 2006; Irie et al., 2008a; Clemer et al., 2010).

From the fourth line in Table 1 it appears that there is an unwanted dependence of the algorithm to the tropospheric NO<sub>2</sub> column itself. This can be solved by making the algorithm iterative, with the tropospheric NO<sub>2</sub> column as an additional dimension of the

**MAX-DOAS aerosol  
corrected  
tropospheric NO<sub>2</sub>  
columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



look-up table, but this step was not applied in this study.

Finally, Table 1 shows that the asymmetry parameter, single scattering albedo, surface albedo and polarization have a relatively small effect on the retrieved tropospheric NO<sub>2</sub> column.

Table 1 was calculated for a relative azimuth of 180° and a solar zenith angle of 60°. Additional studies showed that the values in the table are representative for other solar positions except when the instrument is viewing in a direction close to the sun (either in viewing direction  $\alpha$ , or in the zenith direction) where simulated quantities depend more critically on the aerosol model parameters. Since the Mini MAX-DOAS instrument was pointed towards the North-East, the sun was at relatively large angular distance for most of the time.

### 3.3 Error sources

There are many possible sources of error in the retrieval of NO<sub>2</sub> tropospheric columns from MAX-DOAS observations. Two types of error may be distinguished: observational errors and modeling errors. The term observational error is used here for all factors leading to an incorrect value for the differential slant column and/or relative intensity.

Systematic error contributions to the observational error are: (1) errors caused by incorrect knowledge of the actual field-of-view, which may be caused by incorrect aiming of the instrument (e.g. when it is unattended after periods of heavy winds) or by imprecise knowledge of the offset in the field-of-view as described in Sect. 2.1, (2) incorrect electronic offset correction of raw spectra, (3) errors in the differential cross-sections of NO<sub>2</sub> (e.g. temperature dependency). The DOAS spectral-fitting error represents a mixture of systematic and random errors: incorrect wavelength calibration of spectra, inaccurate knowledge of instrument slit function, incorrect dark current correction, unknown absorbers, and low signal-to-noise.

The DOAS fitting was performed with a cross section at a fixed temperature (see Sect. 2.3.2). This introduces an error in the differential slant column of NO<sub>2</sub> that is proportional to the temperature difference between the fixed temperature of the cross

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



section used in the fit and the effective temperature of the tropospheric NO<sub>2</sub>. Although the NO<sub>2</sub> cross section  $\sigma_{\text{NO}_2}$  itself is not strongly temperature dependent, the differential cross section  $\Delta\sigma_{\text{NO}_2}$  shows a much stronger temperature dependence: a change in temperature of 20 degrees corresponds to a change in  $\sigma_{\text{NO}_2}$  of 1.6% and to a change in  $\Delta\sigma_{\text{NO}_2}$  of 7.2%. An estimate of the effective NO<sub>2</sub> temperature is needed to correct for this effect (see Sect. 4.4).

The modeling errors that affect the differential AMF are introduced in two ways: first by not including parameters in the radiative transfer modeling that do have an effect on the observations, and second by choosing inappropriate values for parameters that are included in the model. Some examples of parameters that were not included in the model are: humidity, clouds, horizontal gradients in all parameters of which a vertical profile is prescribed, and vertical profile shapes other than those described in Sect. 3.1.2.

The effect of inaccurate estimation of the included model parameters was studied in Sect. 3.2. The algorithm described in this study is most sensitive to the boundary layer height, especially to the vertical profile shape of NO<sub>2</sub>. This is a consequence of the decrease in sensitivity to NO<sub>2</sub> with increasing height, especially for low elevations (see e.g. Wittrock et al., 2004, their Fig. 4).

In this study, we estimate the uncertainty  $\varepsilon$  on the retrieved average NO<sub>2</sub> tropospheric column (Eq. 5) as the difference between the maximum and minimum NO<sub>2</sub> tropospheric column that is retrieved for 4°, 8° and 16° elevation:

$$\varepsilon = \max\left(N_{[4^\circ, 8^\circ, 16^\circ]}^{\text{Tr}}\right) - \min\left(N_{[4^\circ, 8^\circ, 16^\circ]}^{\text{Tr}}\right), \quad (17)$$

where the three tropospheric columns are interpolated to the same point in time. Since the tropospheric columns are derived for each elevation independent of the others, this difference, or spread, gives an important indication of the internal consistency of the retrieval. A small spread indicates a consistent retrieval. The spread increases for measurement conditions that differ from the model atmosphere in the look-up table calculations. For instance, an error in the assumed boundary layer height would

**MAX-DOAS aerosol  
corrected  
tropospheric NO<sub>2</sub>  
columns**

T. Vlemmix et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lead to different systematic errors in the retrieved tropospheric NO<sub>2</sub> columns for each elevation angle (see Table 1). This would result in a systematic error in the derived average tropospheric NO<sub>2</sub> column and an increase in the spread. The definition of the measurement uncertainty includes effects such as: presence of clouds, pointing elevation offsets, horizontal gradients, vertical profile shape of aerosols and NO<sub>2</sub>, and uncertainty due to a low signal-to-noise.

Typical values of  $\varepsilon$  are on average much lower for clear sky than for cloudy conditions (see Sect. 4.1). If data is selected for clear sky conditions, using the criterion that the relative intensity of 4°, 8° and 16° is  $> 1$ , then the median of the measurement uncertainty  $\varepsilon$  is  $2.7 \times 10^{15}$  molecules/cm<sup>2</sup> or 17% relative to a mean tropospheric NO<sub>2</sub> column of  $15.6 \times 10^{15}$  molecules/cm<sup>2</sup>.

## 4 Results

In this section various results of our new two-step algorithm will be shown. Firstly, several steps in the tropospheric NO<sub>2</sub> retrieval algorithm will be illustrated for three selected days, and AOT and tropospheric NO<sub>2</sub> retrievals will be shown for four more days. Secondly, the algorithm is compared to the geometrical approximation, which is the default approach to derive tropospheric columns from MAX-DOAS observations. Finally a comparison with AOT from an AERONET instrument and a comparison with OMI tropospheric NO<sub>2</sub> is shown.

### 4.1 AOT and tropospheric NO<sub>2</sub> for selected days

Figure 10 shows the differential slant columns of NO<sub>2</sub>, relative intensity, aerosol optical thickness and tropospheric NO<sub>2</sub> columns for three days, to illustrate the different steps in the retrieval. The first day (12 October 2008) is selected to illustrate the effect of clouds on the retrieval. Until around 13:00 UTC, clouds were present above the measurement location. This can be seen most clearly from the relative intensity

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations. Under a homogeneous cloud cover, the intensity of scattered light is higher in the zenith direction than in other directions, and is decreasing with decreasing elevation. This results in relative intensities  $< 1$ . After the clouds disappear (around 13:00 UTC) the situation is opposite. Under clear sky conditions the intensity of the scattered sunlight above the horizon is generally higher than the intensity in the zenith; the maximum intensity occurs usually somewhere between  $\alpha=8^\circ$  and  $\alpha=25^\circ$ , depending on e.g. solar position and AOT (see Fig. 7). For this reason relative intensity  $> 1$  is a good first indication for clear sky conditions, although this is not true in general: for small solar zenith angles the zenith sky may be brighter than the horizon.

It is to be expected that relative intensity cannot be used to retrieve AOT under cloudy circumstances since the cloud has a much stronger effect on the relative intensity than the aerosols. As a consequence, AOT values larger than the maximum AOT of the look-up table (0.8) are retrieved before 13:00 UTC on 12 October 2008. At later times the three elevations ( $4^\circ$ ,  $8^\circ$ ,  $16^\circ$ ) give similar AOT values. The effect of too large AOTs can also be seen in the NO<sub>2</sub> tropospheric column retrieval for this day. However, a sudden change between 12:00 and 13:00 UTC cannot be seen in the differential slant columns. This illustrates that differential slant column measurements are mostly sensitive to the lowest kilometers of the atmosphere (whether or not below a cloud). The other two days in Fig. 10 (26 December 2008 and 27 January 2009) are included as an example of both a relatively clean and a relatively polluted day. The clean day, December 26, was a public holiday with temperatures below  $0^\circ\text{C}$ . It shows stable conditions in both the tropospheric NO<sub>2</sub> column and AOT.

Four more examples of retrievals for cloud free days are shown in Fig. 11. The spread in the tropospheric columns of NO<sub>2</sub> between the three different elevations is relatively small, about 10%. Possible causes of the spread are discussed in Sect. 3.3.

## 4.2 Comparison with geometrical air mass factor approximation

The geometrical approximation (GA) of the differential air mass factor for MAX-DOAS observations provides a simple way to determine a first order estimate of the

tropospheric column of NO<sub>2</sub>, or other trace gases that are located primarily in the boundary layer. It is applied to MAX-DOAS observations in e.g. Brinksma et al. (2008) and Hains et al. (2010) in a comparison with other methods to measure tropospheric NO<sub>2</sub>, such as lidar and satellite.

Using the geometric approximation is simple: it does not require an inversion based on radiative transfer modeling. The question remains how accurate this approximation is. This issue has been discussed by e.g. Hönninger et al. (2004), Wittrock et al. (2004), and Pinardi et al. (2008), based primarily on radiative transfer modeling results.

Here we follow a different approach, partly based on observations. Simulations with DAK show that, depending on the boundary layer aerosol load, large differences may occur between the geometrical and modeled differential air mass factors at low elevations:  $\alpha \leq 16^\circ$ , (see Fig. 7, right plot). Therefore the GA should not be used for these elevations. For higher elevations, the difference becomes much smaller. It seems from Fig. 7 that the radiative transfer model and the GA have almost the same differential air mass factor for  $30^\circ$  elevation. However, it can be seen from Fig. 12 that even for this high elevation, the difference between the GA and the model may become as large as 25%, depending on the relative position of the sun. At smaller relative azimuths this relative difference is even higher.

The effect of the difference between the geometrical approximation and radiative-transfer based and aerosol corrected differential air mass factors can also be observed in the measurements. Figure 13 shows that the tropospheric column of NO<sub>2</sub> derived from the GA for  $\alpha=30^\circ$ , is not in agreement with the tropospheric column derived with the algorithm using the mean of  $\alpha=4^\circ$ ,  $8^\circ$  and  $16^\circ$ . The algorithm finds a NO<sub>2</sub> tropospheric column that is significantly different from the geometrical approximation, with differences up to 20%.

To find out whether this difference could be due to the difference in differential slant columns, the tropospheric NO<sub>2</sub> column is calculated from the differential slant column at  $30^\circ$  (that were only used before in combination with the GA) with the differential air mass factor calculated according to the two-step algorithm. Thus for this specific study,

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





the algorithm is applied to 30° elevation alone. Figure 13 shows that this method to calculate the tropospheric NO<sub>2</sub> vertical column is better in agreement with the tropospheric columns derived with the standard algorithm than the GA.

It can be concluded that the geometrical approximation is not sufficiently accurate for tropospheric NO<sub>2</sub> retrieval. However, it can be used for a quick indicative conversion from the differential slant column at 30° to a tropospheric column. A disadvantage in using this (relatively high) elevation is the reduced sensitivity to the lower troposphere.

### 4.3 Verification of AOT with AERONET data

As a verification of the AOT retrieved by the new two-step algorithm, a comparison is shown in Fig. 14 between the AOT data from the MAX-DOAS observations of relative intensity at De Bilt and the AOT data from the Aerosol Robotic Network (AERONET, see Dubovik and King, 2000) instrument at Cabauw (22 km from De Bilt). For all days in the operational period (see Sect. 2.2) level 1.5 AERONET AOT data at 440 nm was selected (if available) and compared with MAX-DOAS AOT data if the uncertainty in this value – i.e. the spread between AOTs derived from  $\alpha=4^\circ$ ,  $8^\circ$  and  $16^\circ$  – was below 0.1. This selection resulted in 1251 data points with a correlation of 0.85. The mean difference between the AERONET and MAX-DOAS AOT was  $-0.01$  with a standard deviation of 0.08. Next, a linear regression was done that minimizes the sum of the squared orthogonal distances. From this regression a slope of 1.01 was found and an offset of  $-0.01$ . Considering the difference in observation method (direct sunlight versus scattered sunlight), and the distance between the two sites, this agreement is satisfying.

### 4.4 Comparison with satellite observations

As a first application of the time series of tropospheric NO<sub>2</sub> columns derived from the Mini MAX-DOAS measurements in De Bilt, an inter-comparison was made with tropospheric NO<sub>2</sub> data (DOMINO-product, see Boersma et al., 2007) from the Ozone

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Monitoring Instrument (OMI, see Levelt et al., 2006). The DOMINO data selection was based on the following criteria: the satellite cloud fraction should be below 0.2 and the NO<sub>2</sub> slant column should be below  $2 \times 10^{17}$  molec/cm<sup>2</sup>. Pixels affected by the OMI row-anomaly were removed. A coincident ground and satellite observation was defined as a measurement where the ground site was within the satellite pixel (which has an approximate rectangular shape, ranging from  $13 \times 24$  km<sup>2</sup> at nadir to  $26 \times 135$  km<sup>2</sup> at the edge of the swath (Dobber et al., private communication)). The ground-based tropospheric column was averaged over 15 min around the time of satellite overpass.

A correction was applied to account for the fixed temperature dependence of the NO<sub>2</sub> cross section (see Sect. 2.3.2). Two steps were taken to estimate the effective NO<sub>2</sub> temperature at the time of observation.

First the difference was determined between the surface temperature and the effective NO<sub>2</sub> temperature  $T_{\text{NO}_2}^{\text{eff}}$ :

$$T_{\text{NO}_2}^{\text{eff}} = \frac{\int T(z)m(z)n(z)dz}{\int m(z)n(z)dz}, \quad (18)$$

where  $m(z)$  is the the height-dependent differential air mass factor for an AOT of 0.2 (see Fig. 8),  $n(z)$  is a block profile for NO<sub>2</sub> from 0–1 km, and  $T(z)$  is a standard temperature profile. A difference of about  $-2.5$  °C was found between the surface temperature and the effective NO<sub>2</sub> temperature for all three elevations 4°, 8° and 16°.

Then surface temperature data were taken from KNMI temperature observations in De Bilt, and from this 2.5 °C was subtracted to determine the effective NO<sub>2</sub> temperature. The ratio of  $\Delta\sigma$  (see Fig. 6) at the temperature of the NO<sub>2</sub> cross section in the DOAS-fit and  $\Delta\sigma$  at the effective NO<sub>2</sub> temperature was applied as a temperature correction factor to the MAX-DOAS tropospheric NO<sub>2</sub> columns.

Figure 15 shows a scatter plot of the comparison. Three different selections of the MAX-DOAS data were made, based on different constraints on the (relative) spread of the ground-based tropospheric column of NO<sub>2</sub> (see Table 2). From 362 days of ground-based observations (see Sect. 2.2) only 123 data points remain after applying

the selection criteria on the satellite data (including coincidence with the ground-based observation). Only 17 ground-based observations pass the 10% threshold, as described in Table 2.

The table shows that the correlation between the ground-based and satellite data improves with a more strict selection of the ground-based observations. However, when the estimated uncertainty (spread) of the ground-based observations is less than 10%, the standard deviation of the differences between the two data sets is about 25% ( $3.8 \times 10^{15}$  molec/cm<sup>2</sup> for a mean value of about  $15 \times 10^{15}$  molec/cm<sup>2</sup>).

To test if the spread between the OMI and the MAX-DOAS tropospheric NO<sub>2</sub> columns is dominated by the estimated retrieval errors from both data sets, a  $\chi^2$  test was applied:

$$\chi_v^2 = \frac{1}{N} \sum_{i=1}^N \frac{(x_i - y_i)^2}{\varepsilon_{x_i}^2 + \varepsilon_{y_i}^2}, \quad (19)$$

where  $N$  is the number of data points  $(x_i, y_i)$ ,  $x_i$  is an OMI tropospheric NO<sub>2</sub> column measurement, with retrieval error  $\varepsilon_{x_i}$ , and  $y_i$  is a MAX-DOAS tropospheric NO<sub>2</sub> column measurement with retrieval error  $\varepsilon_{y_i}$  estimated from the spread as described in Eq. (17). For the three selections of the data, as described in Table 2,  $\chi_v^2$  is between 2.5 and 3. Assuming an average difference of zero between the data sets and normal error distributions, the probability of exceeding these values for  $\chi_v^2$  is less than 0.1%. Hence, the spread is larger than can be expected from the estimated retrieval errors alone.

A possible explanation for the part of the spread that is not explained by the retrieval errors, is the difference in observation techniques. First, the spatial representativity of the two types of observation is quite different. The horizontal footprint of an OMI pixel is different from the horizontal domain of the MAX-DOAS observation. Whereas OMI samples a domain of  $> 300$  km<sup>2</sup>, the MAX-DOAS has a horizontal range – in one direction – of roughly 10 km for the lowest elevation of 4°. Also the vertical range that

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**T. Vlemmix et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

contributes to the tropospheric NO<sub>2</sub> signal is different for the satellite and the MAX-DOAS observations. The sensitivity of MAX-DOAS to NO<sub>2</sub> decreases quickly with increasing height of NO<sub>2</sub>, especially for high aerosol loads (see Fig. 8), whereas the satellite has increasing sensitivity with increasing NO<sub>2</sub> height (Eskes and Boersma, 2003). Consequently, the relative contribution of the free-tropospheric NO<sub>2</sub> to the total tropospheric NO<sub>2</sub> is different for the satellite than for MAX-DOAS.

Separation of the free tropospheric and boundary layer contribution to the tropospheric NO<sub>2</sub> column requires accurate knowledge of the NO<sub>2</sub> profile shape, both for the satellite and the MAX-DOAS retrieval. Lack of knowledge of the vertical distribution of NO<sub>2</sub> thus complicates the comparison of satellite and MAX-DOAS NO<sub>2</sub>-data as well as the interpretation of satellite retrievals in terms of surface concentrations.

More observations and understanding of vertical profiles of NO<sub>2</sub> are needed to study the variety of circumstances under which differences between satellite and ground-based observations occur.

## 5 Conclusions

We described a new two-step algorithm to retrieve aerosol corrected tropospheric NO<sub>2</sub> columns from MAX-DOAS observations.

We used relative intensity observations performed with a mini MAX-DOAS instrument to estimate the aerosol optical thickness of the boundary layer. Based on this AOT-estimation, aerosol corrected differential air mass factors for NO<sub>2</sub> were determined to convert differential slant columns of NO<sub>2</sub> to tropospheric columns.

Relative intensity measurements have a strong dependence on boundary layer AOT and almost no dependence on boundary layer height although this dependence increases slightly with decreasing elevation.

Since relative intensity observations are very sensitive to clouds, the retrieval method can only be applied to cloud-free conditions. Under cloud-free conditions, the main source of error is the assumption of a fixed boundary layer height for NO<sub>2</sub> in the

---

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

---

T. Vlemmix et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

algorithm. An uncertainty estimate in the results is derived from the spread in the tropospheric NO<sub>2</sub> columns that are derived independently for each elevation (4°, 8° and 16°). A low spread indicates a consistent retrieval of the tropospheric NO<sub>2</sub> column for the three elevations.

5 The use of the relatively low elevations makes the retrieval method more sensitive to trace gases in the boundary layer than the often used geometrical approximation that can only be applied at higher elevations (30°). The geometrical air mass factor approximation for 30° elevation gives a good first estimate of the tropospheric column. However, even for this high elevation, the differential air mass factor of the geometrical  
10 approach may differ up to 25% from the differential air mass factors using radiative transfer modeling. For relative azimuths smaller than 40°, this difference may even be larger.

The relatively low-cost (Mini) MAX-DOAS instruments are capable of generating long time-series of tropospheric composition data in an automated fashion. MAX-DOAS observations are particularly valuable for the purpose of satellite validation of tropospheric trace gases due to the horizontal and vertical range where trace gases can be detected, which is on the order of 1 to 4 km in the vertical and 5 to 10 km in the horizontal, depending on the AOT. Accurate comparison of satellite and ground-based observations requires knowledge of the NO<sub>2</sub> profile shape, in order to account for differences in the sensitivity to NO<sub>2</sub> at different heights.  
20

The two-step retrieval scheme presented here was applied to cloud-free periods in a twelve month data set of MAX-DOAS observations in De Bilt, the Netherlands, between Autumn 2007 and Spring 2009 (summer of 2008 not included). For cloud free periods, the average tropospheric NO<sub>2</sub> column was  $15.6 \times 10^{15}$  molecules/cm<sup>2</sup>. The median of the estimated relative uncertainty was 17%.  
25

In a comparison with AERONET (Cabauw site, 22 km from De Bilt) a mean difference in AOT (AERONET minus MAX-DOAS) of  $-0.01 \pm 0.08$  was found, and a correlation of 0.85.

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tropospheric NO<sub>2</sub> columns were compared with OMI satellite tropospheric NO<sub>2</sub>. Only satellite pixels over De Bilt were selected. The spread in the tropospheric NO<sub>2</sub> columns retrieved (semi)-simultaneously from the different MAX-DOAS elevations was used as a selection criterion. For ground-based observations restricted to a spread below 10%, a correlation of 0.88 was found, and no significant difference. The spread between OMI and MAX-DOAS tropospheric NO<sub>2</sub> column measurements is larger than can be expected from the estimated retrieval errors alone. This may be due to differences in the spatial representativity of the two observation techniques.

*Acknowledgements.* The Authors would like to thank M. Van Roozendaal, C. Fayt and G. Pinardi from the Belgian Institute for Space and Aeronomy (IASB/BIRA) for providing the Qdoas software that was used for the DOAS analysis, and for giving valuable support and advice.

Furthermore we would like to thank S. Kraus and T. Lehmann of the Institute for Environmental Physics at the University of Heidelberg for providing the DOASIS software package.

We would like to thank K. F. Boersma, J. de Haan, M. Allaart and M. Dobber for useful discussions on this study.

We acknowledge the free use of tropospheric NO<sub>2</sub> column data from the OMI sensor from [www.temis.nl](http://www.temis.nl).

We acknowledge the efforts of the AERONET team, and in particular the TNO team lead by G. de Leeuw for the measurement and provision of AOT data.

Finally, we acknowledge the support of the European Commission through the GEOmon (Global Earth Observation and Monitoring) Integrated Project under the 6th Framework Program (contract number FP6-2005-Global-4-036677).

This work has been financed by User Support Program Space Research via the project “Atmospheric chemistry instrumentation to strengthen satellite validation of CESAR” (EO-091).

## References

- Blond, N., Boersma, K. F., Eskes, H. J., van der A, R. J., Van Roozendael, M., De Smedt, I., Bergametti, G., and Vautard, R.: Intercomparison of SCIAMACHY nitrogen dioxide observations, in situ measurements and air quality modeling results over Western Europe, *J. Geophys. Res.*, 112, D10311, doi:10.1029/2006JD007277, 2007. 2319
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO<sub>2</sub> retrieval from space, *J. Geophys. Res.*, 109, D04311, doi:10.1029/2003JD003962, 2004. 2319
- Boersma, K. F., Eskes, H. J., Veefkind, J. P., Brinksma, E. J., van der A, R. J., Sneep, M., van den Oord, G. H. J., Levelt, P. F., Stammes, P., Gleason, J. F., and Bucsela, E. J.: Near-real time retrieval of tropospheric NO<sub>2</sub> from OMI, *Atmos. Chem. Phys.*, 7, 2103–2118, doi:10.5194/acp-7-2103-2007, 2007. 2340
- Boersma, K. F., Jacob, D. J., Eskes, H. J., Pinder, R. W., Wang, J., and van der A, R. J.: Intercomparison of SCIAMACHY and OMI tropospheric NO<sub>2</sub> columns: Observing the diurnal evolution of chemistry and emissions from space, *J. Geophys. Res.*, 113, D16S26, doi:10.1029/2007JD008816, 2008. 2319
- Bogumil, K., Orphal, J., Homann, T., Voigt, S., Spietz, P., Fleischmann, O. C., Vogel, A., Hartmann, M., Kromminga, H., Bovensmann, H., Frerick, J., and Burrows, J. P.: Measurements of Molecular Absorption Spectra with the SCIAMACHY Pre-Flight Model: Instrument Characterization and Reference Data for Atmospheric Remote-Sensing in the 230–2380 nm Region, *J. Photochem. Photobiol. A.*, 157, 167–184, 2003. 2326
- Brinksma, E. J., Pinardi, G., Volten, H., Braak, R., Richter, A., Schoenhardt, A., Van Roozendael, M., Fayt, C., Hermans, C., Dirksen, R. J., Vlemmix, T., Berkhout, A. J. C., Swart, D. P. J., Oetjes, H., Wittrock, F., Wagner, T., Ibrahim, O., de Leeuw, G., Moerman, M., Curier, R. L., Celarier, E. A., Cede, A., Knap, W. H., Veefkind, J. P., Eskes, H. J., Allaart, M., Rothe, R., Pipers, A. J. M., and Levelt, P. F.: The 2005 and 2006 DANDELIONS NO<sub>2</sub> and aerosol intercomparison campaigns, *J. Geophys. Res.*, 113, D16S46, doi:10.1029/2007JD008988, 2008. 2319, 2322, 2339
- Clémer, K., Van Roozendael, M., Fayt, C., Hendrick, F., Hermans, C., Pinardi, G., Spurr, R., Wang, P., and De Mazière, M.: Multiple wavelength retrieval of tropospheric aerosol optical properties from MAXDOAS measurements in Beijing, *Atmos. Meas. Tech. Discuss.*, 3, 111–145, doi:10.5194/amtd-3-111-2010, 2010. 2322, 2334
- De Haan, J. F., Bosma, P. B., and Hovenier, J. W.: The adding method for multiple scattering

AMTD

3, 2317–2366, 2010

### MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


calculations of polarized light, *Astron. Astrophys.*, 183, 371–393, 1987. 2328

Dubovik, O. and King, M. D.: A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and sky radiance measurements, *J. Geophys. Res.*, 105(D16), 673–696, doi:10.1029/2000JD900282, 2000. 2340

5 Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285–1291, doi:10.5194/acp-3-1285-2003, 2003. 2343

Fayt, C. and Van Roozendaal, M.: WinDOAS 2.1: Software User Manual, <http://www.oma.be/BIRA-IASB/Molecules/BrO/WinDOAS-SUM-210b.pdf>, 2001. 2326

10 Friess, U., Monks, P. S., Remedios, J. J., Rozanov, A., Sinreich, R., Wagner, T., and Platt, U.: MAX-DOAS O<sub>4</sub> measurements: A new technique to derive information on atmospheric aerosols: 2. Modeling studies, *J. Geophys. Res.*, 111, D14203, doi:10.1029/2005JD006618, 2006. 2320, 2322, 2334

15 Hains, J., Boersma, K. F., Kroon, M., and Dirksen, R. J.: Testing and Improving OMI DOMINO Tropospheric NO<sub>2</sub> Using Observations from the DANDELIONS and INTEX-B Validation Campaigns, *J. Geophys. Res.*, 115, D05301, doi:10.1029/2009JD012399, 2010. 2319, 2322, 2339

Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO<sub>2</sub> column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, *J. Geophys. Res.*, 20 114, D13307, doi:10.1029/2009JD011848, 2009. 2320

Hönninger, G., von Friedeburg, C., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS), *Atmos. Chem. Phys.*, 4, 231–254, doi:10.5194/acp-4-231-2004, 2004. 2320, 2321, 2323, 2334, 2339

25 Irie, H., Kanaya, Y., Akimoto, H., Iwabuchi, H., Shimizu, A., and Aoki, K.: First retrieval of tropospheric aerosol profiles using MAX-DOAS and comparison with lidar and sky radiometer measurements, *Atmos. Chem. Phys.*, 8, 341–350, doi:10.5194/acp-8-341-2008, 2008a. 2320, 2322, 2334

Irie, H., Kanaya, Y., Akimoto, H., Tanimoto, H., Wang, Z., Gleason, J. F., and Bucsela, E. J.: Validation of OMI tropospheric NO<sub>2</sub> column data using MAX-DOAS measurements deep inside the North China Plain in June 2006: Mount Tai Experiment 2006, *Atmos. Chem. Phys.*, 30 8, 6577–6586, doi:10.5194/acp-8-6577-2008, 2008b. 2319

Kurucz, R. L., Furenliid, I., and Testerman, L.: Solar Flux Atlas from 296 to 1300 nm, Technical Report, National Solar Observatory, 1984. 2326



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Leigh, R. J., Corlett, G. K., Frieß, U., and Monks, P. S.: Spatially resolved measurements of nitrogen dioxide in an urban environment using concurrent multi-axis differential optical absorption spectroscopy, *Atmos. Chem. Phys.*, 7, 4751–4762, doi:10.5194/acp-7-4751-2007, 2007. 2320
- 5 Levelt, P. F., van den Oord, G. H. J., Dobber, M., Mälkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J., and Saari, H.: The Ozone Monitoring Instrument, *IEEE Trans. Geo. Rem. Sens.*, 44, 1093–1101, 2006. 2341
- Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO<sub>2</sub> columns, *J. Geophys. Res.*, 108, D17, doi:10.1029/2003JD003453, 2003. 2319
- 10 Melo, S. M. L., Strong, K., Bassford, M. R., Preston, K. E., McElroy, C. T., Rozanov, E. V., and Egorova, T.: Retrieval of Stratospheric NO<sub>2</sub> Vertical Profiles from Ground-Based Zenith-Sky DOAS Measurements: Results for the MANTRA 1998 Field Campaign, *Atmosphere-Ocean*, 43, 340–350, 2005. 2320
- 15 Mijling, B., van der A, R. J., Boersma, K. F., Van Roozendael, M., Desmedt, I., and Kelder, H. M.: Reductions of NO<sub>2</sub> Detected from Space During the 2008 Beijing Olympic Games, *Geophys. Res. Lett.*, 36, L13801, doi:10.1029/2009GL038943, 2009. 2319
- Pikelnaya, O., Hurlock, S. C., Trick, S., and Stutz, J.: Intercomparison of multi-axis and long-path optical absorption spectroscopy measurements in the marine boundary layer, *J. Geophys. Res.*, 112, D10S01, doi:10.1029/2006JD007727, 2007. 2320, 2334
- 20 Pinardi, G., Hendrick, F., Clemer, K., Lambert, J. C., Bai, J., and Van Roozendael, M.: On the use of the MAX-DOAS technique for the validation of tropospheric NO<sub>2</sub> column measurements from satellite, *Proc. Eumetsat Conf.*, 2008. 2321, 2339
- Platt, U. and Stutz, J.: *Differential Optical Absorption Spectroscopy*, Springer-Verlag Berlin Heidelberg, 2008. 2320
- 25 Richter, A., Burrows, J. P., Nuess, H., Ganier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129–132, 2005. 2319
- Sinreich, R., Friess, U., Wagner, T., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS) of gas and aerosol distributions, *Faraday Discuss.*, 130, 153–164, 2005. 2320, 2322, 2323
- 30 Stammes, P.: Spectral radiance modeling in the UV-visible range, *IRS2000: Current problems in atmospheric radiation*, Eds. W.L. Smith and Y.M. Timofeyev, A. Deepak Publ., Hampton (VA), 1, 385–388, 2001. 2329



**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Stammes, P., de Haan, J. F., and Hovenier, J. W.: The polarized internal radiation field of a planetary atmosphere, *Astron. Astrophys.*, 225, 239–259, 1989. 2329
- van der A, R. J., Peters, D. H. M. U., Eskes, H., Boersma, K. F., Van Roozendael, M., DeSmedt, I., and Helder, H. M.: Detection of the trend and seasonal variation in tropospheric NO<sub>2</sub> over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594, 2006. 2319
- 5 Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., Merienne, M. F., Jenouvrier, A., and Coquart, B.: Measurements of the NO<sub>2</sub> Absorption Cross-section from 42 000 cm<sup>-1</sup> to 10 000 cm<sup>-1</sup> (238–1000 nm) at 220 K and 298 K, *J. Quant. Spectr. Radiat. Transfer*, 59, 171–184, 1997. 2326
- 10 Wagner, T., Dix, B., von Friedeburg, C., Friess, U., Sanghavi, S., Sinreich, R., and Platt, U.: MAX-DOAS O<sub>4</sub> measurements: A new technique to derive information on atmospheric aerosols - Principles and information content, *J. Geophys. Res.*, 109, D22205, doi:10.1029/2004JD004904, 2004. 2320, 2322
- 15 Wagner, T., Burrows, J. P., Deutschmann, T., Dix, B., von Friedeburg, C., Frieß, U., Hendrick, F., Heue, K.-P., Irie, H., Iwabuchi, H., Kanaya, Y., Keller, J., McLinden, C. A., Oetjen, H., Palazzi, E., Petritoli, A., Platt, U., Postlyakov, O., Pukite, J., Richter, A., van Roozendael, M., Rozanov, A., Rozanov, V., Sinreich, R., Sanghavi, S., and Wittrock, F.: Comparison of box-air-mass-factors and radiances for Multiple-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) geometries calculated from different UV/visible radiative transfer models, *Atmos. Chem. Phys.*, 7, 1809–1833, doi:10.5194/acp-7-1809-2007, 2007. 2329, 2334
- 20 Wagner, T., Ibrahim, O., Shaiganfar, R., and Platt, U.: Mobile MAX-DOAS observations of tropospheric trace gases, *Atmos. Meas. Tech.*, 3, 129–140, doi:10.5194/amt-3-129-2010, 2010. 2322
- 25 Wittrock, F., Oetjen, H., Richter, A., Fietkau, S., Medeke, T., Rozanov, A., and Burrows, J. P.: MAX-DOAS measurements of atmospheric trace gases in Ny-Ålesund – Radiative transfer studies and their application, *Atmos. Chem. Phys.*, 4, 955–966, doi:10.5194/acp-4-955-2004, 2004. 2320, 2321, 2334, 2336, 2339
- 30 Zhou, Y., Brunner, D., Boersma, K. F., Dirksen, R., and Wang, P.: An improved tropospheric NO<sub>2</sub> retrieval for OMI observations in the vicinity of mountainous terrain, *Atmos. Meas. Tech.*, 2, 401–416, doi:10.5194/amt-2-401-2009, 2009. 2319

**Table 1.** Sensitivity study of six parameters affecting relative intensity and the tropospheric NO<sub>2</sub> differential air mass factor (see Sect. 3.2): aerosol optical thickness (AOT), boundary layer height for NO<sub>2</sub> and aerosols (BLH), boundary layer column of NO<sub>2</sub> (N), asymmetry parameter of aerosols (ASY), single scattering albedo of aerosols (SSA), surface albedo (ALB), and polarization (POL). Each parameter was changed in the DAK model from case 1 (reference value) to case 2, with all other parameters unchanged. For the elevations 4°, 8° and 16°, the effect of this change in percent is given for the relative intensity ( $I^{\text{rel}}$ ), for the differential air mass factor ( $\Delta M$ ) and for the differential air mass factor calculated by the two-step algorithm ( $\Delta M^{\text{alg}}$ ). The percentage was calculated as follows:  $[P(\text{case 1})-P(\text{case 2})]/P(\text{case 2})\times 100\%$ , where  $P(\text{case 2})$  is the model simulation of quantity  $P$  for parameter setting case 2. Values were calculated for a solar zenith angle of 60° and a relative azimuth of 180°.

param.	case 1 case 2		$\alpha=4^\circ$ change (%) in:			$\alpha=8^\circ$ change (%) in:			$\alpha=16^\circ$ change (%) in:		
			$I^{\text{rel}}$	$\Delta M$	$N_\alpha^{\text{Tr}}$	$I^{\text{rel}}$	$\Delta M$	$N_\alpha^{\text{Tr}}$	$I^{\text{rel}}$	$\Delta M$	$N_\alpha^{\text{Tr}}$
AOT	0.2	0.4	54	55	0	60	29	0	40	7.4	0
BLH aer.&NO <sub>2</sub> [km]	1.0	1.5	-6.5	6.1	-11	-3.2	4.2	-4.7	-1.1	1.9	-1.6
BLH NO <sub>2</sub> [km]	1.0	1.5	-1.3	23	-24	-0.6	12	-12	-0.1	5.1	-4.7
$N$ [ $10^{15}$ molec/cm <sup>2</sup> ]	20	2	-5.7	0.1	-3.9	-4.3	-2.7	1.8	-2.4	-7.7	7.8
ASY	0.70	0.75	-4.8	-3.7	0.15	-5.0	-3.0	1.6	-4.1	-3.1	2.8
SSA	0.92	0.95	-2.1	-0.2	-1.4	-1.2	0.1	-0.5	-0.4	-0.6	0.6
ALB	0.06	0.03	2.7	-0.5	-3.1	1.5	0.1	-0.7	0.7	0.2	-0.7
POL	on	off	2.4	0.5	1.9	2.7	0.4	0.9	-2.1	-0.4	0.0

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.




**Fig. 1.** Illustration of MAX-DOAS observation. The Mini MAX-DOAS instrument of this study can rotate only in one vertical plane, i.e. it has a fixed azimuth. The viewing direction is referred to as the elevation, which is the angle ( $\alpha$ ) with the horizontal. For both the DOAS method and relative intensity observations, an observation at elevation  $\alpha$  is always combined with a zenith observation ( $\alpha = 90^\circ$ ). Two examples of photon paths are shown in white. MAX-DOAS observations are relatively insensitive to the trace gases in the stratosphere, as long as the zenith and non-zenith observations are taken within a time frame where the solar zenith angle changes only little: the sensitivity of the MAX-DOAS observation to NO<sub>2</sub> in a particular horizontal layer is proportional to the difference in (detected) photon path length through that layer, between the observations pointed at the zenith and at elevation  $\alpha$ .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.




**Fig. 2.** Measurements with the Mini MAX-DOAS instrument of a Mercury line source at 407.8 [nm], at four different temperatures.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.




**Fig. 3.** Set-up of the field-of-view characterization experiment with a light source in the distance. The angle  $\alpha$  denotes the rotation in the vertical plane relative to a horizontal starting position of  $0^\circ$ . At this starting position the light source was at the same height above the table as the lens. The experiment was repeated including a black tube in front of the lens (not drawn), which is used normally to block stray light. Adding the tube did not change the results.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.



**Fig. 4.** Field-of-view of the Mini MAX-DOAS instrument, measured with the set-up of Fig. 3. The black curve represents the measured signal at elevation  $\alpha$ , and the red curve is the normalized surface integral of the black curve. The FWHM is  $0.4^\circ$ . The center of the field-of-view is defined as the elevation where the red curve equals one half. In this case the pointing offset is  $-2.16^\circ$ .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.





Fig. 5. Flow chart of the two-step retrieval algorithm of tropospheric NO<sub>2</sub> columns.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.




**Fig. 6.** Spectrum of the NO<sub>2</sub> absorption cross-section convoluted with the instrumental slit function. The three wavelengths  $\lambda_1$ ,  $\lambda_2$  and  $\lambda_3$  were used in the radiative transfer simulation of the differential AMF and relative intensity. The wavelength range of this figure corresponds to the DOAS spectral fitting window (see Sect. 2.3.2).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.



**Fig. 7.** Radiative transfer simulations with the DAK model of relative intensity (left) and the differential air mass factor of NO<sub>2</sub> (right) at 428.22 nm as a function of the elevation, for different values of the AOT. Note the logarithmic scale on the x-axis. The black line in the right plot gives the differential air mass factor for the geometrical approximation (GA). Solar zenith angle = 50°, relative azimuth angle = 180°.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.



**Fig. 8.** Radiative transfer simulations with the DAK model of height-dependent differential AMF (differential box-AMF) for AOT=0.2 (left) and AOT=0.6 (right), at 428.22 nm. Settings in the simulations: boundary layer height for aerosols (block profile): 1.0 km, solar zenith angle = 60°, relative azimuth angle = 180°,  $\lambda=428$  nm.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.



**Fig. 9.** Difference between the modeled relative intensity for a range of AOT and BLH values, and the reference value of relative intensity for AOT=0.2 and a boundary layer height of 1 km (red dot). Two elevations were used in the calculations: 4° (top) and 30° (bottom). Solar zenith angle = 50°, relative azimuth angle = 180°,  $\lambda=428$  nm.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.




**Fig. 10.** Tropospheric NO<sub>2</sub> column retrieval for three selected days using the two-step algorithm. All plots have the same time scale on the horizontal axis and in each plot the colors correspond to the elevations indicated by the legend on the upper left. The first row shows the measured NO<sub>2</sub> differential slant columns in [ $10^{16}$  molecules/cm<sup>2</sup>]. The second row shows the relative intensities, and the third row the retrieved AOT. The last row shows the retrieved tropospheric column of NO<sub>2</sub> in [ $10^{15}$  molecules/cm<sup>2</sup>].

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.




**Fig. 11.** Retrieved aerosol optical thickness (upper row) and tropospheric NO<sub>2</sub> column (lower row) for four selected days using the two-step algorithm. The red lines represent the average value of the three elevations: 4°, 8°, and 16°. The spread is indicated by the grey lines.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.



**Fig. 12.** Difference between the differential air mass factor simulated with the DAK radiative transfer model, and the differential air mass factor of the geometrical approximation (GA) for 30° elevation, as a function of solar position. AOT = 0.2,  $\lambda = 428.22$  nm. As an example, the red curve represents the path of the sun through the sky above De Bilt on 21 March 2009, relative to the viewing azimuth of the instrument, which was 46° East from North (sunrise is indicated by “R”, sunset by “S”).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns

T. Vlemmix et al.



**Fig. 13.** Retrieved tropospheric NO<sub>2</sub> column in units of  $[10^{15} \text{ molec/cm}^2]$  for 21 March 2009 (left graph). The different colors represent variations of the retrieval method. The green line is based on the geometrical approximation (GA), applied to the differential slant column measurements at 30° elevation. The blue line is derived from the same measurements, but with a differential air mass factor calculated with the DAK model and the settings described in Sect. 3.1.2, where the AOT was derived from the relative intensity measurements at this elevation. The red line is the average tropospheric column as derived by the two-step algorithm, i.e. from the elevations 4°, 8° and 16°. The spread in these values is shown in grey. The noisy character of the retrievals based on the 30° elevation is due to the low sensitivity to NO<sub>2</sub> at 30°, relative to the other elevations. The right graph gives the difference in percent between the blue and the green line. See also the red line in Fig. 12.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.




**Fig. 14.** Comparison between AOT derived from MAX-DOAS relative intensity observations in De Bilt and AERONET AOT (Cabauw). Both sites were 22 km apart. The selection criteria described in Sect. 4.3 were applied to all measurements in the observation period (see Sect. 2.2). The red line represents a linear regression, where the squared orthogonal distance of all points to the 1:1 line was minimized.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**MAX-DOAS aerosol corrected tropospheric NO<sub>2</sub> columns**

T. Vlemmix et al.



**Fig. 15.** Comparison between tropospheric NO<sub>2</sub> columns from MAX-DOAS observations in De Bilt and OMI (DOMINO product). The selection of data-points is described in Table 2 (MAX-DOAS), and Sect. 4.4 (OMI), and was applied to all measurements in the observation period (see Sect. 2.2). The selection described in the first row of Table 2 includes all points in the plot, the selection described in the second row includes all black and red points, and the selection of the last row includes only the red points.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)