

**A comparison of
OEM CO retrievals
from the IASI and
MOPITT instruments**

S. Illingworth et al.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

**S. M. Illingworth¹, J. J. Remedios¹, H. Boesch¹, S.-P. Ho², D. P. Edwards²,
P. I. Palmer³, and S. Gonzi³**

¹Earth Observation Science, Department of Physics and Astronomy, University of Leicester, Leicester, UK

²Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, Colorado, USA

³School of GeoSciences, University of Edinburgh, Edinburgh, UK

Received: 20 September 2010 – Accepted: 22 October 2010 – Published: 10 November 2010

Correspondence to: S. Illingworth (smi3@le.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

Observations of atmospheric CO can only be made on global and regional scales by remote sensing instruments situated in space. One such instrument is the Infrared Atmospheric Sounding Interferometer (IASI), producing spectrally resolved, top-of-atmosphere radiance measurements from which CO vertical layers and total columns can be retrieved. This paper presents the first intercomparison between an IASI CO data set, in this case that produced by the University of Leicester IASI Retrieval Scheme (ULIRS), and the V3 and V4 operationally retrieved CO products from the Measurements Of Pollution In The Troposphere (MOPITT) instrument. The comparison is performed for a localised region of Africa. By comparing both the standard data and a special version of MOPITT data retrieved using the ULIRS a priori for CO, it is shown that standard intercomparisons of CO are strongly affected by the differing a priori data of the retrievals, and by the differing sensitivities of the two instruments. In particular, the differing a priori profiles for MOPITT V3 and V4 data result in systematic retrieved profile changes as expected. Application of averaging kernels is used to derive a difference quantity which is much less affected by smoothing error and hence more sensitive to systematic error. This technique is used to show that the systematic bias between MOPITT V4 and ULIRS IASI data, at MOPITT vertical resolution, is less than 7%. These conclusions are confirmed by simulations with model profiles for the same region.

1 Introduction

Carbon monoxide (CO) in the troposphere acts as a marker, or tracer, of incomplete combustion processes, and through its reactions with the hydroxyl free radical OH, the concentration of CO is related to the oxidising capacity of the troposphere (Thompson, 1992); investigations into perturbations of the sources, sinks and net surface fluxes of CO are therefore of significant importance.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

5 Whilst ground-based and aircraft-mounted instruments are able to make precise measurements of the tropospheric concentrations of CO, they are not able to provide global coverage. Only observations from space allow for fully (in the absence of cloud) global measurements of CO concentrations to be made over a reasonably short time period. Over the past decade the MOPITT (Measurements of Pollution in the Troposphere; Deeter et al., 2003), IMG (Interferometer Monitor for greenhouse Gases) (Kobayashi et al., 1999), AIRS (Atmospheric InfraRed Sounder) (McMillan et al., 2005), and TES (Tropospheric Emission Spectrometer) (Rinsland et al., 2006) instruments have all successfully exploited observations in the 4.7 μm spectral band to increase the vertical information content of profiles and also global coverage. The IASI (Infrared Atmospheric Sounding Interferometer) is the latest instrument in the TIR suite of tropospheric sounders, and the University of Leicester IASI Retrieval Scheme (ULIRS) has been developed to convert IASI Top Of Atmosphere (TOA) radiances into an atmospheric CO product (Illingworth et al., 2010).

15 The objective of this paper is to assess the consistency between the MOPITT and ULIRS retrieved IASI data (hereafter referred to simply as the IASI data or product), and to investigate some of the probable causes of differences that are observed. This work follows on from other studies which have investigated the intercomparison of CO from different satellite products, such as Luo et al. (2007); Ho et al. (2009); Warner et al. (2007); Kopacz et al. (2010), and this paper is organised as follows. In Sect. 2 we summarise the characteristics of the IASI instrument, and discuss the set-up of the ULIRS. Section 3 introduces the MOPITT instrument and presents in detail the retrieval algorithm of the MOPITT CO product, in particular highlighting the similarities and differences between the V3 and V4 operational products. Section 4 presents the results of a standard intercomparison between the IASI and MOPITT instruments and their retrieved CO products, with Sect. 5 performing the analysis with a consistent set of a priori statistics. This section also compares the observed differences between instruments with systematic differences computed using modelled profiles. Finally in Sect. 6 we apply the methodology of Rodgers and Connor (2003) to undertake a comparison



with reduced smoothing error, as demonstrated using synthetic CO fields as for Sect. 5. The conclusions of this work are summarised in Sect. 7.

2 IASI CO retrievals

2.1 IASI

5 The IASI instrument is a high-resolution Michelson interferometer which was launched in 2007, onboard EUMETSAT's (European Organisation for the Exploitation of Meteorological Satellites) MetOp-A satellite, with an equator crossing time of 09:30 LST (local solar time) and an instantaneous Field of View (IFOV) that is approximately 12 km in diameter at nadir; it covers the spectral range between 645 and 2760 cm^{-1} , with a spectral sampling of 0.25 cm^{-1} and a nominal apodised spectral resolution of 0.5 cm^{-1} (Blumstein et al., 2004). A more detailed description of the IASI instrument is given in Clerbaux et al. (2009); here we briefly discuss why the IASI is well suited for providing detailed information about the global distribution of CO, on both short and long term timescales.

15 The IASI instrument's spectral range and low noise in the 4.7 μm region, mean that it is able to observe the CO spectral band centred on 2140 cm^{-1} (4.7 μm); the large swath width (2200 km) enables IASI to achieve a twice daily global coverage (~99%), and as the first of a series of three instruments to be launched every five years, the IASI will allow for the monitoring of long-term climatological trends at a very high temporal resolution.

20 Illingworth et al. (2009) have demonstrated that the likely accuracy in band 1 of the IASI instrument is < 0.1 K at 11 μm . By considering work done by Larar et al. (2010), and also the internal radiometric accuracy of the IASI instrument, as reported by Blumstein et al. (2004), the IASI instrument is likely radiometrically accurate to < 0.3 K in the 25 4.7 μm spectral region.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Retrieval setup

The ULIRS scheme (Illingworth et al., 2010) has been developed to retrieve CO from IASI measured TOA radiances, utilising an Optimal Estimation Method (OEM) (Rodgers, 2000) to constrain the ill-conditioned nature of the retrieval problem. The ULIRS algorithm is based on the retrieval of a VMR state vector, and a CO a priori profile and covariance matrix are constructed using profiles from the Toulouse Off-line Model of Chemistry And Transport (TOMCAT) Chemical Transport Model (CTM) (Chipperfield, 2006), which incorporates a wide variety of CO scenarios. A constant a priori product, i.e. one which is the same for every retrieval, is used to ensure that any spatial or temporal features observed in the retrieved CO product are not symptomatic of features in the a priori. For this particular study a key point is that the a priori product has been constructed using over 8000 TOMCAT profiles from the Southern Africa region (over a grid box bounded longitudinally from -20 to 50° E, and latitudinally from -30 to 30° N) for the year 2004. The ULIRS incorporates the Oxford RFM (Dudhia, 2000) as a forward model, and the retrieval scheme makes use of the spectral interval 2143 to 2181 cm^{-1} , for reasons outlined in Barret et al. (2005). The ULIRS also utilises a spatially precise ($30''$ resolution) surface elevation (USGS, 1998) and emissivity (0.05° resolution) (Seemann et al., 2008), as well as incorporating a quantified back-scattered solar radiation component, which makes use of the Atmospheric Chemistry Experiment (ACE) Fourier Transform Spectrometer (FTS) atlas of the infrared solar spectrum (Hase et al., 2010). The importance of these three parameters, as well as full details of the ULIRS are discussed in Illingworth et al. (2010). A linear error analysis of the ULIRS data suggests that typical random errors over the African region relating to the profiles are found to be $\sim 10\%$ at 5 and 12 km, and on the total columns to be $\sim 12\%$.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 MOPITT CO retrievals

3.1 MOPITT

A summary of the main characteristics of the MOPITT instrument is given below; for a full description of the instrument see e.g. Drummond and Mand (1996). MOPITT is a nadir sounding instrument which measures upwelling infrared radiation in both the 4.7 μm and 2.4 μm spectral bands; it uses gas correlation spectroscopy using Pressure Modulation Cells (PMC) and Length Modulation Cells (LMCs) to calculate total column amounts and profiles of CO in the lower atmosphere. MOPITT was launched on board the Terra satellite in 1999, with an equator crossing time of 10:30 LST \pm 26 min, and a total scanning angle of $\pm 26^\circ$ in each swath, combined with a 22 by 22 km horizontal resolution allows MOPITT to generate a global map of CO once every three days. For the purposes of this work we are interested in the measurements made in the 4.7 μm region, which provide the most information in regards to a retrieved profile of CO. Whilst MOPITT utilises a slightly different technique to IASI in its measurements of TOA radiances, the sensitivity of the two instruments are determined by the same factors.

3.2 Retrieval setup

The MOPITT “version 3” (V3) product first became available in 2002, and was the first data set to offer truly long-term and global coverage about the distribution of tropospheric CO. The characteristics of this product are given in detail by Deeter et al. (2003), but to summarise they include: (1) an OEM retrieval algorithm, which utilises an operational radiative transfer model (MOPFAS) (Edwards et al., 1999) based on prelaunch laboratory measurements of instrument parameters as its forward model; (2) a fixed 7-level pressure grid (floating surface level, 850 hPa, 700 hPa, 500 hPa, 350 hPa, 250 hPa, 150 hPa); (3) a global a priori profile and covariance matrix for all

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



retrievals; and (4) a state vector which consists of a CO profile, given in terms of a Volume Mixing Ratio (VMR), a surface temperature and a surface emissivity.

A full description of the MOPITT “version 4” (V4) product is given by Deeter et al. (2010). As with the V3 product, the V4 algorithm is based on an OEM retrieval technique, using MOPFAS as the forward model. There are however significant differences between the V3 and V4 algorithms, and some of these are now discussed. Whereas the V3 state vector represented the CO vertical profile as a set of VMR values, the V4 state vector represents the CO profile as a set of log(VMR) values; it was found that the use of a VMR probability distribution function in V3 occasionally resulted in unrealistic negative VMR values, and so by representing the CO profile in log(VMR) space in the V4 product, these negative values were eliminated. The motivation for using the log(VMR) state vector for V4 was thus to avoid these negative values, and to provide a greater consistency with observations. In addition to this, the V4 state vector expresses the CO profile on a 10-level pressure grid (floating surface level followed by nine uniformly spaced levels from 900 to 100 hPa) instead of the seven-level grid used for V3; this change to an equidistant pressure grid was implemented because a retrieval grid with uniform grid spacing simplifies the physical interpretation of the retrieval.

For the V4 product some changes to the MOPFAS radiative transfer model were also made in comparison to that used in the V3 algorithm. In extremely polluted conditions, V3 retrievals sometimes failed because they exceeded the upper CO concentration limit of MOPFAS. For the V4 product the MOPFAS forward model was therefore modified to allow for retrievals with significantly higher values, with the number of training profiles expanded from 58 (Edwards et al., 1999) to 116. As part of its processing, MOPFAS also incorporates models of the physical states of the MOPITT LMCs and PMCs, and for the V4 algorithm, both the PMC and LMC models have been updated for consistency with the actual on-orbit cell pressure and temperature values observed during the mission. Specifically, the pressure and temperature values that are used to model the LMCs in MOPFAS are now time-mean values, whilst the shapes and relative phases of the PMC pressure and temperature cycles remain unchanged. These

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



updated models also reflect instrumental modifications performed after the failure of one of MOPITT's two coolers which occurred on 7 May 2001.

One of the most significant differences between the MOPITT V3 and V4 retrieval algorithms is the choice of the a priori profiles and covariance matrices. In the MOPITT V3 product, a global a priori profile was employed for all retrievals; as discussed in Sect. 2.2, this ensures that any features in the retrieved CO actually correspond to information in the measurements. However, it is also true that using a global a priori can sometimes yield large systematic differences between the “true” CO concentration and the retrieved value, at levels where the weighting functions exhibit low sensitivity; this is especially true for the large smoothing length (500 hPA, calculated by computing the delta-pressure for which the off-diagonal element of the covariance matrix was found to be $1/e^2$ times the corresponding diagonal element) of the MOPITT V3 a priori covariance matrix. In an attempt to reduce these a priori-related errors, V4 a priori profiles are based on a monthly climatology from the global CTM MOZART-4 (Model for OZone And Related chemical Tracers, version 4) (Emmons et al., 2010), where for each retrieval, the climatology is spatially and temporally interpolated to the time and location of the observation. As in V3, the V4 algorithm uses a global a priori covariance matrix, but sets the diagonal elements to a fractional VMR variability of 30%, and assumes a smoothing length of 100 hPA (Deeter et al., 2010); these values have been chosen based on analyses of aircraft in situ data sets at individual MOPITT validation sites. This relatively small value for the smoothing length acts to reduce the projection of information from levels where the MOPITT weighting functions are relatively strong (e.g., the mid-lower troposphere) to levels where the weighting functions are relatively weak (e.g., the surface). An example of the differences between the V3 and V4 a priori profiles and covariances matrices is plotted in Fig. 1.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Comparison of MOPITT V3 and V4 averaging kernels

The averaging kernel matrix \mathbf{A} is a representative of the sensitivity of the retrieved state to the true state:

$$\mathbf{A} = \frac{\partial \hat{\mathbf{x}}}{\partial \mathbf{x}} \quad (1)$$

where \mathbf{x} is the “true” state vector, and $\hat{\mathbf{x}}$ is the retrieved state vector. The rows of \mathbf{A} are generally peaked functions, which have a half-width that is representative of the spatial resolution of the observing system. An ideal observing system would have δ -function averaging kernels, peaking at the various levels over which the retrieval was performed, and no noise. The Degrees of Freedom for Signal (DFS) are a measurement of the information content of the retrieval, and are defined as the trace of \mathbf{A} (Rodgers, 2000).

Figure 2 shows a plot of the averaged daytime and oceanic MOPITT V3 and V4 averaging kernels (\mathbf{A}^{MOP3} and \mathbf{A}^{MOP4}) for 1 September 2007 over the region shown in Fig. 4 (hereafter referred to as the Southern Africa region), at each of the pressure levels for the relevant retrieval grid, and illustrates how the MOPITT V3 and V4 measurements contribute to the retrieved CO profiles. It is important to note that whilst Fig. 2 gives a good indication as to the sensitivities of the MOPITT V3 and V4 averaging kernels, \mathbf{A}^{MOP3} and \mathbf{A}^{MOP4} do not represent the same quantity: \mathbf{A}^{MOP3} is $\partial \hat{\mathbf{x}} / \partial \mathbf{x}$, whereas \mathbf{A}^{MOP4} is $\partial \log(\hat{\mathbf{x}}) / \partial \log(\mathbf{x})$. To the first order the V3 and V4 averaging kernels can be converted using the following relationship:

$$\frac{\partial \log(\hat{\mathbf{x}})}{\partial \log(\mathbf{x})} = \left(\frac{\mathbf{x}}{\hat{\mathbf{x}}} \right) \frac{\partial \hat{\mathbf{x}}}{\partial \mathbf{x}} \quad (2)$$

Because of the large variations in the averaging kernel matrices between day and night, and over land and ocean, it is important to consider each of these cases separately. If these different scenarios are not analysed individually then there are too many factors that must be accounted for, including but not limited to: varying thermal contrast; varying CO distributions; varying a priori; and varying surface pressures, to

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



permit a meaningful analysis. Similarly, significant latitudinal effects which make tropical retrievals quite different than polar retrievals, mean that any conclusions will be more justifiable if a specific latitudinal zone or region, rather than a global data set, is analysed.

The main reason for performing this analysis over Southern Africa was because this is the region over which the ULIRS was characterised (Illingworth et al., 2010). Over Southern Africa, especially during the fire season (which typically lasts from late July to early November; Giglio et al., 2006), there is also a large variety in the different CO atmospheric scenarios; Southern Africa thus represents a region over which a wide variety of CO profiles can be observed, but which should not be adversely affected by latitudinal effects. It has been shown previously (see e.g. Deeter et al., 2007b) that the thermal contrast has a large effect on the MOPITT averaging kernels, therefore the main comparison has been carried out during daytime over the ocean, where the effects of the thermal contrast are minimised, and where surface emissivity variations can be neglected. From a technique point of view, a consistent use of all the data sets is important.

As the averaging kernels are strongly dependent upon the different pressure grids that are used in each of the retrieval schemes, to make a consistent comparison, the pressure-layer-normalised averaging kernels are generated using the following equation, taken from Ho et al. (2009):

$$\mathbf{A}_N^{i,j} = \frac{\mathbf{A}^{i,j}}{\Delta P^i}, \quad (3)$$

where i and j are indexes of column and row elements of \mathbf{A} and \mathbf{A}_N , and ΔP^i is the pressure thickness of the layer corresponding to column index i . It is also important to note that \mathbf{A}^{MOP3} and \mathbf{A}^{MOP4} do not represent the same quantity: \mathbf{A}^{MOP3} is $\partial \hat{x} / \partial x$, whereas \mathbf{A}^{MOP4} is $\partial \log(\hat{x}) / \partial \log(x)$. To the first order the V3 and V4 averaging kernels can be converted using the following relationship:

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

$$\frac{\partial \log(\hat{x})}{\partial \log(x)} = \left(\frac{x}{\hat{x}}\right) \frac{\partial \hat{x}}{\partial x}. \quad (4)$$

3.4 Comparison of MOPITT V3 and V4 retrievals

The mean CO profiles of the MOPITT V3 and MOPITT V4 retrieval algorithms, over the ocean and for the daytime of 1 September 2007 over the Southern Africa region (see Fig. 4) are shown in Fig. 3. It can be seen that the largest differences between the MOPITT V3 and V4 CO concentrations occur at the surface (approximately 70 ppbv), whilst the smallest differences occur between 400 and 600 hPa.

The differences between the MOPITT V3 and MOPITT V4 retrieved profiles shown in Fig. 3 are consistent with the results of Deeter et al. (2010), with the mean V3 and V4 retrieved profiles similar in the upper troposphere, and differing greatly in the lower troposphere. One of the major reasons for the observed differences between the MOPITT V3 and V4 profiles is the use of different a priori (profiles and covariance matrices), and the effect of the smoothing length on the retrievals. In particular, for large correlation lengths, retrievals at pressure levels that are insensitive to CO can be strongly influenced by more sensitive levels. One example of where this “false influence” can occur is for scenes with a low-thermal contrast, and hence a lack of sensitivity to the surface; in such scenes a large correlation length can result in the projection of CO features from the mid-troposphere, where there is an increased sensitivity, to the surface. Such a phenomena could well be responsible for the results seen in Fig. 3, with the large differences at the surface being a direct result of the MOPITT V3 being falsely influenced by CO features above 850 hPa.

Another possible cause of the differences between the V3/V4 retrieved profiles is the use of lognormal statistics in the MOPITT V4 retrieval algorithm. Deeter et al. (2007a) showed that the assumption of Gaussian VMR variability in the MOPITT V3 retrieval algorithm is inconsistent with in-situ data sets, and leads to positive retrieval bias in especially clean conditions (e.g., VMR values less than 60 ppbv). As V4 retrievals use

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a state vector based on log(VMR) they are therefore not subject to this effect. A further source of the difference between the MOPITT V3 and V4 data sets is the change in retrieval bias due to drifts in the MOPITT LMCs and PMCs (see Sect. 3.2), which are corrected for in the V4 retrieval algorithm, but not in the V3.

4 Comparison of IASI and MOPITT CO

MOPITT is on board the Terra satellite, which is on a different orbit to MetOp, it is therefore not possible to find exact coincidences between IASI and MOPITT measurements. Thus, to ensure a reasonable coincidence criteria and sample size, only IASI and MOPITT retrievals that correspond spatially to within 50 km, and which differ on a temporal timescale by at most 90 min are considered for this intercomparison; 90 min was chosen, because it allowed for the maximum amount of data to be collated whilst minimising the effects of transportation. The IASI retrievals were further filtered by using the Root Mean Square (RMS) difference between simulated and observed spectra, as well as the χ^2 value of the retrieval. Only retrievals which had an RMS difference of less than $3.5 \text{ nW/cm}^2/\text{cm}^{-1}/\text{sr}$ and a χ^2 of less than 3.5 were considered for the comparison. Any IASI and MOPITT matches where the surface pressures used in the retrieval process differed by more than 20 hPa were also neglected.

The region considered for comparison, along with the retrieved CO total column densities for each of the algorithms is shown in Fig. 4. As was discussed in Sect. 3.3 it is important to consider the day/night and land/ocean retrievals separately, and the remainder of the figures in this study are for the comparison between daytime retrievals over the ocean.

The IASI and MOPITT retrieval algorithms are set up on pressure grids which differently sample the troposphere, thus the IASI retrievals which were on the pressure levels closest to those of the corresponding MOPITT pressure levels were selected for comparison. The MOPITT V3 (\mathbf{A}^{MOP3}) and IASI averaging kernels (\mathbf{A}^{IASI}) at retrieval pressure levels close to those of MOPITT V3 are shown in Fig. 5, whilst the

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOPITT V4 (\mathbf{A}^{MOP4}) and IASI averaging kernels (\mathbf{A}^{IASI}) at retrieval pressure levels closest to those of MOPITT V4 are shown in Fig. 6. These mean averaging kernels (\mathbf{A}^{MOP3} , \mathbf{A}^{MOP4} and \mathbf{A}^{IASI}) correspond to daytime and ocean averages from MOPITT and IASI for 1 September 2007 over the Southern Africa region shown in Fig. 4. As with the differences between \mathbf{A}^{MOP3} and \mathbf{A}^{MOP4} , the different magnitudes between \mathbf{A}^{MOP} and \mathbf{A}^{IASI} are mainly due to the different pressure layer thicknesses of the retrieval grids used by MOPITT V3/V4 (7/10 layers) and IASI (30 layers). Recall that, whilst the IASI and MOPITT V3 averaging kernels both represent $\partial\hat{x}/\partial x$, the MOPITT V4 averaging kernels correspond to $\partial\log(\hat{x})/\partial\log(x)$.

The MOPITT V3 and IASI mean CO profiles and standard deviation relative to their means are shown in Fig. 7a, at MOPITT V3 pressure levels for the 1 September 2007, over the Southern Africa region for the daytime over the ocean. As can be seen from Fig. 7c, x^{IASI} overestimates x^{MOP3} at pressure levels greater than 500 hPa, and is an underestimate at lower pressures. The results from the intercomparison of IASI (x^{IASI}) and MOPITT V3 (x^{MOP3}) CO are summarised in Table 1.

Figure 7b shows the MOPITT V4 and IASI mean CO profiles and standard deviation relative to their means, at MOPITT V4 pressure levels for the 1 September 2007 over the Southern Africa region. The differences between the two products are depicted in Fig. 7d, with large apparent discrepancies between the mean IASI and MOPITT V4 CO profiles below 800 hPa symptomatic of the difference in the relative correlation lengths of the two different retrieval algorithms. As was discussed in Sect. 3.4 the correlation length used in the MOPITT V4 algorithm is 100 hPa and the a priori profile is based on modelled data that is spatially and temporally interpolated to the retrieval scene. As the IASI retrieval scheme uses a constant a priori profile, and has a smoothing length of approximately 400 hPa it has a greater sensitivity to mid-tropospheric events than it does to the surface (as can be seen from Figs. 5b and 6b). The results from the intercomparison of IASI (x^{IASI}) and MOPITT V4 (x^{MOP4}) CO are summarised in Table 2.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[⏴](#)[⏵](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The results presented in Fig. 7 indicate that with respect to the effect of projection of information in the mid-troposphere to the surface, the IASI retrievals more closely resemble the MOPITT V3 than the V4 retrievals. This can in part be explained by the similarity in the smoothing lengths of the IASI and MOPITT V3 a priori covariance matrices (400 and 500 hPa, respectively, in comparison to 100 hPa for the MOPITT V4 algorithm), as well as the fact that the diagonal elements of the a priori covariance matrix are similar for IASI and MOPITT V3, in comparison to those for MOPITT V4 (see horizontal error bars in Fig. 1). It should also be noted that the differences between the MOPITT V3 and V4 retrieved profiles that can be inferred from Fig. 7 are the same as those shown in Fig. 3.

5 Comparison of IASI and MOPITT CO using the same a priori

It is instructive to investigate whether the differences observed between x^{IASI} and x^{MOP} are due to the choice of a priori used by each of the retrieval schemes. A useful way to examine this problem, is to use the same a priori profile and covariance for both the MOPITT and IASI retrievals. In order to do this the IASI a priori profiles and covariance matrices were integrated into the MOPITT V3 and V4 retrieval algorithms.

The differences between the IASI (x^{IASI}) and “adjusted” MOPITT ($x^{\text{MOP3'}}$) profiles shown in Fig. 8c are similar to the differences between the IASI and MOPITT V3 profiles (Fig. 7c), which is to be expected given the similarities in the smoothing length and a priori covariance matrices of the IASI and MOPITT V3 retrieval algorithms. Figure 8d also demonstrates that x^{IASI} and $x^{\text{MOP4'}}$ are in better agreement, particularly in the lower troposphere, compared to the differences between x^{IASI} and x^{MOP4} (Fig. 7d). This result is mainly explained by the fact that both retrievals now use the same a priori statistics, in particular utilising an identical correlation length, meaning that the surface retrievals are equally affected by the influence of mid-tropospheric events. The results from the intercomparison between the IASI and adjusted MOPITT V3 and V4 CO

products are summarised in Tables 1 and 2 respectively. This better agreement corresponds well with other studies (e.g. Luo et al., 2007; Ho et al., 2009; Warner et al., 2007), which also investigated the effect of minimising the differences in the a priori statistics, in the intercomparison of different retrieval products.

5.1 Smoothing bias

Whilst the IASI and MOPITT products have now been retrieved using the same a priori information, there still exist differences in the measurement sensitivity, i.e. weighting functions and noise, of the retrievals. The retrieved state vector can be written as a weighted mean of the true profile (\mathbf{x}_{true}) and the a priori profile (\mathbf{x}_a) (Rodgers and Connor, 2003). In the case of IASI retrievals this can be written as:

$$\mathbf{x}^{\text{IASI}} = \mathbf{A}^{\text{IASI}} \left(\mathbf{x}_{\text{true}} - \mathbf{x}_a^{\text{IASI}} \right) + \mathbf{x}_a^{\text{IASI}} + \epsilon^{\text{IASI}}, \quad (5)$$

whilst for the MOPITT retrievals (V3 and V4):

$$\mathbf{x}^{\text{MOP}} = \mathbf{A}^{\text{MOP}} \left(\mathbf{x}_{\text{true}} - \mathbf{x}_a^{\text{MOP}} \right) + \mathbf{x}_a^{\text{MOP}} + \epsilon^{\text{MOP}}, \quad (6)$$

where ϵ represents the error in the retrieved profiles due to both random and systematic errors in the measured signal and in the algorithm's forward model (Rodgers and Connor, 2003). By processing the MOPITT retrieval algorithms using the IASI a priori, the modified MOPITT retrieved CO profile, $\mathbf{x}^{\text{MOP}'}$, can be written as:

$$\mathbf{x}^{\text{MOP}'} = \mathbf{A}^{\text{MOP}'} \left(\mathbf{x}_{\text{true}} - \mathbf{x}_a^{\text{IASI}} \right) + \mathbf{x}_a^{\text{IASI}} + \epsilon^{\text{MOP}'}, \quad (7)$$

where $\mathbf{A}^{\text{MOP}'}$ represents the adjusted MOPITT (V3 and V4) averaging kernel matrices.

Differences between \mathbf{x}^{IASI} and $\mathbf{x}^{\text{MOP}'}$ can be characterised from Eqs. (5) and (7). Further insight into differences in the smoothing effect of the two retrievals (the smoothing

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bias) can be obtained by examining what happens when the averaging kernels for each instrument are applied to a true profile (thus simulating the effect of retrieval characteristics on the retrieved profiles).

For this study the truth is represented by gridded output from the GEOS-Chem (v7.04.10) CTM (Bey et al., 2001 and <http://www.GEOS-Chem.org/>), driven by meteorological observations from the Goddard Earth Observing System v5 (GEOS-5) from the Global Modeling and Assimilation Office Global Circulation model based at NASA Goddard. The GEOS-Chem global 3-D CTM relates surface emissions of CO to global 3-D distributions of CO concentration. Primary sources of CO include biomass burning, fossil and biofuel combustion. Secondary sources are from the oxidation of co-emitted Volatile Organic Compounds (VOCs). We use surface biomass burning emission estimates from the Global Fire Emission Database (GFED version 2) (Van Der Werf et al., 2006), and the model is used in tagged CO tracer mode at a horizontal resolution of $2 \times 2.5^\circ$, with 47 sigma levels that span the surface to 0.01 hPa. The 3-D meteorological data is updated every six hours, and boundary layer and tropopause heights are updated every three hours. We use GEOS-Chem CO profiles, x^{GEOS} , with the closest x^{GEOS} to the collocated IASI and adjusted MOPITT CO profiles being used for this comparison; the GEOS-Chem total column densities for the studied region are shown in Fig. 4d.

$x_{\text{MODEL}}^{\text{IASI}}$ and $x_{\text{MODEL}}^{\text{MOP}'}$ have been calculated from Eqs. (5) and (7), respectively, with x_{true} given by x^{GEOS} . In the absence of any other biases, $x_{\text{MODEL}}^{\text{MOP}'}$ and $x_{\text{MODEL}}^{\text{IASI}}$ represent the “best possible” retrieval of the GEOS-Chem profiles by the adjusted MOPITT and IASI algorithms, respectively. The results from the intercomparison between the expected smoothing biases δ_{MODEL} MOPITT V3 and V4 CO products are summarised in Tables 1 and 2 respectively.

As can be seen from Figs. 8 and 9, and also from Tables 1 and 2, the differences between $x_{\text{MODEL}}^{\text{MOP}'}$ and $x_{\text{MODEL}}^{\text{IASI}}$ are of a very similar magnitude to those that are observed between x^{IASI} and $x^{\text{MOP}'}$. Retrieving GEOS-Chem modelled profiles in this manner has

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



therefore shown that many of the observed differences between the IASI and adjusted MOPITT retrievals can be explained by the smoothing bias.

6 Comparison with reduced smoothing error

The results of Sect. 5.1 indicate that the differences between $x^{\text{MOP}'}$ and x^{IASI} are a result of the smoothing bias. The direct comparison between the products includes a contribution from smoothing error, even when the same a priori is used for the retrievals of both instruments to be compared. This arises from the non-identical weighting functions and error covariances of the two instruments. Rodgers and Connor (2003) proposed a methodology, also employed by Luo et al. (2007); Warner et al. (2007); Ho et al. (2009), in which the averaging kernels of one retrieval scheme are used to smooth the retrieved data set from the other set of measurements, providing that both retrievals utilise the same set of a priori statistics. With the caveat that this “double smoothing” of the truth will serve to smooth away some of the intrinsic information contained within the retrieved data set, this section discusses such a method, the justification for its use, and the results that are obtained. In particular we note that the real importance here is to show that quantities can be calculated which minimise smoothing contributions.

6.1 Methodology

Rodgers and Connor (2003) showed that the effect of smoothing error in a comparison can be reduced if the retrieval of one instrument is simulated using the retrieval of another. For profiles used in this study, the number of DFS for the MOPITT retrievals were found to be comparable to, yet usually smaller than, that of IASI. The DFS for IASI over the oceanic Southern Africa region during the daytime were found to range from 1.34 to 2.53, whereas those corresponding to $\mathbf{A}^{\text{MOP3}'}$ and $\mathbf{A}^{\text{MOP4}'}$ were found to range from 0.99 to 2.29 and from 0.95 to 2.00, respectively.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Providing that common a priori statistics are used in the retrievals, Rodgers and Connor (2003) states that:

$$\mathbf{x}^{\text{IASI}'} = \mathbf{A}^{\text{MOP}''} \left(\mathbf{x}_{\text{IASI}} - \mathbf{x}_a^{\text{IASI}} \right) + \mathbf{x}_a^{\text{IASI}}, \quad (8)$$

where $\mathbf{A}^{\text{MOP}''}$ is $\mathbf{A}^{\text{MOP}'}$, but converted onto the same pressure grid as that used by IASI, and $\mathbf{x}^{\text{IASI}'}$ now represents the retrieved IASI profile, smoothed using $\mathbf{A}^{\text{MOP}''}$. The IASI retrieval should be optimal with respect to the comparison ensemble, which it is expected to be because the IASI a priori were specifically constructed for the comparison ensemble.

From Eqs. (5), (7), and (8), the difference between $\mathbf{x}^{\text{IASI}'}$ and $\mathbf{x}^{\text{MOP}'}$ can be written as:

$$\mathbf{x}^{\text{MOP}'} - \mathbf{x}^{\text{IASI}'} = \left(\mathbf{A}^{\text{MOP}'} - \mathbf{A}^{\text{MOP}'} \mathbf{A}^{\text{IASI}} \right) \left(\mathbf{x}_{\text{true}} - \mathbf{x}_a^{\text{IASI}} \right) + \mathbf{e}^{\text{MOP}'} - \mathbf{A}^{\text{MOP}'} \mathbf{e}^{\text{IASI}}. \quad (9)$$

Providing that the $\mathbf{A}^{\text{MOP}'} - \mathbf{A}^{\text{MOP}'} \mathbf{A}^{\text{IASI}}$ term is small, Eq. (9) should represent the difference between the MOPITT systematic error and the vertical IASI smoothed systematic error. As can be seen from Figs. 10 and 11, the DFS for $\mathbf{A}^{\text{MOP}'} - \mathbf{A}^{\text{MOP}'} \mathbf{A}^{\text{IASI}}$ are on average 0.07 and 0.03 for the MOPITT V3 and V4 retrievals, respectively. As this difference quantity is larger for the MOPITT V3 comparisons, it makes the results of $\mathbf{x}^{\text{MOP}'} - \mathbf{x}^{\text{IASI}'}$ more difficult to interpret than for the corresponding IASI and V4 analysis.

6.2 Results

Application of the above process to a representative true profile is important in order to demonstrate that the smoothing error terms have been very much reduced. Again, GEOS-Chem CO profiles, were used for this comparison, with the smoothing bias now represented by $\delta_{\text{MODEL}'}$. It should be noted that $\delta_{\text{MODEL}'}$ represents the expected smoothing bias for $\mathbf{x}^{\text{IASI}'}$ and $\mathbf{x}^{\text{MOP}'}$, whilst δ_{MODEL} represents the expected smoothing bias for \mathbf{x}^{IASI} and \mathbf{x}^{MOP} . As can be seen from Fig. 13, the smoothing error across

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the profiles, in the case of both the MOPITT V3 and V4 comparisons is small. By comparing Figs. 13 and 9 it can be seen that the application of Eq. (8) has significantly reduced the expected smoothing bias.

Figure 12 compares the mean values for $x^{\text{IASI}'}$, $x^{\text{MOP3}'}$, and $x^{\text{MOP3}' - x^{\text{IASI}'}$, for each MOPITT V3 pressure level for 1 September 2007 for the daytime over the oceanic Southern Africa region. An offset between the total column densities of 6.32% was also computed, and a correlation coefficient between the IASI and MOPITT V3 total columns of 0.86 is comparable to the correlation coefficient between the IASI operational and MOPITT V3 column amounts of 0.87 that was observed by George et al. (2009). Over ocean and during the daytime, the MOPITT V3 CO profile appears to be an overestimate of the IASI retrieved profile in the mid-lower troposphere.

Figure 12 also compares the mean values for $x^{\text{IASI}'}$, $x^{\text{MOP4}'}$, and $x^{\text{MOP4}' - x^{\text{IASI}'}$, for each MOPITT V4 pressure level. As can be seen from this figure, there is excellent agreement between $x^{\text{IASI}'}$ and $x^{\text{MOP4}'}$ across the profile. The results of the comparison between the adjusted MOPITT V4 and the smoothed IASI products are summarised in Table 2. The agreement between the smoothed IASI and adjusted MOPITT V4 total column densities is also high (with a mean absolute difference of 4.21%), and there is a correlation coefficient between the column amounts for $x^{\text{IASI}'}$ and $x^{\text{MOP4}'}$ of 0.86. The results of this comparison indicate that for the data studied, over ocean and during the daytime, the IASI and MOPITT V4 data sets are in very good agreement.

Tables 3 and 4 summarise the values for $x^{\text{MOP}' - x^{\text{IASI}'}$, along with the expected smoothing bias ($\delta_{\text{MODEL}'}$) for the 1 September 2007 over the region shown in Fig. 4 for the daytime/ocean, nighttime/ocean, and nighttime/land scenarios. The results for the daytime/land scenario are not shown as there were insufficient coincident data points from which to draw conclusions from. It should also be noted that only profiles for which there were retrieved values at each of the fixed MOPITT pressure levels were chosen, and as much of the Southern Africa land region has a surface pressure of less than 900 hPa (but greater than 850 hPa) this resulted in a large discrepancy in

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

the number of MOPITT V3 and V4 comparisons over land during the nighttime. The comparisons show that for V4 data, the anticipated differences due to smoothing error are very small. Therefore, any biases observed are likely to be intrinsic, non-retrieval, systematic biases between MOPITT and IASI, which on average across the profile appear to be less than 4%. The results for V3 are less easy to interpret but would be consistent with systematic biases of up to 10%. As the MOPITT V4 product has been shown to be more reliable than the MOPITT V3 product (see e.g. Deeter et al., 2010), it is encouraging that ULIRS agrees to such a large extent with the MOPITT V4 data set.

7 Conclusions

In this paper, possible causes of the differences between the retrievals of the IASI and MOPITT V3/V4 CO products have been investigated. It was observed that the differences between the standard products were significantly affected by the differing a priori data (both mean profiles, covariance errors, and error correlation lengths) used in the retrievals. To account for this different a priori information in IASI and MOPITT retrievals, IASI a priori profiles and covariance matrices were applied to a modified operational MOPITT retrieval algorithm. The residual differences between the two products were believed to arise from differences in the measurement sensitivity, and this was demonstrated by calculating a smoothing bias, using a set of GEOS-Chem modelled data to represent the truth.

A further development to account for the remaining differences in the retrieved CO products was also presented, in which a difference quantity is calculated which is much more sensitive to systematic error than smoothing error. In effect, MOPITT averaging kernels (from the adjusted MOPITT retrieval scheme) were used to smooth IASI CO retrievals. This methodology, as first proposed by Rodgers and Connor (2003) and applied by Ho et al. (2009), was justified by examining the theory for optimal retrievals and by applying the process also to the model data; analysis of data with an a priori



A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

developed specifically for the region exploited in the comparison is particularly suitable for this purpose. Differences between the IASI data and the MOPITT V4 data, were shown to be less than 7% at the MOPITT V4 vertical resolution. For the comparison to MOPITT V3 data, differences were shown to be of the order of 10% although this result must be treated with caution because of residual smoothing error contributions to the selected difference quantity. This offset is comparable with the results of Deeter et al. (2010).

As yet no other work has been published regarding the intercomparison of IASI and MOPITT V4 data products. As well as demonstrating that the ULIRS CO product is in good agreement with an independent satellite data set, the work presented in this paper can also be seen as an initial independent verification of the offset between the MOPITT V3 and V4 products.

The next step in the continuation of this work is to extend the intercomparison over a larger temporal range, and to different regions. Initially the intercomparison should be widened to incorporate dates from the different seasons, as it would be interesting to see how the IASI and MOPITT CO products compared during a period that was not affected by biomass burning. Such a study would also enable the identification of any inter-annual variability between the two data sets. The intercomparison could then be extended in the spatial domain to include regions other than Africa, especially over land masses which are relatively flat. However, care must be taken when the analysis is performed over larger regions because of potential variations in the a priori specifications and larger variations in smoothing error contributions. Any such analysis should be performed separately for day and night data, and over the land and ocean.

Whilst one day (1 September 2007) has been selected for this study, over the Tropical African region, the results and their interpretations have shown the importance in understanding the constituent parts of any retrieval scheme, and the effects that any of these underlying inputs may have on the retrieved product. In particular any set of profiles and total column amounts should always be provided with a set of matching a priori and averaging kernels, so that the data can be correctly interpreted, and if



needs be compared with another product. The purpose of this paper was to assess the consistency between the MOPITT and IASI data, and to investigate the probable causes of differences between these two different instruments. In doing so a greater understanding as to the correct interpretation of both the MOPITT and the ULIRS CO retrieved products has been achieved.

Acknowledgements. IASI has been developed and built under the responsibility of the Centre National d'Etudes Spatiales (CNES, France). It is flown onboard the Metop satellites as part of the EUMETSAT Polar System. The IASI L1 data are received through the EUMETCast near real time data distribution service.

The authors would like to thank the NEODC for access to ECMWF data, the NERC for funding S. M. Illingworth, and EUMETSAT for access to the IASI data. SATSCAN-IR is a project selected by EUMETSAT/ESA under the first EPS/Metop RAO.

References

- Barret, B., Turquety, S., Hurtmans, D., Clerbaux, C., Hadji-Lazaro, J., Bey, I., Auvray, M., and Coheur, P.-F.: Global carbon monoxide vertical distributions from spaceborne high-resolution FTIR nadir measurements, *Atmos. Chem. Phys.*, 5, 2901–2914, doi:10.5194/acp-5-2901-2005, 2005. 4893
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.-Atmos.*, 106, 23073–23095, 2001. 4904
- Blumstein, D., Chalon, G., Carlier, T., Buil, C., Hébert, P., Maciaszek, T., Ponce, G., Phulpin, T., Tournier, B., Siméoni, D., Astruc, P., Clauss, A., Kayal, G., and Jegou, R.: IASI instrument: Technical overview and measured performances, in: *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, edited by: Strojnik, M., vol. 5543, 2004. 4892
- Chipperfield, M. P.: New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer experiments, *Q. J. Roy. Meteor. Soc.*, 132(617), 1179–1203, 2006. 4893
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, doi:10.5194/acp-9-6041-2009, 2009. 4892

Deeter, M., Edwards, D., Gille, J., Emmons, L., Francis, G., Ho, S.-P., Mao, D., Masters, D., Worden, H., and Drummond, J.: The MOPITT Version 4 CO Product: Algorithm Enhancements, Validation, and Long-Term Stability, *J. Geophys. Res.-Atmos.*, 115, D07306, doi:10.1029/2009JD013005, 2010. 4895, 4896, 4899, 4908, 4909

Deeter, M. N., Emmons, L. K., Francis, G. L., Edwards, D. P., Gille, J. C., Warner, J. X., Khatatov, B., Ziskin, D., Lamarque, J.-F., Ho, S.-P., Yudin, V., Attié, J.-L., Packman, D., Chen, J., Mao, D., and Drummond, J. R.: Operational carbon monoxide retrieval algorithm and selected results for the MOPITT instrument, *J. Geophys. Res.-Atmos.*, 108, 1–11, 2003. 4891, 4894

Deeter, M. N., Edwards, D. P., and Gille, J. C.: Retrievals of carbon monoxide profiles from MOPITT observations using lognormal a priori statistics, *J. Geophys. Res.-Atmos.*, 112, D11311, doi:10.1029/2006JD007999, 2007a. 4899

Deeter, M. N., Edwards, D. P., Gille, J. C., and Drummond, J. R.: Sensitivity of MOPITT observations to carbon monoxide in the lower troposphere, *J. Geophys. Res.-Atmos.*, 112, D24306, doi:10.1029/2007JD008929, 2007b. 4898

Drummond, J. R. and Mand, G. S.: The measurements of pollution in the troposphere (MOPITT) instrument: Overall performance and calibration requirements, *J. Atmos. Ocean. Tech.*, 13, 314–320, 1996. 4894

Dudhia, A.: Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) Reference Forward Model (RFM), Software User's Manual, 2000. 4893

Edwards, D. P., Halvorson, C. M., and Gille, J. C.: Radiative transfer modeling for the EOS Terra satellite Measurement of Pollution in the Troposphere (MOPITT) instrument, *J. Geophys. Res.-Atmos.*, 104, 16755–16775, 1999. 4894, 4895

Emmons, L. K., Walters, S., Hess, P., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 43–67, 2010. 4896

George, M., Clerbaux, C., Hurtmans, D., Turquety, S., Coheur, P.-F., Pommier, M., Hadji-Lazaro, J., Edwards, D. P., Worden, H., Luo, M., Rinsland, C., and McMillan, W.: Carbon monoxide distributions from the IASI/METOP mission: evaluation with other space-borne remote

**A comparison of
OEM CO retrievals
from the IASI and
MOPITT instruments**S. Illingworth et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- sensors, *Atmos. Chem. Phys.*, 9, 8317–8330, doi:10.5194/acp-9-8317-2009, 2009. 4907
- Giglio, L., Csiszar, I., and Justice, C. O.: Global distribution and seasonality of active fires as observed with the Terra and Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) sensors, *J. Geophys. Res.-Biogeo.*, 111, G02016, doi:10.1029/2005JG000142, 2006. 4898
- 5 Hase, F., Wallace, L., McLeod, S. D., Harrison, J., and Bernath, P.: The ACE-FTS atlas of the infrared solar spectrum, *J. Quant. Spectrosc. Ra.*, 111, 521–528, 2010. 4893
- Ho, S. ., Edwards, D. P., Gille, J. C., Luo, M., Osterman, G. B., Kulawik, S. S., and Worden, H.: A global comparison of carbon monoxide profiles and column amounts from Tropospheric Emission Spectrometer (TES) and Measurements of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.-Atmos.*, 114, D21307, doi:10.1029/2009JD012242, 2009. 4891, 4898,
- 10 4903, 4905, 4908
- Illingworth, S. M., Remedios, J. J., and Parker, R. J.: Intercomparison of integrated IASI and AATSR calibrated radiances at 11 and 12 μm , *Atmos. Chem. Phys.*, 9, 6677–6683, doi:10.5194/acp-9-6677-2009, 2009. 4892
- 15 Illingworth, S. M., Remedios, J. J., Boesch, H., Moore, D. P., Sembhi, H., Dudhia, A., and Walker, J. C.: A new optimal estimation retrieval scheme for carbon monoxide using IASI spectral radiances - Part 1: Sensitivity analysis, error budget and simulations, *Atmos. Meas. Tech. Discuss.*, 3, 3747–3802, doi:10.5194/amtd-3-3747-2010, 2010. 4891, 4893, 4898
- Kobayashi, H., Shimota, A., Kondo, K., Okumura, E., Kameda, Y., Shimoda, H., and Ogawa, T.: Development and evaluation of the interferometric monitor for greenhouse gases: A high-throughput Fourier-transform infrared radiometer for nadir Earth observation, *Appl. Optics*,
- 20 38, 6801–6807, 1999. 4891
- Kopacz, M., Jacob, D. J., Fisher, J. A., Logan, J. A., Zhang, L., Megretskaia, I. A., Yantosca, R. M., Singh, K., Henze, D. K., Burrows, J. P., Buchwitz, M., Khlystova, I., McMillan, W. W., Gille, J. C., Edwards, D. P., Eldering, A., Thouret, V., and Nedelec, P.: Global estimates of CO sources with high resolution by adjoint inversion of multiple satellite datasets (MOPITT, AIRS, SCIAMACHY, TES), *Atmos. Chem. Phys.*, 10, 855–876, doi:10.5194/acp-10-855-2010, 2010. 4891
- 25 Larar, A. M., Smith, W. L., Zhou, D. K., Liu, X., Revercomb, H., Taylor, J. P., Newman, S. M., and Schlüssel, P.: IASI spectral radiance validation inter-comparisons: case study assessment from the JAIVEx field campaign, *Atmos. Chem. Phys.*, 10, 411–430, doi:10.5194/acp-10-411-2010, 2010. 4892
- 30 Luo, M., Rinsland, C. P., Rodgers, C. D., Logan, J. A., Worden, H., Kulawik, S., Eldering, A.,

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Goldman, A., Shephard, M. W., Gunson, M., and Lampel, M.: Comparison of carbon monoxide measurements by TES and MOPITT: Influence of a priori data and instrument characteristics on nadir atmospheric species retrievals, *J. Geophys. Res.-Atmos.*, 112, D09303, doi:10.1029/2006JD007663, 2007. 4891, 4903, 4905

5 McMillan, W. W., Barnet, C., Strow, L., Chahine, M. T., McCourt, M. L., Warner, J. X., Novelli, P. C., Korontzi, S., Maddy, E. S., and Datta, S.: Daily global maps of carbon monoxide from NASA's Atmospheric Infrared Sounder, *Geophys. Res. Lett.*, 32, 1–4, 2005. 4891

Rinsland, C. P., Luo, M., Logan, J. A., Beer, R., Worden, H., Rider, D., Osterman, G., Gunson, M., Eldering, A., Goldman, A., Shephard, M., Clough, S. A., Rodgers, C., Lampel, M., and Chiou, L.: Nadir measurements of carbon monoxide distributions by the Tropospheric Emission Spectrometer instrument onboard the Aura Spacecraft: Overview of analysis approach and examples of initial results, *Geophys. Res. Lett.*, 33, L22806, doi:10.1029/2006GL027000, 2006. 4891

15 Rodgers, C.: *Inverse Methods for Atmospheric Sounding: Theory and Practice*, World Scientific, 2000. 4893, 4897

Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys. Res.-Atmos.*, 108, ACH13–1–ACH13–14, 2003. 4891, 4903, 4905, 4906, 4908

20 Seemann, S. W., Borbas, E. E., Knuteson, R. O., Stephenson, G. R., and Huang, H.: Development of a global infrared land surface emissivity database for application to clear sky sounding retrievals from multispectral satellite radiance measurements, *J. Appl. Meteorol. Clim.*, 47, 108–123, 2008. 4893

Thompson, A. M.: The oxidizing capacity of the earth's atmosphere: Probable past and future changes, *Science*, 256, 1157–1165, 1992. 4890

25 USGS: US Geological Survey: GTOPO30 Global 30 Arc-second Digital Elevation Model, Tech. rep., <http://edcwww.cr.usgs.gov/landdaac/gtopo30/gtopo30.html>, last access: November 2010, 1998. 4893

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006. 4904

30 Warner, J., Comer, M. M., Barnet, C. D., McMillan, W. W., Wolf, W., Maddy, E., and Sachse, G.: A comparison of satellite tropospheric carbon monoxide measurements from AIRS and MOPITT during INTEX-A, *J. Geophys. Res.-Atmos.*, 112, D12S17, doi:10.1029/2006JD007925, 2007. 4891, 4903, 4905

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Table 1. Mean and percentage biases for IASI and MOPITT V3 CO profiles, as well as the mean absolute total column biases between the two products, for daytime over the oceanic Southern Africa region on 1 September 2007. N represents the number of retrievals, and the parentheses denote the standard deviations relative to the mean. Note also that the percentage biases represent the ratio to the mean IASI value.

	$x^{\text{MOP3}} - x^{\text{IASI}}$ ($N = 987$)	$x^{\text{MOP3}'} - x^{\text{IASI}'}$ ($N = 974$)	δ_{MODEL} ($N = 975$)	$x^{\text{MOP3}'} - x^{\text{IASI}'}$ ($N = 972$)	$\delta_{\text{MODEL}'}$ ($N = 972$)
850 hPa					
Mean (ppbv)	-12.8 (61.4)	-14.1 (62.9)	-30.3 (18.5)	11.2 (39.1)	7.3 (14.6)
Bias (%)	-7.7 (36.9)	-8.5 (37.8)	-18.7 (11.4)	8.0 (27.8)	5.9 (11.8)
500 hPa					
Mean (ppbv)	14.5 (19.1)	8.1 (20.5)	7.0 (7.1)	5.7 (12.8)	3.6 (5.5)
Bias (%)	15.6 (20.5)	8.8 (22.1)	8.0 (8.1)	6.0 (13.4)	4.1 (6.3)
250 hPa					
Mean (ppbv)	13.9 (16.8)	17.9 (18.3)	17.2 (10.2)	1.5 (8.8)	1.5 (2.1)
Bias (%)	18.2 (22.0)	23.6 (24.1)	23.3 (13.8)	1.6 (9.5)	1.7 (2.1)
Total column					
Absolute Bias (%)	1.2 (20.3)	2.4 (20.4)	1.8 (6.1)	6.4 (17.5)	4.0 (7.3)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Table 2. Same as Table 1, but for IASI and MOPITT V4 CO retrievals.

	$x^{\text{MOP4}} - x^{\text{IASI}}$ ($N = 1041$)	$x^{\text{MOP4}'} - x^{\text{IASI}'}$ ($N = 1038$)	δ_{MODEL} ($N = 1042$)	$x^{\text{MOP4}'} - x^{\text{IASI}'}$ ($N = 1042$)	$\delta_{\text{MODEL}'}$ ($N = 1038$)
900 hPa					
Mean (ppbv)	-74.4 (72.8)	-28.9 (66.7)	-41.9 (35.9)	3.1 (26.9)	3.5 (8.1)
Bias (%)	-42.4 (41.5)	-16.5 (38.0)	-24.3 (20.8)	2.2 (18.7)	2.8 (6.3)
500 hPa					
Mean (ppbv)	22.2 (25.4)	2.0 (19.8)	3.2 (4.2)	2.1 (12.0)	4.2 (6.3)
Bias (%)	23.6 (27.0)	2.2 (21.0)	3.6 (4.8)	2.3 (12.7)	4.8 (7.2)
200 hPa					
Mean (ppbv)	-1.4 (18.6)	13.5 (18.6)	14.3 (11.2)	0.5 (5.8)	2.4 (3.5)
Bias (%)	-1.9 (25.6)	18.5 (25.5)	20.2 (15.9)	0.5 (6.8)	2.9 (4.2)
Total column					
Absolute Bias (%)	14.0 (22.0)	5.6 (20.3)	18.4 (9.3)	2.9 (14.7)	2.5 (3.2)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Table 3. δx_{BIAS} for IASI and MOPITT V3 CO retrievals for different scenarios.

	day and ocean		night and ocean		night and land	
	$x^{\text{MOP3}'} - x^{\text{IASI}'}$ ($N = 972$)	$\delta_{\text{MODEL}'}$ ($N = 972$)	$x^{\text{MOP3}'} - x^{\text{IASI}'}$ ($N = 908$)	$\delta_{\text{MODEL}'}$ ($N = 908$)	$x^{\text{MOP3}'} - x^{\text{IASI}'}$ ($N = 711$)	$\delta_{\text{MODEL}'}$ ($N = 710$)
850 hPa						
Mean (ppbv)	11.2 (39.1)	7.3 (14.6)	11.5 (38.8)	1.2 (7.4)	28.7 (44.1)	19.1 (29.7)
Bias (%)	8.0 (27.8)	5.9 (11.8)	9.7 (32.9)	1.1 (7.0)	15.8 (24.3)	11.4 (17.7)
500 hPa						
Mean (ppbv)	5.7 (12.8)	3.6 (5.5)	12.0 (8.1)	1.2 (2.8)	8.6 (13.1)	-0.6 (5.6)
Bias (%)	5.8 (13.4)	4.1 (6.3)	15.2 (10.2)	1.6 (3.7)	8.3 (12.5)	-0.7 (6.0)
250 hPa						
Mean (ppbv)	1.5 (8.8)	1.5 (2.1)	6.3 (13.4)	-0.1 (1.5)	7.5 (11.0)	-1.5 (4.9)
Bias (%)	1.6 (9.5)	1.7 (2.4)	7.6 (16.3)	-0.1 (1.8)	8.0 (11.8)	-1.7 (5.7)
Total column						
Absolute Bias (%)	6.4 (17.5)	4.0 (7.3)	10.4 (14.4)	0.8 (4.4)	13.0 (14.9)	3.7 (6.7)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.

Table 4. δx_{BIAS} for IASI and MOPITT V4 CO retrievals for different scenarios.

	day and ocean		night and ocean		night and land	
	$x^{\text{MOP4'}} - x^{\text{IASI'}}$ ($N = 1038$)	$\delta_{\text{MODEL'}}$ ($N = 1038$)	$x^{\text{MOP4'}} - x^{\text{IASI'}}$ ($N = 925$)	$\delta_{\text{MODEL'}}$ ($N = 925$)	$x^{\text{MOP4'}} - x^{\text{IASI'}}$ ($N = 211$)	$\delta_{\text{MODEL'}}$ ($N = 211$)
900 hPa						
Mean (ppbv)	3.1 (26.9)	3.5 (8.1)	-3.4 (30.0)	-2.2 (4.4)	-4.9 (48.9)	1.41 (5.6)
Bias (%)	2.2 (18.7)	2.8 (6.3)	-2.7 (24.0)	-1.8 (3.6)	-2.7 (27.3)	1.0 (4.2)
500 hPa						
Mean (ppbv)	2.1 (12.0)	4.2 (6.3)	3.9 (8.0)	0.7 (1.9)	3.9 (12.4)	0.6 (3.7)
Bias (%)	2.3 (12.7)	4.8 (7.2)	4.9 (10.0)	0.8 (2.4)	3.8 (12.0)	0.7 (4.4)
200 hPa						
Mean (ppbv)	0.5 (5.8)	2.4 (3.5)	4.4 (11.5)	0.5 (1.6)	5.7 (11.0)	0.1 (2.6)
Bias (%)	0.5 (6.8)	2.9 (4.2)	5.5 (14.6)	0.6 (2.1)	6.8 (13.1)	0.1 (3.2)
Total column						
Absolute Bias (%)	2.9 (14.7)	2.5 (3.2)	4.5 (14.4)	0.1 (2.1)	7.4 (16.2)	2.1 (4.3)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 1. IASI and MOPITT V3 a priori profile and associated errors (i.e. the diagonal of the a priori covariance matrix), as well as the MOPITT V4 a priori profile averaged over the region shown in Fig. 4 for 1 September 2007. The profiles have all been interpolated onto the MOPITT V3 pressure grid for ease of reference.

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 2. Averaged daytime and ocean averaging kernels for 1 September 2007 over Southern Africa, for: **(A)** MOPITT V3 (A^{MOP3}), **(B)** MOPITT V4 (A^{MOP4}), **(C)** pressure-layer-normalised averaging kernels for MOPITT V3 (A_N^{MOP3}), and **(D)** pressure-layer-normalised averaging kernels for MOPITT V4 (A_N^{MOP4}). The MOPITT V3 averaging kernels are in normal space, whereas the MOPITT V4 averaging kernels relate to $\log(\text{VMR})$ values, and the units of the pressure-layer-normalised averaging kernels are hPa^{-1} .

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A comparison of
OEM CO retrievals
from the IASI and
MOPITT instruments**

S. Illingworth et al.




Fig. 3. MOPITT V3 and V4 mean CO profiles (solid red line and blue line, respectively) and standard deviation relative to their means (dashed horizontal lines) for 1 September 2007 over the Southern Africa region, for the daytime and over the ocean. The MOPITT V3 a priori profile (dashed turquoise line), and mean MOPITT V4 a priori profile (dashed yellow line) used in the retrievals are also shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**A comparison of
OEM CO retrievals
from the IASI and
MOPITT instruments**

S. Illingworth et al.




Fig. 4. CO total column density over Southern Africa during the daytime of 1 September 2007: (A) IASI, (B) MOPITT V3, (C) MOPITT V4, and (D) GEOS-Chem.




Fig. 5. Daytime and ocean averaging kernels over the Southern Africa region for 1 September 2007, for: **(A)** MOPITT V3 (\mathbf{A}^{MOP3}) at MOPITT V3 pressure levels, **(B)** IASI (\mathbf{A}^{IASI}) at the IASI pressure levels closest to the MOPITT V3 pressure levels, **(C)** pressure-layer-normalised averaging kernels for MOPITT V3 ($\mathbf{A}_N^{\text{MOP3}}$), and **(D)** pressure-layer-normalised averaging kernels for IASI ($\mathbf{A}_N^{\text{IASI}}$). The units of the pressure-layer-normalised averaging kernels are hPa^{-1} .




Fig. 6. Daytime and ocean averaging kernels over the Southern Africa region for 1 September 2007, for: **(A)** MOPITT V4 (\mathbf{A}^{MOP4}) at MOPITT V4 pressure levels, **(B)** IASI (\mathbf{A}^{IASI}) at the IASI pressure levels closest to the MOPITT V4 pressure levels, **(C)** pressure-layer-normalised averaging kernels for MOPITT V4 ($\mathbf{A}_N^{\text{MOP4}}$), and **(D)** pressure-layer-normalised averaging kernels for IASI ($\mathbf{A}_N^{\text{IASI}}$). The units of the pressure-layer-normalised averaging kernels are hPa^{-1} .

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 7. (A) MOPITT V3 and IASI CO mean profiles (solid red line and blue line, respectively) and standard deviation relative to their means (dashed lines), (B) MOPITT V4 and IASI CO mean profiles (solid red line and blue line, respectively) and standard deviation relative to their means (dashed lines), (C) the mean difference between MOPITT V3 CO and IASI CO (solid green line) and the standard deviation (dashed green line) relative to the mean, and (D) the mean difference between MOPITT V4 CO and IASI CO (solid green line) and the standard deviation (dashed green line) relative to the mean. The IASI CO retrievals closest to the MOPITT pressure levels are used here, and all comparisons are for over the ocean and during the daytime of 1 September 2007, over the Southern Africa region.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 8. Comparison of retrievals performed with the same a priori (ULIRS profile and error covariances), $\mathbf{x}^{\text{MOPITT}}$ and \mathbf{x}^{IASI} : **(A)** adjusted MOPITT V3 and IASI CO mean profiles; **(B)** adjusted MOPITT V4 and IASI CO mean profiles; **(C)** the mean difference between adjusted MOPITT V3 CO and IASI CO profiles; and **(D)** the mean difference between adjusted MOPITT V4 CO and IASI CO profiles.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 9. Same as Fig. 8, except for $x_{\text{MODEL}}^{\text{MOPITT}}$ and $x_{\text{MODEL}}^{\text{IASI}}$: **(A)** adjusted MOPITT V3 and IASI CO mean profiles; **(B)** adjusted MOPITT V4 and IASI CO mean profiles; **(C)** the mean difference between adjusted MOPITT V3 CO and IASI CO profiles; and **(D)** the mean difference between adjusted MOPITT V4 CO and IASI CO profiles. (A) and (B) are in relation to the retrieval of a true profile, as provided by GEOS-Chem.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 10. Daytime and ocean averaging kernels over the Southern Africa region for 1 September 2007, for: **(A)** Adjusted MOPITT V3 averaging kernels converted onto the IASI pressure grid $A^{MOP3''}$; **(B)** IASI averaging kernels at pressure levels closest to the MOPITT V3 pressure levels A^{IASI} ; **(C)** $A^{MOP3''} - A^{IASI}$; and **(D)** $A^{MOP3''} - A^{MOP3''} \cdot A^{IASI}$.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 11. Same as Fig. 10, but for MOPITT V4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 12. with reduced smoothing error using the derived quantity in Eq. 8, x^{MOPIT} and x^{IASI} : **(A)** adjusted MOPITT V3 and MOPITT-smoothed IASI CO mean profiles; **(B)** adjusted MOPITT V4 and MOPITT-smoothed IASI CO mean profiles; **(C)** the mean difference between adjusted MOPITT V3 CO and MOPITT-smoothed IASI CO profiles; and **(D)** the mean difference between adjusted MOPITT V4 CO and MOPITT-smoothed IASI CO profiles.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

A comparison of OEM CO retrievals from the IASI and MOPITT instruments

S. Illingworth et al.




Fig. 13. Same as Fig. 12, but for $x_{\text{MODEL}}^{\text{MOPITT}}$ and $x_{\text{MODEL}}^{\text{IASI}}$: **(A)** adjusted MOPITT V3 and MOPITT-smoothed IASI CO mean profiles; **(B)** adjusted MOPITT V4 and MOPITT-smoothed IASI CO mean profiles; **(C)** the mean difference between adjusted MOPITT V3 CO and MOPITT-smoothed IASI CO profiles; and **(D)** the mean difference between adjusted MOPITT V4 CO and MOPITT-smoothed IASI CO profiles. (A) and (B) are in relation to the retrieval of a true profile, as provided by GEOS-Chem.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)