

## ***Interactive comment on “A sea surface reflectance model for (A)ATSR, and application to aerosol retrievals” by A. M. Sayer et al.***

**A. M. Sayer et al.**

sayer@atm.ox.ac.uk

Received and published: 26 May 2010

The authors thank the three anonymous reviewers for their comments on the manuscript. Our response to this review is given below.

Review 1.

“My only a big question on this paper is that the authors mentioned (second paragraph in page 1053) “The BRDF can take values from around  $10^{-5}$  to 1 dependent on the wavelength and location with respect to the sun-glint region.” In principle, the BRDF should take value from 0 to infinite, and this can be derived from equations 17 and/or

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



19, for example, when  $\cos(\theta_v) - - > 0$ ,  $\cos(\theta_s) - - > 0$  or ,  $\cos(\beta) - - > 0$ , the  $\rho_{gl} - - > infinite$ "

We agree with this statement. The sentence was intended to convey that the sea BRDF in this spectral region is modelled as taking values between around  $10^{-5}$  to 1 (although as stated BRDF can be between zero and infinite), and this has been clarified in the revised version of the manuscript.

---

Interactive comment on Atmos. Meas. Tech. Discuss., 3, 1023, 2010.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

