

**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi¹, T. Neubert¹, S. Syndergaard², and J. Nielsen²

¹DTU Space, National Space Institute, Copenhagen, Denmark

²DMI, Danish Meteorological Institute, Copenhagen, Denmark

Received: 27 January 2011 – Accepted: 17 February 2011 – Published: 28 February 2011

Correspondence to: R. Biondi (ribi@space.dtu.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The tropical deep convection affects the radiation balance of the atmosphere changing the water vapor mixing ratio and the temperature of the upper troposphere lower stratosphere. The aim of this work is to better understand these processes and to investigate if severe storms leave a significant signature in radio occultation profiles in the tropical tropopause layer. Using tropical cyclone best track database and data from different GPS radio occultation missions (COSMIC, GRACE, CHAMP, SACC and GPSMET), we selected 1194 profiles in a time window of 3 h and a space window of 300 km from the eye of the cyclone. We show that the bending angle anomaly of a GPS radio occultation signal is typically larger than the climatology in the upper troposphere and lower stratosphere and that a double tropopause during deep convection can easily be detected using this technique. Comparisons with co-located radiosondes, climatology of tropopause altitudes and GOES analyses are also shown to support the hypothesis that the bending angle anomaly can be used as an indicator of convective towers. The results are discussed in connection to the GPS radio occultation receiver which will be part of the Atomic Clock Ensemble in Space (ACES) payload on the International Space Station.

1 Introduction

Deep convective systems influence global circulation and the mass exchange between the troposphere and stratosphere (Fueglistaler et al., 2009), and they also have been recognized as a crucial process for the understanding of the climate changes. Temperature and water vapour variation in the upper troposphere and lower stratosphere (UTLS) are closely connected to the convection (Danielsen, 1982; Sherwood et al., 2003; Chae et al., 2010;) but the difficulty to measure those parameters at such altitudes, with high vertical resolution and sensitivity still raises many questions on their behaviour (Forster and Shine, 1999). The stratospheric water vapour, like the CO₂, is

AMTD

4, 1371–1395, 2011

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a greenhouse gas contributing to the global warming (Manabe and Wetherald, 1967; Soden and Fu, 1995; Forster and Shine, 2002; Solomon et al., 2010) and the variation of temperature in the tropical tropopause layer (TTL) during the convection is strictly linked to the variation of water vapor overshooted by the storm (Kim, 2005). The tropical cyclones (TCs) are severe weather events often leading to deep convective activity and persisting for long time. They alter the physics and chemistry of the TTL and they play a fundamental role on the atmospheric circulation and troposphere-stratosphere exchange (Ray and Rosenlof, 2007; Cairo et al., 2008; Fueglistaler et al., 2009; Roms and Kuang, 2009). The GPS Radio Occultation (RO) technique (Kursinki et al., 1997) is useful to study severe weather phenomena because the GPS signal is independent on the weather conditions and it allows profiling atmospheric parameters with high vertical resolution and accuracy. The past and ongoing GPS RO missions, Global Positioning System/Meteorology (GPS/MET) satellite (Rocken et al., 1997), the CHALLENGING Minisatellite Payload (CHAMP) satellite (Wickert et al., 2001), the Satellite de Aplicaciones Cientificas-C (SAC-C) satellite (Hajj et al., 2004), the Constellation Observing System for Meteorology, Ionosphere and Climate (COSMIC) six-satellite constellation (Anthes et al., 2008), and the Gravity Recovery And Climate Experiment (GRACE) twin satellites (Beyerle et al., 2005), have a denser number of measurements at mid and high latitude due to the inclination of the orbits, but they do not provide a good coverage of the tropical area (Fig. 1). A new advanced GPS receiver is planned to be launched in the near future on board of the International Space Station (ISS) as part of the Atomic Clock Ensemble in Space (ACES) payload (Svehla et al., 2006). It will be mounted on the external platform of the Columbus laboratory module, viewing towards the limb in the forward direction. The inclination of ISS orbit (51.6°) will allow the ACES GPS receiver to monitor the major convective regions of the Earth contributing to increase the number of GPS RO within the tropical regions (Fig. 2). We have selected all the RO profiles from the above-mentioned missions during the period 1995–2009, analysed them, and compared them with co-located data along the TC best tracks. The Geostationary Operational Environmental Satellite (GOES) brightness temperature (BT) were

analysed near the location of the RO profiles in order to detect the overshooting during the TCs. The GPS RO bending angle and the temperature profiles show a clear signature of the convection in the UTLS which is confirmed by the comparison with co-located radiosonde (RAOB) data.

2 Data description

The TCs best tracks were collected from different sources: US National Hurricane Center (Atlantic, Caribbean, Gulf of Mexico and Eastern Pacific), Australian Government Bureau of Meteorology (Western, Northern and Eastern Australia), Japan Meteorological Agency (Western and Northern Pacific), Meteo France (Southwest Indian Basin) and Unysis Weather (North Indian Ocean Basin, Bay of Bengal and Arabian Sea). All these institutes provide information every 6 h, including at least the name, dates, coordinates, minimum pressure, maximum wind speed and category of the tropical cyclone. In some cases, during the hurricane hunters missions, additional information is available, such as eye dimension, moving cyclone direction and speed.

The RAOBs in FSL (Forecast System Laboratory) format, are part of the National Oceanic and Atmospheric Administration Earth System Research Laboratory (NOAA-ESRL) radiosonde database, and they are used as in situ validation of RO temperature and water vapour mixing ratio profiles.

Tropopause climatology is used as reference to detect if the convection reaches the TTL or not. For this purpose we used the climatological tropopause height from National Centers for Environmental Prediction (NCEP) analyses, containing monthly mean zonally averaged tropopause heights. This encompasses 40 years of data starting in January 1958 with latitudinal resolution of 2.5 degrees.

The GPS ROs that we have analyzed were downloaded from the COSMIC Data Analysis and Archive Center (CDAAC) website. We collected all the GPS/MET, SAC-C, CHAMP, GRACE and COSMIC profiles as shown in Table 1. The results of this work are obtained using level 2 data products in netCDF format:

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- atmPrf, which is an atmospheric profile without moisture information, containing coordinates and mean sea level altitude of the perigee point, dry pressure and temperature (derived assuming no water vapour), refractivity, bending angle and impact parameter.
- 5 – wetPrf, which is an interpolated product obtained using 1DVar (one dimensional variational) technique with a vertical resolution of 100 m, containing latitude and longitude of the perigee point, pressure, temperature, water vapour pressure, refractivity and mean sea level altitude of the perigee point.
- sonPrf, which is radiosonde data co-located with the GPS RO (time window of 6 h and space window of 400 km) containing latitude, longitude, pressure, tempera-
10 ture, water vapour pressure, refractivity and mean sea level geometric height.

The GOES brightness temperature analyses were used as a detector of overshoots. We downloaded from the Comprehensive Large Array Data Stewardship System (CLASS) the GOES VARIable IMAger data (GVAR_IMG) in “area” format with native
15 spatial resolution acquired by GOES11 and GOES12. Using the BT at 6.8 microns (band 3), which is sensitive to the water vapour, and the BT at 10.7 microns (band 4), it is possible to detect convective systems (Schmetz et al., 1997; Chaboureau et al., 2007). In clear sky conditions and low level clouds, the BT in the water vapour channel is lower than the BT at 10.7 microns. When the overshoot is present, there is an in-
20 version and the BT at 6.8 microns becomes larger by a few degrees. GOES data were also used to detect if the RO and the RAOB were acquired in the same meteorological conditions.

3 Methods

The GPS receivers measure the phase and amplitude of two L-band signals. Using
25 these it is possible to get the bending angle, which is used to compute the refractivity

**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

with the co-located bending angle anomaly shows two positive spikes at the same altitude of the two temperature minima. Similar features are often evident also in the RO temperature profile, but sometimes it is not as clearly seen as it is in the bending angle anomaly.

5 Figure 7 illustrates the situation during the TC Bill. The RO was acquired on 22 August 2009 at 08:27:00 UTC in the Atlantic Ocean when Bill was weakening from hurricane category 3 to hurricane category 2. The radiosonde was launched at 11:00:00 UTC and the distance between the radiosonde station (Bermuda airport) and the RO tangent point at 15 km altitude was 98.32 km.

10 Figure 8 shows the difference of the BT at 6.8 microns and the BT at 10.7 microns from GOES11 in the same region where the RO and the RAOB were acquired. An inversion is present; consequently some overshoot should be expected.

The RAOB temperature profile shows a double minimum, the lowest exactly at the standard tropopause altitude and the second one a couple of kilometers above (Fig. 9). The local temperature variation is larger than 5 K and this defines an evident double tropopause. The bending angle anomaly, as usual during TCs, has two local maxima, the lowest one 10% larger than the climatology in the same area (at 17 km of altitude), and the second one a couple of kilometers above (Fig. 10). Within the uncertainty and limited vertical resolution of the RAOB, and considering that the two measurements are not exactly coincident, the spikes in the bending angle and temperature anomalies, as well as the minima in the RO temperature profile (although not as distinct) corresponds reasonably well to the minima in the RAOB temperature profile. The water vapor mixing ratio profile from the RO (1DVar approach) does not show any variation, but the RAOB profile shows an increase at the altitude of the local temperature maximum (Fig. 11).

25 The double spike of the bending angle anomaly at these altitudes is a common feature of the UTLS during TCs. So far, we have selected about 300 cases. In all the cases the lowest coldest point (and the warming about 1 km above) is clearly recognizable in the RO temperature profiles but it is sometimes difficult to recognize a clear double tropopause.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The double local minima detected during our studies, confirms the results of Danielsen (Danielsen, 1982, 1993) explaining the lowest one as the mean tropopause and the highest one as the cold stratospheric anvil produced by the deep convection. The same temperature trend has been clearly recognized with in situ measurements during the Tropospheric/Stratospheric Water Vapor Exchange Experiment (TSWVEE) (Kley et al., 1982) and the Tropical Convection, Cirrus and Nitrogen Oxides (TROCCI-NOX) (Chaboureau, 2007).

6 Discussions and conclusions

Our results show that the bending angle from the GPS RO signal contains interesting information on the tropopause during TCs. In general the bending angle anomaly relative to the climatology is positive during tropical cyclone season in the UTLS and the magnitude of the anomaly increases during a TC. This variation of anomaly is not enough to detect convective towers since it could be due to a simple reduction of the tropopause temperature, but it is anyway a clear signature of the TC. Extending the analysis to using RAOBs, during strong convective events a double tropopause is often recognized and the bending angle anomaly can be used to detect it. In this way the bending angle can be used as an indicator of convective towers. Our results suggest that ACES missions on board of ISS will increase the RO measurements in the tropics providing important support to the study of TCs and the study of overshoots.

Acknowledgement. We would like to thank G. Beyerle for the ACES simulator.

References

- Anthes, R. A., Bernhardt, P. A., Chen, Y., Cucurull, L., Dymond, K. F., Ector, D., Healy, S. B., Ho, S.-H., Hunt, D. C., Kuo, Y.-H., Liu, H., Manning, K., McCormick, C., Meehan, T. K., Randel, W. J., Rocken, C., Schreiner, W. S., Sokolovskiy, S. V., Syndergaard, S., Thompson, D. C., Trenberth, K. E., Wee, T.-K., Yen, N. L., and Zeng, Z.: The COSMIC/Formosat/3 mission: early results, *B. Am. Meteorol. Soc.*, 89, 313–333, doi:10.1175/BAMS-89-3-313, 2008.
- Beyerle, G., Schmidt, T., Michalak, G., Heise, S., Wickert, J., and Reigber, C.: GPS radio occultation with GRACE: atmospheric profiling utilizing the zero difference technique, *Geophys. Res. Lett.*, 32, L13806, doi:10.1029/2005GL023109, 2005.
- Biondi, R., Neubert, T., Syndergaard, S., and Nielsen, J.: Measurements of the upper troposphere and lower stratosphere during tropical cyclones using the GPS radio occultation technique, *Adv. Space Res.*, 47, 348–355, doi:10.1016/j.asr.2010.05.031, 2011.
- Cairo, F., Buontempo, C., MacKenzie, A. R., Schiller, C., Volk, C. M., Adriani, A., Mitev, V., Matthey, R., Di Donfrancesco, G., Oulanovsky, A., Ravegnani, F., Yushkov, V., Snels, M., Cagnazzo, C., and Stefanutti, L.: Morphology of the tropopause layer and lower stratosphere above a tropical cyclone: a case study on cyclone Davina (1999), *Atmos. Chem. Phys.*, 8, 3411–3426, doi:10.5194/acp-8-3411-2008, 2008.
- Chaboureau, J.-P., Cammas, J.-P., Duron, J., Mascart, P. J., Sitnikov, N. M., and Voessing, H.-J.: A numerical study of tropical cross-tropopause transport by convective overshoots, *Atmos. Chem. Phys.*, 7, 1731–1740, doi:10.5194/acp-7-1731-2007, 2007.
- Chae, J. H., Wu, D. L., Read, W. G., and Sherwood, S. C.: The role of tropical deep convective clouds on temperature, water vapor, and dehydration in the tropical tropopause layer (TTL), *Atmos. Chem. Phys. Discuss.*, 10, 8963–8994, doi:10.5194/acpd-10-8963-2010, 2010.
- Danielsen, E. F.: A dehydration mechanism for the stratosphere, *Geophys. Res. Lett.*, 9, 605–608, doi:10.1029/GL009i006p00605, 1982.
- Danielsen, E. F.: In situ evidence of rapid, vertical, irreversible transport of lower tropospheric air into the lower tropical stratosphere by convective cloud turrets and by larger-scale upwelling in tropical cyclones, *J. Geophys. Res.*, 98, 8665–8681, doi:10.1029/92JD02954, 1993.
- Forster P. M. and Shine, K. P.: Stratospheric water vapour changes as a possible contributor to observed stratospheric cooling, *Geophys. Res. Lett.*, 26, 3309–3312, doi:10.1029/1999GL010487, 1999.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Forster, P. M. and Shine, K. P.: Assessing the climate impact of trends in stratospheric water vapor, *Geophys. Res. Lett.*, 29, 101–102, doi:10.1029/2001GL013909, 2002.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009.
- 5 Hajj, G. A., Ao, C. O., Iijima, B. A., Kuang, D., Kursinski, E. R., Mannucci, A. J., Meehan, T. K., Romans, L. J., de La Torre Juarez, M., and Yunck, T. P.: CHAMP and SAC-C atmospheric occultation results and intercomparisons, *J. Geophys. Res.*, 109, D06109, doi:10.1029/2003JD003909, 2004.
- Kim, H. C.: The effect of deep convection on temperatures in the tropical tropopause layer and its implications to the regulation of tropical lower stratospheric humidity, PhD Thesis, University of Houston, Houston, TX, USA, 2005.
- 10 Kley, D., Schmeltekopf, A. L., Kelly, K., Winkler, R. H., Thompson, T. L., and McFarland, M.: Transport of water through the tropical tropopause, *Geophys. Res. Lett.*, 9, 617–620, doi:10.1029/GL009i006p00617, 1982.
- 15 Kuo, Y. H., Wee, T. K., Sokolovskiy, S., Rocken, C., Schreiner, W., Hunt, D., and Anthes, R. A.: Inversion and error analysis of GPS radio occultation data, *J. Meteor. Soc. Japan*, 82, 507–531, 2004.
- Kursinski, E. R., Hajj, G. A., Schofield, J. T., Linfield, R. P., and Hardy, K. R.: Observing Earth's atmosphere with radio occultation measurements using the Global Positioning System, *J. Geophys. Res.*, 102, 23429–23465, 1997.
- 20 Manabe, S. and Wetherald, R. T.: Thermal equilibrium of the atmosphere with a given distribution of relative humidity, *J. Atmos. Sci.*, 24, 241–259, 1967.
- Ray, E. A. and Rosenlof, K. H.: Hydration of the upper troposphere by tropical cyclones, *J. Geophys. Res.*, 112, D12311, doi:10.1029/2006JD008009, 2007.
- 25 Rocken, C., Anthes, R., Exner, M., Hunt, D., Sokolovskiy, S., Ware, R., Gorbunov, M., Schreiner, W., Feng, D., Herman, B., Kuo, Y.-H., and Zou, X.: Analysis and validation of GPS/MET data in the neutral atmosphere, *J. Geophys. Res.*, 102, 29849–29860, 1997.
- Romps, D. M. and Kuang, Z.: Overshooting convection in tropical cyclones, *Geophys. Res. Lett.*, 36, L09804, doi:10.1029/2009GL037396, 2009.
- 30 Schmetz, J., Tjembs, S. A., Gube, M., and Van de Berg, L.: Monitoring deep convection and convective overshooting with Meteosat, *Adv. Space Res.*, 19, 433–441, doi:10.1016/S0273-1177(97)00051-3, 1997.
- Sherwood, S. C., Horinouchi, T., and Zeleznik, H. A.: Convective impact on temperatures ob-

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

served near the tropical tropopause, *J. Atmos. Sci.*, 60, 1847–1856, 2003.

Soden, B. J. and Fu, R.: A satellite analysis of deep convection, upper tropospheric humidity and the greenhouse effect, *J. Climate*, 8, 2333–2351, 1995.

Solomon, S., Rodenlof, K., Portmann, R., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G. K.: Contribution of stratospheric water vapor to decadal changes in the rate of global warming, *Science*, 327, 1219–1223, doi:10.1126/science.1182488, 2010.

Svehla, D., Rothacher, M., Ziebart, M., and Salomon, C.: Galileo on board International Space Station and synergy with the ACES clock ensemble, EGU General Assembly, 2006.

Wickert, J., Reigber, C., Beyerle, G., König, R., Marquardt, C., Schmidt, T., Grundwaldt, L., Galas, R., Meehan, T. K., Melbourne, W. G., and Hocke, K.: Atmosphere sounding by GPS radio occultation: first results from CHAMP, *Geophys. Res. Lett.*, 28, 3263–3266, doi:10.1029/2001GL013117, 2001.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Table 1. Number of profiles (second column) analysed and number of coincidences between GPS radio occultation and tropical cyclones in a time window of 3 h and a space window of 300 km (third column).

Mission	# profiles	RO-TC
GPSMET (1995–1997)	5002	1
SAC-C (2001–2002)	60 354	40
CHAMP (2001–2008)	397 193	180
COSMIC (2006)	249 923	179
COSMIC (2007)	614 201	194
COSMIC (2008)	641 440	335
COSMIC (2009)	644 035	211
GRACE (2007–2009)	113 567	54
Total	2 725 715	1194

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.

Table 2. Number of coincidences between GPS radio occultation and tropical cyclones in a time window of 3 h and a space window of 300 km, for each year and each mission.

	GPSMET	SACC	CHAMP	COSMIC	GRACE	Total
1995	1					1
2001		13	9			22
2002		27	28			55
2003			29			29
2004			40			40
2005			33			33
2006			12	179		191
2007			9	194	18	221
2008			20	335	22	377
2009				211	14	225
Total	1	40	180	919	54	1194

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. GPS radio occultation coverage. Total number of occultations (from GPSMET, SACC, CHAMP, COSMIC and GRACE covering the period 1995–2009) gridded with 1 degree of resolution.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.




Fig. 2. Histogram of 1 month (July 2008) simulated ACES radio occultations vs. latitude.

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[⏪](#) [⏩](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.




Fig. 3. Averaged bending angle anomaly profile for 1194 TC cases from 1995 to 2009 (black line) \pm with the standard deviation of the mean (grey lines).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.




Fig. 4. Distribution of positive peak bending angle anomalies between 14 and 18 km during TCs.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Radio occultation
bending angle
anomalies during
tropical cyclones**

R. Biondi et al.




Fig. 5. Distribution of mean negative bending angle anomalies between 14 and 18 km during TCs.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.




Fig. 6. Distribution of mean negative bending angle anomalies between 14 and 18 km during normal conditions, i.e., in the absence of TCs.

[Title Page](#)
[Abstract](#) | [Introduction](#)
[Conclusions](#) | [References](#)
[Tables](#) | [Figures](#)
[⏪](#) | [⏩](#)
[◀](#) | [▶](#)
[Back](#) | [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)






Fig. 7. Tropical cyclone Bill on 22 August 2009 with the co-located radiosonde (white balloon) and radio occultation (red line).

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 8. Event Bill 22 August 2009, 08:27:00 UTC, (Latitude $32^{\circ}30'$ and Longitude $-65^{\circ}42'$ at 15 km of altitude): Difference between the brightness temperature (BT) at 6.8 microns and the BT at 10.7 microns (BT in Kelvin) from GOES 11. The red line indicates the projection of the RO tangent points.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.




Fig. 9. Event Bill 22 August 2009, 08:27:00 UTC: temperature profile between 13 and 20 km of altitude. The blue line is the RAOB and the red line is the RO. The horizontal black lines show the minimum and maximum standard tropopause altitude at those latitudes from NCEP 40 years dataset.

[Title Page](#) | [Abstract](#) | [Introduction](#) | [Conclusions](#) | [References](#) | [Tables](#) | [Figures](#)

[⏪](#) | [⏩](#) | [◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 10. Event Bill 22 August 2009, 08:27:00 UTC: bending angle percentage anomaly (green) and corresponding temperature percentage anomaly (blue), between 13 and 20 km.

Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radio occultation bending angle anomalies during tropical cyclones

R. Biondi et al.




Fig. 11. Event Bill 22 August 2009, 08:27:00 UTC: water vapour mixing ratio from the RAOB (blue) and the RO/1DVar (red), between 13 and 20 km. The horizontal black lines show the minimum and maximum standard tropopause altitude at those latitudes from NCEP 40 years dataset.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪
⏩

◀
▶

Back	Close
------	-------

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

