Atmos. Meas. Tech. Discuss., 4, C823–C824, 2011 www.atmos-meas-tech-discuss.net/4/C823/2011/ © Author(s) 2011. This work is distributed under the Creative Commons Attribute 3.0 License.



Interactive comment on "Detection of HO₂ by laser-induced fluorescence: calibration and interferences from RO₂ radicals" *by* H. Fuchs et al.

H. Fuchs et al.

h.fuchs@fz-juelich.de

Received and published: 10 June 2011

We apologize for not having answered this question in our first response.

Comment: It is unclear from the paper whether the hydrocarbons are added to the calibration gas mixture before the Hg lamp or after it. In Fuchs et al 2008, in which the RO2 measurement technique which uses the conversion of RO_2 into HO_2 followed by detection of HO2 using the chemical conversion method with NO is described, it would seem that the parent hydrocarbons are added before the lamp. If this is the case, the authors should discuss possible photolysis of the alkenes in this present study which may lead to the formation of RO_2 or HO_2 and impact on the interference measurements. Ethene for example might be expected to undergo some photolysis to generate an H atom which then generates HO_2 quickly in air. Previous studies (e.g. Giroux et al.,

C823

Can. J. Chem. 67, 1166, 1989, and many others) discuss yields of H atoms at 193 nm, and there have been studies at 185 nm. It may be difficult to distinguish between an HO_2 interference from RO_2 decomposition and photolytic sources of HO_2 .

Response: The hydrocarbons were added to the gas before it reached the photolysis region, as mentioned in Fuchs et al. 2008. In that paper we discussed and excluded the relevance of photolytical radical production from the 185 nm photolysis of isoprene. In the present work, we tested all hydrocarbons listed in Table 2 by feeding the calibration source with a mixture of the corresponding hydrocarbon and dry air, avoiding radical production from the case, a significant radical production from hydrocarbon photolysis could be detected by LIF.

We will add in the manuscript on page 1265 line 26: "Since hydrocarbons were added upstream of the photolysis region of the radical source, photolysis of these hydrocarbons at 185nm may occur (Demers et al. 1978, Giroux et al. 1989). In order to test for effects of photolysis on experiments conducted here, the fluorescence signal was measured, when hydrocarbons were mixed in dry air, so that radicals are only produced by photolysis of the hydrocarbon. No significant radical production from hydrocarbon photolysis could be detected by LIF."

Interactive comment on Atmos. Meas. Tech. Discuss., 4, 1255, 2011.