

Depolarization ratio of Polar Stratospheric Clouds in coastal Antarctica: Comparison analysis between ground-based Micro Pulse Lidar and space-borne CALIOP observations

C. Córdoba-Jabonero¹, J.L. Guerrero-Rascado^{2,3}, D. Toledo¹, M. Parrondo¹, M. Yela¹, M. Gil¹, H.A. Ochoa⁴

[1]{Instituto Nacional de Técnica Aeroespacial (INTA), Atmospheric Research and Instrumentation Branch, Torrejón de Ardoz, 28850, Madrid, Spain}

[2]{Andalusian Center for Environmental Research (CEAMA), Group of Atmospheric Physics, Universidad de Granada-Junta de Andalucía, Granada, Spain}

[3]{Universidad de Granada (UGR), Department of Applied Physics, Granada, Spain}

[4]{Dirección Nacional del Antártico/Instituto Antártico Argentino (DNA/IAA), Buenos Aires, Argentina}

Correspondence to: C. Córdoba-Jabonero (cordobajc@inta.es)

Abstract

Polar Stratospheric Clouds (PSCs) play an important role in polar ozone depletion, since they are involved in diverse ozone destruction processes (chlorine activation, denitrification). The degree of that ozone reduction is depending on the type of PSCs, and hence on their occurrence. Therefore PSC characterization, mainly focused on PSC-type discrimination, is widely demanded. The backscattering (R) and volume linear depolarization (δ^V) ratios are the parameters usually used in lidar measurements for PSC detection and identification. In this work, an improved version of the standard NASA/Micro Pulse Lidar (MPL-4), which includes a built-in depolarization detection module, has been used for PSC observations above the coastal Antarctic Belgrano II station (Argentina, 77.9°S 34.6°W, 256 m a.s.l.) since 2009. Examination of the MPL-4 δ^V feature as a suitable index for PSC-type discrimination is based on the analysis of the two-channel data, i.e. the parallel (p-) and perpendicular (s-) polarized MPL signals. This study focuses on the comparison of coincident δ^V -profiles as obtained from ground-based MPL-4 measurements during three Antarctic winters with those reported from the space-borne lidar CALIOP aboard the CALIPSO satellite in the same period (83 simultaneous cases are analysed for 2009-2011 austral winter times). Three different approaches are considered for the comparison analysis between both lidar profile datasets in order to test the degree of agreement: the correlation coefficient (CC), as a measure of the relationship between both PSC vertical structures; the mean differences together with their root mean square (RMS) values found between datasets; and the percentage differences ($BIAS$), parameter also used in profiling comparisons between CALIOP and other ground-based lidar systems. All of them are examined as a function of the CALIPSO ground-track distance from the Belgrano II station. Results represent a relatively good agreement between both ground-based MPL-4 and space-borne CALIOP profiles of the volume linear depolarization ratio δ^V for PSC events, once the MPL-4 depolarization calibration parameters are applied. Discrepancies between CALIOP and MPL-4 profiles in vertical layering structure are enhanced from 20 km

up, likely due to a decrease of the signal-to-noise ratio (SNR) for both lidars systems at those altitudes. Regarding the results obtained from as the mean as the percentage differences found between MPL-4 and CALIOP δ^V profiles, a predominance of negative values is also observed, indicating a generalized underestimation of the MPL-4 depolarization as compared to that reported by CALIOP. However, absolute differences between those δ^V -profile datasets are no higher than a $10 \pm 11\%$ in average. Moreover, the degree of agreement between both lidar δ^V datasets is slightly dependent on the CALIPSO ground-track overpass distance from the Belgrano II station. That is, small discrepancies are found when CALIPSO ground-track distance is as close as rather far from the ground-based station. These results would indicate that MPL-4 depolarization observations would reflect relatively well the PSC field that CALIOP can detect at relatively large distances from the ground-based station. As a consequence, PSC properties can be statistically similar in average over large volumes, and hence the present weak disagreement found between both the lidar δ^V datasets can be likely dominated by small spatial PSC inhomogeneities along the CALIPSO separation from the station. This statement is based on the fact that Belgrano II is a station located well inside the stable Antarctic polar vortex, allowing determined thermodynamic conditions leading to a very low variability in the PSC field, and then in their properties.

1 Introduction

The polar stratosphere in both hemispheres is characterized by very low temperatures during winter leading to the formation of Polar Stratospheric Clouds (PSC). Heterogeneous chemical reactions occur on the surface of the PSCs, activating destructive compounds of ozone. Ozone destruction processes, through direct chlorine activation or indirectly through denitrification, are directly linked to the presence of a given type of PSC, and hence influencing the degree of ozone depletion (Solomon, 1999). In Polar regions PSCs start to form during winter at stratospheric temperatures below the condensation threshold of the Nitric Acid Trihydrate (NAT), depending on the water vapour and nitric acid partial pressure (Hanson and Mauersberger, 1988). PSCs are classified in three main groups depending on their composition, and then on their temperature formation threshold (i.e., see the review on PSC microphysics and chemistry by Lowe and MacKenzie, 2008, and references therein): type Ia (PSC-Ia) are nitric acid trihydrate (NAT) clouds formed above the frost point ($T_{NAT}=194$ K at 30 hPa), type Ib (PSC-Ib) are super-cooled ternary (H_2SO_4 , HNO_3 , H_2O) solution (STS, liquid particles) clouds, and type II (PSC-II) are water ice clouds ($T_{ice}=185$ K at 30 hPa).

Arctic temperatures are close to the threshold of PSC formation, hence both spatial and temporal PSC distributions present a high variability at daily and yearly scales. In contrast, PSC presence in the Antarctica is almost ubiquitous from the beginning of wintertime to early springtime, since Antarctic temperatures can reach rather lower values than those present in the Arctic (Parrondo et al., 2007), leading to higher occurrence of PSCs over the Antarctic continent.

Lidar measurements have been widely used for PSC classification on the basis of two lidar variables: the backscattering ratio (total backscatter-to-molecular coefficient ratio, R) and the volume linear depolarization ratio (δ^V). Indeed, due to the fact that non-spherical particles change the polarization state of the incident light, unlike spherical particles, both PSC-I (subtype Ia corresponding to solid particle NAT clouds and subtype Ib to liquid STS clouds) and PSC-II (ice clouds), as well as their mixtures, can be detected and identified by using lidar systems with depolarization measurement capabilities. Larger R and δ^V values

are found when PSC-IIIs occur, whereas PSC-Ia and -Ib, and their mixtures, present smaller R and δ^V values (e.g., Adriani et al., 2004; Maturilli et al., 2005; Pitts et al., 2009). PSC-IIs and PSC-IIIs have been detected in both hemispheres; however, the presence of ice clouds over the southern hemisphere is more frequent due to the fact that Antarctic temperatures usually can reach rather lower values than those in the Arctic, as result of a stronger and more stable Antarctic vortex during wintertime (Waugh and Polvani, 2010). However, although ice clouds can provide aerosol surface areas 100 times greater than those of liquid STS (PSC-Ib) or NAT (PSC-Ia) clouds for chlorine activation (e.g., Carslaw et al., 1998; Lowe and MacKenzie, 2008), thus favouring an enhancement of the ozone reduction as compared to that PSC-I clouds would present, their occurrence is rather lower than that for PSC-I (e.g., Adriani et al., 2004; Maturilli et al., 2005; Pitts et al. 2009). Therefore a more relevant role on ozone depletion is actually linked to liquid STS and NAT PSCs, which are the most important PSCs for chlorine activation and denitrification processes, respectively, involved in ozone destruction. Indeed, long-term PSC field monitoring together with PSC-type identification are critical in polar ozone depletion research, and hence directly linked to stratospheric temperature variability.

The Instituto Nacional de Técnica Aeroespacial (INTA, Spain) in collaboration with the Dirección Nacional del Antártico/Instituto Antártico Argentino (DNA/IAA, Argentina) have been performing an extensive program for stratospheric ozone monitoring and research in Antarctica. One of the objectives was the climatology of high clouds in coastal Antarctica to link two highly-correlated fields: PSC formation and ozone depletion (Solomon, 1999). In the frame of the international Polar Year (IPY), an improved version of the standard NASA/Micro Pulse Lidar (MPL v.4, MPL-4, Sigma Space Corp.), which includes a built-in depolarization measurement module, is currently used for PSC observations in the Antarctic Belgrano II station (Argentina, 77.9°S 34.6°W, 256 m a.s.l.) since 2009. The column of air above this station remains well inside the polar vortex during wintertime (Parrondo et al., 2007), as shown in Figure 1 in relation to the Antarctic polar vortex on 24 June 2010 as an example, providing thus an excellent location for PSC observations. Older versions than the MPL-4 have already been deployed in two other Antarctic stations (see Figure 1): Syowa (Japan, 69.0°S 39.5°E), where a PSC type-II single event was reported and attributed to low temperature fluctuations related to inertia gravity waves (Shibata et al., 2003), and remaining usually outside the polar vortex (see Figure 1); and South Pole/Amundsen-Scott (USA, 89.98°S 24.8°E), on the Antarctic Plateau (2835 m a.s.l.), where a 5-year data record was obtained by using the noisier MPL-3 with careful smoothing procedures (Campbell and Sassen, 2008). However, none of them includes polarization measuring capabilities similar to those of the MPL-4.

A good performance of the MPL-4 system for PSC detection was previously achieved in the Arctic (Córdoba-Jabonero et al., 2009), where depolarization data confirmed that all the PSC cases detected during the 2006-2007 winter were related only to PSC-I events, and no PSC-II occurrences with larger δ^V values were found. In particular, the MPL-4 performance for discriminating Type I (Ia and Ib) and Type II PSCs in relation with the δ^V parameter is still to be evaluated. This evaluation is focused on: 1) a more detailed estimation of δ^V from MPL-4 measurements, based on the analysis of the two-channel data, i.e. both parallel (p-) and perpendicular (s-) polarization MPL signals, and 2) the inter-comparison with a reliable lidar located not far from the Belgrano II station.

The space-borne lidar CALIOP on board the CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation, <http://www-calipso.larc.nasa.gov>) has provided valuable PSC information since 2006 at regional scales over both poles (Pitts et al., 2007, 2009, 2011). Therefore, MPL-4 depolarization retrievals of Antarctic PSCs are analyzed in

comparison with the PSC volume linear depolarization ratio δ^V reported from the space-borne lidar CALIOP to test the degree of agreement between both datasets.

Both of the lidar systems and the depolarization data processing are described in Section 2. Section 3 presents results and discussion together with the analysis procedures applied to both δ^V datasets, where three different approaches are considered for that comparison analysis. Finally, the main conclusions are summarized in Section 4.

2 Instrumentation and methods

2.1 Lidar systems

2.1.1 Ground-based lidar: MPL-4

The Micro Pulse Lidar version 4 (MPL-4, Sigma Space Corp.) is an improved version of the standard Micro Pulse Lidar version 3 (MPL-3, SES Inc.) in routine operation within the NASA/MPLNET (Micro-Pulse Lidar Network, <http://mplnet.gsfc.nasa.gov>) (Campbell et al., 2002). The MPL-4 system is small, easy-to-handle with high autonomy and operational in full-time continuous mode. MPL-4 is configured in a zenith, monostatic, coaxial alignment and is based on an eye-safe pulsed Nd:YLF laser emitting at 527 nm with a high repetition-rate (2500 Hz) and low-energy (10 μ J, max.). Its receiver system consists of a Maksutov-Cassegrain 18 cm-diameter telescope, a birefringent polarizer cell, and an avalanche photodiode detector. Backscattered signals are registered with a 1-min integration time and 75-m vertical resolution, commuting at each time the polarization module from parallel- to perpendicular-polarized detection (p- and s-channels, respectively). These 1-min signals registered in alternative mode for each p- and s-channel are hourly-averaged, providing 30-min averaged p- and s-signal profiles in one hour. These hourly-averaged profiles are usually analyzed to study the spatial and temporal variability of the PSC distribution. This MPL-4 configuration allows for probing the atmosphere up to 30 km with a sufficient signal-to-noise ratio (SNR). A full overlap is achieved at altitudes around 4 km up; however, the impact of the incomplete overlap effect on our retrievals is irrelevant because of: (i) its cancellation in the definition of δ^V (see Eq. (1) in Sect. 2.2), and (ii) PSCs frequently appear at higher altitudes where a full overlap is found.

2.1.2 Space-borne lidar: CALIOP

The CALIPSO satellite carries the space-borne lidar instrument CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) which provides horizontally- (along the CALIPSO ground-track) and vertically-resolved measurements for aerosol and clouds distributions at a global scale. CALIOP is based on diode-pumped Nd:YAG laser emitting linearly polarized pulses with a repetition rate of 20.16 Hz and a pulse length of ~ 20 ns, energy per pulse of 220 mJ at 1064 nm and ~ 110 mJ at 532 nm. Its receiver system consists of a 1m-diameter telescope which feeds a three-channel receiver measuring the backscattered intensity at 1064 nm and the two orthogonal polarization components at 532 nm, parallel (p) and perpendicular (s). A full description of the CALIOP system can be found in Winker et al. (2007) and Hunt et al. (2009). CALIOP provides data at 532 nm (the closest wavelength to that of the MPL-4 system) with a different vertical resolution as a function of altitude: 30 m at heights lower than 8.2 km, 60 m at 8.2-20.2 km, 180 m at 20.2-30.1 km, and 300 m at 30.1-40.0 km. In order to improve the SNR, a horizontal averaging over 5-km CALIPSO ground-track and a vertical 7-point adjacent averaging are applied. The latter data smoothing is achieved by using a sliding window of 7 points for averaging through the entire profile. As a result, the final vertical resolution of the CALIOP data keeps unchanged.

2.2 Depolarization data processing

The first lidar measurements of polarization properties were performed in the early 1970s (Schotland et al., 1971; Pal, 1973). It is well known that spherical particles do not change the polarization state of the incident light, while a partial depolarization component is introduced in the 180° backscattered signal after interacting with non-spherical particles. Several definitions are available in the lidar community to describe the depolarization phenomena caused by atmospheric constituents. A review of the most common parameters used in the lidar literature is given by Cairo et al. (1999). In our study, one of the most basic definitions is used, i.e., the volume linear depolarization ratio δ^V defined as follows:

$$\delta^V(z) = \frac{\beta^\perp(z)}{\beta^\parallel(z)} \quad (1)$$

where $\beta^\perp(z)$ and $\beta^\parallel(z)$ are the backscatter (particles plus molecules) coefficients for perpendicular- and parallel-polarization planes, respectively, and z is the height. In general, the term “particles” refers to both cloud and aerosol particles.

2.2.1 Ground-based depolarization measurements

From the practical point of view, the most general expression to calculate the volume linear depolarization ratio δ^V is:

$$\delta^V(z) = K \frac{P^\perp(z)}{P^\parallel(z)} + \chi \quad (2)$$

where $P^\perp(z)$ and $P^\parallel(z)$ are the s- and p-components of the measured MPL signals, respectively, once corrected for intrinsic instrumental factors (Campbell et al., 2002); K is a calibration constant that accounts for the differences of the receiver channel gains; and χ is a correction to account for any slight mismatch in the transmitter and detector polarization planes and any impurity of the laser polarization state (Sassen and Benson, 2001; Sassen, 2005). Because only a single detector is used in the MPL-4, the gain ratio is unity by definition and calibration requirements are vanished, however, at the expense of non-simultaneous measurements of the polarization components (see the review of existing techniques for estimating gain ratio in Álvarez et al. (2006)). Fortunately, the impact of non-simultaneity on our retrievals is negligible due to the rather small PSC variability during the integration time (1 minute) of each measurement. Thus, K is considered to be 1. The remaining correction term χ can be estimated by probing the δ^V values at middle and upper troposphere altitudes under both aerosol- and cloud-free conditions (calibration window). Optimal χ values are obtained by using fitting procedures with molecular backgrounds, considering a molecular volume linear depolarization ratio $\delta_{mol} = 0.0144$. Mean values of χ found for each year are presented in Table 1, showing a data dispersion of 5% among these three years (a total of 778 hourly-averaged profiles were selected for that purpose). Those χ values are then used for calibration of the MPL-4 depolarization measurements following Eq. (2).

Regarding time averaging procedures applied to the MPL-4 measurements, hourly-averaged MPL-4 δ^V profiles, as obtained from those 30-min averaged p- ($P^\parallel(z)$) and s- signal ($P^\perp(z)$) profiles in one hour (see Sect. 2.1.1 for details), are those used in the comparison with CALIOP data instead of instantaneous 1-min profiles (δ^V_{1-min}). As aforementioned (see Sect. 2.1.1), this improves the SNR of the lidar measurements at Belgrano II station. Indeed, the level of noise decreases as the time averaging increases, as shown in Figure 2 (for instance, data on 1 July 2009). δ^V variations depending on the time averaging (5-, 10-

, 15- and 30-min averaged profiles are shown in Figure 2) reveal that the vertical δ^V structure presents a clearly enhanced SNR when the time averaging is higher than 15 min. In particular, an additional PSC feature at around 25 km height can be identified with enough SNR only for 15-min and 30-min averaging of the MPL-4 data (see Figure 2).

Moreover, MPL-4 δ^V fluctuations along that hour are also studied by examining the differences between instantaneous 1-min ($\delta^V_{1-\text{min}}$) and hourly-averaged (δ^V) profiles within the same hour. Mean differences and their RMS values are shown in Figure 3 (for instance, data on 1 July 2009). A height-averaged value of -0.005 ± 0.013 is obtained for these mean differences, and their RMS values show that temporal δ^V fluctuations are lower than 0.05, 0.1 and 0.25 up to altitudes of 18 km, 23 km and 30 km height, respectively.

In summary, a high SNR is achieved for a time averaging of 30 minutes applied to MPL-4 p- and s- signal profiles (i.e., hourly-averaged δ^V profiles) in our comparison analysis, and temporal δ^V fluctuations per hour are lower than 0.1 up to altitudes where PSC features more frequently appear.

An example of this calibration procedure performed on 01 July 2009 is illustrated in Figure 4, where the volume linear depolarization ratio δ^V (left panel) together with the backscattering ratio R (centre panel) and the closest temperature profile provided from the local radiosounding (29 June 2009 at 11:00 UTC, right panel) are also represented. R , the total backscatter-to-molecular coefficient ratio (or the normalised R , $R_{\text{norm}} = 1 - 1/R$) usually used for PSC detection, is obtained by using a lidar ratio (extinction-to-backscatter coefficient ratio) of 30 sr in the Klett-Fernald inversion algorithm for backscatter coefficient retrieval (Klett, 1981; Fernald, 1984). A reference height is fixed when the condition of $R = 1$ ($R_{\text{norm}} = 0$, the strict case for an aerosol- and cloud- free molecular atmosphere) is found in the calibration window range considered. On this particular day, a mean value of $\chi = -0.055 \pm 0.009$ is computed from the 24 daily profiles by using a calibration window from 5 to 7 km a.g.l. (grey band in Figure 4). This value is similar to that reported for 2009 (see Table 1), which was applied to the 2009 winter dataset.

A similar procedure is applied to all other uncalibrated MPL-4 δ^V profiles. In particular, PSC depolarization features observed on 01 July 2009, once the δ^V profile is calibrated, present δ^V values larger than 0.2 at altitudes from 18.5 to 22 km and from 24 km height upwards, where a maximum $\delta^V = 0.5$ at 21-22 km height is identified (see Figure 4, left), and $R \cong 10$ ($R_{\text{norm}} \cong 0.9$) is found (see Figure 4, centre). This combination of R and δ^V values is typical for PSC-II events, as reported for several studies (e.g., Adriani et al. 2004; Maturilli et al. 2005; Pitts et al., 2009). This PSC-II feature is also confirmed by radiosonde data (see Figure 4, right), reporting temperatures at those altitudes between 19.5 and 26.5 km height lower than the threshold for PSC-II formation, i.e. where $T < T^{\text{PSC-II}}$ (see Figure 4, right, grey-shaded bands). These threshold temperatures for PSC-I ($T^{\text{PSC-I}}$, dashed line in Figure 4) formation have been calculated according to the parameterisations of Hanson and Mauersberger (1988) assuming 5-ppmv H_2O and 10-ppbv HNO_3 ; and for PSC-II ($T^{\text{PSC-II}}$, solid line in Figure 4) they were estimated from Marti and Mauersberger (1993) with the same H_2O amount. These proportions were obtained from Maturilli et al. (2005), where these values were reported as typical for this month.

However, $T^{\text{PSC-I}}$ and $T^{\text{PSC-II}}$ can vary depending on the real amounts of HNO_3 and H_2O . Therefore, additionally, as the local radiosounding water vapour profile is available, the

same calculations have been performed by using the in-situ H₂O concentration (radiosonde data) instead of a constant value. These new threshold temperatures for PSC-I and PSC-II formation are also shown in Figure 4 (thick dashed and solid lines, respectively), and differences are clearly found respect to those previously obtained. In particular, no PSC-II region ($T < T^{\text{PSC-II}}$) would be observed, differing the lidar results found as indicated by both the R and δ^V parameters. These discrepancies can be likely due to the no simultaneous lidar measurements with local available radiosounding data, at least for this day. Hence, a further study must be addressed on the relation between the PSC-type features and the stratospheric temperature variability, regarding the temperature thresholds for PSC formation. Nevertheless, this is out of the scope of this work.

2.2.2 Space-borne depolarization measurements

CALIPSO provides Level 1, Level 2 and Level 3 products. Level 1 products include lidar calibrated and geo-located profiles of attenuated-backscatter coefficient at 1064 nm, and total and perpendicular-polarized attenuated backscatter coefficient at 532 nm. Level 2 products include cloud layer, aerosol layer and aerosol profiles at different horizontal resolutions. Level 3 products are monthly-averaged profiles of aerosol optical properties on a uniform grid in the tropospheric region for altitudes below 12 km height. A detailed description of products Levels 1 and 2 can be seen in Powell et al. (2010). The Level 1 V3-01 (version 3.01, validated stage 1) attenuated-backscatter profile products at 532 nm (total and perpendicular-polarized) are used in this study. The attenuated-backscatter coefficient profile is defined as the volume backscatter coefficient β multiplied by the two-way atmospheric transmission T^2 (Hostetler et al., 2006). The total linear depolarization ratio δ_{total} is defined as:

$$\delta_{total}(z) = \frac{\beta^{\perp}(z)}{\beta^{total}(z)} \quad (3)$$

where $\beta^{total} = \beta^{\perp} + \beta^{\parallel}$, $\beta^{\perp}(z)$ and $\beta^{\parallel}(z)$ are the backscatter (particles and molecules) coefficients for s- and p-polarized components, respectively, and z is the range. Again, the term "particles" refers to both cloud and aerosol particles. For convenience, Eq. (3) can be multiplied by the term T^2 , allowing for expressing the total linear depolarization ratio δ_{total} in terms of attenuated-backscatter coefficients:

$$\delta_{total}(z) = \frac{\beta^{\perp}(z)}{\beta^{total}(z)} = \frac{\beta^{\perp}(z) \cdot T^2}{\beta^{total}(z) \cdot T^2} = \frac{\beta_{att}^{\perp}(z)}{\beta_{att}^{total}(z)} \quad (4)$$

where $\beta_{att}^{total} = \beta_{att}^{\perp} + \beta_{att}^{\parallel}$, and $\beta_{att}^{\perp}(z)$ and $\beta_{att}^{\parallel}(z)$ are the attenuated-backscatter coefficient for s- and p-polarized components, respectively. β_{att}^{\perp} and β_{att}^{total} are provided by the CALIPSO Level 1 products. Finally, the δ_{total} values are converted into δ^V values using the following relationship (Cairo et al., 1999):

$$\delta^V(z) = \frac{\delta_{total}(z)}{1 - \delta_{total}(z)} \quad (5)$$

Therefore, the volume linear depolarization ratio δ^V can be compared between both MPL-4 and CALIOP datasets.

However, despite smoothing and averaging procedures applied (see Sect. 2.1.2), high negative δ^V values inside the CALIOP dataset were observed, mainly at higher altitudes.

Therefore, in order to avoid the data contamination with such “unrealistic” values, an a-priori filtering of data is proposed. Restriction conditions to those δ^V values are based on the CALIPSO observations reported by Pitts et al. (2009), where most of aerosol depolarization ratio (δ_a) data were shown between -0.1 and 0.8, falling about 14% overall of CALIOP data outside these limits. Note that volume linear depolarization ratio δ^V data are presented here instead of those δ_a . Therefore, the corresponding restriction limits are calculated for δ^V by using the formulation reported by Cairo et al. (1999), linking both these magnitudes. Hence, δ^V can be expressed as a function of the molecular depolarization ratio δ_{mol} , the backscattering ratio R and δ_a , as follows:

$$\delta^V = \frac{\delta_a \times (R \times \delta_{mol} + R - 1) + \delta_{mol}}{\delta_a + (R - 1) \times \delta_{mol} + R} \quad (6)$$

where R values are varied between 1 and 30, δ_a between -0.1 and 0.8 (Pitts et al., 2009), and δ_{mol} takes these two values: 0.00366 (Cabannes scattering) and 0.0144 (total Rayleigh). Among all the possible combinations, the minimal (-0.1) and maximal (0.8) values are computed. Hence, these limits will act as a conservative restriction range for the overall of δ^V values in both lidar datasets, which are the same presented for δ_a in Pitts et al. (2009). In fact, about $14 \pm 11\%$ and $5 \pm 3\%$ overall of CALIOP and MPL-4 data, respectively, falling outside of these limits, have been disregarded for the comparison analysis performed between both lidar datasets.

3 Results and discussion

3.1 Lidar datasets

PSC observations have been performed at the Belgrano II station since 2009 to the present. MPL-4 measurements for the 2009-2011 Antarctic winters, from May to September, are used for this study. Lidar datasets are compared under the following conditions: coincident profiles in time, with simultaneous measurements carried out at time scales lower than two hours around the CALIPSO overpass, and in space, with CALIPSO ground-track separations from the station closer in 55 km distance. During those winter times, a total of 189 CALIPSO overpasses nearby the Belgrano II station were carried out within fewer 55 km distance. Among them, 104 overpasses are coincident events with MPL-4 measurements reporting PSC detection and 48 of them are simultaneously available for comparison. Moreover, 35 more lidar profiles are analyzed in order to examine the influence on spatial scales when data from rather large CALIPSO overpass distances are included in that lidar comparison. Those additional 35 profiles correspond to separations between 70 and 100 km. In general, four predominant distance-ranges are observed: 0-10 km, 20-30 km, 45-55 km and 70-90 km. All these PSC cases, listed by the CALIPSO ground-track distance from the Belgrano II station, are shown in Table 2.

A height-interval from 5 to 30 km is selected for the comparison between lidar profiles. A delineating altitude of 10 km has been conservatively established as the lower limit for the unambiguous presence of PSCs, distinguishing them from other upper tropospheric clouds (mainly Cirrus clouds). The lower limit of 10 km chosen in this work for PSC detection is based on the fact that the tropopause is not clearly delineated by the temperature profile during wintertime in deep Antarctica (Rubin, 1953). Indeed, a traditional tropopause height, denoted by rapidly increasing static stability above it, can be approximated from December

through March in Belgrano II station sounding data around 9 km. During winter months, however, temperatures decrease with height to nearly 23 km. Dynamic coupling between the troposphere and stratosphere is more likely in such conditions. The region from 8 to 10 km is considered a transitional zone, where cloud type cannot be established with any certainty. Although our study is restricted to PSC formation altitudes, the 5-10 km height interval is also considered for contrast as a PSC-free region. Heights above 30 km are disregarded due to a decreasing of the MPL-4 SNR. Finally, along the overall height-interval selected, every 0.5-km layer is averaged for comparing the MPL-4 and CALIOP datasets. Note that negative δ^V values are not disregarded during the smoothing process of CALIOP data profiles (see Sects. 2.1 and 2.2). Only δ^V values falling outside the (-0.1, 0.8) interval are ignored when this 0.5-km vertical averaging is applied to both lidar profiles.

3.2 Comparison analysis

Three different approaches are considered for the comparison between both lidar δ^V datasets in order to test the degree of agreement as a function of the CALIPSO ground-track separation from Belgrano II station:

- 1) the correlation coefficient (CC), as a measure of the relationship between both PSC vertical layering structures;
- 2) the mean differences, $\Delta(z) = \delta^{MPL}(z) - \delta^{CAL}(z)$, between both MPL-4 and CALIOP δ^V profiles, together with their root mean square (RMS) values; and
- 3) the percentage difference, $BIAS(z)$, since this parameter is also used in profiling comparisons between CALIOP and other ground-based lidar systems (Mamouri et al., 2009; Mona et al., 2009), and defined as:

$$BIAS(z) = \frac{100 \times [\delta^{MPL}(z) - \delta^{CAL}(z)]}{\delta^{CAL}(z)} \quad (7)$$

For all these three approaches, $\delta^{MPL}(z)$ and $\delta^{CAL}(z)$ are the 0.5-km averaged volume linear depolarization ratio δ^V profiles for MPL-4 and CALIOP, respectively.

3.2.1 Comparison analysis I: Linear regression and correlation coefficient (CC)

The correlation coefficient (CC) between the 0.5-km averaged MPL-4 and CALIOP profiles, was calculated for 83 cases over the total altitude range (see Table 2). Among these cases, 12, 24, 12 and 35 of them, respectively, correspond to the four predominant CALIPSO distance-ranges: < 10 km, 20-30 km, 45-55 km and 70-90 km, denoted as D1, D2, D3 and D4, respectively. This analysis is also performed as a function of different altitude intervals in order to examine similarities and/or discrepancies between different PSC layers as observed by both MPL-4 and CALIOP in each case. This procedure can reveal the degree of agreement between both vertical layering structures as the height increases. Five height intervals for correlation fitting are selected: 5-10 km, 5-15 km, 5-20 km, 5-25 km and 5-30 km. Table 2 summarizes the results. They are based on the significance of these correlation coefficients obtained. Therefore, a value of CC = 0.5 is considered statistically significant depending on the number of data points deemed into the calculation, where a p-value less than the chosen significance level α of 0.05 must be reported. In particular, for the case of a height interval of 5-15 km with a smaller number of data points than that for other height intervals considered in this study, a test statistic $t = 2.09$ and CC = 0.46 are obtained reporting p-values lower than 0.05. Consequently, for other height intervals with larger number of data points, p-values must be also lower than α , and therefore those obtained CC can be regarded as statistically significant in this study.

A few dissimilarities can be observed between each year, but in general the correlation is better with a higher number of cases presenting CC values higher than 0.5 (referred as the frequency in %) for the fitting height interval of 5-20 km, being slightly lower for those four others. In particular, among those 83 MPL-4/CALIOP profile coincidences examined in total during the 2009-2011 winter periods, 49 out of them (59%) present $CC > 0.5$, at least in that fitting 5-20 km height interval (see Table 2). In fact, the lowest value (35%) of that frequency (cases with $CC > 0.5$) is found when the overall height interval of 5-30 km is considered for correlation fitting. This result indicates that discrepancies between CALIOP and MPL-4 in vertical layering structure are enhanced from 20 km up, likely due to a decrease of the SNR for both lidars systems at those altitudes. However, PSC formation occurs indeed more frequently at altitudes lower than those heights. Therefore, these results represent a relatively good agreement for the PSC volume linear depolarization ratio δ^V between MPL-4 and CALIOP profiles.

Regarding the dependence of the correlation between lidar datasets on the distance of the CALIPSO overpass from Belgrano II station, that frequency (cases with $CC > 0.5$) for all three 2009-2011 winters, and individually for 2009, 2010 and 2011 wintertime periods is shown in Figure 5 as a function of the four predominant distance-ranges. Results confirm a weak dependence on that distance for all the years, in general. Indeed, no large differences are unexpectedly observed in the correlation between both vertical structures of depolarization when the CALIPSO ground-track is just at a few kilometres from the station (at 0-10 km far, D1), showing a frequency of 75% with $CC > 0.5$, respect to rather large separations (at 70-90 km far, D4) with a lower frequency of 49% (see Figure 5). Actually, these results indicate that MPL-4 depolarization observations would reflect relatively well the PSC field that CALIOP can detect at large distances from the ground-based station.

3.2.2 Comparison analysis II: Mean differences (Δ) and their root mean square (RMS) values

In order to complement that previous comparison analysis the differences, $\Delta = \delta^{MPL} - \delta^{CAL}$, between MPL-4 and CALIOP 0.5-km averaged profiles, δ^{MPL} and δ^{CAL} , respectively, are calculated for the four predominant CALIPSO distance-ranges: < 10 km (D1), 20-30 km (D2), 45-55 km (D3) and 70-90 km (D4). Mean differences between those profiles fulfilling that their corresponding CALIPSO separation is within a given distance-range, $\Delta_d(z)$, together with their root mean square (RMS) values are shown in Figure 6. In general, slightly discrepancies are observed for as the mean differences as the RMS values depending on the CALIPSO separation (see Figure 6). Results indicate that $\Delta_d(z)$ are mostly negatives with a tendency to values close to zero or even positives at the 20-25 km height interval, reaching larger negatives values with higher data dispersion at altitudes from 25 km height up. In addition, RMS values are no larger than 0.3 in overall, being lower than 0.15 at altitudes up to 20 km, in general for all the distance-ranges. This result shows that only a slightly data dispersion is found at high altitudes. Variations of the RMS values are rather small depending on the distance-range.

Moreover, these Δ_d profiles have been averaged, for simplicity, in 5-km layers thick, Δ_d^{5-km} , in order to examine the behaviour of these mean differences Δ_d in relation with the CALIPSO ground-track distances (data also shown in Figure 6). As previously, slightly discrepancies are observed at all the distance-ranges, and a predominance of negative values is found among all the 5-km averaged layers. Similar height-averaged values are obtained at altitudes up to 20 km height, being their mean value -0.011 ± 0.006 within all the distance-ranges. In addition, absolute Δ_d^{5-km} values are no higher than 0.1, but with a

larger data dispersion at altitudes higher than 20 km. However, Δ_d (in absolute value) unexpectedly seems to slightly decrease as distance increases up to 70-90 km far.

3.2.3 Comparison analysis III: Percentage differences ($BIAS$)

This procedure has previously been used for ground-based lidar profiles in comparison with CALIOP data in the case of the tropospheric attenuated backscatter coefficients at middle-latitude regions (Mamouri et al., 2009; Mona et al., 2009). Hence, it is applied for the PSC volume linear depolarization ratio δ^V for the first time in this work. Both 0.5-km averaged MPL-4 and CALIOP datasets are used to calculate the $BIAS$ (see Eq. (7)) as an altitude-dependent parameter, obtained for comparison analysis between both lidar datasets.

Due to the large $BIAS$ data dispersion obtained in general, a determined constraint condition is applied to the $BIAS$ profiles: only $BIAS$ values within a given interval ($-50\% < BIAS < +50\%$) are regarded. This constrained selection is done to evaluate $BIAS$ values with a given realistic significance, ignoring values rather higher than $\pm 50\%$. From a statistical point of view, a height-averaged $BIAS$, $BIAS_z$, is calculated from the 'constrained' profiles, also considering the number of data points in each profile fulfilling that condition. 'Constrained' $BIAS_z$ values for all the cases are shown in Table 2 together with their standard deviation (SD), including the percentage of number of data points fulfilling that condition (N^{BIAS} , in %) for each case.

Results show that δ^{MPL} is in general underestimated respect to CALIOP δ^{CAL} values with a clear predominance of negative $BIAS_z$ values: 74% out of those 83 cases available for comparison (see Table 2). In addition, the number of data points fulfilling the constraint ($-50\% < BIAS < +50\%$) condition in average $\langle N^{BIAS} \rangle$ is $25 \pm 13\%$, thus showing a large data dispersion with $BIAS$ values higher than $\pm 50\%$. Despite this result, mean absolute differences between δ^{MPL} and δ^{CAL} are no higher than a $10 \pm 7\%$ respect to CALIOP values with a median value of 9%. It is worth mentioning that the calculated percentage differences, $BIAS$, are rather large at PSC-free altitudes, mostly in the 5-10 km height interval, due to relatively low δ^{CAL} values close to the molecular one in that region, and as a consequence $BIAS$ considerably increases following Eq. (7). Additional calculations have been performed and $BIAS$ values have been averaged within an smaller height-range from 10 to 20 km height in order to avoid that particular PSC-free region and also altitudes higher than 20 km with a larger data dispersion (as shown in the previous Sect. 3.2.2). In this case, a 71% out of cases present negatives $BIAS_z$ and $\langle N^{BIAS} \rangle = 34 \pm 23\%$, being the mean absolute differences between δ^{MPL} and δ^{CAL} no higher than a $14 \pm 11\%$ respect to CALIOP values with a median value of 11%. These results are in agreement with those values obtained for the mean differences, as expected.

Moreover, likewise the two previous approaches (see Sects. 3.2.1 and 3.2.2), the $BIAS$ dependence on the CALIPSO ground-track distance from Belgrano II station is also evaluated. $BIAS$ values obtained for each year as a function of those four main distance-ranges (D1: < 10 km, D2: 20-30 km, D3: 45-55 km, and D4: 70-90 km) are shown in Figure 7. A predominance of negative values is observed, showing the previously commented δ^{MPL} underestimation respect to δ^{CAL} . In particular, averaged $BIAS$ on those four distance-ranges, $BIAS_d$, are obtained: $-10 \pm 8\%$, $-6 \pm 8\%$, $-9 \pm 9\%$ and $-8 \pm 11\%$, respectively (see Figure 7). These similar values indicate that the $BIAS$ is independent on CALIPSO separation with practically no increase in those differences between both lidar

datasets as the CALIPSO overpass is far from the station. However, a slightly increasing dispersion of those *BIAS* values can be observed as the distance increases.

3.2.4 Comparison analysis of particular cases

A few cases of particular PSC events are described in more detail below. They depict different comparison features as a function of the CALIPSO ground-track distance from the Belgrano II station (within the four predominant distance-ranges).

Figures 8-11 represents simultaneous 0.5-km averaged profiles of the volume linear depolarization ratio δ^V for MPL-4 (δ^{MPL} , full circles) and CALIOP (δ^{CAL} , open triangles), together with the corresponding differences, $\Delta = \delta^{MPL} - \delta^{CAL}$ (as shown by black stars), found between both lidar δ^V profiles depending on the four predominant (D1 to D4) CALIPSO distance-ranges, respectively. PSC events correspond to those cases with high/moderate and low CC values (left and right panels in each figure, respectively) (see Sect. 3.1 and Table 2).

In particular, PSC event observed on 01 July 2009 when the CALIPSO ground-track overpass is at 3.6 km distance from Belgrano II station (see Figure 8, left) present a high correlation coefficient (0.91) between both lidar δ^V datasets for a fitting height-range of 5-25 km, decreasing as height increases (see Table 2). δ^V values higher than 0.2 are obtained from around 18.5 up to 22.5 km height; this result is also in agreement with those stratospheric temperatures lower than T^{PSC-II} present in the same height-range (see Figure 4 and Sect. 2.2.1 for more detail). However, PSC depolarization features observed on 25 September 2009 (see Figure 8, right) by both lidar systems present high uncorrelated vertical layering structures as indicated by a low CC value (0.12), despite the CALIPSO separation is within the same distance-range as the former case (see Table 2). In addition, absolute differences are mostly lower 0.15 on 01 July 2009 (case with high CC) as compared to those found on 25 September 2009 (low CC) with a higher dispersion of those differences. Height-averaged values of those MPL-4/CALIOP differences $\langle \Delta \rangle$ are also calculated: -0.02 ± 0.12 and -0.16 ± 0.21 are found on 01 July 2009 and 25 September 2009, respectively. As previously stated, these negatives differences indicate an underestimation of the MPL-4 depolarization respect to that for CALIOP, being this underestimation more severe for that case with a low correlation between both lidar datasets, i.e. on 25 September 2009 (see Figure 8).

PSC events observed on 23 August 2011 and 03 July 2010 are the cases presenting a high (CC = 0.75) and low (CC = 0.20) correlation, respectively, within a fitting height-range of 5-25 km between MPL-4 and CALIPSO δ^V -profiles (see Figure 9, left and right, respectively) when CALIPSO ground-track overpass is within the D2 distance-range (20-30 km) from the ground-based station. As previously, the correlation decreases as height increases (see Table 2), with a more severe decreasing on 03 July 2010 (low CC). In particular, $\delta^V > 0.2$ values are mainly found from 9 up to 16.5 km height (a Cirrus/PSC overlapping at 9-10 km height-range is present) and also at altitudes from 19 up to 24 km height on 23 August 2011 (see Figure 9, left), in contrast with those found on 03 July 2010 (see Figure 9, right) with the overall $\delta^V < 0.2$, except for two spike-like PSC features at 15.5 km and 18 km height reported by CALIOP data. In both cases, absolute differences are mostly lower 0.15 up to 25 km height. Height-averaged values $\langle \Delta \rangle = -0.06 \pm 0.16$ and -0.03 ± 0.16 are found on 23 August 2011 and 03 July 2010, respectively. That previously observed MPL-4 δ^V underestimation is also found, but with similar values for both cases (see Figure 9).

Particular cases when CALIPSO ground-track separation is fairly far from the Belgrano II station (within the D3 distance-range: 45-55 km) are selected, corresponding to PSC events observed on 22 August 2011 and 12 July 2010. These two cases present a moderate (0.60) and low (0.31) correlation coefficients, respectively, within a fitting height-range of 5-25 km (see Table 2) between MPL-4 and CALIPSO δ^V -profiles (see Figure 10, left and right, respectively). A decreasing of the degree of correlation is also observed, as in those previous cases, when a larger height interval is examined. In particular, vertical δ^V structures are well correlated up to altitudes lower than 25 km height in both cases, but a more severe disagreement is observed for the low CC case (12 July 2010, see Figure 10, right), as expected. In addition, $\delta^V > 0.2$ values are mainly found from 11 up to 23 km height on 22 August 2011 (see Figure 10, left), being mostly close to 0.2 on 12 July 2010 (see Figure 10, right), except for several spike-like PSC features centred at 19.5 km, 22.5 km and 27 km height as observed by CALIOP. In both cases, absolute differences are mostly lower 0.15 up to 20 km height. Height-averaged values $\langle \Delta \rangle = -0.02 \pm 0.16$ and -0.06 ± 0.23 are found on 22 August 2011 and 12 July 2010, respectively. As aforementioned, a MPL-4 δ^V underestimation is also obtained for both cases, but presenting a higher dispersion for the lower correlated case (see Figure 10).

Figure 11 shows those PSC features observed on 31 July 2010 (left) and 22 June 2011 (right) presenting high (0.86) and low (0.27) correlation coefficients, respectively, within a fitting height-range of 5-25 km (see Table 2). These cases correspond to rather far CALIPSO ground-track separations from the Belgrano II station (D4 distance-range: 70-90 km). In general, the degree of correlation is also reduced at altitude-ranges higher than 25 km up, as previously observed, being more notably observed for the low CC case (22 June 2011, see Figure 11, right), where the correlation drops from 20 km height up. In particular, $\delta^V > 0.2$ values are mainly found from around 9 up to 20 km height on 31 July 2010 (see Figure 11, left), where a Cirrus/PSC overlapping at 8-10 km height-range is present. Besides, a spike-like PSC feature centred at around 26 km height can be only observed by CALIOP data with δ^V values rather higher than 0.2. δ^V values are mostly around 0.2 on 22 June 2011 (see Figure 11, right), presenting also two spike-like PSC features with $\delta^V > 0.2$ centred at around 15 km and 17.5 km height by CALIOP data, and one more at around 25 km height detected by both lidars. Absolute differences are mainly lower 0.15 up to 25 km height on the high CC case, but not for the low CC one with a higher number of points outside this ± 0.15 range. $\langle \Delta \rangle$ values are similar to those obtained for the previous cases corresponding to CALIPSO smaller separations. In particular, -0.03 ± 0.21 and -0.08 ± 0.18 are found on 31 July 2010 and 22 June 2011, respectively. As for the other cases, a MPL-4 δ^V underestimation is also obtained for both these ones (see Figure 11).

These results indicate that particular PSC features are observed when individual simultaneous cases are examined. However, in general, a high degree of correlation between the vertical δ^V structures from MPL-4 and CALIOP datasets is found for those cases presenting large δ^V values. In addition, the correlation decreases at altitudes higher than 25 km height. Moreover, absolute differences between both lidar datasets are lower than 0.15 at altitudes up to 20 km height, in general, for the high CC cases; higher data dispersion with values outside this range is found for those other low CC cases. In particular, height-averaged Δ values are similar independently on the CALIPSO ground-track distance from the station. A generalized underestimation of the MPL-4 depolarization respect to that for CALIOP is obtained, as expected (see previous Sects. 3.2.1-3.2.3).

4. Summary and conclusions

This study appears as a first application of the lidar depolarization technique to Antarctic PSC detection and identification by using an improved version (MPL-4) of the standard NASA/Micro Pulse Lidar. In particular, this work represents a significant advance on PSC-type discrimination studies by using MPL-4 δ^V data.

Calibration parameters for suitable MPL-4 δ^V retrievals have been calculated from MPL-4 measurements. These calibrated δ^V profiles have been compared with coincident CALIPSO data as a reference during 2009-2011 austral winters, from May to September periods, over Belgrano II station (Argentina, 77.9°S 34.6°W, 256 m a.s.l.). Coincident observations are referred to simultaneous measurements between both lidar systems within two hours around the closest CALIPSO overpass to the Belgrano II station. That is, four predominant distance-ranges between the CALIPSO ground-track and the station have been selected in order to examine the dependence of the degree of agreement between both lidar δ^V -profile datasets on that distance from the Belgrano II station. Three analysis procedures for δ^V -profile comparison between both lidar datasets have been presented.

Correlation analysis shows that a 59% out of all the comparison cases present CC values higher than 0.5, at least in the altitude-range from 5 to 20 km height. This frequency (number of cases with CC > 0.5) decreases down to 35% when the overall height interval of 5-30 km is considered for correlation fitting, indicating that discrepancies between CALIOP and MPL-4 in vertical layering structure are enhanced from 20 km up, likely due to a decrease of the SNR for both lidars systems at those altitudes. However, PSC formation occurs indeed more frequently at altitudes lower than those heights. Hence, a relatively good agreement is found between both ground-based MPL-4 and space-borne CALIOP profiles of the volume linear depolarization ratio δ^V for PSC events, once the MPL-4 depolarization calibration parameters are applied. Moreover, no large differences are unexpectedly observed in the correlation analysis between both vertical depolarization structures when the CALIPSO ground-track is just at a few kilometres from the station, showing a frequency of 75% with CC > 0.5, respect to rather large separations with a lower frequency (49%).

Regarding the differences between both lidar δ^V -profile datasets, two related variables are analysed: the mean differences, Δ , within a given CALIPSO distance-range; and the percentage differences, *BIAS* (see Eq. (7)), since, despite its relation with the former Δ , *BIAS* is a parameter also used in profiling comparisons between CALIOP and other ground-based lidar systems.

Slightly discrepancies are observed for the mean differences between those δ^V profiles depending on the CALIPSO separation. Mean differences Δ are mostly negatives in the overall height interval, with higher data dispersion at altitudes from 25 km height up. In addition, absolute Δ values are no higher than 0.1 in average, being lower than 0.05 at altitudes up to 25 km height, with RMS values no larger than 0.3 in overall for all the distance-ranges. The other comparison parameter, *BIAS*, seems to be a less robust indicator of the degree of agreement for lidar δ^V datasets, since rather less than a half of the cases fulfil the selected constraint condition ($-50\% < BIAS < +50\%$), showing a large data dispersion with absolute percentage differences higher than 50%. This is related to the fact that *BIAS* is rather large when relatively low δ^{CAL} values are present within a given CALIPSO profile (as, for instance, those are at PSC-free altitudes), and as a consequence, *BIAS* considerably increases (see Eq. (7)), exceeding the percentage values of that

supposed constraint condition. In addition, that previously observed predominance of negative values is also found, indicating a generalized δ^{MPL} underestimation with respect to CALIOP data. However, absolute differences between δ^{MPL} and δ^{CAL} are no higher than a $10 \pm 11\%$ in average as compared to CALIOP values. As expected, these results are in agreement with those obtained for the mean differences.

Moreover, the degree of agreement between both lidar δ^V datasets is moderately dependent on the CALIPSO ground-track overpass distance from the Belgrano II station, as shown by the results obtained in each one of the comparison analyses carried out: the vertical correlation (CC), and both the mean (Δ) and percentage (*BIAS*) differences. That is, no large discrepancies are found when CALIPSO ground-track distance is as close as < 10 km far as well as rather far (at 70-90 km) from the Belgrano II station.

Actually, these results indicate that MPL-4 depolarization observations would reflect relatively well the PSC field that CALIOP can detect at large distances from the ground-based station. As a consequence, PSC properties would be statistically similar in average over large volumes, and hence the present disagreement found between both the lidar δ^V datasets would be likely related to be dominated by small spatial PSC inhomogeneities along the CALIPSO separation from the station. This statement is based on the fact that Belgrano II is a station located well inside the polar vortex during almost all the wintertime period. Indeed, the Antarctic polar vortex is quite stable to allow determined thermodynamic conditions leading to a very low variability of the PSC field, and then of their properties.

Therefore, a further study, out of the scope of this work, on the correlation between the PSC features and the variability of both the polar potential vorticity and the stratospheric temperature fields would provide an understanding on what the observed discrepancies are really based on. In addition, a detailed going-on 3-year statistical analysis of PSC occurrence over Belgrano II station in terms of both the backscattering, R , and volume linear depolarization, δ^V , ratios from MPL-4 measurements would complete these studies. This will include a PSC-type discrimination assessment over Belgrano II station, a station well inside the Antarctic polar vortex.

Finally, it is worth mentioning these results are useful for PSC detection and classification in both Polar Regions by using this kind of micro pulse lidar that operates in full-time continuous mode, providing a more complete evolution of the PSC field on a daily basis.

Acknowledgements

This work has been supported by the Spanish Ministry for Science and Innovation (MICINN) under grant CGL2010-20353 (project VIOLIN), the Spanish Ministry of Education (MEC) fellowship EX2009-0700 and the Andalusian Regional Government project P10-RNM-6299. CALIPSO data were obtained from the NASA Langley Research Center Atmospheric Science Data Center. Authors specially thank the DNA/IAA teams at Belgrano II station for their valuable assistance and support in the lidar and radiosounding measurements.

References

Adriani, A., Massoli, P., Di Donfrancesco, G., Cairo, F., Moriconi, M.L., and Snels, M.: Climatology of polar stratospheric clouds based on lidar observations from 1993 to 2001 over McMurdo station, Antarctica, *J. Geophys. Res.*, 109, D24211, doi:10.1029/2004JD004800, 2004.

Álvarez, J.M., Vaughan, M.A., Hostetler, C.A., Hunt, W.H., and Winker, D.M.: Calibration technique for polarization-sensitive lidars, *J. Atmos. Ocean. Technol.*, 23, 683-699, 2006.

Cairo, F., Di Donfrancesco, G., Adriani, A., Pulvirenti, L., and Fierli, F.: Comparison of various depolarization parameters measured by lidar, *Appl. Opt.*, 38, 4425-4432, 1999.

Campbell, J.R., and Sassen, K.: Polar Stratospheric Clouds at the South Pole from Five Years of Continuous Lidar Data: Macrophysical, Optical and Thermodynamic Properties, *J. Geophys. Res.*, 113, D20204, doi:10.1029/2007JD009680, 2008.

Campbell, J.R., Hlavka, D.L., Welton, E.J., Flynn, C.J., Turner, D.D., Spinhirne, J.D., Scott, V.S., and Hwang, I.H.: Full-time, eye-safe cloud and aerosol lidar observation at Atmospheric Radiation Measurement program sites: Instruments and data processing, *J. Atmos. Ocean. Technol.*, 19, 431-442, 2002.

Carslaw, K.S., Wirth, M., Tsias, A., Luo, B.P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J.T., Reimer, E., Peter, T.: Increased stratospheric ozone depletion due to mountain-induced atmospheric waves, *Nature*, 391, 675-678, 1998.

Córdoba-Jabonero, C., Gil, M., Yela, M., Maturilli, M., and Neuber, R.: Polar Stratospheric Cloud observations in the 2006/07 Arctic winter by using an improved Micro Pulse Lidar, *J. Atmos. Ocean. Technol.*, 26, 2136-2148, 2009.

Fernald, F.G.: Analysis of atmospheric lidar observations: some comments, *Appl. Optics*, 23, 652-653, 1984.

Hanson, D. R., and Mauersberger K.: Laboratory studies of the nitric acid trihydrate: implications for the south polar stratosphere, *Geophys. Res. Lett.*, 15, 855-858, 1988.

Hostetler, C. A., Liu, Z., Reagan, J., Vaughan, M., Winker, D., Osborn, M., Hunt, W. H., Powell, K. A., and Treppe, C.: CALIOP Algorithm Theoretical Basis Document, Calibration and level 1 Data Products, PC-SCI-201, NASA Langley Research Center, Hampton, VA 23681, USA, 2006.

Hunt, W.H., Winker, D.M., Vaughan, M.A., Powell, K.A., Lucker, P.L., and Weimer, C.: CALIPSO Lidar Description and Performance Assessment, *J. Atmos. Ocean. Technol.*, 26, 1214-1228, 2009.

Innis, J.L., and Klekociuk, A.R.: Planetary wave and gravity wave influence on the occurrence of polar stratospheric clouds over Davis Station, Antarctica, seen in lidar and radiosonde observations, *J. Geophys. Res.*, 111, D22102, doi:10.1029/2006JD007629, 2006.

Klett, J.D.: Stable analytic inversion solution for processing lidar returns, *Appl. Optics*, 20, 211-220, 1981.

Lowe, D., and MacKenzie, A. R.: Polar stratospheric cloud microphysics and chemistry, *J. Atmos. Solar-Terr. Phys.*, 70, 13-40, 2008.

Mamouri, R.E., Amiridis, V., Papayannis, A., Giannakaki, E., Tsaknakis, G., and Balis, D.S.: Validation of CALIPSO space-borne-derived attenuated backscatter coefficient profiles using a ground-based lidar in Athens, Greece, *Atmos. Meas. Tech.*, 2, 513-522, 2009.

Marti, J., and Mauersberger, K.: A survey and new measurements of ice vapor pressure at temperatures between 170 K and 250 K, *Geophys. Res. Lett.*, 20, 363-366, 1993.

Maturilli, M. , Neuber, R. , Massoli, P. , Cairo, F. , Adriani, A. , Moriconi, M.L., and Di Donfrancesco, G.: Differences in Arctic and Antarctic PSC occurrence as observed by lidar in Ny-Alesund [79°N 12°E] and McMurdo [78°S 16.7°E], *Atmos. Chem. Phys.*, 5, 2081-2090, 2005.

- Mona, L., Pappalardo, G., Amodeo, A., D'Amico, G., Madonna, F., Boselli, A., Giunta, A., Russo, F., and Cuomo, V.: One year of CNR-IMAA multi-wavelength Raman lidar measurements in coincidence with CALIPSO overpasses: Level 1 products comparison, *Atmos. Chem. Phys.*, 9, 7213–7228, 2009.
- Pal, S.R., and Carswell, A.I.: Polarization properties of lidar backscattering from clouds, *Appl. Opt.*, 12, 1530–1535, 1973.
- Parrondo, M.C., Yela, M., Gil, M., Von der Gathen, P., and Ochoa, H.: Mid-winter lower stratosphere temperatures in the Antarctic vortex: comparison between observations and ECMWF operational model, *Atmos. Chem. Phys.*, 7, 1–7, 2007.
- Pitts, M.C., Thomason, L.W., Poole, L.R., and Winker, D.M.: Characterization of Polar Stratospheric Clouds with spaceborne lidar: CALIPSO and the 2006 Antarctic season, *Atmos. Chem. Phys.*, 7, 5207–5228, 2007.
- Pitts, M.C., Poole, L.R., and Thomason, L.W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, 2009.
- Pitts, M.C., Poole, L.R., Dörnbrack, A., and Thomason, L.W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, 2011.
- Powell, K., Vaughan, M., Winker, D., Lee, K.-P., Pitts, M., Trepte, C., Detweiler, P., Hunt, W., Lambeth, J., Lucker, P., Murray, T., Hagolle, O., Lifermann, A., Faivre, M., Garnier, A., Pelon, J.: Cloud – Aerosol LIDAR Infrared Pathfinder Satellite Observations (CALIPSO), Data Management System, Data Products Catalog, Document No: PC-SCI-503, Release 3.2, August 2010, NASA Langley Research Center, Hampton, Virginia, USA, 2010.
- Rubin, M. J.: Seasonal variations of the Antarctic tropopause, *Met.*, 10, 127–134, 1953.
- Santacesaria, V., MacKenzie, A.R., and Stefanutti, L.: A climatological study of polar stratospheric clouds (1989–1997) from LIDAR measurements over Dumont d'Urville (Antarctica), *Tellus*, 53B, 306–321, 2001.
- Sassen, K., and Benson, S.: A midlatitude cirrus cloud climatology from the Facility for Atmospheric Remote Sensing. Part II: Microphysical properties derived from lidar depolarization, *J. Atmos. Sci.*, 58, 2103–2112, 2001.
- Sassen, K.: Polarization in Lidar, in *Lidar: Range-Resolved Optical Remote Sensing of the Atmosphere*, Ed. C. Weitkamp, Optical Sci., Springer Ser., Singapore, 2005.
- Schotland, R.M., Sassen, K., and Stone, R.: Observations by lidar of linear depolarization ratios by hydrometeors, *J. Appl. Meteorol.*, 10, 1011–1017, 1971.
- Shibata, T., Sato, K., Kobayashi, H., Yabuki, M., and Shiobara, M.: Antarctic polar stratospheric clouds under temperature perturbation by nonorographic inertia gravity waves observed by micropulse lidar at Syowa Station, *J. Geophys. Res.*, 108, D3, 4105, doi:10.1029/2002JD002713, 2003.
- Solomon, S.: Stratospheric ozone depletion: a review of concepts and history, *Rev. Geophys.*, 37, 275–316, 1999.
- Toon, O.B., Browell, E.V., Kinne, S., and Jordan, J.: An analysis of lidar observations of polar stratospheric clouds, *Geophys. Res. Lett.*, 17, 393–396, 1990.
- Waugh, D.W., and Polvani, L.M.: Stratospheric polar vortices, in *The Stratosphere: Dynamics, Chemistry, and Transport*, Geophys. Monogr. Ser., 190, , edited by L.M. Polvani, A. H. Sobel, and D. W. Waugh, pp. 43–57, AGU, Washington, D. C., 2010.

Winker, D.M., Hunt, W.H., and McGill, M.J.: Initial performance assessment of CALIOP, *Geophys. Res. Lett.*, 34, L19803, doi:10.1029/2007GL030135, 2007.

Table 1. Mean χ values together with their SD (%SD) and the number of profiles used.

| Year | χ | SD (%SD) | Number of profiles |
|------------|--------|--------------|--------------------|
| 2009 | -0.053 | 0.005 (10.4) | 271 |
| 2010 | -0.059 | 0.006 (10.8) | 275 |
| 2011 | -0.053 | 0.008 (15.4) | 232 |
| Mean/Total | -0.055 | 0.003 (5.2) | 778 |

Table 2. Coincident PSC events between MPL-4 and CALIOP measurements at four predominant CALIPSO ground-track distance-ranges from Belgrano II station: Correlation coefficient (CC) for three selected height intervals, and the calculated height-averaged $BIAS$, $BIAS_z$, and their SD together with the number of data points (in %, respect to the total data points) fulfilling the constraint condition ($-50\% < BIAS < +50\%$). Particular analysed cases are marked by asterisks.

| Distance (km) | Date | CC 5-20 km | CC 5-25 km | CC 5-30 km | $BIAS_z$ (%) | SD (%) | N^{BIAS} (%) |
|--|------------|---------------|---------------|---------------|-----------------|--------|-------------------|
| Distance-range (D1): < 10 km (12 cases) | | | | | | | |
| 3.28 | 15/06/2009 | 0.34 | 0.19 | 0.06 | -10 | 23 | 12 |
| * 3.55 | 01/07/2009 | 0.85 | 0.91 | 0.54 | +2 | 31 | 38 |
| 4.72 | 02/08/2009 | 0.77 | 0.66 | 0.52 | -8 | 27 | 22 |
| * 0.34 | 25/09/2009 | 0.25 | 0.12 | 0.18 | -24 | 20 | 22 |
| 1.58 | 08/06/2010 | 0.71 | 0.56 | 0.27 | -22 | 19 | 8 |
| 1.98 | 24/06/2010 | 0.89 | 0.83 | 0.78 | +3 | 23 | 44 |
| 3.75 | 04/07/2010 | -0.23 | -0.21 | 0.00 | -11 | 30 | 10 |
| 1.02 | 27/08/2010 | 0.58 | 0.23 | 0.04 | +1 | 29 | 22 |
| 3.05 | 07/07/2011 | 0.90 | 0.74 | 0.21 | -14 | 22 | 24 |
| 2.48 | 29/07/2011 | 0.70 | 0.57 | 0.32 | -16 | 26 | 36 |
| 2.66 | 14/08/2011 | 0.59 | 0.12 | -0.06 | -7 | 18 | 46 |
| 2.73 | 24/08/2011 | 0.70 | 0.61 | 0.62 | -11 | 26 | 22 |
| Distance-range (D2): 20 – 30 km (24 cases) | | | | | | | |
| 26.9 | 28/06/2009 | 0.46 | 0.22 | -0.07 | 0 | 33 | 16 |
| 25.5 | 30/06/2009 | 0.73 | 0.95 | 0.51 | +5 | 25 | 28 |
| 25.6 | 30/07/2009 | 0.71 | 0.79 | 0.72 | -12 | 26 | 40 |
| 27.0 | 01/08/2009 | 0.51 | 0.34 | 0.35 | -21 | 30 | 26 |
| 22.2 | 25/08/2009 | 0.72 | 0.68 | 0.51 | -16 | 24 | 42 |
| 30.1 | 27/08/2009 | 0.52 | 0.13 | 0.06 | -17 | 29 | 30 |
| 23.3 | 26/09/2009 | 0.13 | -0.11 | 0.23 | -11 | 25 | 24 |
| 29.0 | 28/09/2009 | 0.88 | 0.62 | 0.43 | -6 | 6 | 6 |
| 22.4 | 09/06/2010 | 0.73 | 0.69 | 0.63 | -8 | 31 | 4 |
| 29.8 | 11/06/2010 | 0.51 | 0.44 | 0.41 | +5 | 26 | 24 |
| 22.2 | 25/06/2010 | 0.76 | 0.63 | 0.69 | -6 | 28 | 32 |
| 29.9 | 27/06/2010 | 0.49 | 0.48 | 0.48 | +4 | 29 | 34 |
| 24.4 | 01/07/2010 | 0.39 | 0.39 | 0.18 | +9 | 30 | 12 |

| | | | | | | | |
|--|------------|-------|-------|-------|-----|----|----|
| * 28.0 | 03/07/2010 | 0.10 | 0.20 | 0.16 | +1 | 17 | 6 |
| 29.0 | 13/07/2010 | 0.54 | 0.00 | 0.28 | +5 | 27 | 40 |
| 24.1 | 28/08/2010 | 0.66 | 0.03 | 0.32 | -1 | 24 | 18 |
| 23.5 | 19/09/2010 | -0.03 | -0.16 | -0.19 | -6 | 34 | 10 |
| 29.5 | 29/05/2011 | 0.43 | -0.04 | 0.16 | -15 | 22 | 8 |
| 29.3 | 06/07/2011 | 0.48 | 0.49 | 0.17 | -7 | 31 | 20 |
| 28.9 | 16/07/2011 | 0.65 | 0.50 | 0.34 | -8 | 26 | 30 |
| 23.9 | 30/07/2011 | 0.28 | 0.61 | 0.45 | -4 | 31 | 34 |
| 23.9 | 15/08/2011 | 0.20 | -0.18 | -0.24 | -9 | 29 | 20 |
| 23.1 | 21/08/2011 | 0.85 | 0.85 | 0.85 | -2 | 23 | 52 |
| * 29.4 | 23/08/2011 | 0.76 | 0.75 | 0.53 | -12 | 23 | 60 |
| Distance-range (D3): 45 – 55 km (12 cases) | | | | | | | |
| 50.1 | 29/06/2009 | 0.05 | 0.81 | 0.55 | -2 | 32 | 18 |
| 48.9 | 31/07/2009 | 0.55 | 0.53 | 0.43 | -14 | 28 | 28 |
| 54.0 | 27/09/2009 | 0.05 | -0.07 | -0.10 | -18 | 30 | 8 |
| 55.5 | 10/06/2010 | 0.06 | 0.12 | -0.10 | -18 | 26 | 8 |
| 55.7 | 26/06/2010 | 0.51 | 0.60 | 0.64 | +2 | 23 | 32 |
| 49.7 | 02/07/2010 | 0.53 | 0.60 | 0.62 | -14 | 23 | 32 |
| * 54.9 | 12/07/2010 | 0.25 | 0.31 | 0.15 | -5 | 31 | 42 |
| 50.2 | 20/09/2010 | 0.55 | 0.55 | 0.55 | -14 | 27 | 26 |
| 56.8 | 28/05/2011 | -0.29 | -0.16 | 0.49 | -10 | 26 | 12 |
| 50.3 | 03/06/2011 | -0.05 | -0.18 | -0.03 | +11 | 26 | 12 |
| 56.7 | 15/07/2011 | 0.54 | 0.41 | 0.19 | -10 | 24 | 30 |
| * 50.8 | 22/08/2011 | 0.84 | 0.60 | 0.60 | -17 | 22 | 42 |
| Distance-range (D4): 70 - 90 km (35 cases) | | | | | | | |
| 80.1 | 16/06/2009 | 0.53 | 0.28 | 0.37 | -7 | 30 | 26 |
| 78.8 | 26/06/2009 | 0.61 | 0.03 | 0.65 | +5 | 24 | 20 |
| 80.5 | 02/07/2009 | -0.07 | -0.17 | -0.06 | +1 | 35 | 6 |
| 77.6 | 06/07/2009 | 0.17 | 0.56 | 0.48 | -8 | 27 | 18 |
| 79.3 | 12/07/2009 | -0.10 | 0.38 | 0.59 | -16 | 29 | 30 |
| 81.2 | 18/07/2009 | 0.07 | 0.65 | 0.38 | +7 | 24 | 22 |
| 77.1 | 22/07/2009 | -0.31 | -0.13 | -0.12 | -10 | 40 | 10 |
| 75.6 | 28/07/2009 | -0.16 | 0.49 | 0.13 | -23 | 16 | 8 |
| 80.4 | 13/08/2009 | 0.89 | 0.59 | 0.61 | +3 | 27 | 42 |
| 82.4 | 19/08/2009 | 0.19 | -0.10 | -0.10 | -5 | 28 | 28 |

| | | | | | | | |
|--------|------------|-------|-------|-------|-----|----|----|
| 77.5 | 08/09/2009 | 0.65 | 0.56 | 0.50 | +3 | 25 | 20 |
| 78.0 | 24/09/2009 | 0.85 | 0.75 | 0.68 | -10 | 30 | 18 |
| 82.8 | 05/07/2010 | 0.63 | 0.69 | 0.76 | +3 | 28 | 44 |
| 77.7 | 09/07/2010 | 0.73 | 0.69 | 0.48 | -14 | 28 | 36 |
| 78.5 | 25/07/2010 | 0.68 | 0.24 | 0.10 | -16 | 30 | 12 |
| * 75.2 | 31/07/2010 | 0.87 | 0.86 | 0.56 | +4 | 25 | 46 |
| 78.9 | 10/08/2010 | -0.33 | -0.27 | -0.12 | -15 | 15 | 8 |
| 75.2 | 16/08/2010 | 0.71 | 0.47 | 0.26 | -17 | 23 | 28 |
| 74.7 | 01/09/2010 | 0.51 | 0.33 | 0.17 | -42 | 5 | 8 |
| 83.0 | 07/09/2010 | 0.20 | 0.03 | 0.04 | -6 | 30 | 26 |
| 78.9 | 31/05/2011 | 0.67 | 0.67 | 0.43 | +1 | 24 | 6 |
| 73.0 | 31/05/2011 | -0.07 | -0.46 | 0.35 | +1 | 32 | 18 |
| 78.2 | 16/06/2011 | 0.31 | 0.04 | -0.16 | +14 | 21 | 10 |
| * 83.9 | 22/06/2011 | 0.13 | 0.27 | 0.10 | -14 | 27 | 32 |
| 78.3 | 26/06/2011 | 0.48 | 0.42 | 0.12 | -11 | 20 | 34 |
| 73.2 | 02/07/2011 | 0.42 | 0.77 | 0.55 | -13 | 21 | 28 |
| 84.3 | 08/07/2011 | 0.74 | 0.57 | 0.32 | -17 | 26 | 60 |
| 84.0 | 24/07/2011 | 0.26 | 0.37 | 0.07 | -25 | 20 | 10 |
| 73.4 | 03/08/2011 | 0.17 | 0.13 | 0.08 | -12 | 26 | 28 |
| 78.7 | 13/08/2011 | 0.35 | 0.04 | 0.04 | -18 | 25 | 32 |
| 78.0 | 19/08/2011 | 0.81 | 0.74 | 0.72 | +1 | 21 | 40 |
| 73.0 | 19/08/2011 | 0.32 | 0.31 | 0.10 | -4 | 29 | 18 |
| 84.3 | 25/08/2011 | -0.28 | -0.07 | -0.12 | -8 | 29 | 22 |
| 79.0 | 29/08/2011 | 0.64 | 0.26 | 0.06 | -19 | 32 | 18 |
| 78.9 | 30/09/2011 | 0.51 | 0.42 | 0.34 | -8 | 28 | 18 |

Figure captions

Figure 1. Stable polar vortex over Antarctica represented by the potential vorticity (PV) at 475 K-level on 24 June 2010. The white line indicates the edge of the polar vortex. Also shown are the location of Belgrano II station (Argentina, 77.9°S 34.6°W) together with Syowa (Japan, 69.0°S 39.5°E), which is outside the polar vortex edge, and South Pole/Amundsen-Scott (USA, 89.98°S 24.8°E) stations.

Figure 2. MPL-4 δ^V profiles (grey-lined open circles, in black background of their SD values) on 1 July 2009 depending on the time averaging (see legend inside each panel, from up to down and right to left): 5-, 10-, 15-min and 30-m averaged profiles.

Figure 3. MPL-4 mean ($\delta^V_{1-\min} - \delta^V$) differences along an hour and their RMS values (for instance, data on 1 July 2009).

Figure 4. An example of calibration of the volume linear depolarization ratio δ^V . From left to right: 1h-averaged (01:00-02:00 UTC) δ^V profile on 01 July 2009, once calibrated (thick solid line) and before calibration (thin solid line); the corresponding backscattering ratio R profile; and the closest available temperature radiosounding on 29 June 2009 at 11:00 UTC. The threshold temperatures for PSC-I (T^{PSC-I}) and PSC-II (T^{PSC-II}) formation are also shown assuming: 10-ppbv HNO₃ and 5-ppmv H₂O (thin dashed and solid lines, respectively) as typical values for June (Maturilli et al., 2005), and the in-situ water vapour concentration profile (radiosonde data) (thick dashed and solid lines, respectively). Calculations have been performed by using the parameterizations of Hanson and Mauersberger (1988) for PSC-I, and Marti and Mauersberger (1993) for PSC-II. CALIPSO ground-track distance was 3.6 km far from the Belgrano II station on this day.

Figure 5. Frequency (%) of Correlation Coefficients (CC) higher than 0.5 for four fitting height intervals (see legend) for the four predominant CALIPSO ground-track distance from Belgrano II station (D1: < 10 km, D2: 20 – 30 km, D3: 45 – 55 km, and D4: 70 – 90 km) examined in this work during all three years (2009-2011), and individually for 2009, 2010 and 2011 wintertime periods.

Figure 6. Mean differences ($\Delta = \delta^{MPL} - \delta^{CAL}$) (open symbols, left) and their RMS values (solid line, right) for the four predominant CALIPSO ground-track distance-ranges (from left to right, and from up to down, respectively): < 10 km (D1), 20-30 km (D2), 45-55 km (D3) and 70-90 km (D4). 5-km averaged mean differences (black stars) are also shown.

Figure 7. Height-averaged $BIAS$, $BIAS_z$, in dependence on the CALIPSO ground-track distance from Belgrano II station for 2009 (circles), 2010 (triangles) and 2011 (squares) wintertime periods. Those averaged values $BIAS_d$ (SD is marked by error bars) on each CALIPSO predominant separation (from D1 to D4) are also shown (black stars).

Figure 8. Simultaneous 0.5-km averaged profiles of the volume linear depolarization ratio δ^V for MPL-4 (full circles) and CALIOP (open triangles), and the differences $\Delta = \delta^{MPL} - \delta^{CAL}$ (black stars) found between both lidar δ^V profiles for CALIPSO ground-track distances < 10 km (D1 distance-range) far from Belgrano II station. PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 1 July 2009 (left) and 25 September 2009 (right). A height-averaged Δ value ($\langle \Delta \rangle \pm \text{SD}$) is also shown in each case.

Figure 9. The same as for Fig. 8, but for the CALIPSO D2 distance-range (20-30 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 23 August 2011 (left) and 3 July 2010 (right).

Figure 10. The same as for Fig. 8, but for the CALIPSO D3 distance-range (45-55 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 22 August 2011 (left) and 12 July 2010 (right).

Figure 11. The same as for Fig. 8, but for the CALIPSO D4 distance-range (70-90 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 31 July 2010 (left) and 22 June 2011 (right).




Figure 1. Stable polar vortex over Antarctica represented by the potential vorticity (PV) at 475 K-level on 24 June 2010. The white line indicates the edge of the polar vortex. Also shown are the location of Belgrano II station (Argentina, 77.9°S 34.6°W) together with Syowa (Japan, 69.0°S 39.5°E), which is outside the polar vortex edge, and South Pole/Amundsen-Scott (USA, 89.98°S 24.8°E) stations.




Figure 2. MPL-4 δ^V profiles (grey-lined open circles, in black background of their SD values) on 1 July 2009 depending on the time averaging (see legend inside each panel, from up to down and right to left): 5-, 10-, 15-min and 30-m averaged profiles.




Figure 3. MPL-4 mean ($\delta_{1-\min}^V - \delta^V$) differences along an hour and their RMS values (for instance, data on 1 July 2009).




Figure 4. An example of calibration of the volume linear depolarization ratio δ^V . From left to right: 1h-averaged (01:00-02:00 UTC) δ^V profile on 01 July 2009, once calibrated (thick solid line) and before calibration (thin solid line); the corresponding backscattering ratio R profile; and the closest available temperature radiosounding on 29 June 2009 at 11:00 UTC. The threshold temperatures for PSC-I (T^{PSC-I}) and PSC-II (T^{PSC-II}) formation are also shown assuming: 10-ppbv HNO_3 and 5-ppmv H_2O (thin dashed and solid lines, respectively) as typical values for June (Maturilli et al., 2005), and the in-situ water vapour concentration profile (radiosonde data) (thick dashed and solid lines, respectively). Calculations have been performed by using the parameterizations of Hanson and Mauersberger (1988) for PSC-I, and Marti and Mauersberger (1993) for PSC-II. CALIPSO ground-track distance was 3.6 km far from the Belgrano II station on this day.




Figure 5. Frequency (%) of Correlation Coefficients (CC) higher than 0.5 for four fitting height intervals (see legend) for the four predominant CALIPSO ground-track distance from Belgrano II station (D1: < 10 km, D2: 20 – 30 km, D3: 45 – 55 km, and D4: 70 – 90 km) examined in this work during all three years (2009-2011), and individually for 2009, 2010 and 2011 wintertime periods.




Figure 6. Mean differences ($\Delta = \delta^{MPL} - \delta^{CAL}$) (open symbols, left) and their RMS values (solid line, right) for the four predominant CALIPSO ground-track distance-ranges (from left to right, and from up to down, respectively): < 10 km (D1), 20-30 km (D2), 45-55 km (D3) and 70-90 km (D4). 5-km averaged mean differences (black stars) are also shown.




Figure 7. Height-averaged $BIAS$, $BIAS_z$, in dependence on the CALIPSO ground-track distance from Belgrano II station for 2009 (circles), 2010 (triangles) and 2011 (squares) wintertime periods. Those averaged values $BIAS_d$ (SD is marked by error bars) on each CALIPSO predominant separation (from D1 to D4) are also shown (black stars).




Figure 8. Simultaneous 0.5-km averaged profiles of the volume linear depolarization ratio δ^V for MPL-4 (full circles) and CALIOP (open triangles), and the differences $\Delta = \delta^{MPL} - \delta^{CAL}$ (black stars) found between both lidar δ^V profiles for CALIPSO ground-track distances < 10 km (D1 distance-range) far from Belgrano II station. PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 1 July 2009 (left) and 25 September 2009 (right). A height-averaged Δ value ($\langle \Delta \rangle \pm \text{SD}$) is also shown in each case.




Figure 9. The same as for Fig. 8, but for the CALIPSO D2 distance-range (20-30 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 23 August 2011 (left) and 3 July 2010 (right).




Figure 10. The same as for Fig. 8, but for the CALIPSO D3 distance-range (45-55 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 22 August 2011 (left) and 12 July 2010 (right).




Figure 11. The same as for Fig. 8, but for the CALIPSO D4 distance-range (70-90 km). PSC events correspond to those cases with a high/moderate and low CC values as observed, respectively, on 31 July 2010 (left) and 22 June 2011 (right).