Atmos. Meas. Tech. Discuss., 5, C423–C430, 2012 www.atmos-meas-tech-discuss.net/5/C423/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License.



AMTD

5, C423-C430, 2012

Interactive Comment

Interactive comment on "Areal-averaged trace gas emission rates from long-range open-path measurements in stable boundary layer conditions" by K. Schäfer et al.

K. Schäfer et al.

klaus.schaefer@imk.fzk.de

Received and published: 29 March 2012

Point-by-point reply to referee review

Anonymous Referee #1

General

The MS first considers various techniques available for measuring trace gas fluxes over large spatial scales of order of a ha or more and notes that micrometeorological methods are required, but all of them have problems during stable atmospheric conditions due to low levels of turbulence. The authors also conclude that non-intrusive methods



Printer-friendly Version

Interactive Discussion



are preferable for trace gases emitted from soils because they avoid the influences of the environment on the emission process. They appear to favour long-range open-path concentration measurements, but note that ideally, path-integrated wind and turbulence measurements made in the same control volume as the concentration measurements are required. The authors conclude that unfortunately, present instrumentation cannot provide the necessary accuracies during stable and very stable conditions and move on to consider using open-path gas measurements in a flux-gradient approach. They observe, however, that successful application of this method requires confidence in the gradients in gas concentration and confidence in the applicability of boundarylayer turbulence theory. Noting that under stable, light wind conditions, the continuous 'steady-state' turbulence of the surface boundary layer breaks down and the layer can have phenomena like intermittent and wave-like turbulent structures, they declare that they will deal only with what they call quality-assured determinations of fluxes under low wind, stable or night-time atmospheric conditions. One assumes that guality-assured data have been filtered for outlier values and values of parameters such as u and c. that fall within certain ranges. Provision of a table that shows what data have been filtered and what the acceptable ranges for the various parameters are would be a welcome addition. <Such a table is included now.> Finally, the authors present various flux-gradient methods and examine how well 3 of them succeeded in estimating fluxes in 2 experimental situations: one in a study of N2O fluxes from grassland and the other in an examination of NH3 fluxes from cattle lagoons. One of the methods used a similarity approach that does not assume log-linear relationships between wind speeds or gas concentrations and height while the other 2 used MOST flux-gradient approaches that assume log-linear profiles of wind speed and concentration. The flux calculations were tested by comparing with chamber data in the grassland experiment and integrated horizontal flux calculations in the lagoon experiment. It was concluded that the first of the approaches, the similarity approach, provided the best estimate of gas fluxes in stable conditions while the other methods were of limited value because the wind speed and concentration profiles were not log-linear as assumed by MOST.

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



While I think the topic of the paper of the paper is important and fits well with the journal's scope, I find the MS too long and too discursive in its present form. I feel its length could be reduced by a considerable amount, maybe 40 or 50% with the heaviest reductions in Section 1, the Introduction, and 3, Methods. Those Sections contain interesting information, but much of it is unnecessary. Examples include Table 1 which is a useful reference source in other contexts, but this paper is about open-path fluxgradient techniques, not closed-path techniques and it doesn't probe the strengths and weaknesses of chambers. < Table 1 is deleted. > A lot of Section 3, particularly Section 3.2 on open-path flux-gradient methods is conventional micrometeorology and could be abbreviated. The telling of the story seems to dart off in different directions unexpectedly in these 2 Sections and I suggest that their structures be tightened to more concise and logical forms. < This reduction is done as follows. Sect 1: more than 40 %; Sect. 2: about 30 %; Sect. 3.1.1: about 20 %, new order of three paragraphs; Sect. 3.1.2: about 10 %; Sect. 3.2: more than 50 %, also equations by 50 %; Sect. 3.3: more than 50 %, also all equations; Sect. 3.4: about 30 %; Sect. 3.5: about 30 %; Sect. 3.6: about 30 %; Sect. 4.1: about 30 %; Sect. 4.2: about 10 %; Sect. 4.3: about 20 %, new order of three paragraphs; Sect. 5: about 10 %, new order of one paragraph; Sect. 6: about 20 %, one paragraph shifted to Sect. 4.3. The reduction is in total 15 pages i.e. more than one third of the original text. Some of the below requested corrections were deleted by these abbreviations.> I am concerned that there is no really convincing demonstration either in the paper or in the references that flux-gradient relationships developed for measurements in the vertical plane apply over long horizontal paths of around 100 m. The authors consider the problem, but seem convinced that their guality assurance measures will overcome it. To me, something more convincing seems necessary, although I must admit that I can't say what it is; perhaps more chambers or anemometers or scintillation measurements or better still some demonstration either through measurement or modelling that time integrated means will iron out local perturbations. < This remark is added to the conclusions. > I also feel that a weakness in the paper is the lack of an unequivocal measurement of source strengths. Reliance

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on just 4 chambers (which gave quite different fluxes) and IHF methods to test the flux gradient methods seems rather risky. <We agree.> A good deal of editing for colloquial English is necessary. I have made a number of suggestions in my Specific comments, but other changes still need to be made. <This is done.>

Specific

p.4, line 14: suggest "other" for "further" <ok>

p.4, line 15: delete "the" between "of" and "source" <ok>

p.4, line16: suggest move " and sink" to to previous line after source; delete the end bracket after "2006" and the forward bracket before "Chapuis-Lardy" <ok>

p.5, line 6 : delete "the" <ok>

p.5, line 11: I don't know of an instance where an open-path TDLAS instrument has been used for eddy covariance. Perhaps give a reference. <improved>

p.5, line 18: suggest insert "the" between "called" and "integrated" <ok>

p.5, line 19: "Lagrangian" for "Lagrange" <ok>

p.5, last line: perhaps "evaluating" for "approximating' <done>

p.6, line 3: suggest "low turbulence levels" for "insufficient turbulent transport" <ok>

p.6, line 12: suggest delete the first "and"; insert "recently" between "and" and "for" <ok>

p.6, line 13: suggest move "of QCL" to between "versions" and "are" <ok>

p.6, lines 13-16: closed-path FTIR deserves a mention here <done>

p.6, line 17: suggest move "path-integrated wind and turbulence measurements" to beginning of line; add "are" after it <ok>

p.7, lines 3 & 4: The statement that non-intrusive flux measurements are available for

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CO2 and H2O only is not true. The open-path Licor 7700 instrument measures CH4 at a point in just the way the authors describe: non-obtrusively & without any enclosure or sampling or measuring cell and is intended for eddy covariance measurements, while open-path lasers are available for all 3 of N2O, CH4 and NH3. <corrected, improved>

p.7, line 2 from bottom: See comment about "quality-assured" in my General comments <improved>

p.8, line 7: suggest reword the sentence to read "The footprint function describes the relationship....." <ok>

p.8, lines 10 & 11: suggest reword the sentence to read "It is the objective of the study....." <done>

p.8, line 13: suggest "of " for "to" and "defining" for "defined" <ok>

p.8, line 15: suggest "However," for "But also,"; "analysis" for "analyses"; "of" for "to": "determining" for "determine" <ok>

p.8, line 17: suggest delete "the" <ok>

p.8, line 18:suggest "analysis" for "analyses" <ok>

p.8 line 19: suggest delete the second "the" <ok>

p.9, lines 4 & 5: suggest "changes" for "change" and "take" for "takes" <ok>

p.9, line 6: suggest delete the 2nd "the" <ok>

p.9, line 8: suggest insert "to occur" after "fluxes" <ok>

p.9, line 16: suggest "with" for "to" <ok>

p.9, line 20: suggest "in the present context" for "here" <ok>

p.10, line 8: suggest delete "the" <ok>

p.10, line 16: "at" for "in" <ok>

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



p.11,line 2: "upon" for "from" <ok>

p.12, line 3: clarify if 1-, 2- or 3-D <done>

p.13, line 17: zF is often set at the geometric mean of z- and z+, e.g., Paulson (1970) <Thank you for this comment. Paulson stated that for the heights 0.5 m and 16 m the error is 6 %. We shortened the text so that this detail is no more described.>

p.13, line 21: "on" for "from" <ok>

p.13, line 22: insert "the" after "as" <ok>

p.14, discussion of .m and .c after Eq.(2): It is usual to equate .m and .c in neutral conditions. Is that what is intended here? <improved>

p.17,line13: insert "at" after "set" <ok>

- p.22, line 11: delete "is applied which" ? <ok>
- p.22, line 5: suggest "used" for "was using" <ok>

p.23, lines 5 to 7: The sentence here needs rewording for clarity <improved>

p.23, lines 10, 12, 14: "at" for "in" <ok>

p.23, line 22: insert "is" after "and" <ok>

p.26 & 27: I wonder if the discussion in the first 2 paragraphs of this new section is necessary <shortened>

p.27, last paragraph: Perhaps justification for the use of IHF to represent the real flux is needed. Reasons could include simplicity of concept, few theoretical assumptions, etc. <Exactly this discussion about fluxes is in the next chapter and shown in the next figure (Fig. 7). Here is a discussion about the turbulence.>

p.28, lines1 & 2: The meaning is unclear to me. Suggest reword to clarify. <improved>

p.30, lines 2 and 14: "at" for "in" <ok>

AMTD

5, C423–C430, 2012

Interactive Comment



Printer-friendly Version

Interactive Discussion



P. 30, line 7: insert "in" after "gaps" <ok>

p.31, lines 1,5, 6: "at" for "in" <ok>

p.31, line 1: suggest delete "was" <ok>

p.31, line 2: insert "was" after "height" <ok>

p.31, lines 12-16: This sentence needs rewording to clarify what is being said <improved>

p.31, line 16: delete "up" <ok>

p.32, line 9: insert "the" before Indiana <ok>

p.32, lines 11 to 13: Higher emissions during near-neutral conditions might also be due to effects on soil and lagoon temperatures, which will change equilibrium gas concentrations <improved by extension with these effects>

p.33, line 7: the range 0 < . =+10 would exclude all unstable cases because . would be < 0. <corrected by writing xsi<=+10>

p.34, line 3: "periods" for "period" <ok>

p.34, line 10: Where is Sect 5.1? Should it be Sect. 3.2? <corrected>

p.35, line 8: "layers' for "layer" <ok>

p.35, line 11: suggest move "calculated" to before "N2O" <ok>

p.35, lines 10-17: There is too much crammed into this sentence. Suggest split it and rewrite. <improved>

p.35, last 3 lines: meaning unclear <re-formulated>

p.38, 1st reference: Year of publication missing <included>

p.38, 2nd reference: 2 years of publication. Maybe one of the years belongs to the 1st

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reference <corrected>

p.48, Table 1: NH3 is a sticky and very soluble gas and chambers are not recommended for measuring NH3 fluxes. <Table is excluded.>

- p.57, Figure 8: What does the curved line represent? <added>
- p.58, Figure 9: What is the distinction between diamonds and squares? <added>
- p. 59, Figure 10: Same comment as Fig. 9 <added>
- p.60, Figure 11: What does the curved line represent? <added>

All changes in the original manuscript are shown in a separate file.

Interactive comment on Atmos. Meas. Tech. Discuss., 5, 1459, 2012.

AMTD

5, C423-C430, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

