

**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

# Assessment of GPS radiosonde descent data

M. Venkat Ratnam<sup>1</sup>, N. Pravallika<sup>2</sup>, S. Ravindra babu<sup>3</sup>, G. Basha<sup>4</sup>, M. Pramitha<sup>1</sup>,  
and B. V. Krishna Murthy<sup>5</sup>

<sup>1</sup>National Atmospheric Research Laboratory, Gadanki, India

<sup>2</sup>Sri Venkateswara University, Tirupati, India

<sup>3</sup>Jawaharlal Nehru Technology University, Hyderabad, India

<sup>4</sup>SRM University, Chennai, India

<sup>5</sup>B1, Ceebros, 47/20, Illrd Main Road, Chennai, India

Received: 7 November 2013 – Accepted: 22 November 2013 – Published: 3 December 2013

Correspondence to: M. Venkat Ratnam (vratnam@narl.gov.in)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Radiosondes are widely used to obtain basic meteorological parameters such as pressure ( $P$ ), temperature ( $T$ ), relative humidity (RH), and horizontal winds during the balloon ascent up to the altitude of balloon burst, usually  $\sim 32\text{--}35$  km. Data from the radiosondes released from Gadanki ( $13.5^\circ$  N,  $79.2^\circ$  E), a tropical station in India, has been collected during the ascent and during the descent as well without attaching any parachute or its equivalent since the year 2008. In the present study an attempt has been made to characterize the radiosonde descent data with the main objective of exploring its usefulness and reliability for scientific purposes. We compared the data obtained during ascent and descent phases of the same sounding. The mean differences in  $T$ , RH and horizontal winds between ascent and descent data are found to be small and are sometimes even within the uncertainty of the measurements and/or expected diurnal variation itself. The very good consistency observed between the ascent and the descent data shows that one more profile of the meteorological parameters can be constructed within 3 h of time of balloon launch practically at no additional cost. Further checks are done by utilizing the 3 hourly radiosonde observations collected during the Tropical Tropopause Dynamics campaign conducted at Gadanki. In the process of checking the consistency between the radiosonde ascent and descent data, several new findings are arrived at and are reported in this study. In general, it has taken more than half-an-hour for the balloon to reach the ground from the burst altitude. It is also observed that the fall velocity is close to  $10\text{ m s}^{-1}$  near the surface. Finally, it is suggested to record also the observations when the balloon is descending as this information is also useful for scientific purposes.

## 1 Introduction

Measurements of different meteorological parameters such as pressure ( $P$ ), temperature ( $T$ ), relative humidity (RH) and horizontal winds are very important for forecasting

# AMTD

6, 10361–10398, 2013

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the weather and for studying the thermal and dynamical state of the atmosphere. Several efforts are made to probe the atmosphere using different in-situ and remote sensing techniques. In the in-situ measurements, the sensors are normally located in the instruments carried aloft in aircraft and rockets or as radiosondes if vertical structure is needed. Instrument intended to be carried by a balloon through the atmosphere, is equipped with devices to measure one or several meteorological variables ( $P$ ,  $T$ , RH, winds etc.), and is provided with a radio transmitter for sending this information to the observing station. Originally named a radio-meteor graph, the instrument is now referred to as a radiosonde, a name apparently derived by *H. Hergesell* from a combination of the words “radio” for the onboard radio transmitter and “sonde”, which is messenger from old English.

The earliest radio soundings were done in the late 1920’s and 1930’s by *V. Väisälä*. However, archived data generally begin in the 1940’s, with significant expansion of the network at the time of the International Geophysical Year (1957–1958). Currently, the global radiosonde network includes about 900 upper-air stations and about two-thirds of these conduct the experiment twice daily (at 00:00 and 12:00 UTC) (Durre, 2006; Immler et al., 2010). The network is predominantly land-based and favors the Northern Hemisphere as it is covered mostly by land. Measurements at every 12 h from these locations may be enough to define the large-scale weather patterns, though not sufficient to define small-scale weather events. However, for many purposes the balloon sounding is an ideal research tool. In addition, some radiosondes are flown with sensing systems for atmospheric constituents, such as ozone concentration and radioactivity. The ranges of meteorological parameters are: 1050 to 5 hPa for  $P$ , 50 to  $-90^\circ\text{C}$  for  $T$  and 100 to 1 % for RH, with the systems being able to sustain continuous reliable operation even when operating in heavy rain or in the vicinity of thunderstorms or in severe icing conditions. See detailed review on the quality of the radiosonde measurements worldwide by Durre et al. (2006) and the WMO inter-comparison report of high quality radiosonde systems (WMO, 2011). Note that Sodell et al. (2004) employed Aerosonde which is a programmable robotic aircraft developed mainly for use over remote and

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



inaccessible regions (like oceans) for meteorological observations and compared them with simultaneous radiosonde observations and found good agreement between the two. However, Aerosonde has not been much in use probably due to the complexities involved and cost factor.

As mentioned earlier, in general, radiosondes are widely used to obtain major meteorological parameters up to an altitude of balloon burst, usually  $\sim 32$ – $35$  km. It costs up to US \$ 200 per each launch including the cost of balloon, gas, manpower etc. However, many observations worldwide are not done up to altitudes higher than tropopause (Durre et al., 2006) or about 25 km because of the higher cost of the balloons and gas necessary to lift the equipment to the low pressures. In general, across the world data are collected only during ascent of balloons and once balloon bursts the data collection is terminated. There may be several reasons for this including the thought of sensing different air masses between the ascent and the descent (due to balloon drifts), sensor response time (descent rate is faster than the ascent rate) etc., Chen et al. (2013) employed a light weight hard ball as a parachute to the descending radiosonde to deduce vertical air motions at altitudes below  $\sim 12$  km. In the present study we obtained the meteorological data from the descending radiosonde without attaching any parachute or its equivalent and explored the usefulness and reliability of the data during the descent. To the best of our knowledge there was no such study earlier. The ascent and descent data has been obtained from the radiosondes released (without any parachute or equivalent) from Gadanki ( $13.5^\circ$  N,  $79.2^\circ$  E), a tropical station in India since the year 2008 for this purpose. The results of this study are presented in this communication.

## 2 Data base

High-resolution GPS radiosonde (Meisei, Japan make, RD-06G) balloons were launched almost regularly from Gadanki during January 2008 to December 2011 around 17:30 IST (IST = UT + 5.5 h). The weight of the sonde is 150 g and is attached to a 600 g Totex balloon. The dimensions of the sonde are  $98(W) \times 155(H) \times 88(D)$  mm.

**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Note that no parachute is attached to the balloons launched from Gadanki. About 1461 profiles of  $P$ ,  $T$ , RH, and horizontal winds reaching an average altitude of  $\sim 30$  km were obtained in different seasons during the aforementioned period. The data of the atmospheric parameters were collected at sampling intervals of 1 s yielding an altitude resolution of 5–6 m. The original response time of the temperature (NTC thermistor) sensor vary from 0.12 to 10 s though the data is recorded for every 1 s. However, the response time for the humidity (capacitance based thin polymer) is 10 s for a step change from 11 % to 93 % RH, respectively. The balloon burst altitude of all the radiosondes launched during 1 January 2008 to 31 December 2011 is shown in Fig. 1a and the minimum altitude reached during descent is also superimposed in the same panel. Number of profiles reaching different altitudes in the ascent and the descent are shown in Fig. 1b and the statistics in different seasons is given in Table 1. Note that the minimum altitude in the descent is not zero (ground) owing to the failure of the radio communication at altitudes close to the ground due to surrounding complex terrain of Gadanki location. The minimum altitude has a seasonal dependence as can be seen from the table. During the monsoon season the profiles reaching down close to the ground is less than that during the other seasons due to blocking of the hills located in the south-west side. Note that radiosonde will drift in the southwestern side of Gadanki during this season due to the prevailing winds in that direction.

As mentioned earlier, there were 1461 soundings out of which 537 soundings were later eliminated due to non-availability of usable descent data mainly because of very short duration of the data for a meaningful assessment. The descent data of radiosondes was not recorded initially in the starting year 2008. Thus, there are 924 soundings in total available for the present study and when they are segregated into seasons the number of soundings are 256, 234, 218, and 216 during winter, pre-monsoon, monsoon, and post-monsoon seasons, respectively. Out of the 924 balloons, only 14 balloons reached less than 16 km altitude in ascent. About 98.2 % of the balloons reached the altitude of tropical tropopause (around 16 km) and 92.6 and 85.5 % reached the altitude of greater than 20 and 25 km, respectively. Note that higher altitudes are attained

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over our station by using quality balloons (Totex, Japan) unlike in the worldwide radiosonde statistics reported by Durre et al. (2006) which reveals that most of the radiosondes terminate near tropopause. Number of balloons reaching close to the surface (0–1 km) in descent is highest in the pre-monsoon season and least in the monsoon season. Note that more than 50% of balloons reached 5 km (in descent) in all the seasons. In contrast to this, number of balloons reaching the maximum altitude is lowest in the pre-monsoon season and highest in the monsoon season.

### 3 Background conditions

#### 3.1 Topographical conditions

Before going into the details of the results, the topography and background atmospheric conditions over the study region are presented which will be useful in interpreting the observed results. The station Gadanki is in a rural environment located at about 120 km northwest of Chennai (Madras) on the east coast of the South Indian peninsula. The location and topography of the Indian subcontinent and more detailed topography of Gadanki are shown in Basha and Ratnam (2009). Note that this station is surrounded by hills with a maximum altitude of 350–400 m above the station, and the station is at an altitude of 375 m above mean sea level (hereinafter all altitudes are mentioned above mean sea level only). The local topography is complex with a number of small hillocks around and a high hill of ~ 1 km about 30 km from the balloon launching site in the northeast direction.

#### 3.2 Background weather conditions

Over the Indian subcontinent, meso-scale phenomena including convection dominate. On the basis of the meteorological conditions over the observational site, the months are grouped into four seasons, namely, winter (December–February), summer/pre monsoon (March–May), monsoon/progressing southwest (SW) monsoon

(June–August), and post monsoon/retreating SW monsoon (September–November). The onset of the SW monsoon will be in the first or second week of June and withdrawal in the last week of September. Besides this, the northeast (NE) monsoon due to northeast circulation occurs over the southern part of India. Normal date of onset of the NE monsoon is in the middle of October, with a deviation of about a week on either side.

### 3.3 Background meteorological conditions obtained during the balloon ascent

In this sub-section background meteorological conditions prevailing over the observational site are briefly described based on the data collected during the balloon ascent. The monthly mean contours of the  $T$ , RH, zonal ( $U$ ) and meridional ( $V$ ) winds are shown in Fig. 2a–d.

The mean zonal velocities are generally westerly during winter (with a maximum velocity of  $10\text{--}12\text{ ms}^{-1}$ ) and pre-monsoon (up to  $14\text{ km}$  with maximum velocity of  $13\text{ ms}^{-1}$ ). There seem to be weak easterlies above the altitude of  $14\text{ km}$  during the pre-monsoon. During the monsoon season low-level westerlies exist below  $7\text{--}8\text{ km}$  and easterlies above. The tropical easterly jet (TEJ) is prevalent over this region in the SW monsoon season, with peak velocity sometimes reaching more than  $40\text{ ms}^{-1}$  (Roja Raman et al., 2009). There exist large vertical shears during monsoon in the zonal wind. In general, meridional velocities are very small and are southward up to  $8\text{ km}$  and northward above in all the seasons, except during monsoon, with peak velocity of  $5\text{ ms}^{-1}$ . During the winter and monsoon, somewhat stronger southerlies and northerlies prevailed, respectively, between  $12$  and  $15\text{ km}$ . A clear annual oscillation can be noticed in both zonal and meridional velocities. Similar variations are also observed by the MST radar located at the same site in between  $4$  and  $20\text{ km}$  (Ratnam et al., 2008; Basha and Ratnam, 2009; Debashis Nath et al., 2009).

Temperature does not show significant variations seasonally in the lower to middle troposphere, but shows variations in the lower stratosphere. There exist significant seasonal variations in the RH. During winter RH is very small (about  $50\%$ ) in the first

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



few kilometers and is almost negligible above. However, during the other seasons, particularly in the monsoon season, large RH concentrations (60–70 %) can be noticed up to middle troposphere.

## 4 Results and Discussion

### 4.1 Maximum horizontal drift of the radiosonde while in ascent and in descent

Before assessing the radiosonde descent data, it is desirable to see the maximum horizontal drift of the balloons from the launch location. Monthly mean drifts of the radiosondes from Gadanki location during ascent time observed in different seasons are depicted in Fig. 3. Figure 4 shows the locations of the balloons at the maximum altitude reached for all the radiosondes launched from Gadanki (shown with start mark) during January 2008 to December 2011 separated according to the season. The radiosonde balloon drift at height level  $z_i$  is computed from the successively reported horizontal positions of the balloon with altitude. This is done by accumulating successive horizontal distances travelled by the balloon as given by Houchi et al. (2010):

$$\text{Drift}(z_i) = \sqrt{\left(\sum_{j=1}^{j=i} dx_j\right)^2 + \left(\sum_{j=1}^{j=i} dy_j\right)^2} \quad (1)$$

where  $dx_j = 0.5(u_j + u_{j+1})dt$  and  $dy_j = 0.5(v_j + v_{j+1})dt$  are the zonal and meridional distances travelled by the radiosonde balloon from one atmospheric level  $j$  to another  $j + 1$  with respect to the launch location (13.5° N, 79.2° E).  $z_i$  is the level height, where  $i$  indicates the level number at which the drift is computed. In general, radiosonde has drifted maximum during the monsoon season mainly due to large winds observed due to prevailing TEJ during this season as shown in Fig. 2. Radiosondes have gone greater than 120 km from the launch station during this season. The radiosonde drift in winter

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

season is a minimum and the maximum distance it has drifted during this season is only 20 km. During post-monsoon and pre-monsoon seasons the radiosonde drifts for maximum of 80–100 km and 40–60 km, respectively. The observed drifts closely match with the background wind velocities as expected, i.e., the higher the wind velocities, the larger the radiosonde drifts and vice versa. Though we cannot directly compare our results with those reported by Houchi et al. (2010) and Seidel et al. (2011) (as they have not included any Indian radiosonde observations) on the balloon drifts, note that drifts are larger at our location in contrast to those reported by them. Seidel et al. (2011) reported that drifts are larger at mid-latitudes than tropical, but even at a tropical latitude station like Gadanki, we could notice large drifts. Another striking difference observed is that summer season drifts are larger than winter unlike those reported by Seidel et al. (2011). These differences are arising mainly due to prevailing strong tropical easterly jets over the Indian region whereas the drifts at the mid latitudes are due to the prevailing sub-tropical jets (Seidel et al., 2011).

The maximum drift of the radiosondes from the balloon burst altitude and the locations of the balloons reaching the lowest altitude for all the radiosondes launched from Gadanki is also estimated for each radiosonde launched during January 2008 to December 2011 (Figure not shown). It is observed that the radiosonde drifted with the background wind in descent also as expected. It is also observed that the mean drifts during descent are small in all the seasons, being less than 20 km. Thus, in principle, it can be concluded that during the ascent and the descent same air masses are probed. Further, McGrath et al. (2010) reported that impact of any errors due to the mean drifts on climate statistics is very small though individual errors may be of significance for weather modelling and for the tuning of satellite data (Sun et al., 2010). However, MacPherson (1995) reported that “while the impact of balloon drift can indeed be detected, it is small compared with the impact from the observation when assimilated with neglect of balloon drift”. We discuss this aspect further in the following.

## 4.2 Mean differences between the ascent and the descent durations

Histograms showing the times taken for the balloon to reach maximum and minimum altitudes during different seasons along with the 4 yr mean ascent and descent duration are depicted in Fig. 5. The histograms showing the ascent and descent durations reveal that it has taken about one and half hours to two hours to reach maximum altitude of 30–35 km and mostly half-an-hour for the descent. There is another peak near 10 min suggesting that quite often it has taken less than 10 min to reach to the ground/near ground. Note that the total ascent duration depends on the maximum altitude reached and not all the balloons reached altitudes greater than 35 km and above. However, it is very interesting to see that about 70 % of the balloons had taken more than 30 min to reach the ground/near surface from the balloon burst altitude. The 4 yr mean ascent and descent rates along with standard deviation shown in Fig. 5d reveal that the mean ascent rate is close to 5–6 ms<sup>-1</sup> in troposphere and reaches slightly higher values in the stratosphere. Whereas the descent rate is as high as 50 ms<sup>-1</sup> near the balloon burst altitudes and it reaches close to 10 ms<sup>-1</sup> near the ground.

We computed the descent rate (fall velocity) using the following equation (Johansson and Bergstrom, 2005) based on the balance between gravity and drag forces suffered by the radiosonde (weight of 150 g) (without the gas but assuming the portion of bursted balloon is still attached).

$$\text{Drag} = \text{mass} \times g = \frac{1}{2} \times \rho \times V_T^2 \times C_d \times \text{area} \quad (2)$$

$$\text{Terminal velocity } V_T = \sqrt{\frac{2 \times \text{mass} \times g}{\rho \times \text{area} \times C_d}} \quad (3)$$

Where  $g$  is acceleration due to gravity,  $\rho$  is the density,  $C_d$  is the drag coefficient and frontal area is calculated using both edge and no-edge wise. The four year mean density profile along with standard deviations used in estimating  $V_T$  is shown in Fig. 6a. By considering the dimensions of the radiosonde as 155 mm × 98 mm × 88 mm, the frontal

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



area is calculated as  $0.015 \text{ m}^2$  (non-edge) or  $0.008 \text{ m}^2$  (edge wise). Taking  $C_d$  as 1.0 for convenience and the radiosonde mass as 150 g, the terminal velocity is calculated both edge wise and non-edge wise and is shown in Fig. 6b. At 10 km altitude ( $\rho \sim 0.4 \text{ kg m}^{-3}$ ) the terminal velocity is close to  $22 \text{ m s}^{-1}$  (edge) or  $29 \text{ m s}^{-1}$  (non-edge). At 30 km, these values ( $\rho$  is  $\sim 0.017$ ) are  $106 \text{ m s}^{-1}$  or  $140 \text{ m s}^{-1}$ . The maximum observed value is  $60 \text{ m s}^{-1}$  (Fig. 5c) suggesting the radiosonde might be tumbling experiencing a much higher drag.

Note that the descent duration also depends on the maximum altitude i.e., higher the maximum altitude, longer the time taken to reach the ground. The time taken to reach different altitudes while descending from the maximum altitude is shown in Table 2. It is interesting to see no difference in the time taken to reach from 30 km to 10 km and 25 km to 10 km suggesting that radiosonde velocity is high when it bursted which can be seen from Fig. 5c. This is also true for the balloons reaching from 30 to 8 km and 25 to 8 km, 30 to 6 km and 25 to 6 km, 30 to 4 km and 25 to 4 km. Note that in winter season it takes less time to reach ground and more in monsoon season.

### 4.3 Differences between the ascent and the descent data

As mentioned earlier, the radiosonde data is available at every 5–6 m (1 s interval). However, data has been smoothed to 100 m resolution so as to remove the errors, if any, arising due to random motion of the balloon. Typical profiles of  $T$ , RH,  $U$ ,  $V$  (from ascent data) and ascent rate observed on 1 January 2008 at 12:00 UT are shown in Fig. 7. Since one of the main objectives is to examine the reliability and usefulness of the descent data, the profiles of  $T$ , RH,  $U$ , and  $V$  from descent data and descent rate are also superimposed in the respective panels. Note that the descent profiles of  $T$ , RH,  $U$  and  $V$  resemble the smoothed version of the ascent profiles. In general, very good consistency can be observed between the ascent and descent data sets revealing that one more profile can be constructed within 3 h of time from the balloon launch.

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The differences between ascent and descent data in  $T$ , RH,  $U$  and  $V$  are estimated and are shown in the middle panels of Fig. 7. Note that the horizontal winds,  $T$  and RH are measured with an accuracy of  $0.15 \text{ ms}^{-1}$ ,  $0.5 \text{ K}$  and  $5\%$ , respectively. The difference in  $T$  between ascent and descent is within  $1 \text{ K}$  up to  $\sim 16 \text{ km}$  ( $\sim$  tropopause) and above the tropopause it is more and can reach as high as  $4 \text{ K}$ . In the case of RH, there is a difference of about  $10\%$  between the ascent and descent data with higher RH during descent phase below  $10 \text{ km}$  and reverse above that altitude. The difference in  $U$  between ascent and descent is within  $4 \text{ ms}^{-1}$  with slightly higher values in the lower stratosphere. Below  $15 \text{ km}$ ,  $V$  during the descent is more and reverses above. Generally ascent rate is close to  $5 \text{ ms}^{-1}$ . The descent rate initially is as high as  $60 \text{ ms}^{-1}$  and drastically decreases with increase of density to almost  $10 \text{ ms}^{-1}$  before it reaches the ground. Wild fluctuations in descent rate can be noticed which probably can be attributed to tumbling of the radiosonde as indicated by the estimated terminal velocities with two different radiosonde frontal areas. The terminal velocity is inversely proportional to the square root of the frontal area. Roughly the frontal area is about one-third edgewise, so that the terminal speed would be 1.7 times the non-edgewise descent. This appears to be consistent with top two rows of Fig. 7e panels. In order to check the variability of the differences in  $T$ , RH,  $U$ , and  $V$  and between the ascent and descent rates, these are shown as histograms in the third row panels of Fig. 7. In general, most of the difference in  $T$ , RH,  $U$ ,  $V$ , and ascent and descent rate are within  $2 \text{ K}$ ,  $5\%$ ,  $2 \text{ ms}^{-1}$ ,  $2 \text{ ms}^{-1}$ , and  $20 \text{ ms}^{-1}$ , respectively.

Similar analysis as mentioned above has been done for the entire 4 yr of data and is shown in the bottom panels of Fig. 7. Large spread in each of the parameters mentioned above is noticed which can be mainly attributed to the day-to-day, seasonal, annual and inter-annual variability in the respective parameters. Relatively less variability is seen in  $T$  than in RH,  $U$  and  $V$ . The difference in  $T$  is within  $3 \text{ K}$  up to  $16 \text{ km}$  and increases to  $5 \text{ K}$  sometimes. In case of RH, there is a difference of about  $30\%$ – $40\%$  between the ascent and the descent data with higher RH during the descent phase below  $10 \text{ km}$ . The difference in  $U$  between the ascent and the descent is seen

up to  $5 \text{ ms}^{-1}$  which increase to  $10 \text{ ms}^{-1}$  in the lower stratosphere. The difference in  $V$  can go as high as  $10 \text{ ms}^{-1}$  between the ascent and the descent data. The descent rate is as high as  $100 \text{ ms}^{-1}$  drastically decreasing with increase of density below to almost  $10 \text{ ms}^{-1}$  before it reaches to the ground. In general, most of the differences in  $T$ ,  $RH$ ,  $U$ ,  $V$ , and ascent and descent rate are within  $5 \text{ K}$ ,  $30 \%$ ,  $5 \text{ ms}^{-1}$ ,  $5 \text{ ms}^{-1}$ , and  $10 \text{ ms}^{-1}$ , respectively. Note that the observed differences seems to be at higher side and is mainly arising due to including the differences immediate after balloon burst, which otherwise the differences will be low and details will be shown in Fig. 9. However, there could be several reasons for these differences within 3 h of observations including the diurnal variation and prevailing waves and are discussed in detail in the following sub-sections.

#### 4.4 Seasonal mean differences in the meteorological parameters between the ascent and the descent data

As there is a time difference of two and half hours to three hours between the ascent and decent profiles, it is quite natural to expect some difference between these profiles because of diurnal variation of the parameter itself. To get a clear picture of the diurnal variation in each parameter, we make use of the 3 hourly radiosonde launchings done during the Tropical Tropopause Dynamics (TTD) campaign (Ratnam et al., 2013) conducted at Gadanki every month for 3 days (total 24 launches) since December 2010. The diurnal variation observed in  $T$ ,  $RH$ ,  $U$  and  $V$  during the above mentioned period separated seasonally is shown in Fig. 8. The diurnal variation in  $T$  below the tropopause is less than  $1 \text{ K}$ . However, strong diurnal variation in the lower stratosphere and near ground reaching as high as  $3 \text{ K}$  can be noticed and there exists some seasonal dependence in the diurnal variation. Diurnal variation with warmer and colder conditions during day and night time can be clearly noticed in the lower troposphere. The diurnal variation in  $RH$  can be as high as  $15 \%$  with high and low values during noon and mid-night hours, respectively, in all the seasons except in winter. In winter, no strong diurnal variation can be noticed in  $RH$ . The diurnal variation observed in  $U$  and  $V$  is

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



close to  $2 \text{ ms}^{-1}$  in the troposphere whereas it is as high as  $4 \text{ s}^{-1}$  in the lower stratosphere. The diurnal variation between 17 IST (ascent time) and 20 IST (descent time) is highlighted with vertical lines in Fig. 8. A difference of 2–3 K, 15–20 %, 3–4  $\text{ms}^{-1}$  in  $T$ , RH and wind components respectively, can be noticed between 17 IST and 20 IST. Note that differences in all the meteorological parameters have been noticed (Fig. 5) between the ascent and descent profiles. It can be seen that the differences in the above mentioned parameters between ascent and descent are consistent with the diurnal variations strongly supporting the contention that the descent data is meaningful and reliable which can be used for scientific studies.

In order to check whether there is seasonal dependence in the observed differences in the meteorological parameters between the ascent and the descent profiles, similar procedure is followed as for the individual cases and for all four years of the data. Figure 9 shows the seasonal mean difference profiles of  $T$ , RH,  $U$  and  $V$  between ascent and descent data along with standard deviations at different altitudes. The seasonal mean differences between the ascent and descent data at different altitudes are shown in Table 3 and for all the four years data in Table 4. It is very interesting to see that the mean difference in  $T$  is less than 0.5 K below 30 km except near tropopause altitude during monsoon season where it reaches 1 K with standard deviation of 2 K. Above 30 km the mean difference reaches 2 K with standard deviation of 4–6 K. The mean RH difference between ascent and descent is less than 10 % with standard deviation of 20–25 % below 12 km. It is again consistent to see higher RH during ascent time than descent time as the atmosphere cools by the time of descent. Since radiosonde is not sensitive to RH below  $-40^\circ\text{C}$  (Miloshevich et al., 2009 and references therein), no conclusion can be reached for altitudes above 12 km and they are erroneous. Differences seen above 25 km in  $T$  must be due to inadequate sensor response during the fast descent so that the values are biased toward higher altitudes during descent.

The mean difference in  $U$  and  $V$  between ascent and descent is less than  $1 \text{ ms}^{-1}$  with standard deviation of about  $5 \text{ ms}^{-1}$ . In general, the mean temperature difference in all the parameters is less in winter season compared to the other seasons. At altitudes

near the balloon burst, since the descent rate is high (Fig. 5c), the response time of sensors is also important, where ambient density is low and the terminal velocity is high (Fig. 6). The sensor will lag behind the ambient value because of the finite response time.

#### 5 4.5 Consistency check for the descent data with immediate balloon ascent data

In general, it takes two to two and half hours to reach the maximum altitude of 35 km and takes only 30 min to reach surface or near surface. To reduce this time difference for a better comparison, we made use of the 3 hourly radiosonde launching done during TTD campaign mentioned above. Using this data we have estimated the difference between the descent data and the ascent data of immediate next radiosonde. In this case note that the time difference is less than half an hour below the tropopause altitude. The profiles of temperature obtained during TTD campaign launched every 3 h during 28–31 December 2010 are shown in Fig. 10. Descent profiles are also superimposed in the same panel. A very good consistency between the ascent and the descent profiles can be noticed. The mean difference between the ascent and the descent time along with the standard deviation shown in the same figure and Table 5 reveals a very low mean difference with maximum standard deviation of 2 K. Detail of differences between the descent and the immediate ascent data for individual profiles are shown in Table 6. Note that this mean difference and standard deviations are small when compared to the ascent and the descent data of the same sounding mentioned in the last sub-section. However, from this figure it is seen that descent temperature data is lower than the immediate ascent data.

#### 4.6 Validation of descent data with immediate balloon ascent data: day and night differences

It is further desirable to check the consistency in the ascent and descent data between the day and night. Here we take again TTD campaign data conducted during

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



21–24 July 2011 to compare the temperature difference between the descent and the immediate ascent profile during daytime and night time separately. Figure 11 shows profile of temperature obtained during TTD campaign launched every 3 h during 21–24 July 2011 separately for the profiles obtained during day and night times. During day time the temperature during the descent is observed to be less than the immediate ascent profile as expected. However, no such systematic difference during night time is noticed.

### 4.7 Background meteorological conditions constructed using descent data

From the above, it can be concluded that descent profiles from the radiosonde observations are reliable and provide useful information similar to that of the profiles obtained from the ascent data. The profiles of all the meteorological parameters are obtained during the descent phase for each of the ascent profiles and the monthly mean  $T$ , RH,  $U$  and  $V$  obtained during these descent profiles are shown in Fig. 12. The major features of the ascent data seen in Fig. 2 can be noticed in Fig. 12 as well except perhaps in RH. This may be due to slower response of RH.

## 5 Summary and conclusions

A systematic study has been made to assess the quality of the radiosonde descent data (without attaching parachute) obtained from high resolution GPS radiosonde launched from a tropical station Gadanki during January 2008 to December 2011 for the first time. The main conclusions drawn from the present study are the following.

1. About 98.2 % of the total balloons reached the altitude of tropical tropopause (around 16 km) and 92.6 % and 85.5 % reached the altitude of greater than 20 km and 25 km, respectively. The number of balloons reaching the maximum altitude is lowest in the pre-monsoon season and highest in the monsoon season.



**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2. More than 50 % of the balloons reached (in the descent) down to 5 km. Number of balloons reaching close to the surface (0–1 km) are high in the pre-monsoon season and least in the monsoon season.
3. In general, radiosonde has drifted maximum during the monsoon season mainly due to large winds observed due to prevailing tropical easterly jet during this season. The drift in the winter season is minimum and the maximum distance it has drifted during this season is only 20 km. Mean drifts during descent is less in all seasons i.e., 20 km from balloon burst location.
4. It roughly takes one and half hours to two hours to reach maximum altitude of 30–35 km in the ascent. It is interesting to see that about 70 % of the balloons had taken more than 30 min to reach the ground/near surface from the maximum altitude. This time is matching well with estimated drag and terminal velocity. As quite often it is taking more than half an hour for descent it appears that the bursted balloon itself is acting like a kind of parachute. Moreover the descent rate while it is close to ground is only  $10 \text{ ms}^{-1}$  (mean is around  $20 \text{ ms}^{-1}$ ) which is allowing us to construct one more profile. Note that no parachute is attached to the balloon. The horizontal winds,  $T$ , and RH are measured with an accuracy of  $0.15 \text{ ms}^{-1}$ , 0.5 K and 5 %, respectively, during ascent. These parameters exhibit significant diurnal variation. It is very interesting to see that the mean difference in  $T$  between ascent and descent is less than 0.5 K below 30 km except near tropopause altitude during monsoon season where it reaches 1 K with standard deviation of 2 K. Above 30 km mean difference reaches 2 K with standard deviation of 4–6 K.
5. The mean RH difference between ascent and descent is less than 10 % with standard deviation of 20–25 % below 12 km. The mean difference in  $U$  and  $V$  between ascent and descent is less than  $1 \text{ ms}^{-1}$  with standard deviation of about  $5 \text{ ms}^{-1}$ .

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

6. The above obtained results are further confirmed using the immediate ascent profile obtained from TTD campaign.

7. The radiosonde descent profile provides reliable data of the meteorological parameters and can be used for scientific studies.

5 Thus one more reliable profile of meteorological profiles can be constructed with in 3 h of time of balloon launch and can be used for scientific purposes. It is suggested to record the descent data whenever and wherever this type of radiosondes are launched across the globe so that one more profile of meteorological parameters can be obtained at practically NO ADDITIONAL COST. Various studies like boundary layer processes, detection of zero degree isotherm, tropopause characteristics, turbulence (Clayson, and Kantha, 2008), and estimation of instability indices using descent data will be presented in a separate communication. However, some caution is advised while using the descent data. Note that descent rate is quite high ( $50\text{--}60\text{ ms}^{-1}$ ) immediate after balloon burst and it takes some time to stabilize. Thus, the data within a few kilometers from the balloon burst may be biased because of improper sampling.

*Acknowledgements.* We wish to thank NARL staff for launching radiosonde every day which is the main source for the preset study.

## References

- 20 Basha, G. and Ratnam, M. V.: Identification of atmospheric boundary layer height over a tropical station using high-resolution radiosonde refractivity profiles: comparison with GPS radio occultation measurements, *J. Geophys. Res.*, 114, D16101, doi:10.1029/2008JD011692, 2009.
- 25 Chen, H., Zhu, Y., Zhang, J., and Xuan, Y.: Vertical air motions derived from a descending radiosonde using a lightweight hard ball as the parachute, *Atmos. Meas. Tech. Discuss.*, 6, 8107–8127, doi:10.5194/amtd-6-8107-2013, 2013.

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Clayson, C. A. and Kantha, L.: On turbulence and mixing in the free atmosphere inferred from high resolution data, *J. Atmos. Ocean. Technol.*, 25, 833–852, doi:10.1175/2007JTECHA992.1, 2008.

Debashis Nath, Venkat Ratnam, M., Jagannadha Rao, V. V. M., Krishna Murthy, B. V., and Vijaya Bhaskara Rao, S.: Gravity wave characteristics observed over a tropical station using high-resolution GPS radiosonde soundings, *J. Geophys. Res.*, 114, D06117, doi:10.1029/2008JD011056, 2009.

Durre, I., Vose, R. S., Russell, S., and Wuertz, D. B.: Overview of the integrated global radiosonde archive, *J. Climate*, 19, 53–68, 2006.

Houchi, K., Stoffelen, A., Marseille, G. J., and de Kloe, J.: Comparison of wind and wind-shear climatologies derived from high-resolution radiosondes and the ECMWF model, *J. Geophys. Res.*, 115, D22123, doi:10.1029/2009JD013196, 2010.

Immler, F. J., Dykema, J., Gardiner, T., Whiteman, D. N., Thorne, P. W., and Vömel, H.: Reference Quality Upper-Air Measurements: guidance for developing GRUAN data products, *Atmos. Meas. Tech.*, 3, 1217–1231, doi:10.5194/amt-3-1217-2010, 2010.

Johansson, C. and Bergström, H.: An auxiliary tool to determine the height of the boundary layer, *Bound.-Lay. Meteorol.*, 115, 423–432, 2005, 2005.

MacPherson, B.: Radiosonde balloon drift – does it matter for data assimilation?, *Meteorol. Appl.*, 2, 301–305, doi:10.1002/met.5060020402, 1995.

McGrath, R., Semmler, T., Sweeney, C., and Wang, S.: Impact of balloon drift errors in radiosonde data on climate statistics, *J. Climate*, 19, 3430–3442, doi:10.1175/JCLI3804.1, 2006.

Miloshevich, L. M., Vömel, H., Whiteman, D. N., and Leblanc, T.: Accuracy assessment and correction of Vaisala RS92 radiosonde water vapor measurements, *J. Geophys. Res.*, 114, D11305, doi:10.1029/2008JD011565, 2009.

Ratnam, M. V., Narendra Babu, A., Jagannadha Rao, V. V. M., Vijaya bhaskar Rao, S., and Narayana Rao, D.: Gravity wave climatology observed over a tropical station using Indian MST radar: source mechanisms, *J. Geophys. Res.*, 113, D07109, doi:10.1029/2007JD008986, 2008.

Ratnam, M. V., Sunilkumar, S. V., Parameswaran, K., Krishna Murthy, B. V., Geetha Ramkumar, Rajeev, K., Ghouse Basha., Ravindra Babu, S., Muhsin, M., Manoj Kumar Mishra., Hemanth Kumar, A., Akhil Raj, S. T., and Pramitha, M.: Tropical Tropopause Dynamics (TTD)

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Campaigns over Indian region: Overview and Preliminary results, *J. Atmos. Sol. Terr. Phys.*, communicated, 2013.

Roja Raman, M., Jagannadha Rao, V. V. M., Venkat, M., Ratnam., M., Rajeevan, B., Rao, S. V., Narayana Rao, D., and Prabhakara Rao, N.: Characteristics of the Tropical Easterly Jet: Long-term trends and their features during active and break monsoon phases, *J. Geophys. Res.*, 114, D19105, doi:10.1029/2009JD012065, 2009.

Seidel, D. J., Sun, B., Pettey, M., and Reale, A.: Global radiosonde balloon drift statistics, *J. Geophys. Res.*, 116, D07102, doi:10.1029/2010JD014891, 2011.

Soddell, J. R., Kendal McGuffie, G., and Holland, J.: Inter-comparison of atmospheric soundings from the aerosonde and radiosonde, *J. Appl. Meteor.*, 43, 1260–1269, 2004.

Sun, B., Reale, A., Seidel, D. J., and Hunt, D. C.: Comparing radiosonde and COSMIC atmospheric profile data to quantify differences among radiosonde types and the effects of imperfect collocation on comparison statistics, *J. Geophys. Res.*, 115, D23104, doi:10.1029/2010JD014457, 2010.

WMO: WMO inter-comparison of high quality radiosonde systems, Instruments and observing methods, Report No. 107, WMO/TD-No. 1580, 2011.

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

**Table 1.** Statistics of number of balloons reaching the minimum and maximum altitudes while descending and ascending time, respectively, during different seasons together with statistics of 4 yr.

| Minimum altitude | Winter (256) | Pre-monsoon (234) | Monsoon (218) | Post-monsoon (216) | 4 yr (924)   |
|------------------|--------------|-------------------|---------------|--------------------|--------------|
| 0–1 km           | 33 (12.89%)  | 49 (20.94%)       | 2 (0.92%)     | 24 (11.11%)        | 108 (11.69%) |
| 1–2 km           | 79 (30.86%)  | 110 (47.01%)      | 13 (5.96%)    | 66 (30.55%)        | 268 (29%)    |
| 2–3 km           | 108 (42.19%) | 130 (55.56%)      | 45 (20.64%)   | 94 (43.52%)        | 377 (40.80%) |
| 3–4 km           | 126 (49.22%) | 137 (58.55%)      | 69 (31.65%)   | 108 (50.0%)        | 440 (47.62%) |
| 4–5 km           | 131 (51.17%) | 140 (59.83%)      | 78 (35.78%)   | 114 (52.78%)       | 465 (50.32%) |
| 5–10 km          | 149 (58.20%) | 158 (67.52%)      | 106 (48.62%)  | 129 (59.72%)       | 542 (58.66%) |
| Maximum altitude |              |                   |               |                    |              |
| 10–15 km         | 255 (99.61%) | 231 (98.72%)      | 218 (100%)    | 214 (99.07%)       | 918 (99.35%) |
| 15–20 km         | 253 (98.83%) | 231 (98.72%)      | 218 (100%)    | 214 (99.07%)       | 916 (99.13%) |
| 20–25 km         | 242 (94.53%) | 223 (95.30%)      | 208 (95.41%)  | 168 (77.77%)       | 841 (91.02%) |
| 25–30 km         | 224 (87.5%)  | 209 (89.32%)      | 186 (85.32%)  | 138 (63.89%)       | 757 (81.93%) |
| 30–35 km         | 136 (53.12%) | 152 (64.96%)      | 131 (60.09%)  | 9 (41.20%)         | 508 (54.98%) |
| 35–40 km         | 11 (4.30%)   | 7 (2.99%)         | 41 (18.81%)   | 14 (6.48%)         | 73 (7.90%)   |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

**Table 2.** The mean time (hours) along with standard deviation taken for the balloons to reach from maximum altitude to the minimum altitudes during different seasons along with 4 yr mean descent timings.

| Altitude    | Winter      | Pre-Monsoon | Monsoon     | Post-Monsoon | 4 yr        |
|-------------|-------------|-------------|-------------|--------------|-------------|
| 30 to 10 km | 0.21 ± 0.05 | 0.22 ± 0.05 | 0.27 ± 0.08 | 0.24 ± 0.06  | 0.23 ± 0.06 |
| 25 to 10 km | 0.21 ± 0.07 | 0.21 ± 0.06 | 0.26 ± 0.08 | 0.24 ± 0.1   | 0.22 ± 0.07 |
| 20 to 10 km | 0.16 ± 0.05 | 0.17 ± 0.06 | 0.21 ± 0.07 | 0.2 ± 0.09   | 0.18 ± 0.07 |
| 15 to 10 km | 0.09 ± 0.02 | 0.09 ± 0.03 | 0.11 ± 0.03 | 0.11 ± 0.05  | 0.10 ± 0.03 |
| 30 to 8 km  | 0.25 ± 0.06 | 0.26 ± 0.05 | 0.32 ± 0.08 | 0.28 ± 0.06  | 0.27 ± 0.07 |
| 25 to 8 km  | 0.25 ± 0.07 | 0.25 ± 0.07 | 0.31 ± 0.09 | 0.28 ± 0.1   | 0.27 ± 0.08 |
| 20 to 8 km  | 0.21 ± 0.06 | 0.21 ± 0.07 | 0.26 ± 0.08 | 0.26 ± 0.12  | 0.22 ± 0.07 |
| 15 to 8 km  | 0.13 ± 0.03 | 0.14 ± 0.04 | 0.16 ± 0.04 | 0.17 ± 0.08  | 0.14 ± 0.04 |
| 30 to 6 km  | 0.29 ± 0.06 | 0.3 ± 0.05  | 0.35 ± 0.08 | 0.33 ± 0.07  | 0.31 ± 0.07 |
| 25 to 6 km  | 0.3 ± 0.08  | 0.29 ± 0.07 | 0.35 ± 0.09 | 0.33 ± 0.1   | 0.31 ± 0.08 |
| 20 to 6 km  | 0.25 ± 0.07 | 0.25 ± 0.07 | 0.31 ± 0.09 | 0.29 ± 0.1   | 0.27 ± 0.08 |
| 15 to 6 km  | 0.18 ± 0.04 | 0.18 ± 0.04 | 0.2 ± 0.05  | 0.2 ± 0.06   | 0.19 ± 0.05 |
| 30 to 4 km  | 0.34 ± 0.06 | 0.35 ± 0.06 | 0.38 ± 0.07 | 0.38 ± 0.07  | 0.35 ± 0.07 |
| 25 to 4 km  | 0.35 ± 0.09 | 0.34 ± 0.08 | 0.4 ± 0.1   | 0.39 ± 0.11  | 0.36 ± 0.09 |
| 20 to 4 km  | 0.3 ± 0.08  | 0.3 ± 0.08  | 0.36 ± 0.1  | 0.35 ± 0.11  | 0.32 ± 0.09 |
| 15 to 4 km  | 0.23 ± 0.05 | 0.23 ± 0.05 | 0.26 ± 0.06 | 0.26 ± 0.07  | 0.24 ± 0.06 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

**Table 3.** Mean difference between ascent and descent observed in all the parameters at different altitudes in different seasons.

| Temperature (K) Humidity (%) |              |              |              |              |               |               |               |               |
|------------------------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|---------------|
| Altitude                     | Winter       | Pre-Monsoon  | Monsoon      | Post-Monsoon | Winter        | Pre-Monsoon   | Monsoon       | Post-Monsoon  |
| 5 km                         | 0.43 ± 0.78  | 0.42 ± 0.85  | 0.16 ± 1.03  | 0.37 ± 0.82  | -2.84 ± 10.62 | -6.66 ± 17.03 | -4.69 ± 18.86 | -5.52 ± 14.04 |
| 10 km                        | 0.40 ± 0.52  | 0.25 ± 0.61  | 0.040.76     | 0.16 ± 0.62  | -7.57 ± 9.44  | -8.92 ± 15.61 | -8.60 ± 14.52 | -7.39 ± 15.44 |
| 15 km                        | 0.16 ± 1.07  | 0.30 ± 1.16  | -0.28 ± 1.68 | -0.01 ± 1.10 | 15.17 ± 8.02  | 16.37 ± 10.73 | 24.18 ± 14.52 | 17.77 ± 12.47 |
| 20 km                        | 0.38 ± 1.25  | 0.67 ± 1.72  | 0.12 ± 2.04  | 0.09 ± 1.42  | -             | -             | -             | -             |
| 25 km                        | -0.12 ± 1.21 | 0.43 ± 1.75  | 0.40 ± 1.61  | 0.04 ± 1.03  | -             | -             | -             | -             |
| 30 km                        | -1.71 ± 2.24 | -1.18 ± 2.43 | -0.44 ± 2.21 | -0.91 ± 1.26 | -             | -             | -             | -             |
| 35 km                        | -1.34 ± 2.38 | -1.78 ± 2.28 | -1.80 ± 2.51 | -0.49 ± 3.17 | -             | -             | -             | -             |

| $U$ ( $\text{ms}^{-1}$ ) $V$ ( $\text{ms}^{-1}$ ) |              |              |              |              |              |              |              |              |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Altitude  | Winter       | Pre-Monsoon  | Monsoon      | Post-Monsoon | Winter       | Pre-Monsoon  | Monsoon      | Post-Monsoon |
| 5 km  | 0.51 ± 1.90  | 0.27 ± 2.17  | 1.29 ± 4.47  | 0.40 ± 1.97  | 0.03 ± 2.94  | 0.36 ± 2.22  | -0.66 ± 2.61 | 0.04 ± 1.81  |
| 10 km   | 0.04 ± 2.11  | 0.10 ± 2.25  | 0.97 ± 5.46  | -0.15 ± 2.79 | -0.26 ± 2.80 | -0.27 ± 2.84 | 0.38 ± 3.42  | -0.20 ± 2.55 |
| 15 km   | -0.33 ± 3.93 | -0.66 ± 5.57 | -2.48 ± 6.51 | -3.01 ± 5.25 | 0.86 ± 4.61  | 0.13 ± 3.90  | -1.05 ± 4.86 | 0.73 ± 4.80  |
| 20 km   | 1.22 ± 4.41  | -0.44 ± 5.22 | -0.79 ± 5.89 | -0.89 ± 5.54 | 0.77 ± 4.20  | -0.03 ± 3.95 | 0.20 ± 3.71  | -0.25 ± 4.73 |
| 25 km   | 1.00 ± 4.69  | -0.02 ± 4.83 | -1.26 ± 5.45 | -1.74 ± 4.95 | 0.47 ± 3.75  | 0.08 ± 3.56  | 0.11 ± 4.37  | -0.06 ± 3.88 |
| 30 km   | -0.22 ± 5.08 | -0.61 ± 4.06 | -0.51 ± 4.62 | -1.82 ± 5.23 | -0.46 ± 4.25 | 0.20 ± 4.07  | 0.42 ± 4.36  | -1.10 ± 3.12 |
| 35 km   | 0.09 ± 2.67  | 0.79 ± 7.92  | 0.70 ± 3.74  | 0.15 ± 3.41  | 0.47 ± 3.55  | -0.07 ± 4.80 | 1.05 ± 3.96  | -0.53 ± 4.64 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

**Table 4.** Mean difference along with standard deviation observed in all the parameters between ascent and descent at different altitudes obtained during January 2008 to December 2011.

| 4 yr  | $T$ (K)          | RH (%)            | $U$ ( $\text{ms}^{-1}$ ) | $V$ ( $\text{ms}^{-1}$ ) |
|-------|------------------|-------------------|--------------------------|--------------------------|
| 5 km  | $0.37 \pm 0.86$  | $-4.98 \pm 15.15$ | $0.54 \pm 2.62$          | $0.01 \pm 2.44$          |
| 10 km | $0.23 \pm 0.63$  | $-8.12 \pm 13.90$ | $0.20 \pm 3.24$          | $-0.12 \pm 2.89$         |
| 15 km | $0.07 \pm 1.26$  | $18.00 \pm 11.88$ | $-1.51 \pm 5.44$         | $0.22 \pm 4.56$          |
| 20 km | $0.35 \pm 1.64$  | –                 | $-0.13 \pm 5.29$         | $0.21 \pm 4.14$          |
| 25 km | $0.18 \pm 1.47$  | –                 | $-0.28 \pm 5.05$         | $0.18 \pm 3.86$          |
| 30 km | $-1.13 \pm 2.20$ | –                 | $-0.68 \pm 4.72$         | $-0.17 \pm 4.07$         |
| 35 km | $-1.45 \pm 2.59$ | –                 | $0.47 \pm 4.08$          | $0.49 \pm 4.07$          |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

**Table 5.** Mean difference (descent – ascent) in temperature between descent profiles and immediate ascent profiles observed during 28–31 December 2010.

| Altitude region | Difference in $T$ |
|-----------------|-------------------|
| 1–5 km          | $-1.12 \pm 0.29$  |
| 5–10 km         | $-0.99 \pm 0.21$  |
| 10–15 km        | $-0.81 \pm 0.14$  |
| 15–20 km        | $-1.04 \pm 0.41$  |
| 20–25 km        | $-0.62 \pm 0.39$  |
| 25–30 km        | $-0.34 \pm 0.50$  |
| 30–35 km        | $1.73 \pm 0.70$   |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 6.** Mean temperature difference between descent profile and immediate ascent profile below and above the tropopause altitude.

| Date                  | 0–18 km          | 18-maximum altitude |
|-----------------------|------------------|---------------------|
| 28 Dec 2010, 11:00 LT | $-1.42 \pm 1.24$ | $-0.51 \pm 1.75$    |
| 28 Dec 2010, 14:00 LT | $-0.66 \pm 1.18$ | $-1.44 \pm 1.12$    |
| 28 Dec 2010, 20:00 LT | $-0.01 \pm 1.04$ | $-0.94 \pm 1.54$    |
| 29 Dec 2010, 08:00 LT | –                | $2.2 \pm 0$         |
| 29 Dec 2010, 11:00 LT | $-1.23 \pm 1.09$ | $-0.24 \pm 1.19$    |
| 29 Dec 2010, 14:00 LT | $-2.07 \pm 0.84$ | $-1.16 \pm 1.34$    |
| 29 Dec 2010, 17:00 LT | $-0.30 \pm 0.90$ | $-0.13 \pm 1.37$    |
| 29 Dec 2010, 20:00 LT | $-0.23 \pm 0.65$ | $-0.05 \pm 1.18$    |
| 30 Dec 2010, 02:00 LT | $-0.15 \pm 0.70$ | $0.53 \pm 1.09$     |
| 30 Dec 2010, 05:00 LT | $-0.74 \pm 0.89$ | $-0.65 \pm 1.49$    |
| 30 Dec 2010, 08:00 LT | $-2.52 \pm 0.80$ | $-0.95 \pm 1.31$    |
| 30 Dec 2010, 11:00 LT | $-0.82 \pm 1.09$ | $-0.72 \pm 2.13$    |
| 30 Dec 2010, 14:00 LT | $-2.08 \pm 0.59$ | $-1.11 \pm 2.19$    |
| 30 Dec 2010, 17:00 LT | $-0.26 \pm 0.60$ | –                   |
| 30 Dec 2010, 20:00 LT | $-0.16 \pm 1.45$ | –                   |
| 31 Dec 2010, 05:00 LT | $-1.51 \pm 1.00$ | $-1.59 \pm 1.47$    |

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.




**Fig. 1. (a)** Balloon burst altitude (ascent) and the minimum altitude (in descent) of all the radiosondes launched from Gadanki during 1 January 2008 to 31 December 2011. **(b)** Number of balloons reaching maximum and minimum altitudes while in ascent and in descent, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.




**Fig. 2.** Time – altitude section of monthly mean (a)  $T$ , (b) RH, (c)  $U$  and (d)  $V$  observed over Gadanki using GPS radiosonde observations during January 2008 to December 2011 using balloon ascent data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Assessment of GPS radiosonde descent data


M. Venkat Ratnam et al.



**Fig. 3.** Monthly mean drifts of the radiosonde from Gadanki location during ascent time observed during **(a)** Winter, **(b)** Pre-monsoon, **(c)** Monsoon and **(d)** Post-monsoon seasons.

**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.




**Fig. 4.** The locations of the balloons reaching the maximum altitude for all the radiosonde launched from Gadanki (shown with start mark) during January 2008 to December 2011 separated according to the season.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.



**Fig. 5.** Histograms showing the time taken for the balloons to reach **(a)** maximum and **(b)** minimum altitudes during different seasons along with 4 yr mean ascent and descent timings. **(c)** Mean ascent and descent rates along with standard deviations obtained while averaging over 4 yr.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Assessment of GPS  
radiosonde descent  
data**

M. Venkat Ratnam et al.




**Fig. 6.** (a) Mean profiles of density along with standard deviation observed over Gadanki averaged over four years (2008–2011). (b) Mean terminal velocity calculated for radiosonde edge and non-edge (see text for details) sections along with standard deviations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.




**Fig. 7.** (First panels) Typical profiles of  $T$ , RH,  $U$ ,  $V$ , and ascent rate observed on 1 January 2008 at 12:00 UT at Gadanki. The profiles obtained during descent are also superimposed in respective panels. (Second panels) The difference observed in  $T$ , RH,  $U$ ,  $V$ , and ascent rate and descent rate between ascent and descent phase. (Third panels) Histograms showing the variability in the difference of  $T$ , RH,  $U$ ,  $V$ , and ascent rate between ascent and descent profiles. (Fourth panels) Same as third panels but for complete 4 yr period (January 2008 to December 2011).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.



**Fig. 8.** Diurnal variation observed seasonally during December 2010 to May 2012 in (a)  $T$ , (b) RH, (c)  $U$ , and (d)  $V$ . The vertical lines are drawn at 17:00LT and 20:00LT to highlight the variability during ascent and descent times.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Assessment of GPS radiosonde descent data


M. Venkat Ratnam et al.



**Fig. 9.** Mean difference in  $T$ ,  $RH$ ,  $U$  and  $V$  between ascent and descent times observed during different seasons along with 4 yr mean difference. Horizontal bars represent the standard deviation.

**Assessment of GPS  
radiosonde descent  
data**


M. Venkat Ratnam et al.



**Fig. 10.** (Left panel) Profiles of temperature obtained during Tropical Tropopause Dynamics campaign launched every 3 h during 28–31 December 2010. Descent profiles are also superimposed in same panel. (Right panel) Mean difference between descent and immediate ascent profiles along with standard deviation.

## Assessment of GPS radiosonde descent data


M. Venkat Ratnam et al.



**Fig. 11.** (Left panels) Profiles of temperature obtained during tropical tropopause dynamics campaign launched during daytime (top panel) and night time (bottom panel) every 3 h during 21–24 July 2011. Descent profiles are also superimposed in same panels. Ascent and descent cases are abbreviated as A and D, respectively. (Right panels) Mean difference between descent and immediate ascent profiles along with standard deviation.

## Assessment of GPS radiosonde descent data

M. Venkat Ratnam et al.



**Fig. 12.** Time – altitude section of monthly mean during descending **(a)**  $T$ , **(b)** RH, **(c)**  $U$  and **(d)**  $V$  observed over Gadanki using GPS Radiosonde observations during January 2008 to December 2011 using balloon descent data.