

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

SI²N overview paper: ozone profile measurements: techniques, uncertainties and availability

B. Hassler^{1,2}, I. Petropavlovskikh^{1,3}, J. Staehelin⁴, T. August⁵, P. K. Bhartia⁶, C. Clerbaux⁷, D. Degenstein⁸, M. De Mazière⁹, B. M. Dinelli¹⁰, A. Dudhia¹¹, G. Dufour¹², S. M. Frith¹³, L. Froidevaux¹⁴, S. Godin-Beekmann¹⁵, J. Granville⁹, N. R. P. Harris¹⁶, K. Hoppel¹⁷, D. Hubert⁹, Y. Kasai¹⁸, M. J. Kurylo¹⁹, E. Kyrölä²⁰, J.-C. Lambert⁹, P. F. Levelt²¹, C. T. McElroy²², R. D. McPeters⁶, R. Munro⁵, H. Nakajima²³, A. Parrish²⁴, P. Raspollini²⁵, E. E. Remsberg²⁶, K. H. Rosenlof², A. Rozanov²⁷, T. Sano²⁸, Y. Sasano²⁹, M. Shiotani²⁸, H. G. J. Smit³⁰, G. Stiller³¹, J. Tamminen²⁰, D. W. Tarasick³², J. Urban³³, R. J. van der A²¹, J. P. Veefkind²¹, C. Vigouroux⁹, T. von Clarmann³¹, C. von Savigny³⁴, K. A. Walker^{35,36}, M. Weber²⁷, J. Wild³⁷, and J. Zawodny²⁶

¹CIRES, University of Colorado at Boulder, Boulder, Colorado, USA

²NOAA/ESRL, Chemical Sciences Division, Boulder, Colorado, USA

³NOAA/ESRL, Global Monitoring Division, Boulder, Colorado, USA

⁴ETH-Zürich, Zürich, Switzerland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁵EUMETSAT, Darmstadt, Germany

⁶NASA Goddard Space Flight Center, Greenbelt, MD, USA

⁷UPMC Univ. Paris 06, Université Versailles St-Quentin, CNRS/INSU, LATMOS-IPSL, Paris, France

⁸University of Saskatchewan, Saskatoon, Saskatchewan, Canada

⁹Belgian Institute for Space Aeronomy (IASB-BIRA), Brussels, Belgium

¹⁰ISAC-CNR, Bologna, Italy

¹¹AOPP, Physic Department, University of Oxford, Oxford, UK

¹²LISA, UMR CNRS 7583, Université Paris-Est Créteil et Université Paris-Diderot, Créteil, France

¹³Science Systems and Applications, Inc., NASA GSFC, Greenbelt, Maryland, USA

¹⁴Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

¹⁵Observatoire de Versailles Saint-Quentin-en-Yvelines, Guyancourt Cedex, France

¹⁶University of Cambridge, Chemistry Department, Cambridge, UK

¹⁷Remote Sensing Division, Naval Research Laboratory, Washington, DC, USA

¹⁸National Institute of Information and Communications Technology, Tokyo, Japan

¹⁹Universities Space Research Association, Goddard Earth Sciences, Technology, and Research, NASA Goddard Space Flight Center, Greenbelt, MD, USA

²⁰Finnish Meteorological Institute, Earth Observation, Helsinki, Finland

²¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, the Netherlands

²²Department of Earth and Space Science and Engineering (Lassonde School of Engineering), York University, Toronto, Canada

²³National Institute for Environmental Studies, Tsukuba, Japan

²⁴Department of Astronomy, University of Massachusetts, Amherst, MA, USA

²⁵Istituto di Fisica Applicata “N. Carrara” (IFAC) del Consiglio Nazionale delle Ricerche (CNR), Florence, Italy

²⁶NASA Langley Research Center, Hampton, Virginia, USA

²⁷Institute of Environmental Physics and Remote Sensing (IUP/IFE), University of Bremen, Bremen, Germany

²⁸Research Institute for Sustainable Humanosphere, Kyoto University, Kyoto, Japan

²⁹Association of International Research Initiatives for Environmental Studies, Tokyo, Japan

³⁰ Forschungszentrum Jülich, Institut für Chemie der belasteten Atmosphäre (ICG2), Jülich, Germany

³¹ Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, Karlsruhe, Germany

³² Environment Canada, Downsview, Ontario, Canada

³³ Chalmers University of Technology, Department of Earth and Space Sciences, Göteborg, Sweden

³⁴ Institut für Physik, Ernst-Moritz-Arndt-Universität Greifswald, Greifswald, Germany

³⁵ Department of Physics, University of Toronto, Toronto, Ontario, Canada

³⁶ Department of Chemistry, University of Waterloo, Waterloo, Ontario, Canada

³⁷ Innovim, LLC, NOAA/NWS/NCEP/Climate Prediction Center, College Park, Maryland, USA

Received: 21 October 2013 – Accepted: 24 October 2013 – Published: 14 November 2013

Correspondence to: B. Hassler (birgit.hassler@noaa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Peak stratospheric chlorofluorocarbon (CFC) and other ozone depleting substance (ODS) concentrations were reached in the mid to late 1990s. Detection and attribution of the expected recovery of the stratospheric ozone layer in an atmosphere with reduced ODSs as well as efforts to understand the evolution of stratospheric ozone in the presence of increasing greenhouse gases are key current research topics. These require a critical examination of the ozone changes with an accurate knowledge of the spatial (geographical and vertical) and temporal ozone response. For such an examination, it is vital that the quality of the measurements used be as high as possible and measurement uncertainties well quantified.

In preparation for the 2014 United Nations Environment Programme (UNEP)/World Meteorological Organization (WMO) Scientific Assessment of Ozone Depletion, the SPARC/IO₃C/IGACO-O₃/NDACC (SI²N) initiative was designed to study and document changes in the global ozone profile distribution. This requires assessing long-term ozone profile data sets in regards to measurement stability and uncertainty characteristics. The ultimate goal is to establish suitability for estimating long-term ozone trends to contribute to ozone recovery studies. Some of the data sets have been improved as part of this initiative with updated versions now available.

This summary presents an overview of stratospheric ozone profile measurement data sets (ground- and satellite-based) available for ozone recovery studies. Here we document measurement techniques, spatial and temporal coverage, vertical resolution, native units and measurement uncertainties. In addition, the latest data versions are briefly described (including data version updates as well as detailing multiple retrievals when available for a given satellite instrument). Archive location information is for each data set is also given.

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

As man-made ozone depleting substances (ODSs) decline in the stratosphere in response to the 1987 Montreal Protocol and its subsequent Amendments and Adjustments, the ozone layer is expected to recover globally. A small increase in stratospheric ozone abundance over Northern mid-latitudes was first reported in the 2010 WMO Ozone assessment (WMO, 2011). Although supported by a variety of ground-based, balloon-born and satellite observations, statistical analysis could not attribute this change exclusively to the decline of ODSs due to observational uncertainty, geophysical and dynamical variability and changes in ozone due to increasing greenhouse gases that cool the stratosphere. Global total ozone remained significantly lower than during the early 1980s, and surface ultraviolet radiation measurements were too variable to strongly support ozone recovery. Indications of the onset of stratospheric ozone recovery have been noted since the publication of the last ozone assessment. There are several recent publications (e.g. Maeder et al., 2010; Nair et al., 2013; Kuttippurath et al., 2013) supporting the statement that the second stage of ozone recovery is ongoing (as defined by the WMO, 2007 statement that “occurrence of statistically significant increases in ozone above previous minimum values due to declining equivalent effective stratospheric chlorine”) other publications show that ozone recovery in Antarctica should be detectable within the next decade (Hassler et al., 2011). The additional four years of data available since analysis was done for the 2010 WMO Ozone assessment as well as improvements in the homogenization of ozone data records should improve the signal to noise in the ozone time series making trend analyses results more robust.

“SPARC Report No.1 – Trends in the vertical distribution of ozone” (SPARC, 1998) provided a comprehensive summary of available data and methods for trend analysis at that time. Its main objectives were to understand the limitations of the data, to assess the consistency between various data records and create a reference database for the analysis of long-term ozone changes as a function of altitude. One of the main reasons

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for this painstaking approach was to avoid misinterpreting instrument drifts as actual ozone changes.

Trend analysis is made more complex by the addition of new satellite records (e.g. Aura, Envisat, Odin, and SCISAT) where each covers a shorter time period than the traditionally used long-term records (e.g. ozonesondes, Umkehr, lidars, SAGE, HALOE). Adding to the problem, some of the long-term platforms have ceased taking measurements (e.g. SAGE, HALOE, instruments on Envisat), while records from other satellite platforms (NOAA/2) are difficult to combine due to drifting orbits and different instrument generations (i.e. NIMBUS or SBUV retrievals from Aura OMI). For a full long-term analysis of stratospheric ozone throughout the decline due to increases in ODSs and subsequent recovery, multiple instrument records need to be considered. Thus, several different data sources have to be merged to allow a complete view of past ozone changes. Recent work has shown that ozone depletion and greenhouse gas (GHG) increases impact dynamical quantities in an additive manner, with ozone recovery possibly cancelling atmospheric circulation effects associated with increasing GHGs. (e.g. IPCC/TEAP, 2005; McLandress et al., 2010; Polvani et al., 2011). Clearly, an understanding of atmospheric transport, circulation and wave breaking patterns is important for our understanding of ozone recovery.

The consideration of uncertainties and artifacts is crucial in ozone trend analyses, particularly when the ozone recovery signal is small as compared to natural geophysical variability. As methods for trend and attribution analysis become more advanced, the uncertainties and instrumental artifacts buried in currently available datasets start to play a bigger role. Therefore, detailed information about measurement uncertainties, data jumps due to changes in instrumentation and drifts during long-term operation of a given instrument are essential.

There have been several attempts to produce historical global ozone profile data sets; these include satellite-ozonesonde combinations starting in 1979 when the first global satellite measurements became available (e.g. Randel and Wu, 2007; Bodeker et al., 2013), global ozone data sets that extend values to 2100 using model and fit-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ting techniques (Hassler et al., 2009; Cionni et al., 2011) for use in global climate models without stratospheric ozone chemistry and also data sets that extend the historical record backward to 1900 (Brönnimann et al., 2013). Issues of biases between instruments, sampling, noise, gaps and differences in vertical registration (pressure or altitude) have to be addressed when creating combined data records to ensure no artificial trends are introduced.

Uncertainties in calculated ozone trends need to be reduced. In the 2010 WMO Ozone Assessment, inconsistencies in the available ozone profile data prevented a clear overall picture of long-term ozone profile changes. A successful assessment of trends for the upcoming 2014 WMO Ozone Assessment requires addressing the problems of time series artefacts due to changes in instrumentation, homogenizing applied satellite retrievals and reduce and describing measurement uncertainties. The SPARC/IO₃C/IGACO-O₃/NDACC (SI²N) Initiative has brought together scientists dealing with ground and satellite ozone profile measurements in order to consolidate efforts documenting long-term ozone profile changes. SI²N has also defined the time frame to ensure that any dataset improvements are finalized to ensure inclusion in the 2014 WMO Ozone Assessment analysis.

This paper provides a summary of the most important available ozone datasets, describes measurement methods, detection limits and uncertainty estimates, gives details about the vertical and temporal resolution of individual datasets and provides references to relevant published work. In addition, improvements and homogenizations that have occurred to date are discussed. This is the first of three overview papers discussing the results from the SI²N Initiative. The second paper describes the validation of the different available ozone profile measurements (Lambert et al., 2013) and the third paper provides information and an overview about the merging and homogenizing of different data sets and calculation of long-term trends (Harris et al., 2013). Ground-based ozone profiles, ordered according to the number of available stations, are described in Sect. 2. In Sect. 3, satellite-based ozone profiles are covered. The satellite measurements are organized by measurement techniques; these include

solar-/stellar-occultation measurements, limb measurements and nadir measurements. A summary, including discussion of possible temporal and spatial coverage problems for the measurement systems, is presented in Sect. 4.

2 Ground-based measurement systems

5 The global map with location of the ground-based instrument systems (ozonesonde, Dobson/Brewer, lidar, FTIR and microwave) is shown in Fig. 1. Ozonesondes and Dobson/Brewer instruments provide the longest record of stratospheric ozone variability (see Fig. 2). Ozonesondes are all weather sounders, and can measure at night and through clouds. Their data quality is good in the lower stratosphere and troposphere
10 where satellites measurement quality is less reliable, and their vertical resolution is much higher than is achievable from remotely sensed measurement. Ozonesondes are inexpensive and easily deployable. Dobson Umkehr measurements have captured evolution of the upper stratospheric ozone prior to the satellite era and still remain the most inexpensive measurement system requiring minimal manpower. Umkehr measurements with Brewer instruments were added later, but they offer fully automated and remote-controlled operations, and furthermore add spectral UV measurements. Ozone lidar measurements are self-calibrated and provide high vertically resolved measurements with small uncertainties (especially for the low to mid stratosphere). Ground-based high-resolution FTIR solar absorption measurements do not require an absolute calibration either; the ozone absorption lines are very narrow and are therefore self-calibrated with the reference being the surrounding continuum. This technique has also the advantage of providing precise total columns in addition to vertical information. Ground-based microwave measurements can be made with a time resolution of an hour or less throughout a full diurnal cycle; their calibration is based on temperature,
20 the measurements are insensitive to stratospheric aerosols, and the instruments are nearly fully automated and require less personnel time and other resources to operate than most other ground-based instruments.

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 Ozonesondes

Ozonesondes are small, lightweight and compact balloon-borne instruments, developed for measuring the vertical distribution of atmospheric ozone up to an altitude of about 30–35 km (see Table 1a). During flight operation, ozonesondes are coupled with standard meteorological radiosondes providing temperature, pressure and wind measurements. The effective vertical resolution of the ozone profile, which is dependent on the balloon ascent rate, is about 100–150m (see Table 1b and Fig. 3a).

There were 66 stations contributing data in the 2000s (Fig. 1). Ozonesonde records provide the longest time series of the vertical ozone distribution throughout both troposphere and stratosphere; some station records start in the 1960s (see Fig. 2 and Table 1c). Most of the profiles are from the electrochemical concentration cell (ECC)-type ozonesonde (Komhyr, 1969) introduced in the early 1970s and adopted by most stations in the global network by the early 1980s. Almost all data in the most recent decade are from ECC sondes. Remaining data are from Brewer–Mast (B-M) sondes (Brewer and Milford, 1960), Japanese carbon iodine cell sondes (KC96) (Kobayashi and Toyama, 1996), and Indian sondes. Prior to the early 1990s, three stations in Eastern Europe flew the GDR sonde. A majority of the data before 1980 is from BM sondes or similar (both the GDR and Indian sondes are similar in design to the BM sonde). All five types use the reaction of ozone with potassium iodide in an aqueous solution as the method of ozone detection, but the instrumental layouts are significantly different (e.g., Smit, 2002; Tarasick and Slater, 2008).

When properly prepared and handled, ECC sondes yield profiles with random errors of 3–5% ($1 - \sigma$) and overall uncertainties of about 5% in the stratosphere (Kerr et al., 1994; Smit et al., 2007; Deshler et al., 2008; Liu et al., 2009). Two types of ECC sondes are in current use, with minor differences in construction and some variation in recommended concentrations of the sensing solution and of its phosphate buffer. The maximum change in stratospheric response resulting from these systematic differences is on the order of 2–3% (Smit et al., 2007). Other sonde types have somewhat

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



let part of the zenith sky spectrum. Measurements are taken when the sun elevation changes between 60° and 90° Solar Zenith Angle (SZA), and under the assumption of static atmospheric conditions and no clouds in the zenith viewing area. The vertical resolution of an Umkehr ozone profile is ≈ 10 km. Ozone profiles (in Dobson Units (DU)) are historically provided in 10 pressure layers, where pressure at the top of the layer is half of the pressure at the bottom, while the tropospheric measurement is one thick layer below 250 hPa (see Table 1a, b). The uncertainty in the retrieved ozone profile is a combination of measurement error (increases with increasing SZA) and smoothing error (about 10–20 % in the troposphere and low stratosphere and ≈ 5 % in middle and upper stratosphere).

The Umkehr record length varies by station; the longest record is from Arosa, Switzerland (starting in 1956; see Table 1c). The latest version of the ozone profile algorithm for processing of Dobson Umkehr data (UMK04) is described in Petropavlovskikh et al. (2005). UMK04 is optimized for retrievals to be used as monthly averages in long-term trend analysis and has essentially no a priori dependence. The forward model used to process Brewer Umkehr data (O3BUmkehr v.2.6, <http://www.o3soft.eu/o3bumkehr.html>) was adapted from the UMK04 model taking into account the different optical characteristics of the Brewer instrument (Petropavlovskikh et al., 2011, see Table 1d).

The Umkehr method uses a technique that minimizes systematic errors of instrumental drifts by normalizing to one of its measurements. Other methods are used to track instrument stability, such as once a month checks of wavelength registration by using a standard discharge lamps, ratio of measurements at several spectral channels (for Brewer instruments, daily), and optical wedge calibration produced during Dobson intercomparisons (every 4 yr). These methods track degradation of the optical system. Effects of ozone cross section values and uncertainties of stray light effects on Dobson and Brewer Umkehr ozone profiles are discussed in Petropavlovskikh et al. (2011) and WMO (2008). The replacement of the instrument at the station can cause a step change in the Umkehr record due to difference in the out-of-band contribution that

is specific to each instrument. Thus, additional homogenization of the record may be required (Zanis et al., 2006; Miyagawa et al., 2009). In addition, interference from optically thick aerosol loading (volcanic) at Umkehr spectral channels results in systematic errors that can be as large as 10–15%. Therefore, about two years of data are typically removed after the Pinatubo (1991) and El Chichon (1982) volcanic eruptions. Data from 28 stations with current Umkehr measurements and of 58 stations with historical Umkehr measurements are archived at WOUDC, Toronto, Canada (see Table 1d for the URL).

2.3 Lidar

Lidar (light detection and ranging) is an active, remote-sensing instrument that uses the interaction between a laser beam and atmospheric molecules and particles. Ozone lidar measurements are performed according using the Differential Absorption Lidar method (DIAL), which is based on the absorption of ultraviolet radiation by ozone and requires the emission of two laser wavelengths (so called “on” and “off” wavelengths) characterized by different ozone absorption cross-sections (Pelon et al., 1986; Godin et al., 1989; McDermid et al., 1990). The use of pulsed lasers provides range resolved measurements. Ozone DIAL systems include one or two lasers, depending on the technique used for the generation of the off wavelength, an optical receiving system that includes a telescope for the collection of the laser light, a spectrometer or filters for the separation of the received wavelengths and an electronic data acquisition system. In the case of stratospheric ozone, photon-counting is generally used for the acquisition of the lidar signal to provide high sensitivity and low noise. The ozone number density is commonly retrieved from the difference of the slopes of the logarithm of the lidar signal corrected from the background noise and saturation (for large signals originating from lower altitudes). A correction is applied corresponding to differential terms linked to molecular and aerosol scattering, and absorption by other atmospheric constituents (e.g. NO₂, SO₂).

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The DIAL technique does not require any calibration since the laser wavelengths are chosen such that the correction term represents less than 10 % of the ozone number density derived from the slope of the lidar signal in the altitude range of interest. When stratospheric aerosol loading is large, such as after cataclysmic volcanic eruptions, this condition is not fulfilled and DIAL measurements are locally perturbed at the altitude of the aerosols. In such conditions, two additional wavelengths corresponding to the first Stokes vibrational Raman scattering of the laser radiation by atmospheric nitrogen can be detected and the Raman lidar signal is used to retrieve ozone (McGee et al., 1993).

Most lidar measurements are performed during the night and averaged over several hours, resulting in a spatial resolution of about 100 km, depending on atmospheric conditions. The vertical resolution decreases as a function of altitude ranging from several hundred meters at lower altitudes to several kilometers above 40 km (Table 1a, b; Fig. 3a). The systematic uncertainty ranges from a few percent below 20 km to more than 10–15 % above 45 km (Godin-Beekmann et al., 2003). Ozone lidar measurements are self-calibrated. However, measurement stability can be affected by instrumental artifacts such as alignment error, changes in the intensity of the lidar signals due to varying laser power, or changes in weather conditions (Godin-Beekmann et al., 2003).

There are twelve stratospheric ozone lidar instruments operational globally at this point (see Fig. 1), with the longest continuous records starting in the late 1980s (Steinbrecht et al., 2009). Data are archived in the NDACC database (see Table 1d for the URL).

2.4 FTIR

Ground-based FTIR solar absorption measurements are performed over the 600–4500 cm^{-1} spectral range, under clear-sky conditions using primarily high resolution spectrometers such as the Bruker 120M (or 125M) or Bruker 120HR (or 125HR). The spectrometers can achieve a spectral resolution of 0.0035 and 0.002 cm^{-1} respectively. The advantage of the FTIR technique is that, for atmospheric gases such as ozone which have very narrow lines, an absolute calibration is not needed; the ozone

absorption signatures are self-calibrated with the reference being the surrounding continuum. Nevertheless, for the determination of vertical trends, especially in the upper stratosphere, it is important to regularly monitor the instrumental line-shape. This is achieved with gas cell measurements (Hase et al., 1999).

In addition to total columns, low-vertical-resolution profiles can be obtained from the temperature and pressure dependence of the absorption line shapes. Two profile retrieval algorithms are widely used, PROFITT9 (Hase, 2000) and SFIT2 (Pougatchev et al., 1995); both are based on the optimal estimation method (Rodgers, 2000). A European FTIR network has been optimized by using a common retrieval strategy to derive ozone trends (1995–2005) in 4 layers (Vigouroux et al., 2008), one in the troposphere and three in the stratosphere, up to about 45 km (Table 1a, b). Thus, a high degree of freedom for the signal (DOFS \approx 4.5) is achieved with the use of the 1000–1005 cm^{-1} window (Barret et al., 2002). The random error on total columns is about 2–4 % (García et al., 2012; Vigouroux et al., 2008), the leading error source being the temperature. The estimated random error on the ozone partial columns can vary a little depending on the station (different instrument, ozone natural variability impacting the smoothing error). For example, at Kiruna the random error for the layers ground–10 km, 10–18 km, 18–27 km, and 27–42 km are 10, 7, 9, and 6 %, respectively (Vigouroux et al., 2008). The smoothing error is dominant in the two lowest layers, whereas the temperature error dominates in the middle and upper stratosphere. Further improvement (on-going research, not applied to the NDACC network yet) can be achieved retrieving a simultaneous temperature profile, reducing the total column random error to less than 1 %, and the upper stratospheric error to 3 % (García et al., 2012). The systematic error is dominated by the uncertainties on the spectroscopic parameters (line intensity and air broadening coefficient). The systematic error due to the line intensity uncertainty was estimated to 2–3 % (Barret et al., 2003; Schneider et al., 2008) on total and partial columns. The error due to the air broadening coefficient uncertainty is negligible for total columns but can be as large as 4 % when partial columns are concerned (Barret et al., 2003; Schneider et al., 2008). Other sources of systematic errors, such

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as the temperature or the instrumental line-shape (if not correctly determined), should also be considered in the upper layers (García et al., 2012).

The ozone trends in Vigouroux et al. (2008) were calculated with a bootstrap re-sampling method, applied to the daily mean partial columns of the ozone time-series.

This work was extended up to 2009 in the WMO 2010 Ozone Assessment report (WMO, 2011) for five European stations (28° N to 79° N). A current update up to 2012 is ongoing, involving additional stations, including stations in the Southern Hemisphere. The spectroscopic database has been changed to HITRAN 2008 (Rothman et al., 2009), and an additional effort of homogenization has been made in contrast to the work described in Vigouroux et al. (2008); all stations are using a priori information from the model WACCM.

The length of the FTIR ozone time-series varies by station (see Fig. 1 for the geographical distribution of the stations); the longest time-series available starts in 1995, and the shortest in 2002 (Fig. 2; Table 1c). In the coming years, more recently established observatory stations (e. g. Reunion Island: long-term measurements started in 2009; Toronto: regular data containing the 1000–1005 cm⁻¹ range started in 2010) or stations that have not yet reprocessed their retrievals using the optimized common strategy (e.g.: Eureka, Bremen) could be included in ozone trends studies.

The FTIR data not yet available on NDACC database (see Table 1d for the URL) can be acquired by direct contact with the responsible Principal Investigator. There are currently 17 active long-term NDACC FTIR sites. An additional long-term site (Kitt Peak, AZ, USA (31.9° N, 111.6° W)) has data from 1978–2005, but operations have unfortunately ceased.

2.5 Microwave

The NDACC microwave ozone profiling instruments measure the spectra of emission lines produced by thermally excited, purely rotational ozone transitions at millimeter wavelengths. All are based on the same principles but differ in technical details. All use a sensitive heterodyne downconverter (receiver) that produces a replica of the

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spectrum it receives from the sky at a much lower so-called intermediate frequency where it can be processed by, a filter bank or digital FFT spectrometer. In all systems, the spectral intensity scale is established by substituting the thermal radiation from two black body sources for the radiation from the sky at the receiver input. One source is at ambient temperature, the second is typically chilled using liquid nitrogen. The ozone altitude distribution is retrieved from the details of the pressure broadened line shape, typically using various implementations of the optimal estimation method of Rodgers (1976). The attenuation of the ozone signal in the troposphere is determined by measuring the tropospheric thermal emission and relating the tropospheric opacity to its emission using a radiative transfer model. Stratospheric aerosols do not affect microwave measurements because the wavelength is large compared to the aerosol particle size. The altitude range is between about 20 and 55 to 72 km, depending on the instrument (Table 1a, b). The fundamental native measurement units are mixing ratio vs. pressure. The instruments operate continuously, and profile retrievals are obtained in weather ranging from clear to some overcast conditions. Temporal resolution can be hourly or less, so diurnal variations of ozone in the stratosphere and mesosphere can be observed.

The instruments at Lauder (New Zealand), Bern and Payerne (Switzerland), and Mauna Loa/Hawaii (USA) have operated essentially continuously from 1992, 1994, 2000, and 1995 to the present respectively. Ny Ålesund (Norway) has been making measurements since 1994, with some extended breaks. These are NDACC Primary Instruments listed in http://www.iapmw.unibe.ch/research/collaboration/ndsc-microwave/instruments/index_spe.html. Other instruments, also listed on the above site, have operated over periods of months to a decade (Table 1c). Instrument and measurement details for Lauder and Mauna Loa were published in Parrish et al. (1992), for Payerne in Maillard-Barras et al. (2009) and Hocke et al. (2007), for Bern in Studer et al. (2013), for Ny Ålesund in Palm et al. (2010). Data from these instruments are archived on the NDACC webpage (see Table 1d for URL).

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Theoretical investigations of microwave ozone measurement errors and vertical resolution have been reported by Connor et al. (1995) and Palm et al. (2010). The former is applicable to the Lauder and Mauna Loa measurements, the latter to the Ny Ålesund measurements. As derived values vary substantially from one instrument to another, the vertical resolution shown in Fig. 3a is an ensemble average; see papers associated with each instrument for its resolution. Because instrument calibration is based on temperature it can theoretically be kept stable over long periods of time. In practice, a more important issue is that the details of the spectral line shape must be measured with a precision that is technically challenging to achieve. The types of spectral errors most commonly encountered tend to propagate into profile errors in the lower to middle stratosphere. Despite these challenges, the profiles of the Mauna Loa and Lauder instruments were found to agree with others to within $< 5\%$ from ≈ 22 to 65 km during formal blind intercomparison campaigns involving several types of instruments (McDermid et al., 1998a, b; McPeters et al., 1999). Differences between the Bern (Studer et al., 2013) instrument and others mostly fall within $\approx 7\%$, as do those at Payerne up to ≈ 50 km. Relative drifts between the Lauder and MLO measurements and those made with lidar, ozonesonde, SAGE II, and HALOE were typically $< 0.5\% \text{ yr}^{-1}$ from 22 to 60 km over the period from the mid-1990s to the mid-2000s (Boyd et al., 2007). While no decadal scale drift tests have been reported for the other stations, Steinbrecht et al. (2009) found that the Bern measurements were very consistent with others over the period from 1994 to 2008.

3 Satellite-based measurement systems

In the following paragraphs several satellite-based systems measuring ozone profiles are described. Systems are grouped according to measurement technique. The solar occultation technique provides spectral measurements with very high signal-to-noise ratio allowing the detection of species with low concentrations without the need for averaging measurements. This self-calibrating method is less susceptible to changes

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in instrument performance over the mission lifetime because of the reference exo-atmospheric measurements used in the retrieval. Stellar occultation measurements are relative by definition and therefore they are less sensitive to instrumental degradation. Due to the multitude of stars, global coverage is obtained. Since stars are point sources the pointing information is excellent (uncertainty < 30 m) and the vertical resolution very good, about 2 km around the ozone maximum and below. The limb viewing technique offers high sensitivity to a number of trace gases with vertical resolution between 1 and 3 km (depending on instrument and wavelength) and provides day and night retrievals, with dense sampling along the sub-orbital tracks and hundreds to a few thousands of profiles per day. Self-calibration is typically provided by reference views to (stable) cold space and onboard blackbody targets. Nadir viewing instruments provide full global coverage of the sunlit Earth on a daily basis, with 6 km vertical resolution for the stratospheric ozone profile, and up to tenth of thousands of profiles per day.

Temporal coverage of each measurement system is shown in Fig. 2. Average spatial coverage for the solar-occultation instruments is shown in Fig. 4 since this is very distinct for each of the systems. Spatial coverage for the stellar-occultation, limb and nadir systems described in the following sections is shown in detail in Toohey et al. (2013), and is therefore not explicitly repeated here. Figure 3 b-d show in detail the vertical resolution of each of the satellite measurement systems that are described below.

3.1 Solar-occultation instruments

3.1.1 SAGE

All of the Stratospheric Aerosol and Gas Experiment (SAGE) instruments (SAGE I on the Applications Explorer Mission-B (AEM-B), SAGE II on the Earth Radiation Budget Satellite (ERBS) and SAGE III on the Meteor-3M spacecraft), used solar occultation to take spectral transmission measurements along a slant path through the atmosphere. SAGE III added lunar occultation measurements. The SAGE instruments scan a small

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



field of view across the solar disk every 2 s. The wavelength-dependent atmospheric transmission is determined by taking the ratio of a measurement looking through the atmosphere to one taken when the sun or moon is above the atmosphere as seen from the spacecraft. The systematic uncertainty of the transmission is independent of instrument characteristics as long as the instrument is stable over the brief few minutes spanned by each occultation event. Ozone is inferred from spectral measurements near 600 nm, at the peak of the Chappuis band and nearby channels are used primarily to characterize the spectral aerosol extinction. The small field-of-view and high signal-to-noise ratio enabled by looking at such bright sources enables the retrieval of the ozone profile at 1 km vertical resolution with low random uncertainty ($\approx 1\%$) from just below the tropopause to well above the stratopause (Table 1a, b).

SAGE I made measurements over a 33 month period, starting in February 1979 (Fig. 2). SAGE II made measurements for 21 yr starting in October 1984 (Table 1c) covering the time when ozone depletion was at its peak as well as during the transition to lower chlorine loading in response to the Montreal Protocol. SAGE III made measurements from December 2001 to December 2005; it provided complementary high latitude coverage (Fig. 4).

The solar occultation technique produces a very stable long-term data set. The ratio between scans across the solar disk above the atmosphere and scans through the atmosphere to determine the spectral extinction vs. altitude are calculated. The instrumental radiometric properties divide out in this ratio. Scanning across, rather than staring at, the Sun provides accurate knowledge of the altitude registration of each measurement.

The most recent version of the SAGE data is version 7. This is the first version that will unify the algorithm, ancillary data and spectroscopy across all three SAGE missions (Damadeo et al., 2013). Data are archived at the Atmospheric Science Data Center, NASA Langley Research Center in Hampton, Virginia (see Table 1d for URL).

3.1.2 HALOE

The Halogen Occultation Experiment (HALOE) was launched aboard the Upper Atmosphere Research Satellite (UARS) in 1991 and operated successfully for fourteen years until November 2005. HALOE made global observations verifying the effects of chlorine compounds on the chemical loss of ozone in the upper stratosphere. In addition to HCl and ozone, HALOE provided profile measurements of H₂O, CH₄, NO, NO₂, HF, temperature and aerosol extinction using the technique of solar occultation. Its measurements of HF, HCl, CH₄, and NO were obtained using gas filter correlation radiometry. HALOE ozone transmission profiles were measured by the more traditional method of broadband filter radiometry (Russell et al., 1993), using a broadband channel from 9.2 to 10.4 micrometers. Overall measurement uncertainties range from 9–25 % in the lower stratosphere and from 9–20 % in the upper stratosphere and mesosphere.

The HALOE retrieval algorithm employed a top-down, onion-peeling approach and iterated the calculated, tangent-layer transmissions to achieve a match with the measured transmissions from each of its channels. The one exception was the method for obtaining the temperature profile, which began with the assignment of the co-located National Center for Environmental Prediction (NCEP) temperature and pressure (or $T(p)$ value) at 31.5 km to the HALOE-measured CO₂ channel transmission profile at that same level. Then, a first-guess $T(p)$ profile was iterated and proceeded upward by matching the calculated transmission value in each tangent layer with that observed in the HALOE CO₂ channel centered at 2.8 μ m. The measured transmission profiles for all the species, including ozone, were then registered vs. the pressure profiles obtained in this way, prior to final retrievals of the species volume mixing ratio profiles (Table 1a–c). The HALOE Level 2 profiles can be downloaded from the NASA Goddard Earth Sciences Data & Information Services Center (GES DISC) or the HALOE webpage (see Table 1d for the URLs).

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1.3 ILAS and ILAS-II

The Improved Limb Atmospheric Spectrometer (ILAS) was a satellite-borne solar-occultation sensor on board the Japanese Advanced Earth Observing Satellite (ADEOS) (Sasano, 2002). ILAS consisted of an infrared spectrometer that covers the wavelength region from about 6 to 12 μm with a detector array of 44 elements and a visible spectrometer from 753 to 784 μm with a detector array of 1024 elements. ADEOS was successfully launched in a sun-synchronous orbit on 17 August 1996. After a 3 month initial checkout period, continuous operation of ILAS started on 30 October, lasting until 29 June 1997 when a failure in the satellite's solar battery system occurred (Fig. 2). For a period of about 8 months, ILAS made solar-occultation measurements at 57–71° N and 64–88° S, and collecting \approx 6700 profiles of O₃, HNO₃, NO₂, N₂O, CH₄, H₂O and aerosol extinction coefficients at 780 nm.

ILAS-II was a satellite-borne solar-occultation sensor on board the satellite ADEOS-II (Nakajima, 2006). ILAS-II consisted of four spectrometers: an infrared spectrometer (between 6.21 μm to 11.76 μm); a mid-infrared spectrometer (between 3.0 μm and 5.7 μm); a high-resolution infrared spectrometer (between 12.78 μm and 12.85 μm); and a visible spectrometer (between 753 nm and 784 nm). ADEOS-II was successfully launched on 14 December 2002, also in a sun-synchronous orbit. After a 3 month initial checkout period, continuous operation of ILAS-II started on 2 April 2003, lasting until 24 October 2003 when the satellite's solar battery system malfunctioned (Fig. 2, Table 1c). For a period of about 7 months, ILAS-II made solar-occultation measurements at 57–71° N and 64–88° S, and gathered \approx 5700 profiles of O₃, HNO₃, NO₂, N₂O, CH₄, H₂O, ClONO₂, N₂O₅, CFC-11, CFC-12 and aerosol extinction coefficients at 780 nm.

Ozone and other minor gas vertical profiles were retrieved by applying both the onion-peeling method and the nonlinear least squares fitting method. Version 5.20 ILAS and version 1.4 ILAS-II ozone profiles were validated by comparing with various data sources (Sugita et al., 2002, 2006). These validations showed that ILAS ozone data agreed with other data sets within $\pm 10\%$ with a few exceptions between 11 and 64 km

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Table 1a, b). ILAS-II ozone data agreed with other data sets within $\pm 10\%$ at altitudes between 11 and 40 km for the Northern Hemisphere. Above 41 km, a negative bias between the NH ILAS-II ozone data and the other data increased, reaching 30% at 61–65 km. Southern Hemisphere ILAS-II ozone data agreed with other data sets within $\pm 10\%$ at altitudes between 11 and 60 km. Since the solar-occultation technique is used with both ILAS and ILAS-II, the instruments are self-calibrated by measuring 100% and 0% intensity of the sun each time before and after the atmospheric measurement. ILAS ozone data are archived at the ILAS and ILAS-II webpages (see Table 1d for URLs).

3.1.4 POAM II and POAM III

The Polar Ozone and Aerosol Measurement II (POAM II) instrument was developed by the Naval Research Laboratory (NRL) to measure the vertical distribution of atmospheric ozone, water vapor, nitrogen dioxide, aerosol extinction and temperature using with the solar-occultation technique. POAM II measured solar extinction in nine narrow band channels, covering the spectral range from approximately 350 to 1060 nm. POAM II was launched aboard the French SPOT-3 satellite on 26 September 1993 into a Sun synchronous polar orbit. The POAM II mission was ended by the failure of the SPOT-3 satellite in November of 1996 (Fig. 2).

The POAM III instrument was a satellite-based nine-channel, visible/near-infrared photometer for making measurements of ozone, aerosol extinction, H₂O and NO₂ in the polar stratosphere using the solar occultation technique (Lucke et al., 1999). The POAM III instrument was developed by NRL and measured solar extinction in nine narrow band channels, covering the spectral range from approximately 350 to 1020 nm. POAM III was carried by the French SPOT-4 spacecraft, which launched on an Ariane-4 rocket in March 1998. The POAM III instrument failed on 5 December 2005.

POAM made 14–15 measurements per day in each hemisphere around a circle of latitude. In the Northern Hemisphere the measurement latitude ranged from 55° N to 73° N, and in the Southern Hemisphere, 63° S to 88° S. The POAM III retrieval version 4 includes the following profile data: ozone concentration (5–60 km), NO₂ concentration

(20–40 km), aerosol extinction at 6 wavelengths (5–25 km), H₂O (5–45 km) and UK Met Office temperature, pressure, and potential vorticity (0–60 km), all with a ≈ 1 km vertical resolution. POAM II has a slightly smaller altitude range for the retrieved profiles than POAM III (Table 1a, b). The estimate for random error on the retrieved concentrations is approximately 3–5 % above 20 km and increasing at lower altitudes. For a complete error analysis see Lumpe et al. (2002). As solar occultation instruments, both POAM systems are largely self-calibrating. No drift or degradation of the POAM III ozone retrievals has been observed.

The most recent version of the POAM retrieved profiles is version 4 (pdf document on the POAM ftp archive, see Table 1d for URL), which differs only very little from version 3 (Lumpe et al., 2002). Data is available either through the NASA Langley Atmospheric Science Data Center or through the Naval Research Laboratory (see Table 1d for URLs).

3.1.5 ACE-MAESTRO

The Atmospheric Chemistry Experiment (ACE) Measurement of Aerosol Extinction in the Stratosphere and Troposphere (MAESTRO) occultation instrument (McElroy et al., 2007; Nowlan et al., 2007), on board the Canadian SCISAT satellite, is a dual, diode-array spectrometer measuring in the UV-VIS-NIR spectral range with a nominal wavelength range of 285–1015 nm (525 and 1010 nm for its primary ozone measurement mode in solar occultation). SCISAT was designed to measure the composition of the Earth's atmosphere from space, with particular emphasis on the middle atmosphere ozone distribution, and related trace gases in the Arctic. The satellite was launched on 12 August 2003 and designed for a 2 yr mission, however it is still operating (Fig. 2). It is in a 74° inclined circular orbit at an altitude of 650 km.

MAESTRO makes measurements in the solar occultation mode and shares a common suntracker and optical viewing direction with the Atmospheric Chemistry Experiment Fourier Transform Spectrometer instrument on the same satellite. Therefore, they observe almost exactly the same slant column of air, albeit with different fields of view,

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and with only a slight difference resulting from differential refractive effects. MAESTRO has a relatively small (0.02°) optical resolution which provides a resolution of ≈ 1 km in altitude (Table 1b).

The primary purpose in adding MAESTRO to the satellite payload was to obtain wavelength-dependent, aerosol extinction information. However, in order to remove the effects of gaseous absorbers, a spectral fitting analysis is done which produces ozone, NO_2 and water vapor slant column amounts. Vertical profiles of these species in volume mixing ratio are also produced, covering the altitude range of 5–35 km (Table 1a). Overall uncertainty on the retrieved profiles is $\approx 10\%$, with smaller uncertainties ($\approx 3\%$) in the mid- to upper stratosphere (Dupuy et al., 2009; Kar et al., 2007).

Because MAESTRO is measuring in occultation mode, a reference spectrum is collected before or after each occultation. Therefore, the optical depths calculated are almost independent of instrument sensitivity. The UV half of MAESTRO has lost sensitivity steadily throughout the mission, with the loss of sensitivity migrating from shorter to longer wavelengths to the point where the instrument has producing no useful data for the last five years. However, the visible-near-IR spectrophotometer has shown very little change in sensitivity. Based on the behavior of the two instruments in the overlap region of the spectrum it is likely that the loss of sensitivity is due to attenuation taking place outside the instrument.

MAESTRO data can be obtained from the ACE Science Team at the University of Waterloo, Canada (see Table 1d for URL).

3.1.6 ACE-FTS

The Atmospheric Chemistry Experiment-Fourier Transform Spectrometer (ACE-FTS), on board the SCISAT satellite, is a solar occultation instrument measuring in the mid-infrared spectral region (Bernath, 2006). SCISAT began routine measurements on 21 February 2004 (Fig. 2).

ACE-FTS is a high-resolution (0.02 cm^{-1}) FTS that measures the full spectral range between 750 and 4400 cm^{-1} (Bernath et al., 2005). Each day ACE-FTS measures ap-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



proximately 15 sunrise and 15 sunset occultations and achieves nearly complete global latitude coverage over a period of three months (e.g., one “season”; Table 1c). These mid-infrared spectral measurements extend from the cloud tops to 150 km and ozone is retrieved up to 95 km (Table 1a). The vertical spacing between each 2 s ACE-FTS measurement varies between 1.5 and 6 km depending on the satellite’s orbit geometry. The vertical sampling is 3–4 km based on the field of view of the instrument (Table 1b).

Constituent profiles are retrieved from the ACE-FTS transmission spectra as volume mixing ratios (VMR) using an unconstrained non-linear least squares global fitting approach (Boone et al., 2005). Initially, CO₂ transitions are used to determine the pressure and temperature as a function of altitude between 12 and 120 km. Below 12 km, meteorological results from the Canadian Meteorological Centre (CMC) operational weather analysis and forecast system are used (note that in version 3.5 of the ACE-FTS retrievals, the CMC pressure and temperature results are used below 15 km, Boone et al., 2013). The resulting temperature and pressure profiles are the basis for retrieving VMR profiles of ozone and over 30 trace gas species. For the ozone retrieval, spectral features between 984 and 1190 cm⁻¹ are used.

The spectroscopic parameters on which the calculations are based are derived from the HIgh-resolution TRANsmission (HITRAN) molecular absorption database 2004 line list (Rothman et al., 2005). The version 2.2 ozone updated (herein v2.2) dataset was validated extensively by Dupuy et al. (2009). In comparisons with ozonesondes and satellite limb sounding instruments, a small positive bias of +1 to +8 % (+5 % on average) was found between 16 and 44 km. At altitudes above 45 km, larger differences of up to +40 % (+20 % on average) are found for the v2.2 ozone profiles. The current version of the ACE-FTS dataset is version 3.5 (Boone et al., 2013). Waymark and colleagues (personal communication) have evaluated the differences between the well-validated v2.2 and new v3.5 profiles and found that there is a reduction of ≈ 5 % at altitudes above 35 km in v3.5 compared to v2.2. There is no error budget available for the ACE-FTS ozone product at present. However, there is an associated fitting uncertainty provided for each measurement.

The ACE-FTS instrument is self-calibrating. A series of reference exo-atmospheric spectra (“high sun” measurements) are collected during each orbit and are used with the atmospheric measurements to calculate the transmittance spectra that are then used in the retrieval process. Data are made available by the ACE team on request or via webpage of the ACE Science Team at the University of Waterloo, Canada (see Table 1d for more detail).

3.2 Stellar occultation instruments

3.2.1 GOMOS

The Global Ozone Monitoring by Occultation of Stars (GOMOS) instrument is one of the three atmospheric instruments on-board the European Space Agency’s Environmental Satellite (Envisat) satellite. The satellite was launched in March 2002 into a polar sun synchronous orbit with a descending node at 10 p.m. GOMOS measurements ceased in April 2012 when contact with Envisat was lost (Fig. 2). About 410 000 night-time ozone profiles were measured by GOMOS during its 10 yr period of operation.

The GOMOS spectrometer covers the wavelength range from ultraviolet, visible and near infrared (UV-VIS-NIR). It uses the stellar occultation technique for observing trace gas and aerosol profiles from the upper troposphere to the mesosphere (Bertaux et al., 2010). Approximately 160 stars are bright enough for atmospheric profiling, providing global coverage from pole to pole. The altitude range is typically from the top of the clouds to 100 km (Table 1a). Geophysical validations have shown that the night-time measurements of ozone are accurate with a negligible $\pm 2\%$ bias in the stratosphere between 20–40 km (e.g. Meijer et al., 2004; van Gijsel et al., 2010). The sampling resolution of GOMOS is 0.5–1.7 km depending on the geometry of the occultation (Table 1b). Due to refraction the sampling resolution improves below 40 km. The stars are point sources, which ensures very good knowledge of the measurement position (< 30 m).

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The retrieval of ozone and other trace gases using stellar occultation measurement is relatively straightforward (Kyrölä et al., 2010). The present retrieval assumes known Rayleigh extinction, which is estimated using the European Centre for Medium-Range Weather Forecasts (ECMWF) and MSIS90 models for neutral density profiles. Only a minimal amount of a priori information on ozone is needed and it is included in the retrieval by using Tikhonov regularization method assuming smoothness of the ozone profiles. The vertical resolution of the ozone profiles is thus lower than the actual sampling resolution: 2 km in the stratosphere and 3 km in the mesosphere. Ozone is retrieved together with aerosols, NO₂, and NO₃ using wide wavelength band 250–675 nm. The stellar characteristics (temperature and brightness) are linked to signal-to-noise ratio resulting in a varying random uncertainty of the data depending on the stars used. The random uncertainty of ozone measurements in the stratosphere ranges from 0.5–4 %, except in the lowermost stratosphere where values are about 10 %, and 2–10 % in the mesosphere. The largest systematic error contribution is due to uncertainties in aerosol modeling and is about 20 % below 20 km, 1–5 % at 20–25 km and negligible above 25 km (Tamminen et al., 2010).

As an occultation instrument GOMOS is, in principle, less sensitive to instrument degradation. However, careful evaluation of possible drifts is needed since the signal to noise ratio is low with dim stars and the increasing amount of dark charge is known to cause problems despite the self-calibrating measurement principle. These problems are minimized by careful screening of the data for dim and cool stars.

In addition to the nighttime occultation measurements discussed above, GOMOS has daytime occultation measurements (Table 1c). Presently GOMOS daytime occultations are of poor quality (e.g. van Gijssel et al., 2010). Daytime ozone profiles can, however, be derived from GOMOS measurements of scattered solar light (Tukiainen et al., 2011). These data are processed by the Finnish Meteorological Institute (FMI) and ozone profiles are available from the ESA SPIN project. All occultation measurements are available via the ESA EARTHNET webpage (see Table 1d for more detail).

3.3 Limb emission instruments

3.3.1 LIMS

The Limb Infrared Monitor of the Stratosphere (LIMS), an infrared, limb-emission instrument, operated on Nimbus 7 following its launch in late October 1978 through late May 1979, at which time its solid cryogen coolant was depleted (Fig. 2). The quality of the original LIMS Version 5 dataset is described in Gille and Russell (1984) and references therein. The LIMS retrieval algorithm uses a top-down onion-peeling approach and iterates to achieve a match of the calculated and measured limb radiances for its wide and narrow CO₂ radiometer channels in the 15-micrometer region. A constant CO₂ volume mixing ratio profile of 325 ppmv is assumed for the radiance calculations in the upper stratosphere, where the forward models for the wide and narrow CO₂ channels are aligned and iterated. Radiance profiles for the LIMS species channels are registered against the pressure profiles that are obtained in this manner.

In order for the LIMS dataset to be more compatible with those from instruments of the UARS, Envisat, and the EOS satellite programs of more recent decades, LIMS profiles were retrieved anew using updated spectral line parameters for CO₂ and other species, including ozone, from the 1992 or 1996 HITRAN line lists (Remsberg et al., 2004). LIMS ozone profiles were obtained from infrared limb radiance measurements at 9 to 10 micrometers. The resultant LIMS Version 6 Level 2 profiles of temperature, species and height are tabulated at 18 levels per decade of pressure or at a spacing of 0.88 km (Table 1a). They have an effective vertical resolution of 3.7 km, in order to account properly for the effects of the instrument on the observed radiances and of the retrieved temperatures on the retrieved ozone profiles (Table 1b). The Version 6 dataset was processed in 2002 and the Level 2 and 3 products are archived at NASA Goddard (GES DISC; see Table 1d for URL). A more detailed description of the LIMS V6 ozone and its quality is provided in Remsberg et al. (2007). Overall uncertainties for LIMS ozone range from 20–30 % in the lower stratosphere, to approximately 12 % in the mid- to upper stratosphere.

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



capabilities to account for horizontal volume mixing ratio gradients. Version 5 was the most recent public release of UARS MLS O₃ data (see Livesey et al., 2003) and the work by Froidevaux et al. (1996) on an earlier data version for more details and validation results).

UARS MLS ozone profiles are recommended from 100 to 0.22 hPa and are retrieved on the same vertical grid as for Aura MLS v2.2 O₃; the vertical resolution is 3.5–5 km in the stratosphere and 5–8 km in the mesosphere (Table 1a, b). Estimated random errors are 0.2–0.3 ppmv and typical systematic error estimates (2σ) are $\approx 6\%$, degrading to 15% (or more) near 68–100 hPa, where some high biases vs. correlative data were noted. The official public distribution location for UARS MLS data is at the NASA GES-DISC Mirador website (see Table 1d for URL).

3.3.3 Aura MLS

NASA's Earth Observing System (EOS) Microwave Limb Sounder (MLS) instrument was launched on the Aura satellite on 15 July 2004 (Fig. 2). This improved sounder (and successor to UARS MLS) measures a variety of atmospheric species (and temperature); O₃ retrievals are provided from the upper troposphere to the mesosphere. Aura MLS performs limb scans ahead of the Aura satellite, which is in a sun-synchronous, near-polar orbit. Aura MLS measures ozone microwave emission for its primary (standard) product near 240 GHz. Retrieved profiles occur at the same latitude every orbit and are separated by ≈ 170 km along the sub-orbital track. This yields nearly 3500 profiles every day (day and night) from 82° S to 82° N (Table 1c). Waters et al. (2006) describe the instrument and its measurements.

The Aura MLS retrievals are obtained via optimal estimation (Rodgers, 1976, 2000). Gauss–Newton iteration is used, with a second-order Tikhonov constraint (Tikhonov, 1963). This constraint is applied to the profile second derivatives (vertically and horizontally). Specific retrieval aspects include adaptation to a two-dimensional system, using the line-of-sight measurements from several scans to derive multi-profile information (Livesey and Read, 2000). The Level 2 data include screening flags regard-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing instrument/retrieval status, quality of fit and retrieval convergence (see Froidevaux et al., 2008). Retrieval details are provided by Livesey et al. (2006) and references therein.

Version 2.2 O₃ is retrieved on a pressure grid with 6 levels per decade up to 0.1 hPa (and coarser vertical resolution above this); v3.3 data are provided on a finer grid and down to 261 hPa, but are not recommended overall over v2.2 because of larger vertical oscillations in the upper troposphere and lower stratosphere (UTLS) at low latitudes (Table 1a). Aura MLS v2.2 single profiles are recommended for scientific use in the 215 hPa to 0.02 hPa range; the vertical resolution is \approx 3 km (or slightly less) in the stratosphere, degrading to 4–6 km for pressures of 0.1 hPa or less (Table 1b). Estimated random errors are 0.1 to 0.3 ppmv (\approx 2–15 %) and systematic error estimates (2σ) are 5–10 % for most of the stratosphere and lower mesosphere (for more details and validation, see Froidevaux et al., 2008, and references therein). The official public distribution location for Aura MLS data is at the NASA GES-DISC Mirador website (see Table 1d for URL).

3.3.4 MIPAS

The Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) is a limb emission, Fourier transform spectrometer (Fischer et al., 2008). Looking horizontally through the atmosphere at various elevation angles, it measured high-resolution, infrared emission spectra of atmospheric constituents, thus providing altitude-resolved information on these constituents. MIPAS was one of the atmospheric chemistry instruments on ESA's Envisat research satellite, which was launched into its sun-synchronous polar orbit on 1 March 2002. In March 2004, problems with the interferometer slide occurred and operation was interrupted, however on 27 January 2005 operation in the nominal mode was resumed with reduced spectral but improved spatial resolution (optimized resolution measurements). MIPAS provided data until 8 April 2012 when contact to the satellite was lost (Fig. 2).

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There exist multiple MIPAS level-2 retrievals. One version was generated with the research processor developed and operated by the Institute of Meteorology and Climate Research (IMK), KIT, Karlsruhe, Germany, in cooperation with the Instituto de Astrofísica de Andalucía (IAA), CSIC, Granada, Spain (von Clarmann et al., 2003, 2009).

The retrieval was performed on a fixed altitude grid usually finer than the tangent altitude spacing, and was regularized by a Tikhonov first order approach that leads to smoothing of the profiles without bias towards an a priori. IMK high-spectral-resolution ozone data (2002–2004) were validated by Steck et al. (2007) and the first validation of the reduced spectral resolution data was published by Stiller et al. (2012).

Frequent gain and offset calibration by means of deep space and blackbody measurements minimize drifts of MIPAS measurements. There is a higher order effect due to slowly decreasing detector non-linearity that is supposedly overestimated towards the end of the MIPAS measurement period. This explains a predominantly negative drift of MIPS-IMK ozone vs. MLS-Aura ozone, which can reach $-0.2 \text{ ppmv decade}^{-1}$ around 40 km (Eckert et al., 2013).

Another MIPAS processor is run by ESA (Ridolfi et al., 2000; Raspollini et al., 2006, 2013). It is based on the algorithm ORM (Optimized Retrieval Model) developed by an European Consortium led by IFAC-CNR (Istituto di Fisica Applicata “Nello Carrara”, part of the Consiglio Nazionale delle Ricerche). This processor was specifically designed for operating in near real time, and hence for working automatically in different atmospheric conditions, and uses the minimum amount of a priori information that may introduce a bias in the profiles. To this purpose, it uses the regularizing Levenberg Marquardt regularization inside the iterations and an a posteriori Tikhonov regularization with a self-adapting constraint that ensure that minimum constraints are used and the best vertical resolution obtainable from the measurements is achieved in all atmospheric conditions. ESA spectral resolution ozone data (2002–2004) were validated by Cortesi et al. (2007).

A third MIPAS retrieval algorithm was developed at Oxford University (Moore and Remedios, 2010) and is based on the Optimal Estimation approach (Rodgers, 2000).

the measurements have remained relatively stable (equator crossing times between 6 a.m.–7 a.m. and 6 p.m.–7 p.m.). The Odin sun-synchronous orbit leads to daytime observations in the summer hemisphere and nighttime observations in the winter hemisphere.

5 The Sub-Millimeter Radiometer (SMR) on board the Odin satellite, launched in 2001, makes limb measurements of thermal emission lines of ozone in several different spectral bands. The 501.8 GHz version –2.1 level-2 product provides stratospheric ozone data in the ≈ 12 –60 km range (above 17–18 km at mid-latitudes) with 2.5–3.5 km vertical resolution and single-profile random uncertainty of about 20 % (Jones et al., 2007; 10 Jégou et al., 2008, Table 1a, b). The systematic error is estimated to be smaller than 0.75 ppmv (Urban et al., 2005, 2006).

Measurements in this observation mode were carried out on every 3rd day until April 2007 and on every other day thereafter (Fig. 2). The thermal emission technique allows ozone to be measured during day and night and global fields between $\approx 83^\circ$ S and $\approx 83^\circ$ N are produced during typically 14–15 orbits per observation day based on up to 65 limb-scans per orbit.

The calibration for SMR is done with a standard hot-cold radiance calibration scheme, based on the internal hot load (viewed at the end of each scan, i.e. every 90 s) and cold sky (viewed each tangent-view, i.e. every 0.85 s). This leads to an excellent long-term stability of the measurements. There are no known signs of instrument degradation so far.

Individual ozone profiles of the 501.8 GHz band are quite noisy since they are derived from a relatively weak line, but averages agree reasonably well with correlative measurements (Jones et al., 2009). Other SMR ozone products such as from the strong ozone line in the 544.6 GHz band (version –2.0) are less noisy but have larger (known) 25 biases. Data are available from the Odin-SMR webpage (see Table 1d for URL).

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Limb scattering instruments

3.4.1 SCIAMACHY

The Scanning Imaging Absorption spectroMeter for Atmospheric CHartographyY (SCIAMACHY) was a payload on Envisat launched in March 2002 (Bovensmann et al., 1999, 2011). Due to the failure of communication to the Envisat platform, SCIAMACHY measurements ended in April 2012. SCIAMACHY is one of the new-generation spaceborne instruments capable of performing spectrally resolved measurements in several different modes: alternate nadir and limb observations of the solar radiation scattered in the atmosphere or reflected by the Earth's surface as well as observations of the light transmitted through the atmosphere during solar or lunar occultation. SCIAMACHY was a passive imaging spectrometer comprising eight spectral channels covering a wide spectral range from 214 to 2386 nm. For the limb-scatter ozone profile retrieval selected wavelengths in the Hartley (ultraviolet) and Chappuis (visible) ozone bands are used (Sonkaew et al., 2009).

In the limb-scatter geometry, SCIAMACHY observed the atmosphere tangentially to the Earth's surface starting at ≈ 3 km below the horizon, i.e., when the Earth's surface is still within the field of view of the instrument, and then scanned vertically up to the top of the neutral atmosphere (about 100 km tangent height; Table 1a). The useful vertical range for ozone retrieval is about 10 to 60 km altitude and the vertical resolution of the retrieved ozone profiles is about 4 km (Table 1b). In the nominal mode about 100 measurements per orbit with 14 orbits per day were performed. Global coverage of the sunlit part of the earth was achieved in 6 days (Table 1c).

Similar to other limb-scattering instruments, pointing uncertainty is a major error source (von Savigny et al., 2005). The systematic uncertainty of the pointing for the whole limb scan is estimated to be about 200 m. The relative pointing error between different tangent heights is negligible. Measurements at lower tangent heights were affected by clouds, hence ozone retrievals were not possible in the presence of clouds in the instrument field of view. Above 20 km measurements have uncertainties up to

AMTD

6, 9857–9938, 2013

SI²N: ozone profile
measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



15 % (for a single profile), with a possible 8 % systematic error. Extensive validation of the SCIAMACHY limb-scatter ozone profiles by comparisons with other satellites has been reported for an earlier retrieval version in Mieruch et al. (2012). A detailed error budget is provided in Rahpoe et al. (2013) and ozone trends during the last decade are reported in Gebhardt et al. (2013). Instrument calibration effects are largely removed by using ratios of the limb spectra at different tangent heights. SCIAMACHY version 2.5 data are available from the webpage of the Institute of Environmental Physics, University of Bremen, Germany (see Table 1d for URL).

An alternative retrieval of ozone profiles from SCIAMACHY limb observations is provided by the European Space Agency/DLR (Doicu et al., 2007).

3.4.2 Odin-OSIRIS

The Canadian-built, Optical Spectrograph and InfraRed Imaging System (OSIRIS) (Llewellyn et al., 2004) has been in routine operation on the Swedish spacecraft Odin (Murtagh et al., 2002) since the fall of 2001 (Fig. 2). OSIRIS measures vertically resolved profiles of spectrally dispersed limb scattered sunlight from 280 nm to 800 nm with approximately 1 nm resolution. Typically the Odin spacecraft rotates to scan the OSIRIS single line of sight through a tangent altitude range that extends from 10 km to 60 km (Table 1a). In the sunlit part of the orbit, a single vertical ozone profile is retrieved from a single Odin scan that typically takes 90 s. This results in as many as 60 scans per orbit (Table 1c). The Odin scan speed along with the OSIRIS imaging rate and instantaneous field of view combine to give an approximately 1.5 km sampling resolution in the UTLS and 2 km at higher altitudes (Table 1b).

The retrieval scheme used to infer vertical ozone profiles from limb-scattered sunlight was developed at the University of Saskatchewan (Degenstein et al., 2009) and involves an adapted Multiplicative Algebraic Reconstruction Technique and the SASKTRAN (Bourassa et al., 2007) radiative transfer model. The vertical ozone profile within the SASKTRAN model is relaxed in a multiplicative fashion until the model and measurements agree to within a certain tolerance. This method is extremely stable, and

because the OSIRIS noise values are very low, the retrieved data quality is very good. Overall uncertainties range from 5 % between 25 km and 50 km, 15 % below 20 km and 7 % between 50 km and 55 km.

Studies have shown that OSIRIS ozone compares well with coincident SAGE II ozone profiles with biases less than 3 % from 18 to 50 km (Degenstein et al., 2009). OSIRIS retrievals are self-calibrating as the radiance data is normalized against both higher altitudes and non-absorbing wavelengths. These normalizations stabilize the retrieval to the majority of known factors; the exception being upwelling radiation from the ground and lower troposphere. Data are available from the Odin-OSIRIS webpage (see Table 1d for URL).

3.5 Nadir viewing instruments

3.5.1 SBUV

The Solar Backscatter Ultraviolet instruments are nadir-viewing instruments that infer total column ozone and the ozone vertical profile by measuring sunlight scattered from the atmosphere in the middle ultraviolet. The instruments use Ebert–Fastie type double monochromators and measure at twelve discrete wavelengths from 252–340 nm. The ratios of the backscattered radiance to direct solar irradiance are used to infer ozone concentration as a function of altitude. Optimal Estimation techniques are used to invert the measurements and estimate ozone amounts as a function of pressure (Table 1a). The vertical resolution of SBUV is 6 km in the upper stratosphere (≈ 3 hPa) and gets progressively worse at higher and lower altitudes, particularly in the tropics (Table 1b). Figure 9 in Bhartia et al. (2013) provides more details.

The SBUV record is the longest available for satellite-based profile ozone measurements from a single instrument type (McPeters et al., 2013). The record began with the launch of Nimbus-4 BUV in 1970. An improved version, SBUV, was launched in October 1978 on the Nimbus-7 satellite. In 1984 the first of seven SBUV/2 instruments

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was launched on the NOAA-9 satellite, with subsequent launches on the NOAA 11, 14, 16, 17, 18, and 19 satellites (Fig. 2).

The instruments are flown in sun synchronous polar orbits to obtain global coverage. Since the measurements rely on backscattered sunlight, data are taken only on the day lit side of each orbit and no data are taken during polar winter. The NOAA polar orbiting satellites were not exactly sun-synchronous and consequently drifted from near-noon orbits with NOAAs 9–14 drifting the most. NOAA-11 for example, drifted from an initial 1.30 p.m. equator crossing time to a 5.00 p.m. orbit in six years, returning as a morning orbit for the later part of the data record. Starting with NOAA-16 the orbital drift was partially mitigated, providing ≈ 5 yr of data before significant drift began. As the orbit drifts the solar zenith angle at which the measurements are made increases and latitude coverage decreases, which can be a problem for trend determination (Table 1c).

The most recent retrieval algorithm is Version 8.6 (Bhartia et al., 2013). Significant changes from Version 8 include new ozone (McPeters and Labow, 2012) and cloud climatologies (Joiner and Vasilkov, 2006), a change in ozone cross-sections to Brion, Daumont, and Malicet, and updated instrument calibration (DeLand et al., 2012). The absolute calibrations of the individual instruments are adjusted at the radiance level to improve the consistency over the multi-instrument period. This is accomplished using coincident measurements during periods of overlap referred to as the “no local time” correction. Local solar time coincidence is required as well as spatial coincidence to account for real diurnal variations in ozone (DeLand et al., 2012). The instruments have been cross-calibrated during two time periods. The calibration of NOAA-17 SBUV/2 is used as the standard for NOAA-14, NOAA-16, and NOAA-18, while the NOAA-11 base calibration is used as the reference for the Nimbus-7 and NOAA 9 absolute calibrations. No instrument overlaps Nimbus-4 BUV; its absolute calibration is based on comparisons with ground-based instruments and has not changed in Version 8.6.

The primary data product for time series analysis is the monthly zonal-mean data, available at the NASA Goddard DISC webpage (see Table 1d for URL). Validation of SBUV monthly zonal mean profiles (Kramarova et al., 2013b) shows that the mean

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



biases and standard deviations are mostly within 5% in the stratosphere between 25 and 1 hPa. The SBUV vertical resolution decreases in the lower stratosphere and troposphere, and as a result the SBUV smoothing error, defined as the component of vertical ozone variability that the observation system cannot measure, increases from < 1% in the middle stratosphere up to 10–15% in the lower stratosphere and troposphere. Combining layers in the troposphere/lower stratosphere and above 1 hPa is recommended to limit smoothing errors, as described in Kramarova et al. (2013a).

In order to make the SBUV measurements accessible for trend analysis, it is necessary to combine the data from the different NOAA satellites. Two versions of a “merged” data set are available (NASA Goddard, S. Frith, personal communication, 2013; NOAA CPC, J. Wild, personal communication, 2013). Both are described in more detail in the second and third SI²N overview papers (Lambert et al., 2013; Harris et al., 2013).

3.5.2 GOME/GOME-2

The Global Ozone Monitoring Experiment (GOME) is a nadir-scanning ultraviolet and visible (UV-VIS) spectrometer. Its launch on board the European Remote Sensing 2 satellite in April 1995 started a new era of measuring Earth-reflected solar radiation with relatively high spectral resolution for UV-VIS wavelengths (Fig. 2). SCIAMACHY, GOME-2 and OMI followed the GOME heritage. The UV-VIS spectrum between 270 nm and 340 nm is used for deriving global, height-resolved information on the ozone distribution in the atmosphere (Munro et al., 1998). As for SBUV, Optimal Estimation techniques (Rodgers, 2000) are used for the retrievals. GOME ozone profiles were extensively analyzed by Meijer et al. (2006) and when the OMI measurements became available these were validated by Kroon et al. (2011).

A common feature of nadir UV-VIS ozone profiles is their limited vertical resolution, estimated typically at 7–15 km, with a considerable amount of vertical smoothing (Table 1b). Smoothing of the retrieved ozone profile is described by the averaging kernel, which is part of the retrieval results. Uncertainty of the retrieved ozone profile is determined by measurement error, a priori error and by smoothing error (about 30% in

the troposphere, and $\approx 10\%$ in the stratosphere). The ozone profiles retrieval depends on the absolute calibration of the instruments and therefore it is sensitive to degradation effects. An offline degradation correction is needed in case of degradation of the instrument.

5 Within the Climate Change Initiative (CCI) program of ESA a new algorithm has been developed to generate high-quality data for nadir ozone profiles from GOME, SCIAMACHY and GOME-2. Data are available from the EUMETSAT archive, and it is planned to make it accessible via CCI web portal (see Table 1d for URLs). Furthermore, GOME-2 profiles are retrieved on an operational basis by KNMI within the framework
10 of EUMETSAT's Ozone and Atmospheric Chemistry Monitoring SAF (O3MSAF) (e.g. Mijling et al., 2010; van Peet et al., 2009) that can be accessed via the Eumetcast dissemination (see Table 1d for URL).

3.5.3 OMI

15 The Ozone Monitoring Instrument (OMI) is a nadir viewing UV-VIS spectrometer on board of the Aura satellite mission (Levelt et al., 2006a, b). Due to the application of two-dimensional detectors OMI combines daily global coverage with a relative high spatial resolution of $13\text{ km} \times 24\text{ km}$ at nadir for wavelengths from 308–500 nm and $13\text{ km} \times 48\text{ km}$ for wavelengths below 308 nm. The OMI science data record currently extends from October 2004 to present (Fig. 2). The OMI instrument is extremely
20 stable radiometrically, with only a few percent of degradation even in the shortest ultraviolet channels, making it a unique data set for ozone monitoring. Since the beginning of 2009 part the OMI swath is affected by the so-called row anomaly (<http://www.knmi.nl/omi/research/product/rowanomaly-background.php>).

25 The OMI instrument will be succeeded by the TROPOMI instrument on board of the Sentinel-5 Precursor satellite, planned for launch late 2015 (Veefkind et al., 2012). TROPOMI uses the same observation technique as OMI, and includes several improvements, including the signal-to-noise ratio, the spatial resolution and additional

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spectral bands. TROPOMI will have a separate spectral band from 270 to 320 nm, which is expected to lead to improved ozone profile retrievals.

The ozone profile retrieval for OMI is based on the optimal estimation method (Rodgers, 2000). As discussed above in the GOME/GOME-2 section (Sect. 3.5.2), the vertical information is limited, especially in the UTLS and the troposphere (Table 1a–c). The OMI ozone profile data are available at the NASA Goddard DISC (see Table 1d for URL). The validation status of this product is described in Kroon et al. (2011) based on comparisons with ozonesondes and several satellite data products. The OMI row anomaly has several effects on the data quality and some of these depend on the position in the orbit. Currently the effect of the row anomaly on the ozone profile quality in the Northern Hemisphere has not yet been fully quantified. In the future, further improvements to the OMI ozone profile product are planned using a new version of the retrieval algorithm including accurate flagging of the row anomaly. In addition to the product available at the NASA DISC, there is also a research product (Liu et al., 2010), with a strong focus on the retrieval of tropospheric ozone information.

3.5.4 IASI

The Infrared Atmospheric Sounding Interferometer (IASI) instrument is a nadir-viewing, Fourier transform spectrometer. It was first launched on board the sun synchronous polar orbiting MetOp-A satellite on 19 October 2006 (Fig. 2). A second instrument was launched on 17 September 2012 on board Metop-B and, together with the forthcoming Metop-C, will ensure products continuity for at least 15 yr.

IASI scans across track in 30 successive steps (four pixels are acquired simultaneously) either side of nadir, with a total swath of around 2000 km (Table 1c). This allows for global coverage twice a day. Each field of view has a typical footprint size of 12 km at nadir. IASI measures spectra in the thermal infrared between 645 and 2760 cm^{-1} at a spectral sampling of 0.25 cm^{-1} (0.5 cm^{-1} resolution after apodization). The 8461 channels are distributed in 3 bands, which, due to their spectral coverage, allow for the retrieval of surface and cloud properties, temperature and humidity profiles as well as

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ozone and a number of other atmospheric constituents (e.g. CO, CH₄, CO₂, N₂O). The ozone profiles are retrieved in near-real time, using the FORLI-O3 (Fast Optimal Retrievals on Layers for IASI) software (Hurtmans et al., 2012), as part of the IASI Level 2 product processing facility (August et al., 2012, Table 1a–c). Validation was performed using ground-based (Antón et al., 2011; Dufour et al., 2012), aircraft (Parrington et al., 2012; Pommier et al., 2012), balloon (Gazeaux et al., 2013) and other satellite data (Scannell et al., 2012).

IASI is auto-calibrated, with high radiometric accuracy. Retrieval methods for profiles measurements use standard Optimal Estimation Method (OEM) or regularization techniques. Overall uncertainties in the ozone profiles range from about 20–30 % in the altitude range 0–6 km, and are about 10–20 % in the stratosphere and upper troposphere. At the moment, IASI ozone data are available from the IASI team on request. Data should be available via Eumetcast in 2014 (see Table 1d for more detail).

3.5.5 SMILES

The Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) was developed and operated under the cooperation of the Japan Aerospace Exploration Agency (JAXA) and the National Institute of Information and Communications Technology (NICT) (Kikuchi et al., 2010). SMILES conducted its atmospheric observation from 12 October 2009 to 21 April 2010 (Fig. 2) on the Japanese Experiment Module (JEM) aboard the International Space Station (ISS). SMILES used three observation bands in the frequency regions of 624.32–626.32 GHz and 649.12–650.32 GHz. This instrument measured about 1600 profiles per day (both daytime and nighttime) within the latitude range of 38° S – 65° N. Owing to the non-sun-synchronous orbit of the ISS, measured the atmosphere at different local times. This dataset provides unprecedented opportunities to study diurnal ozone variations throughout the stratosphere (Imai et al., 2013; Sakazaki et al., 2013; Kasai et al., 2013; Kreyling et al., 2013).

The SMILES mission has two products for ozone. One is the ISAS product (Imai et al., 2013) and the other is the NICT product (Kasai et al., 2013). Version 3.0 (118-12-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0602) of O₃ profiles in SMILES ISAS product are retrieved based on the combination of Optimal Estimation method (Rodgers, 2000) with Tikhonov Regularization method (L2 constraint). Degradation of the instrument is corrected within engineering process from raw data to brightness temperature spectrum with some parameters obtained on-orbit calibration (Ochiai et al., 2013). The SMILES ozone product provides ozone concentration as a value of volume mixing ratio. The nominal retrieved altitude range is from 8 to 120 km, and the typical “useful altitude range” is from 16 to 95 km (Table 1a). The retrieval vertical grid step is 2.5 km throughout the retrieval altitude range (Table 1b). The information on the convergence status of iteration in retrieval process, the validity of the observation altitude range, and the FOV interference onboard the ISS, are represented as the “screening flags” and these flags are stored in the “status field” of each HDF file.

The NICT products of ozone is called SMILES NICT Level-2 (L2) version 2.1.5 product, which used the version 007 calibrated Level-1b (L1b) spectra. The detail of the evaluation of the L1b spectrum, retrieval procedure including the theoretical error analysis, and comparison/evaluation of the ozone vertical profile was shown in Kasai et al. (2013).

Degradation of the instrument is corrected within the engineering process from raw data to the brightness temperature spectrum with some parameters obtained on-orbit calibration. The public SMILES L2 and L3 (SMILES climatology) data are available from the Data ARchives and Transmission System (DARTS) in the Institute of Space and Astronautical Science (ISAS) of JAXA and from the NICT SMILES page (see Table 1d for URLs).

3.6 Additional instruments

Beyond the instruments/missions described in detail above, several other satellite instruments also provide ozone data. These will only be mentioned briefly in the following paragraphs since these instruments were either not used in any SI²N activity (e.g. included in merged data sets, or updated and improved retrievals) or have their main fo-

SI²N: ozone profile measurements

B. Hassler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

cus on tropospheric rather than stratospheric measurements. However, for the sake of completeness we include them here. The TES (Tropospheric Emission Sounder) instrument on board of Aura is a nadir-looking high-resolution infrared-imaging fourier transform spectrometer observing in nadir, which provides profiles of ozone and many other trace species in the troposphere and lowermost stratosphere. TES provides about 1000 observations a day, with a nadir footprint of 5 km × 8 km. The sensitivity of the measurements is highest in the mid-troposphere. The data record has been thoroughly validated (Worden et al., 2007).

The HIRDLS (High Resolution Dynamics Limb Sounder) instrument is a 21 channel (from 6.12 to 17.76 microns) limb scanning infrared radiometer measuring emission from the Earth's limb onboard Aura. These measurements are inverted to yield global distributions of temperature, clouds, aerosols, and concentrations of ozone and other trace gases at high vertical and horizontal resolution from the upper troposphere into the mesosphere. After launch, activation of the HIRDLS instrument revealed that the optical path was blocked so that 20% of the aperture could view the Earth's atmosphere. Nevertheless, measurements at high vertical resolution could be made at one scan angle. Geographically the data extend from 63° S to 80° N with 1° spacing, and cover the period from 29 January 2005 to 17 March 2008, with very few gaps. The estimated total accuracy is 1–10%, varying with altitude and latitude, and 10 to > 50% in the tropical UTLS (Nardi et al., 2008).

The Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) instrument provides volume mixing ratio profiles of ozone from 15–100 km retrieved from mid-infrared limb emission measurements (Russell et al., 1999). The validation of the v1.07 ozone profiles in the middle and upper atmosphere (\approx 15–70 km) are described in Rong et al. (2009).

4 Summary

This paper describes long-term ozone profile data derived from five of the most common ground-based systems and 19 satellite-based instruments. The physical principles of measurements, specifics of data retrievals and uncertainties of measurements have been described in more detail in various publications and reports with relevant references given in this paper. Therefore, the goal of this paper is to provide an overview of well-established ozone records, while also offering updated information on recently reprocessed and newly available data sources. The summary includes links to data archives, lists uncertainties of measurements and retrievals, provides spatial and temporal characteristics of the datasets described above and indicates the latest released data version (see Table 1d, the numbers given are indicative or typical values). This overview does not represent a complete list of all available ozone profile measurements, but aims to provide the community with information about the latest improvements and recalculations of the most commonly used ozone profile data time series that are most relevant in the context of trend analysis, bundled in one place for easy access and comparison.

Although according to Figs. 2 and 3 (also Fig. 1 in Toohey et al., 2013), it appears that the ozone profile measurements are available globally for the entire 1960–2000 period, one needs to keep in mind that there are regions of the globe and times when data coverage is sparse. A small number of quasi-continuous ground-based measurements by the Umkehr method began in the early 1960s and only a few ozonesonde locations provide continuous records going back to the late 1960s; there is thus limited Northern Hemisphere mid-latitude information about ozone changes in the 1960s and throughout most of the 1970s and virtually no information for other parts of the globe. At the beginning of the 1990s high quality ozone profile measurement at NDACC sites started providing reliable long-term information at selected locations. However, ground-based measurement stations are not uniformly distributed around the globe. The coverage of the ozone variability at high latitudes and, to some extent, in the tropics, were improved

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by various polar research projects and with the SHADOZ network. Nevertheless, there are still large regions with low measurement frequency and sparse coverage.

More complete information for the global analysis of ozone changes only became available with the launch of the first satellite instruments. A BUUV instrument operated for a limited period in the early 1970s with quasi-continuous satellite monitoring starting in the late 1970s. The period between the end of the SAGE I measurements and the beginning of the SAGE II measurements (see Fig. 2) has proven to be one of the most challenging obstacles for reliable, well vertically resolved, global ozone trend detection, especially with the altitude registration problems reported for SAGE I measurements (Wang et al., 1996). SBUV measurements are available during this time, however, their coarse vertical resolution compared to the SAGE measurements severely limits their utility in tying the two SAGE records together, especially in the lower stratosphere. It is evident that the largest ozone decreases in the extra tropics occurred during the 1980s and, therefore, the beginning of this decade is often used as the starting point for trend analyses. However, correctly anchoring trends in this period is complicated and represents a challenge for long-term trend analysis (McPeters et al., 1994; Randel et al., 1999; Cunnold et al., 2000; Wang et al., 2002; Terao and Logan, 2007).

Another period of sparse global measurements resulted from the eruption of Mt. Pinatubo in June 1991. The increased aerosol loading in the stratosphere resulted in retrieval problems for several satellite instruments (and some ground-based instruments) measuring at that time (SAGE II, HALOE, SBUV), resulting in recommendations by some instrument science teams that some ozone data for multiple months after the Pinatubo eruption be discarded (Yu and She, 1995; Cunnold et al., 1996). This reduces the available lower stratospheric ozone measurements during that period to primarily ground-based instruments (ozonesondes).

One goal of the SI²N Initiative is to make the measurement community aware of the increased demand for high quality and consistent data sets with well-defined uncertainty estimates to fulfill requirements for advanced statistical methods used to detect small changes in stratospheric ozone variability. It is clear that ODS reductions in re-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sponse to the Montreal Protocol has led to a reversal of the negative column ozone trends in the extra tropics (e.g. Ziemke and Chandra, 2012), however the documentation of the beneficial effect on ozone profile changes is still challenging (in particular, in an atmosphere where greenhouse gas increases may also be playing a role). As stated earlier, this is the first of three initiative overview papers and provides a summary of available ozone profile measurements. The second overview paper (Lambert et al., 2013) focuses on comparison and validation of the different data sets and the different retrievals for the various instruments that are introduced here. A quantitative understanding of the offsets and drifts among data sets and their limitations is essential when merging them for time series analyses. The third overview paper (Harris et al., 2013) focuses on describing and characterizing data sets that have been derived from a combination of different data sources (multiple satellite instruments). Several of these combined data sets are compared to each other and a detailed evaluation of them is given.

Together, the three overview papers provide the scientific community with a source of information about the latest activities of ozone profile measurement groups, with the intention of providing needed background information and guidance for future ozone profile trend analysis. However, at this point it is also important to point out that there remains a significant need for high-quality stratospheric ozone measurements. Even with the reduction of ODSs in the stratosphere following the Montreal Protocol and its Amendments and Adjustments an exact attribution of stratospheric ozone recovery is not possible due to changes associated with increasing greenhouse gas concentrations. Only with highly stable and accurate profile ozone measurements will it be possible to separate and correctly attribute future ozone changes.

Acknowledgements. We would like to thank the different space agencies that support missions with instruments that measure stratospheric ozone profiles (ESA, NASA, NOAA, JAXA, NICT, CSA-ASC). We also would like to thank the several data bases where ground-based measurements are stored and made accessible (NDACC, WOUDC, SHADOZ). The Atmospheric Chemistry Experiment (ACE) is a Canadian-led mission mainly supported by the Canadian Space Agency and the Natural Sciences and Engineering Research Council of Canada.

References

- Antón, M., Loyola, D., Clerbaux, C., López, M., Vilaplana, J. M., Bañón, M., Hadji-Lazaro, J., Valks, P., Hao, N., Zimmer, W., Coheur, P. F., Hurtmans, D., and Alados-Arboledas, L.: Validation of the Metop-A total ozone data from GOME-2 and IASI using reference ground-based measurements at the Iberian Peninsula, *Remote Sens. Environ.*, 115, 1380–1386, 2011. 9898
- Attmannspacher, A. and Dütsch, H. U.: International ozone sonde intercomparison at the Observatory Hohenpeissenberg, *Ber. Dtsch. Wetterdienstes*, 120, 1–85, 1970. 9866
- Attmannspacher, A. and Dütsch, H. U.: Second international ozone sonde intercomparison at the Observatory Hohenpeissenberg, *Ber. Dtsch. Wetterdienstes*, 157, 1–64, 1980. 9866
- August, T., Klaes, D., Schlüssel, P., Hultberg, T., Crapeau, M., Arriaga, A., O’Carroll, A., Coppens, D., Munro, R., and Calbet, X.: IASI on Metop-A: operational level 2 retrievals after five years in orbit, *J. Quant. Spectrosc. Ra.*, 113, 1340–1371, doi:10.1016/j.jqsrt.2012.02.028, 2012. 9898
- Barath, F. T., Chavez, M. C., Cofield, R. E., Flower, D. A., Frerking, M. A., Gram, M. B., Harris, W. M., Holden, J. R., Jarnot, R. F., Kloezeman, W. G., Klose, G. J., Lau, G. K., Loo, M. S., Maddison, B. J., Mattauch, R. J., McKinney, R. P., Peckham, G. E., Pickett, H. M., Siebes, G., Soltis, F. S., Suttie, R. A., Tarsala, J. A., Waters, J. W., and Wilson, W. J.: The upper atmosphere research satellite microwave limb sounder instrument, *J. Geophys. Res.*, 98, 10751–10762, doi:10.1029/93JD00798, 1993. 9885
- Baron, P., Urban, J., Sagawa, H., Möller, J., Murtagh, D. P., Mendrok, J., Dupuy, E., Sato, T. O., Ochiai, S., Suzuki, K., Manabe, T., Nishibori, T., Kikuchi, K., Sato, R., Takayanagi, M., Murayama, Y., Shiotani, M., and Kasai, Y.: The Level 2 research product algorithms for the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES), *Atmos. Meas. Tech.*, 4, 2105–2124, doi:10.5194/amt-4-2105-2011, 2011. 9934
- Barret, B., De Mazière, M., and Demoulin, P.: Retrieval and characterization of ozone profiles from solar infrared spectra at the Jungfraujoch, *J. Geophys. Res.*, 107, 4788, doi:10.1029/2001JD001298, 2002. 9870
- Barret, B., De Mazière, M., and Demoulin, P.: Correction to “Retrieval and characterization of ozone profiles from solar infrared spectra at the Jungfraujoch”, *J. Geophys. Res.*, 108, 4372, doi:10.1029/2003JD003809, 2003. 9870

AMTD

6, 9857–9938, 2013

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bernath, P.: Atmospheric Chemistry Experiment (ACE): analytical chemistry from orbit, TRAC-Trend. Anal. Chem., 25, 647–654, 2006. 9880
- Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P.-F., Colin, R., DeCola, P., DeMaziere, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn, E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nichitui, F., Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelton, R., Sloan, J. J., Soucy, M.-A., Strong, K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry Experiment (ACE): mission overview, Geophys. Res. Lett., 32, L15S01, doi:10.1029/2005GL022386, 2005. 9880
- Bertaux, J. L., Kyrölä, E., Fussen, D., Hauchecorne, A., Dalaudier, F., Sofieva, V., Tamminen, J., Vanhellefont, F., Fanton d'Andon, O., Barrot, G., Mangin, A., Blanot, L., Lebrun, J. C., Pérot, K., Fehr, T., Saavedra, L., Leppelmeier, G. W., and Fraisse, R.: Global ozone monitoring by occultation of stars: an overview of GOMOS measurements on ENVISAT, Atmos. Chem. Phys., 10, 12091–12148, doi:10.5194/acp-10-12091-2010, 2010. 9882
- Bhartia, P. K., McPeters, R. D., Flynn, L. E., Taylor, S., Kramarova, N. A., Frith, S., Fisher, B., and DeLand, M.: Solar Backscatter UV (SBUV) total ozone and profile algorithm, Atmos. Meas. Tech., 6, 2533–2548, doi:10.5194/amt-6-2533-2013, 2013. 9893, 9894, 9934
- Bodeker, G. E., Hassler, B., Young, P. J., and Portmann, R. W.: A vertically resolved, global, gap-free ozone database for assessing or constraining global climate model simulations, Earth Syst. Sci. Data, 5, 31–43, doi:10.5194/essd-5-31-2013, 2013. 9862
- Boone, C. D., Nassar, R., Walker, K. A., Rochon, Y., McLeod, S. D., Rinsland, C. P., and Bernath, P. F.: Retrievals for the Atmospheric Chemistry Experiment Fourier-Transform Spectrometer, Appl. Optics, 44, 7218–7231, 2005. 9881, 9933
- Boone, C. D., Walker, K. A., and Bernath, P. F.: Version 3 retrievals for the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), in: The Atmospheric Chemistry Experiment ACE at 10: a Solar Occultation Anthology, A. Deepak Publishing, Hampton, Virginia, USA, 2013. 9881
- Bourassa, A. E., Degenstein, D. A., Llewellyn, E. J., and SASKTRAN: A spherical geometry radiative transfer code for efficient estimation of limb scattered sunlight, J. Quant. Spectrosc. Ra., 109, 52–73, doi:10.1016/j.jqsrt.2007.07.007, 2007. 9892

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noel, S., Rozanov, V. V., Chance, K. V., and Goede, A. H. P.: SCIAMACHY – mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, 1999. 9891

5 Bovensmann, H., Aben, I., van Roozendaal, M., Kühl, S., Gottwald, M., von Savigny, C., Buchwitz, M., Richter, A., Frankenberg, C., Stammes, P., de Graaf, M., Wittrock, F., Sinnhuber, M., Sinnhuber, B.-M., Schönhardt, A., Beirle, S., Gloudemans, A., Schrijver, H., Bracher, A., Rozanov, A. V., Weber, M., and Burrows, J. P.: SCHIAMACHY's view of the changing earth's environment, in: SCIAMACHY – Exploring the Changing Atmosphere, Springer, Dordrecht, 175–216, doi:10.1007/978-90-481-9896-2_10, 2011. 9891

10 Boyd, I. S., Parrish, A. D., Froidevaux, L., von Clarmann, T., Kyrölä, E., Russell III, J. M., and Zawodny, J. M.: Ground-based microwave ozone radiometer measurements compared with Aura-MLS v2.2 and other instruments at two network for detection of atmospheric composition change sites, *J. Geophys. Res.*, 112, D24S33, doi:10.1029/2007JD008720, 2007. 9873

15 Boyd, I. S., Parrish, A. D., Zawodny, J., and Froidevaux, L.: Using NDACC groundbased microwave ozone measurements to homogenize long-term SAGE II and Aura MLS time-series in 2 zonal regions, in preparation, 2013. 9933

Brewer, A. W., and Milford, J. R.: The Oxford–Kew ozone sonde, *Proc. R. Soc. Lon.-Ser. A*, 256, 470–495, 1960. 9865

20 Brönnimann, S., Bhend, J., Franke, J., Flückiger, S., Fischer, A. M., Bleisch, R., Bodeker, G., Hassler, B., Rozanov, E., and Schraner, M.: A global historical ozone data set and prominent features of stratospheric variability prior to 1979, *Atmos. Chem. Phys.*, 13, 9623–9639, doi:10.5194/acp-13-9623-2013, 2013. 9863

25 Brühl, C., Drayson, S. R., Russel III, J. M., Crutzen, P. J., McInerney, J. M., Purcell, P. N., Claude, H., Gernandt, H., McGee, T. J., McDermid, I. S., and Gunson, M. R.: Halogen Occultation Experiment ozone channel validation, *J. Geophys. Res.*, 101, 10217–10240, 1996. 9933

Carlotti, M., Brizzi, G., Papandrea, E., Prevedelli, M., Ridolfi, M., Dinelli, B. M., and Magnani, L.: GMTR: two-dimensional multi-target retrieval model for MIPAS-ENVISAT observations, *Appl. Optics*, 45, 716–727, 2006. 9889

30 Casadio, S., Castelli, E., Dinelli, B. M., De Laurentis, M., Papandrea, E., Burini, A., and Carlotti, M.: MIPAS2D, MLS and ACE-FTS inter-comparison of stratospheric profiles, in: ESA Living Planet Symposium, Edinburgh, Scotland, 9–13 September 2013, 2-P-300, 2013. 9889

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cionni, I., Eyring, V., Lamarque, J. F., Randel, W. J., Stevenson, D. S., Wu, F., Bodeker, G. E., Shepherd, T. G., Shindell, D. T., and Waugh, D. W.: Ozone database in support of CMIP5 simulations: results and corresponding radiative forcing, *Atmos. Chem. Phys.*, 11, 11267–11292, doi:10.5194/acp-11-11267-2011, 2011. 9863
- 5 Connor, B. J., Parrish, A., Tsou, J. J., and McCormick, M. P.: Error analysis for the ground-based microwave ozone measurements during STOIC, *J. Geophys. Res.*, 100, 9283–9291, 1995. 9873
- Cortesi, U., Lambert, J. C., De Clercq, C., Bianchini, G., Blumenstock, T., Bracher, A., Castelli, E., Catoire, V., Chance, K. V., De Mazière, M., Demoulin, P., Godin-Beekmann, S.,
10 Jones, N., Jucks, K., Keim, C., Kerzenmacher, T., Kuellmann, H., Kuttippurath, J., Iarlori, M., Liu, G. Y., Liu, Y., McDermid, I. S., Meijer, Y. J., Mencaraglia, F., Mikuteit, S., Oelhaf, H., Piccolo, C., Pirre, M., Raspollini, P., Ravegnani, F., Reburn, W. J., Redaelli, G., Remedios, J. J., Sembhi, H., Smale, D., Steck, T., Taddei, A., Varotsos, C., Vigouroux, C., Waterfall, A., Wetzel, G., and Wood, S.: Geophysical validation of MIPAS-ENVISAT operational ozone data, *Atmos. Chem. Phys.*, 7, 4807–4867, doi:10.5194/acp-7-4807-2007, 2007. 9888
- 15 Cunnold, D. M., Wang, H., Chu, W. P., and Froidevaux, L.: Comparison between Stratospheric Aerosol and Gas Experiment II and microwave limb sounder ozone measurements and aliasing of SAGE II ozone trends in the lower stratosphere, *J. Geophys. Res.*, 101, 10061–10075, 1996. 9902
- 20 Cunnold, D. M., Wang, H. J., Thomason, L. W., Zawodny, J. M., Logan, J. A., and Megretskaia, I. A.: SAGE (version 5.96) ozone trends in the lower stratosphere, *J. Geophys. Res.*, 105, 4445–4457, 2000. 9902
- Damadeo, R. P., Zawodny, J. M., Thomason, L. W., and Iyer, N.: SAGE version 7.0 algorithm: application to SAGE II, *Atmos. Meas. Tech. Discuss.*, 6, 5101–5171, doi:10.5194/amtd-6-5101-2013, 2013. 9875, 9933
- 25 Degenstein, D. A., Bourassa, A. E., Roth, C. Z., and Llewellyn, E. J.: Limb scatter ozone retrieval from 10 to 60 km using a multiplicative algebraic reconstruction technique, *Atmos. Chem. Phys.*, 9, 6521–6529, doi:10.5194/acp-9-6521-2009, 2009. 9892, 9893, 9934
- DeLand, M. T., Taylor, S. L., Huang, L. K., and Fisher, B. L.: Calibration of the SBUV version
30 8.6 ozone data product, *Atmos. Meas. Tech.*, 5, 2951–2967, doi:10.5194/amt-5-2951-2012, 2012. 9894, 9934
- Deshler, T., Mercer, J., Smit, H. G. J., Stubi, R., Levrat, G., Johnson, B. J., Oltmans, S. J., Kivi, R., Davies, J., Thompson, A. M., Witte, J., Schmidlin, F. J., Brothers, G., and Sasaki, T.:

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric comparison of electrochemical cell ozonesondes from different manufacturers, and with different cathode solution strengths: the Balloon Experiment on Standards for Ozonesondes, *J. Geophys. Res.*, 113, D04307, doi:10.1029/2007JD008975, 2008. 9865, 9866

5 Dinelli, B. M., Arnone, E., Brizzi, G., Carlotti, M., Castelli, E., Magnani, L., Papandrea, E., Prevedelli, M., and Ridolfi, M.: The MIPAS2D database of MIPAS/ENVISAT measurements retrieved with a multi-target 2-dimensional tomographic approach, *Atmos. Meas. Tech.*, 3, 355–374, doi:10.5194/amt-3-355-2010, 2010. 9889, 9934

10 Dinelli, B. M., Arnone, E., and Castelli, E.: MIPAS2D – 10 years of MIPAS/Envisat measurements analyzed with a 2-D tomographic approach, in: *ESA publications SP-708 Advances in Atmospheric Science and Applications*, Noordwijk, ESA Communications, ISBN 978-92-9092-272-8, 2012. 9889, 9934

15 Doicu, A., Schreier, F., Hilgers, S., von Bargaen, A., Slijkhuis, S., Hess, M., and Aberle, B.: An efficient inversion algorithm for atmospheric remote sensing with application to UV limb observations, *J. Quant. Spectrosc. Ra.*, 103, 193–208, doi:10.1016/j.jqsrt.2006.05.007, 2007. 9892

20 Dufour, G., Eremenko, M., Griesfeller, A., Barret, B., LeFlochmoën, E., Clerbaux, C., Hadji-Lazaro, J., Coheur, P.-F., and Hurtmans, D.: Validation of three different scientific ozone products retrieved from IASI spectra using ozonesondes, *Atmos. Meas. Tech.*, 5, 611–630, doi:10.5194/amt-5-611-2012, 2012. 9898, 9934

25 Dupuy, E., Walker, K. A., Kar, J., Boone, C. D., McElroy, C. T., Bernath, P. F., Drummond, J. R., Skelton, R., McLeod, S. D., Hughes, R. C., Nowlan, C. R., Dufour, D. G., Zou, J., Nichitiu, F., Strong, K., Baron, P., Bevilacqua, R. M., Blumenstock, T., Bodeker, G. E., Borsdorff, T., Bourassa, A. E., Bovensmann, H., Boyd, I. S., Bracher, A., Brogniez, C., Burrows, J. P., Catoire, V., Ceccherini, S., Chabrillat, S., Christensen, T., Coffey, M. T., Cortesi, U., Davies, J., De Clercq, C., Degenstein, D. A., De Mazi Dupuy, E., Walker, K. A., Kar, J., Boone, C. D., McElroy, C. T., Bernath, P. F., Drummond, J. R., Skelton, R., McLeod, S. D., Hughes, R. C., Nowlan, C. R., Dufour, D. G., Zou, J., Nichitiu, F., Strong, K., Baron, P., Bevilacqua, R. M., Blumenstock, T., Bodeker, G. E., Borsdorff, T., Bourassa, A. E., Bovensmann, H., Boyd, I. S., Bracher, A., Brogniez, C., Burrows, J. P., Catoire, V., Ceccherini, S., Chabrillat, S., Christensen, T., Coffey, M. T., Cortesi, U., Davies, J., De Clercq, C., Degenstein, D. A., De Maziere, M., Demoulin, P., Dodion, J., Firanski, B., Fischer, H., Forbes, G., Froidevaux, L., Fussen, D., Gerard, P., Godin-Beekmann, S., Goutail, F., Granville, J., Grif-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fith, D., Haley, C. S., Hannigan, J. W., Höpfner, M., Jin, J. J., Jones, A., Jones, N. B., Jucks, K., Kagawa, A., Kasai, Y., Kerzenmacher, T. E., Kleinböhl, A., Klekociuk, A. R., Kramer, I., Küllmann, H., Kuttippurath, J., Kyrölä, E., Lambert, J.-C., Livesey, N. J., Llewellyn, E. J., Lloyd, N. D., Mahieu, E., Manney, G. L., Marshall, B. T., McConnell, J. C., McCormick, M. P., McDermid, I. S., McHugh, M., McLinden, C. A., Mellqvist, J., Mizutani, K., Murayama, Y., Murtagh, D. P., Oelhaf, H., Parrish, A., Petelina, S. V., Piccolo, C., Pommereau, J.-P., Randall, C. E., Robert, C., Roth, C., Schneider, M., Senten, C., Steck, T., Strandberg, A., Strawbridge, K. B., Sussmann, R., Swart, D. P. J., Tarasick, D. W., Taylor, J. R., Tétard, C., Thomason, L. W., Thompson, A. M., Tully, M. B., Urban, J., Vanhellemont, F., Vigouroux, C., von Clarmann, T., von der Gathen, P., von Savigny, C., Waters, J. W., Witte, J. C., Wolff, M., and Zawodny, J. M.: Validation of ozone measurements from the Atmospheric Chemistry Experiment (ACE), *Atmos. Chem. Phys.*, 9, 287–343, doi:10.5194/acp-9-287-2009, 2009. 9880, 9881

Eckert, E., von Clarmann, T., Kiefer, M., Stiller, G. P., Lossow, S., Glatthor, N., Degenstein, D. A., Froidevaux, L., Godin-Beekmann, S., Leblanc, T., McDermid, S., Pastel, M., Steinbrecht, W., Swart, D. P. J., Walker, K. A., and Bernath, P. F.: Drift-corrected trends and periodic variations in MIPAS IMK/IAA ozone measurements, *Atmos. Chem. Phys. Discuss.*, 13, 17849–17900, doi:10.5194/acpd-13-17849-2013, 2013. 9888, 9889, 9934

Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., von Clarmann, T., Delbouille, L., Dudhia, A., Ehhalt, D., Endemann, M., Flaud, J. M., Gessner, R., Kleinert, A., Koopman, R., Langen, J., López-Puertas, M., Mosner, P., Nett, H., Oelhaf, H., Perron, G., Remedios, J., Ridolfi, M., Stiller, G., and Zander, R.: MIPAS: an instrument for atmospheric and climate research, *Atmos. Chem. Phys.*, 8, 2151–2188, doi:10.5194/acp-8-2151-2008, 2008. 9887

Froidevaux, L., Read, W. G., Lungu, T. A., Cofield, R. E., Fishbein, E. F., Flower, D. A., Jarnot, R. F., Ridenoure, B. P., Shippony, Z., Waters, J. W., Margitan, J. J., McDermid, I. S., Stachnik, R. A., Peckham, G. E., Braathen, G., Deshler, T., Fishman, J., Hofmann, D. J., and Oltmans, S. J.: Validation of UARS Microwave Limb Sounder ozone measurements, *J. Geophys. Res.*, 101, 10017–10060, 1996. 9886

Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sumnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Pe-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



run, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113, D15S20, doi:10.1029/2007JD008771, 2008. 9887

5 García, O. E., Schneider, M., Redondas, A., González, Y., Hase, F., Blumenstock, T., and Sepúlveda, E.: Investigating the long-term evolution of subtropical ozone profiles applying ground-based FTIR spectrometry, *Atmos. Meas. Tech.*, 5, 2917–2931, doi:10.5194/amt-5-2917-2012, 2012. 9870, 9871

10 Gazeaux, J., Clerbaux, C., George, M., Hadji-Lazaro, J., Kuttippurath, J., Coheur, P.-F., Hurtmans, D., Deshler, T., Kovilakam, M., Campbell, P., Guidard, V., Rabier, F., and Thépaut, J.-N.: Intercomparison of polar ozone profiles by IASI/MetOp sounder with 2010 Concordiasi ozonesonde observations, *Atmos. Meas. Tech.*, 6, 613–620, doi:10.5194/amt-6-613-2013, 2013. 9898

15 Gebhardt, C., Rozanov, A., Hommel, R., Weber, M., Bovensmann, H., Burrows, J. P., Degenstein, D., Froidevaux, L., and Thompson, A. M.: Stratospheric ozone trends and variability as seen by SCIAMACHY during the last decade, *Atmos. Chem. Phys. Discuss.*, 13, 11269–11313, doi:10.5194/acpd-13-11269-2013, 2013. 9892

Gille, J. and Russell III, J.: The limb infrared monitor of the stratosphere (LIMS): experiment description, performance and results, *J. Geophys. Res.*, 89, 1525–1540, 1984. 9884

20 Godin, S., Mégie, G., and Pelon, J.: Systematic lidar measurements of the stratospheric ozone vertical distribution, *Geophys. Res. Lett.*, 16, 547–550, 1989. 9868

Godin-Beekmann, S., Porteneuve, J., and Garnier, A.: Systematic DIAL ozone measurements at Observatoire de Haute-Provence, *J. Environ. Monitor.*, 5, 57–67, 2003. 9869

25 Harris, N. R. P., Hassler, B., Tummon, F., Bodeker, G. E., Petropavlovskikh, I., Fioletov, V. E., Steinbrecht, W., Davis, S. M., Wang, H. J., Froidevaux, L., Frith, S. M., Wild, J., Kyrölä, E., Sioris, C., and Zawodny, J.: SI²N Overview paper: Analysis and Interpretation of Changes in the Vertical Distribution of Ozone, in preparation, 2013. 9863, 9895, 9903

Hase, F.: Inversion von Spurengasprofilen aus hochaufgelösten bodengebundenen FTIR-Messungen in Absorption, *Wissenschaftliche Berichte Forschungszentrum Karlsruhe, FZKA 6512*, Karlsruhe, Forschungszentrum Karlsruhe GmbH, ISSN 0947-8620, 2000. 9870

30 Hase, F., Blumenstock, T., and Paton-Walsh, C.: Analysis of the instrumental line shape of high-resolution Fourier transform IR spectrometers with gas cell measurements and new retrieval software, *Appl. Optics*, 38, 3417–3422, 1999. 9870

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hassler, B., Bodeker, G. E., Cionni, I., and Dameris, M.: A vertically resolved, monthly mean, ozone database from 1979 to 2100 for constraining global climate model simulations, *Int. J. Remote Sens.*, 30, 4009–4018, 2009. 9863

Hassler, B., Daniel, J. S., Johnson, B. J., Solomon, S., and Oltmans, S. J.: An assessment of changing ozone loss rates at South Pole: twenty-five years of ozonesonde measurements, *J. Geophys. Res.*, 116, D22301, doi:10.1029/2011JD016353, 2011. 9861

Hocke, K., Kämpfer, N., Ruffieux, D., Froidevaux, L., Parrish, A., Boyd, I., von Clarmann, T., Steck, T., Timofeyev, Y. M., Polyakov, A. V., and Kyrölä, E.: Comparison and synergy of stratospheric ozone measurements by satellite limb sounders and the ground-based microwave radiometer SOMORA, *Atmos. Chem. Phys.*, 7, 4117–4131, doi:10.5194/acp-7-4117-2007, 2007. 9872

Hurtmans, D., Coheur, P.-F., Wespes, C., Clarisse, L., Scharf, O., Clerbaux, C., Hadji-Lazaro, J., George, M., and Turquety, S.: FORLI radiative transfer and retrieval code for IASI, *J. Quant. Spectrosc. Ra.*, 113, 1391–1408, 2012. 9898, 9934

Imai, K., Manago, N., Mitsuda, C., Naito, Y., Nishimoto, E., Sakazki, T., Fujiwara, M., Froidevaux, L., von Clarmann, T., Stiller, G., Murtagh, D., Rong, P.-P., Mlyniczak, M., Walker, K. A., Kinnison, D., Akiyoshi, H., Nakamura, T., Miyasaka, T., Nishibori, T., Mizobuchi, S., Kikuchi, K., Ozeki, H., Takahashi, C., Hayashi, H., Sano, T., Suzuki, M., Takayanagi, M., and Shiotani, M.: Validation of ozone data from the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES), *J. Geophys. Res.*, 118, 5750–5769, doi:10.1002/jgrd.50434, 2013. 9898, 9934

Intergovernmental Panel on Climate Change/Technology and Economic Assessment Panel (IPCC/TEAP): Safeguarding the Ozone Layer and the Global Climate System: Issues Related to Hydrofluorocarbons and Perfluorocarbons, Cambridge University Press, UK, 478 pp., 2005. 9862

Jégou, F., Urban, J., de La Noë, J., Ricaud, P., Le Flochmoën, E., Murtagh, D. P., Eriksson, P., Jones, A., Petelina, S., Llewellyn, E. J., Lloyd, N. D., Haley, C., Lumpe, J., Randall, C., Bevilacqua, R. M., Catoire, V., Huret, N., Berthet, G., Renard, J. B., Strong, K., Davies, J., Mc Elroy, C. T., Goutail, F., and Pommereau, J. P.: Technical Note: Validation of Odin/SMR limb observations of ozone, comparisons with OSIRIS, POAM III, ground-based and balloon-borne instruments, *Atmos. Chem. Phys.*, 8, 3385–3409, doi:10.5194/acp-8-3385-2008, 2008. 9890

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Joiner, J. and Vasilkov, A. P.: First results from the OMI rotational Raman scattering cloud pressure algorithm, *IEEE T. Geosci. Remote*, 44, 1272–1282, 2006. 9894
- Jones, A., Murtagh, D., Urban, J., Eriksson, P., and Rösevall, J.: Inter-comparison of Odin/SMR ozone measurements with MIPAS and balloon sonde data, *Can. J. Phys.*, 85, 1111–1123, 2007. 9890
- 5 Jones, A., Urban, J., Murtagh, D. P., Eriksson, P., Brohede, S., Haley, C., Degenstein, D., Bourassa, A., von Savigny, C., Sonkaew, T., Rozanov, A., Bovensmann, H., and Burrows, J.: Evolution of stratospheric ozone and water vapour time series studied with satellite measurements, *Atmos. Chem. Phys.*, 9, 6055–6075, doi:10.5194/acp-9-6055-2009, 2009. 9890
- 10 Kar, J., McElroy, C. T., Drummond, J. R., Zhou, J., Nichitiu, F., Walker, K. A., Randall, C. E., Nowlan, C. R., Dufour, D. G., Boone, C. D., Bernath, P. F., Treppe, C. R., Thomason, L. W., and McLinden, C.: Initial comparison of ozone and NO₂ profiles from ACE-MAESTRO with balloon and satellite data, *J. Geophys. Res.*, 112, D16301, doi:10.1029/2006JD008242, 2007. 9880
- 15 Kasai, Y., Sagawa, H., Kreyling, D., Dupuy, E., Baron, P., Mendrok, J., Suzuki, K., Sato, T. O., Nishibori, T., Mizobuchi, S., Kikuchi, K., Manabe, T., Ozeki, H., Sugita, T., Fujiwara, M., Irimajiri, Y., Walker, K. A., Bernath, P. F., Boone, C., Stiller, G., von Clarmann, T., Orphal, J., Urban, J., Murtagh, D., Llewellyn, E. J., Degenstein, D., Bourassa, A. E., Lloyd, N. D., Froidevaux, L., Birk, M., Wagner, G., Schreier, F., Xu, J., Vogt, P., Trautmann, T., and Yasui, M.: Validation of stratospheric and mesospheric ozone observed by SMILES from International Space Station, *Atmos. Meas. Tech.*, 6, 2311–2338, doi:10.5194/amt-6-2311-2013, 2013. 9898, 9899, 9934
- 20 Kerr, J. B., Fast, H., McElroy, C. T., Oltmans, S. J., Lathrop, J. A., Kyrö, E., Paukkunen, A., Claude, H., Köhler, U., Sreedharan, C. R., Takao, T., and Tsukagoshi, Y.: The 1991 WMO international ozonesonde intercomparison at Vanscoy, Canada, *Atmos. Ocean*, 32, 685–716, 1994. 9865, 9866
- 25 Kikuchi, K., Nishibori, T., Ochiai, S., Ozeki, H., Irimajiri, Y., Kasai, Y., Koike, M., Manabe, T., Mizukoshi, K., Murayama, Y., Nagahama, T., Sano, T., Sato, R., Seta, M., Takahashi, C., Takayanagi, M., Masuko, H., Inatani, J., Suzuki, M., and Shiotani, M.: Overview and early results of the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES), *J. Geophys. Res.*, 115, D23306, doi:10.1029/2010JD014379, 2010. 9898
- 30

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kobayashi, J. and Toyama, Y.: On various methods of measuring the vertical distribution of atmospheric ozone (III) – carbon iodine type chemical ozonesonde, *Pap. Meteorol. Geophys.*, 17, 113–126, 1996. 9865

Komhyr, W. D.: Electrochemical concentration cells for gas analysis, *Ann. Geophys.*, 25, 203–210, 1969,

<http://www.ann-geophys.net/25/203/1969/>. 9865

Kramarova, N. A., Bhartia, P. K., Frith, S. M., McPeters, R. D., and Stolarski, R. S.: Interpreting SBUV smoothing errors: an example using the quasi-biennial oscillation, *Atmos. Meas. Tech.*, 6, 2089–2099, doi:10.5194/amt-6-2089-2013, 2013a. 9895, 9934

Kramarova, N. A., Frith, S. M., Bhartia, P. K., McPeters, R. D., Taylor, S. L., Fisher, B. L., Labow, G. J., and DeLand, M. T.: Validation of ozone monthly zonal mean profiles obtained from the version 8.6 Solar Backscatter Ultraviolet algorithm, *Atmos. Chem. Phys.*, 13, 6887–6905, doi:10.5194/acp-13-6887-2013, 2013b. 9894, 9934

Kreyling, D., Sagawa, H., Wohltmann, I., Lehmann, R., and Kasai, Y.: SMILES zonal and diurnal variation climatology of strato- and mesospheric trace gases: O₃, HCl, HNO₃, ClO, BrO, HOCl, HO₂, and temperature, *J. Geophys. Res.*, 118, 1–16, doi:10.1002/2012JDO19420, 2013. 9898, 9934

Kroon, M., de Haan, J. F., Veefkind, J. P., Froidevaux, L., Wang, R., Kivi, R., and Hakkarainen, J. J.: Validation of operational ozone profiles from the ozone monitoring instrument, *J. Geophys. Res.*, 116, D18305, doi:10.1029/2010JD015100, 2011. 9895, 9897

Kuttippurath, J., Lefèvre, F., Pommereau, J.-P., Roscoe, H. K., Goutail, F., Pazmiño, A., and Shanklin, J. D.: Antarctic ozone loss in 1979–2010: first sign of ozone recovery, *Atmos. Chem. Phys.*, 13, 1625–1635, doi:10.5194/acp-13-1625-2013, 2013. 9861

Kyrölä, E., Tamminen, J., Sofieva, V., Bertaux, J. L., Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton d'Andon, O., Barrot, G., Guirlet, M., Mangin, A., Blanot, L., Fehr, T., Saavedra de Miguel, L., and Fraisse, R.: Retrieval of atmospheric parameters from GOMOS data, *Atmos. Chem. Phys.*, 10, 11881–11903, doi:10.5194/acp-10-11881-2010, 2010. 9883

Kyrölä, E., Laine, M., Sofieva, V., Tamminen, J., Päivärinta, S.-M., Tukiainen, S., Zawodny, J., and Thomason, L.: Combined SAGE II-GOMOS ozone profile data set for 1984–2011 and trend analysis of the vertical distribution of ozone, *Atmos. Chem. Phys.*, 13, 10645–10658, doi:10.5194/acp-13-10645-2013, 2013.

Laeng, A., Hubert, D., Verhoelst, T., von Clarmann, T., Dinelli, B. M., Dudhia, A., Raspollini, P., Stiller, G., Grabowski, U., Keppens, A., Kiefer, M., Sofieva, V., Froidevaux, L., Walker, K. A.,

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Lambert, J.-C., and Zehner, C.: The ozone climate change initiative: comparison of four level 2 processors for the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *Remote Sens. Environ.*, under review, 2013. 9889, 9934

Lambert, J.-C., Hubert, D., Adams, C., Damadeo, R. P., DeLand, M. T., Eckert, E., Garcia, O. E., Gebhardt, C., Godin-Beekmann, S., Harris, N. R. P., Hassler, B., Kasai, Y., Kirgis, G., Kra-marova, N. A., Kyrola, E., Laeng, A., Leblanc, T., Miyagawa, K., Nair, P. J., Petropavlovskikh, I., Rahpoe, N., Schneider, M., Sofieva, V. F., Staehelin, J., Studer, S., Vigouroux, C., and Weber, M.: SI²N overview paper: synthesis of ozone profiles intercomparisons in the SI²N context, in preparation, 2013. 9863, 9895, 9903

Levelt, P. F., Hilsenrath, E., Leppelmeier, G. W., Van Den Oord, G. H. J., Bhartia, P. K., Tamminen, J., de Haan, J. F., and Veefkind, J. P.: Science objectives of the ozone monitoring instrument, *IEEE T. Geosci. Remote*, 44, 1199–1208, doi:10.1109/TGRS.2006.872336, 2006a. 9896

Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O., and Saari, H.: The ozone monitoring instrument, *IEEE T. Geosci. Remote*, 44, 1093–1101, 2006b. 9896

Liu, G., Tarasick, D. W., Fioletov, V. E., Sioris, C. E., and Rochon, Y. J.: Ozone correlation lengths and measurement uncertainties from analysis of historical ozonesonde data in North America and Europe, *J. Geophys. Res.*, 114, D04112, doi:10.1029/2008JD010576, 2009. 9865

Liu, X., Bhartia, P. K., Chance, K., Spurr, R. J. D., and Kurosu, T. P.: Ozone profile retrievals from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 10, 2521–2537, doi:10.5194/acp-10-2521-2010, 2010. 9897

Livesey, N. J. and Read, W. G.: Direct retrieval of line-of-sight atmospheric structure from limb sounding observations, *Geophys. Res. Lett.*, 27, 891–894, 2000. 9886

Livesey, N. J., Read, W. G., Froidevaux, L., Waters, J. W., Pumphrey, H. C., Wu, D. L., Santee, M. L., Shippony, Z., and Jarnot, R. F.: The UARS Microwave Limb Sounder version 5 dataset: theory, characterization and validation, *J. Geophys. Res.*, 108, 4378, doi:10.1029/2002JD002273, 2003. 9886, 9933

Livesey, N. J., Snyder, W. V., Read, W. G., and Wagner, P. A.: Retrieval algorithms for the EOS Microwave Limb Sounder (MLS) instrument, *IEEE T. Geosci. Remote*, 44, 1144–1155, 2006. 9887

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Llewellyn, E. J., Lloyd, N. D., Degenstein, D. A., Gattinger, R. L., Petelina, S. V., Bourassa, A. E., Wiensz, J. T., Ivanov, E. V., McDade, I. C., Solheim, B. H., McConnell, J. C., Haley, C. S., von Savigny, C., Sioris, C. E., McLinden, C. A., Griffioen, E., Kaminski, J., Evans, W. F. J., Puckrin, E., Strong, K., Wehrle, V., Hum, R. H., Kendall, D. J. W., Matsushita, J., Murtagh, D. P., Brohede, S., Stegman, J., Witt, G., Barnes, G., Payne, W. F., Piché, L., Smith, K., Warshaw, G., Deslauniers, D.-L., Marchand, P., Richardson, E. H., King, R. A., Wevers, I., McCreath, W., Kyrölä, E., Oikarinen, L., Leppelmeier, G. W., Auvinen, H., Mégie, G., Hauchecorne, A., Lefèvre, F., de La Nöe, J., Ricaud, P., Frisk, U., Sjöberg, F., von Schéele, F., and Nordh, L.: The OSIRIS instrument on the Odin Spacecraft, *Can. J. Phys.*, 82, 411–422, 2004. 9892
- Lucke, R. L., Korwan, D. R., Bevilacqua, R. M., Hornstein, J. S., Shettle, E. P., Chen, D. T., Daehler, M., Lumpe, J. D., Fromm, M. D., Debrestian, D., Neff, B., Squire, M., König-Langlo, G., and Davies, J.: The Polar Ozone and Aerosol Measurement (POAM) III instrument and early validation results, *J. Geophys. Res.*, 104, 18785–18799, 1999. 9878
- Lumpe, J. D., Bevilacqua, R. M., Hoppel, K. W., and Randall, C. E.: POAM III retrieval algorithm and error analysis, *J. Geophys. Res.*, 107, 4575, doi:10.1029/2002JD002137, 2002. 9879, 9933
- Mäder, J. A., Staehelin, J., Peter, T., Brunner, D., Rieder, H. E., and Stahel, W. A.: Evidence for the effectiveness of the Montreal Protocol to protect the ozone layer, *Atmos. Chem. Phys.*, 10, 12161–12171, doi:10.5194/acp-10-12161-2010, 2010. 9861
- Maillard-Barras, E., Ruffieux, D., and Hocke, K.: Stratospheric ozone profiles over Switzerland measured by SOMORA, ozonesonde, and MLS/Aura satellite, *Int. J. Remote Sens.*, 30, 4033–4041, 2009. 9872
- McDermid, I. S., Godin, S. M., and Lindquist, L. O.: Ground-based laser DIAL system for long-term measurements of stratospheric ozone, *Appl. Optics*, 29, 3603–3612, 1990. 9868
- McDermid, I. S., Bergwerff, J. B., Bodeker, G., Boyd, I. S., Brinksma, E. J., Connor, B. J., Farmer, R., Gross, M. R., Kimvilakani, P., Matthews, W. A., McGee, T. J., Ornel, F. T., Parrish, A., Singh, U., Swart, D. P. J., Tsou, J. J., Wang, P. H., and Zawodny, J.: OPAL: Network for the detection of stratospheric change ozone profiler assessment at Lauder, New Zealand. II. Intercomparison of revised results, *J. Geophys. Res.*, 103, 28693–28699, 1998a. 9873
- McDermid, I. S., Bergwerff, J. B., Bodeker, G., Boyd, I. S., Brinksma, E. J., Connor, B. J., Farmer, R., Gross, M. R., Kimvilakani, P., Matthews, W. A., McGee, T. J., Ormel, F. T., Parrish, A., Singh, U., Swart, D. P. J., Tsou, J. J., Wang, P. H., and Zawodny, J.: OPAL: Net-

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



work for the detection of stratospheric change ozone profiler assessment at Lauder, New Zealand. I. Blind intercomparison, *J. Geophys. Res.*, 103, 28683–28692, 1998b. 9873

McElroy, C. T., Nowlan, C. R., Drummond, J. R., Bernath, P. F., Barton, D. V., Dufour, D. G., Midwinter, C., Hall, R. B., Ogyu, A., Ullberg, A., Wardle, D. I., Kar, J., Zou, J., Nichitui, F., Boone, C. D., Walker, K. A., and Rowlands, N.: The ACE-MAESTRO instrument on SCISAT: description, performance, and preliminary results, *Appl. Optics*, 46, 20, 4341–4356, 2007. 9879, 9933

McGee, T. J., Gross, M., Ferrare, R., Heaps, W. S., and Singh, U. N.: Raman DIAL measurements of stratospheric ozone in the presence of volcanic aerosols, *Geophys. Res. Lett.*, 20, 955–958, 1993. 9869

McLandress, C., Jonsson, A. I., Plummer, D. A., Reader, M. C., Scinocca, J. F., and Shepherd, T. G.: Separating the dynamical effects of climate change and ozone depletion. Part I Southern Hemisphere stratosphere, *J. Climate*, 23, 5002–5020, doi:10.1175/2010JCLI3586.1, 2010. 9862

McPeters, R. D. and Labow, G.: Climatology 2011: an MLS-derived ozone climatology for satellite retrieval algorithms, *J. Geophys. Res.*, 117, D10303, doi:10.1029/2011JD017006, 2012. 9894

McPeters, R. D., Miles, T., Flynn, L. E., Wellenmeyer, C. G., and Zawodny, J. M.: Comparison of SBUV and SAGE II profiles: implications for ozone trends, *J. Geophys. Res.*, 99, 20513–20524, 1994. 9902

McPeters, R. D., Hofmann, D. J., Clark, M., Froidevaux, L., Gross, M., Johnson, B., Koenig, G., McDermid, S., Murcray, F., Oltmans, S., Parrish, A., Singh, U., Tsou, J. J., and Zawodny, J.: Results from the 1995 stratospheric ozone profile intercomparison at Mauna Loa, *J. Geophys. Res.*, 104, 30505–30514, 1999. 9873

McPeters, R. D., Bhartia, P. K., Haffner, D., Labow, G. J., and Flynn, L.: The version 8.6 SBUV ozone data record: an overview, *J. Geophys. Res.*, 118, 8032–8039, doi:10.1002/jgrd.50597, 2013. 9893, 9934

Meijer, Y. J., Swart, D. P. J., Allaart, M., Andersen, S. B., Bodeker, G., Boyd, I., Braathena, G., Calisesia, Y., Claude, H., Dorokhov, V., von der Gathen, P., Gil, M., Godin-Beekmann, S., Goutail, F., Hansen, G., Karpetchko, A., Keckhut, P., Kelder, H. M., Koelemeijer, R., Kois, B., Koopman, R. M., Lambert, J.-C., Leblanc, T., McDermid, I. S., Pal, S., Kopp, G., Schets, H., Stubi, R., Suortti, T., Visconti, G., and Yela, M.: Pole-to-pole validation of ENVISAT/GOMOS

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ozone profiles using data from ground-based and balloon-sonde measurements, *J. Geophys. Res.*, 109, D23305, doi:10.1029/2004JD004834, 2004. 9882

Meijer, Y. J., Kelder, H. M., van der A, R. J., and van Oss, R. F.: Evaluation of Global Ozone Monitoring Experiment (GOME) ozone profiles from nine different algorithms, *J. Geophys. Res.*, 111, D21306, doi:10.1029/2005JD006778, 2006. 9895

Mieruch, S., Weber, M., von Savigny, C., Rozanov, A., Bovensmann, H., Burrows, J. P., Bernath, P. F., Boone, C. D., Froidevaux, L., Gordley, L. L., Mlynczak, M. G., Russell III, J. M., Thomason, L. W., Walker, K. A., and Zawodny, J. M.: Global and long-term comparison of SCIAMACHY limb ozone profiles with correlative satellite data (2002–2008), *Atmos. Meas. Tech.*, 5, 771–788, doi:10.5194/amt-5-771-2012, 2012. 9892, 9934

Mijling, B., Tuinder, O. N. E., van Oss, R. F., and van der A, R. J.: Improving ozone profile retrieval from spaceborne UV backscatter spectrometers using convergence behaviour diagnostics, *Atmos. Meas. Tech.*, 3, 1555–1568, doi:10.5194/amt-3-1555-2010, 2010. 9896

Mitsuda, C., Suzuki, M., Iwata, Y., Manago, N., Naito, Y., Takahashi, C., Imai, K., Nishimoto, E., Hayashi, H., Shiotani, M., Sano, T., Takayanagi, M., and Taniguchi, H.: Current status of level 2 product of Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES), *Proc. SPIE*, 8176, 81760M, doi:10.1117/12.898135, 2011. 9934

Miyagawa, K., Sasaki, T., Nakane, H., Petropavlovskikh, I., and Evans, R. D.: Reevaluation of long-term Umkehr data and ozone profiles at Japanese stations, *J. Geophys. Res.*, 114, D07108, doi:10.1029/2008JD010658, 2009. 9868

Moore, D. P. and Remedios, J. J.: Seasonality of Peroxyacetyl nitrate (PAN) in the upper troposphere and lower stratosphere using the MIPAS-E instrument, *Atmos. Chem. Phys.*, 10, 6117–6128, doi:10.5194/acp-10-6117-2010, 2010. 9888

Munro, R., Siddans, R., Reburn, W. J., and Kerridge, B. J.: Direct measurement of tropospheric ozone distributions from space, *Nature*, 392, 168–171, doi:10.1038/32392, 1998. 9895

Murtagh, D., Frisk, U., Merino, F., Ridal, M., Jonsson, A., Stegman, J., Witt, G., Eriksson, P., Jiménez, C., Megie, G., de la Nöe, J., Ricaud, P., Baron, P., Hauchcorne, A., Llewellyn, E. J., Degenstein, D. A., Gattinger, R. L., Lloyd, N. D., Evans, W. F. J., McDade, I. C., Haley, C. S., Sioris, C., von Savigny, C., Solheim, B. H., McConnell, J. C., Strong, K., Richardson, E. H., Leppelmeier, G. W., Kyrölä, E., Auvinen, H., and Oikarinen, L.: An overview of the Odin Atmospheric Mission, *Can. J. Phys.*, 80, 309–319, 2002. 9892

Nair, P. J., Godin-Beekmann, S., Kuttippurath, J., Ancellet, G., Goutail, F., Pazmiño, A., Froidevaux, L., Zawodny, J. M., Evans, R. D., Wang, H. J., Anderson, J., and Pastel, M.: Ozone

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- trends derived from the total column and vertical profiles at a northern mid-latitude station, *Atmos. Chem. Phys.*, 13, 10373–10384, doi:10.5194/acp-13-10373-2013, 2013. 9861, 9933
- Nakajima, H.: Preface to special section on ILAS-II: The Improved Limb Atmospheric Spectrometer-II, *J. Geophys. Res.*, 111, D20S90, doi:10.1029/2006JD007412, 2006. 9877
- 5 Nardi, B., Gille, J. C., Barnett, J. J., Randall, C. E., Harvey, V. L., Waterfall, A., Reburn, W. J., Leblanc, T., McGee, T. J., Twigg, L. W., Thompson, A. M., Godin-Beekmann, S., Bernath, P. F., Bojkov, B. R., Boone, C. D., Cavanaugh, C., Coffey, M. T., Craft, J., Craig, C., Dean, V., Eden, T. D., Francis, G., Froidevaux, L., Halvorson, C., Hannigan, J. W., Hoplewhite, C. L., Kinnison, D. E., Khosravi, R., Krinsky, C., Lambert, A., Lee, H., Loh, J.,
- 10 Massie, S. T., McDermid, I. S., Packman, D., Torpy, B., Valverde-Canossa, J., Walker, K. A., Whiteman, D. N., Witte, J. C., and Young, G.: Validation of HIRDLS ozone measurements, *J. Geophys. Res.*, 113, D16S36, doi:10.1029/2007JD008837, 2008. 9900
- Nowlan, C. R., McElroy, C. T., and Drummond, J. R.: Measurements of the O₂ A- and B-bands for determining temperature and pressure profiles from ACE-MAESTRO: forward model and retrieval algorithm, *J. Quant. Spectrosc. Ra.*, 108, 371–388, 2007. 9879
- 15 Ochiai, S., Kikuchi, K., Nishibori, T., Manabe, T., Ozeki, H., Mizobuchi, S., and Irimajiri, Y.: Receiver performance of the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) on the international space station, *IEEE T. Geosci. Remote*, 51, 3791–3802, doi:10.1109/TGRS.2012.2227758, 2013. 9899
- 20 Palm, M., Melsheimer, C., Noël, S., Heise, S., Notholt, J., Burrows, J., and Schrems, O.: Integrated water vapor above Ny Ålesund, Spitsbergen: a multi-sensor intercomparison, *Atmos. Chem. Phys.*, 10, 1215–1226, doi:10.5194/acp-10-1215-2010, 2010. 9872, 9873
- Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D.,
- 25 Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, 2012. 9898
- Parrish, A., Connor, B. J., Tsou, J. J., McDermid, I. S., and Chu, W. P.: Ground-based microwave monitoring of stratospheric ozone, *J. Geophys. Res.*, 97, 2541–2546, 1992. 9872
- 30 Pelon, J., Godin, S., and Megie, G.: Upper stratospheric (30–50 km) lidar observations of the ozone vertical distribution, *J. Geophys. Res.*, 91, 8666–8671, 1986. 9868

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Petropavlovskikh, I., Bhartia, P. K., and DeLuisi, J.: New Umkehr ozone profile retrieval algorithm optimized for climatological studies, *Geophys. Res. Lett.*, 32, L16808, doi:10.1029/2005GL023323, 2005. 9867, 9933

Petropavlovskikh, I., Evans, R., McConville, G., Oltmans, S., Quincy, D., Lantz, K., Disterhoft, P., Stanek, M., and Flynn, L.: Sensitivity of Dobson and Brewer Umkehr ozone profile retrievals to ozone cross-sections and stray light effects, *Atmos. Meas. Tech.*, 4, 1841–1853, doi:10.5194/amt-4-1841-2011, 2011. 9867

Polvani, L. M., Waugh, D. W., Correa, G. J. P., and Son, S. W.: Stratospheric ozone depletion: the main driver of twentieth-century atmospheric circulation changes in the Southern Hemisphere, *J. Climate*, 24, 795–812, doi:10.1175/2010JCLI3772.1, 2011. 9862

Pommier, M., Clerbaux, C., Law, K. S., Ancellet, G., Bernath, P., Coheur, P.-F., Hadji-Lazaro, J., Hurtmans, D., Nédélec, P., Paris, J.-D., Ravetta, F., Ryerson, T. B., Schlager, H., and Weinheimer, A. J.: Analysis of IASI tropospheric O₃ data over the Arctic during POLARCAT campaigns in 2008, *Atmos. Chem. Phys.*, 12, 7371–7389, doi:10.5194/acp-12-7371-2012, 2012. 9898

Pougatchev, N. S., Connor, B. J., and Rinsland, C. P.: Infrared measurements of the ozone vertical distribution above Kitt Peak, *J. Geophys. Res.*, 100, 16689–16697, 1995. 9870

Rahpoe, N., von Savigny, C., Weber, M., Rozanov, A.V., Bovensmann, H., and Burrows, J. P.: Error budget analysis of SCIAMACHY limb ozone profile retrievals using the SCIATRAN model, *Atmos. Meas. Tech. Discuss.*, 6, 4645–4676, doi:10.5194/amtd-6-4645-2013, 2013. 9892, 9934

Randall, C. E., Rusch, D. W., Bevilacqua, R. M., Hoppel, K. W., Lumpe, J. D., Shettle, E., Thompson, E., Deaver, L., Zawodny, J., Kyrö, E., Johnson, B., Kelder, H., Dorokhov, V. M., König-Langlo, G., and Gil, M.: Validation of POAM III ozone: comparisons with ozonesonde and satellite data, *J. Geophys. Res.*, 108, 4367, doi:10.1029/2002JD002944, 2003. 9933

Randel, W. J. and Wu, F.: A stratospheric ozone profile data set for 1979–2005: variability, trends, and comparisons with column ozone data, *J. Geophys. Res.*, 112, D06313, doi:10.1029/2006JD007339, 2007. 9862

Randel, W. J., Wu, F., Russell III, J. M., and Waters, J.: Space-time patterns of trends in stratospheric constituents derived from UARS measurements, *J. Geophys. Res.*, 104, 3711–3727, 1999. 9902

Raspolini, P., Belotti, C., Burgess, A., Carli, B., Carlotti, M., Ceccherini, S., Dinelli, B. M., Dudhia, A., Flaud, J.-M., Funke, B., Höpfner, M., López-Puertas, M., Payne, V., Piccolo, C.,

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Remedios, J. J., Ridolfi, M., and Spang, R.: MIPAS level 2 operational analysis, *Atmos. Chem. Phys.*, 6, 5605–5630, doi:10.5194/acp-6-5605-2006, 2006. 9888

Raspollini, P., Carli, B., Carlotti, M., Ceccherini, S., Dehn, A., Dinelli, B. M., Dudhia, A., Flaud, J.-M., López-Puertas, M., Niro, F., Remedios, J. J., Ridolfi, M., Sembhi, H., Sgheri, L., and von Clarmann, T.: Ten years of MIPAS measurements with ESA Level 2 processor V6 – Part 1: Retrieval algorithm and diagnostics of the products, *Atmos. Meas. Tech.*, 6, 2419–2439, doi:10.5194/amt-6-2419-2013, 2013. 9888, 9934

Reber, C. A., Trevathan, C. E., McNeal, R. J., and Luther, M. R.: The Upper Atmosphere Research Satellite (UARS) mission, *J. Geophys. Res.*, 98, 10643–10647, 1993. 9885

Remsberg, E., Gordley, L. L., Marshall, B. T., Thompson, R. E., Burton, J., Bhatt, P., Harvey, V. L., Lingenfeller, G., and Natarajan, M.: The Nimbus 7 LIMS version 6 radiance conditioning and temperature retrieval methods and results, *J. Quant. Spectrosc. Ra.*, 86, 395–424, doi:10.1016/j.jqsrt.2003.12.007, 2004. 9884

Remsberg, E., Lingenfeller, G., Natarajan, M., Gordley, L., Marshall, B. T., and Thompson, E.: On the quality of the Nimbus 7 LIMS version 6 ozone for studies of the middle atmosphere, *J. Quant. Spectrosc. Ra.*, 105, 492–518, doi:10.1016/j.jqsrt.2006.12.005, 2007. 9884, 9933

Remsberg, E., Fairlie, D., Natarajan, M., Wargan, K., Kim, G., Lingenfeller, G., and Pawson, S.: Impact of assimilating Nimbus 7 LIMS version 6 ozone into NASA's MERRA system, in: *Quadrennial Ozone Symposium (QOS)*, Toronto, Canada, 27–31 August 2012, 5788, 2012. 9885

Ridolfi, M., Carli, B., Carlotti, M., von Clarmann, T., Dinelli, B. M., Dudhia, A., Flaud, J. M., Höpfner, M., Morris, P. E., Raspollini, P., Stiller, G., and Wells, R. J.: Optimized forward model and retrieval scheme for MIPAS near-real-time data processing, *Appl. Optics*, 39, 1323–1340, 2000. 9888

Rodgers, C. D.: Retrieval of atmospheric temperature and composition from remote measurements of thermal radiation, *Rev. Geophys.*, 14, 609–624, doi:10.1029/RG014i004p00609, 1976. 9872, 9886

Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding: Theory and Practice*, Series on Atmospheric, Oceanic and Planetary Physics, Vol. 2, World Scientific Publishing Co., Singapore, 2000. 9870, 9886, 9888, 9895, 9897, 9899

Rong, P. P., Russell III, J. M., Mlynczak, M. G., Remsberg, E. E., Marshall, B. T., Gordley, L. L., and López-Puertas, M.: Validation of Thermosphere Ionosphere Mesosphere Energetics and Dynamics/Sounding of the Atmosphere using Broadband Emission Radiometry

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(TIMED/SABER) v1.07 ozone at 9.6 μm in altitude range 15–70 km, *J. Geophys. Res.*, 114, D04306, doi:10.1029/2008JD010073, 2009. 9900

Rothman, L. S., Jacquemart, D., Barbe, A., Chris Benner, D., Birk, M., Brown, L. R., Carleer, M. R., Chackerian, C., Chance, K., Coudert, L. H., Dana, V., Devi, V. M., Flaud, J.-M., Gamache, R. R., Goldman, A., Hartmann, J.-M., Jucks, K. W., Maki, A. G., Mandin, J.-Y., Massie, S. T., Orphal, J., Perrin, A., Rinsland, C. P., Smith, M. A. H., Tennyson, J., Tolchenov, R. N., Toth, R. A., Vander Auwera, J., Varanasi, P., and Wagner, G.: The HITRAN 2004 molecular spectroscopic database, *J. Quant. Spectrosc. Ra.*, 96, 139–204, doi:10.1016/j.jqsrt.2004.10.008, 2005. 9881

Rothman, L. S., Gordon, I. E., Barbe, A., Chris Benner, D., Bernath, P. F., Birk, M., Boudon, V., Brown, L. R., Campargue, A., Champion, J.-P., Chance, K., Coudert, L. H., Dana, V., Devi, V. M., Fally, S., Flaud, J.-M., Gamache, R. R., Goldman, A., Jacquemart, D., Kleiner, I., Lacombe, N., Lafferty, W. J., Mandin, J.-Y., Massie, S. T., Mikhailenko, S. N., Miller, C. E., Moazzen-Ahmadi, N., Naumenko, O. V., Nikitin, A. V., Orphal, J., Perevalov, V. I., Perrin, A., Predoi-Cross, A., Rinsland, C. P., Rotger, M., Šimečková, M., Smith, M. A. H., Sung, K., Tashkun, S. A., Tennyson, J., Toth, R. A., Vandaele, A. C., and Vander Auwera, J.: The HITRAN 2008 molecular spectroscopic database, *J. Quant. Spectrosc. Ra.*, 110, 533–572, doi:10.1016/j.jqsrt.2009.02.013, 2009. 9871

Russell III, J. M., Gordley, L. L., Park, J. H., Drayson, S. R., Tuck, A. F., Harries, J. E., Cicerone, R. J., Crutzen, P. J., and Frederick, J. E.: The Halogen Occultation Experiment, *J. Geophys. Res.*, 98, 10777–10797, 1993. 9876

Russell III, J. M., Mlynczak, M. G., Gordley, L. L., Tansock Jr., J. J., and Esplin, R.: Overview of the SABER experiment and preliminary calibration results, *Proc. SPIE Int. Soc. Opt. Eng.*, 3756, 277–288, 1999. 9900

Sakazaki, T., Fujiwara, M., Mitsuda, C., Imai, K., Manago, N., Naito, Y., Nakamura, T., Akiyoshi, H., Kinnison, D., Sano, T., Suzuki, M., and Shiotani, M.: Diurnal ozone variations in the stratosphere revealed in observations from the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) on board the International Space Station (ISS), *J. Geophys. Res.-Atmos.*, 118, 2991–3006, doi:10.1002/jgrd.50220, 2013. 9898

Sasano, Y.: Preface, *J. Geophys. Res.*, 107, 8204, doi:10.1029/2002JD002155, 2002. 9877

Scannell, C., Hurtmans, D., Boynard, A., Hadji-Lazaro, J., George, M., Delcloo, A., Tuinder, O., Coheur, P.-F., and Clerbaux, C.: Antarctic ozone hole as observed by IASI/MetOp for 2008–2010, *Atmos. Meas. Tech.*, 5, 123–139, doi:10.5194/amt-5-123-2012, 2012. 9898

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schneider, M., Hase, F., Blumenstock, T., Redondas, A., and Cuevas, E.: Quality assessment of O₃ profiles measured by a state-of-the-art ground-based FTIR observing system, *Atmos. Chem. Phys.*, 8, 5579–5588, doi:10.5194/acp-8-5579-2008, 2008. 9870

Smit, H. G. J.: Ozonesondes, in: *Encyclopedia of Atmospheric Sciences*, edited by: Holton, J., Pyle, J., and Curry, J., Academic Press, London, 1469–1476, 2002. 9865

Smit, H. G. J., Sträter, W., Helten, M., Kley, D., Ciupa, D., Claude, H., Köhler, U., Hoegger, B., Levrat, G., Johnson, B., Oltmans, S. J., Kerr, J. B., Tarasick, D. W., Davies, J., Shitamichi, M., Srivastav, S. K., and Vialle, C.: JOSIE: the 1996 WMO international intercomparison of ozonesondes under quasi-flight conditions in the environmental chamber at Jülich, in: *Atmospheric Ozone: Proceedings of the Quadrennial O₃ Symposium*, l'Aquila, Italy, edited by: Bojkov, R. D. and Visconti, G., *Parco Sci. e Tecnol. d'Abruzzo, Edigrafital, S. Atto*, Italy, 971–974, 1996. 9866

Smit, H. G. J., Sträter, W., Johnson, B., Oltmans, S., Davies, J., Tarasick, D. W., Hoegger, B., Stubi, R., Schmidlin, F., Northam, T., Thompson, A., Witte, J., Boyd, I., and Posny, F.: Assessment of the performance of ECC-ozonesondes under quasi-flight conditions in the environmental simulation chamber: insights from the Jülich Ozone Sonde Intercomparison Experiment (JOSIE), *J. Geophys. Res.*, 112, D19306, doi:10.1029/2006JD007308, 2007. 9865

Smit, H. G. J. and ASOPOS-panel: Quality Assurance and Quality Control for Ozonesonde Measurements in GAW, WMO Global Atmosphere Watch report series, No. 201, World Meteorological Organization, Geneva, available at: http://www.wmo.int/pages/prog/arep/gaw/documents/GAW_201_30_Sept.pdf (last access:8 Novembe 2013), 2012.

Sofieva, V. F., Rahpoe, N., Tamminen, J., Kyrölä, E., Kalakoski, N., Weber, M., Laeng, A., von Clarmann, T., Stiller, G., Lossow, S., Degenstein, D., Bourassa, A., Adams, C., Roth, C., Lloyd, N., Bernath, P., Hargreaves, R. J., Urban, J., Murtagh, D., Hauchecorne, A., Van Roozendaal, M., Kalb, N., and Zehner, C.: Harmonized dataset of ozone profiles from satellite limb and occultation measurements, *Earth Syst. Sci. Data Discuss.*, 6, 189–222, doi:10.5194/essdd-6-189-2013, 2013.

Sonkaew, T., Rozanov, V. V., von Savigny, C., Rozanov, A., Bovensmann, H., and Burrows, J. P.: Cloud sensitivity studies for stratospheric and lower mesospheric ozone profile retrievals from measurements of limb-scattered solar radiation, *Atmos. Meas. Tech.*, 2, 653–678, doi:10.5194/amt-2-653-2009, 2009. 9891

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SPARC: SPARC-IOC-GAW Assessment of Trends in the Vertical Distribution of Ozone, SPARC Report No. 1, WMO Global Ozone Research and Monitoring Project Report No. 43, World Meteorological Organization, Geneva, 1998. 9861, 9866, 9928

5 Steck, T., von Clarmann, T., Fischer, H., Funke, B., Glatthor, N., Grabowski, U., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Milz, M., Stiller, G. P., Wang, D. Y., Allaart, M., Blumenstock, Th., von der Gathen, P., Hansen, G., Hase, F., Hochschild, G., Kopp, G., Kyrö, E., Oelhaf, H., Raffalski, U., Redondas Marrero, A., Remsberg, E., Russell III, J., Stebel, K., Steinbrecht, W., Wetzol, G., Yela, M., and Zhang, G.: Bias determination and precision validation of ozone profiles from MIPAS-Envisat retrieved with the IMK-IAA processor, *Atmos. Chem. Phys.*, 7, 3639–3662, doi:10.5194/acp-7-3639-2007, 2007. 9888

10 Steinbrecht, W., Claude, H., Schönenborn, F., McDermid, I. S., Leblanc, T., Godin-Beekmann, S., Keckhut, P., Hauchecorne, A., van Gijssel, J. A. E., Swart, D. P. J., Bodeker, G. E., Parrish, A., Boyd, I. S., Kämpfer, N., Hocke, K., Stolarski, R. S., Frith, S. M., Thomason, L. W., Remsberg, E. E., von Savigny, C., Rozanov, A., and Burrows, J. P.: Ozone and temperature trends in the upper stratosphere at five stations of the Network for the Detection of Atmospheric Composition Change, *Int. J. Remote Sens.*, 30, 3875–3886, doi:10.1080/01431160902821841, 2009. 9869, 9873

15 Stiller, G. P., Kiefer, M., Eckert, E., von Clarmann, T., Kellmann, S., García-Comas, M., Funke, B., Leblanc, T., Fetzer, E., Froidevaux, L., Gomez, M., Hall, E., Hurst, D., Jordan, A., Kämpfer, N., Lambert, A., McDermid, I. S., McGee, T., Miloshevich, L., Nedoluha, G., Read, W., Schneider, M., Schwartz, M., Straub, C., Toon, G., Twigg, L. W., Walker, K., and Whiteman, D. N.: Validation of MIPAS IMK/IAA temperature, water vapor, and ozone profiles with MOHAVE-2009 campaign measurements, *Atmos. Meas. Tech.*, 5, 289–320, doi:10.5194/amt-5-289-2012, 2012. 9888

20 Studer, S., Hocke, K., Pastel, M., Godin-Beekmann, S., and Kämpfer, N.: Intercomparison of stratospheric ozone profiles for the assessment of the upgraded GROMOS radiometer at Bern, *Atmos. Meas. Tech. Discuss.*, 6, 6097–6146, doi:10.5194/amtd-6-6097-2013, 2013. 9872, 9873

25 Sugita, T., Yokota, T., Nakajima, H., Kanzawa, H., Nakane, H., Gernandt, H., Yushkov, V., Shibasaki, K., Deshler, T., Kondo, Y., Godin, S., Goutail, F., Pommereau, J.-P., Camy-Peyret, C., Payan, S., Jeseck, P., Renard, J.-B., Bösch, H., Fitzenberger, R., Pfeilsticker, K., von König, M., Bremer, H., Küllmann, H., Schlager, H., Margitan, J. J., Stachnik, B., Toon, G. C., Jucks, K., Traub, W. A., Johnson, D. G., Murata, I., Fukunishi, H., and Sasano, Y.:

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Validation of ozone measurements from the Improved Limb Atmospheric Spectrometer, *J. Geophys. Res.*, 107, 8212, doi:10.1029/2001JD000602, 2002. 9877

Sugita, T., Nakajima, H., Yokota, T., Kanzawa, H., Gernandt, H., Herber, A., von der Gathen, P., König-Langlo, G., Sato, K., Dorokhov, V., Yushkov, V. A., Murayama, Y., Yamamori, M., Godin-Beekmann, S., Goutail, F., Roscoe, H. K., Deshler, T., Yela, M., Taalas, P., Kyrö, E., Oltmans, S. J., Johnson, B. J., Allaart, M., Litynska, Z., Klekociuk, A., Andersen, S. B., Braathen, G. O., De Backer, H., Randall, C. E., Bevilacqua, R. M., Taha, G., Thomason, L. W., Irie, H., Ejiri, M. K., Saitoh, N., Tanaka, T., Terao, Y., Kobayashi, H., and Sasano, Y.: Ozone profiles in the high-latitude stratosphere and lower mesosphere measured by the Improved Limb Atmospheric Spectrometer (ILAS)-II: comparison with other satellite sensors and ozonesondes, *J. Geophys. Res.*, 111, D11S02, doi:10.1029/2005JD006439, 2006. 9877

Tamminen, J., Kyrölä, E., Sofieva, V. F., Laine, M., Bertaux, J.-L., Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton-d'Andon, O., Barrot, G., Mangin, A., Guirlet, M., Blanot, L., Fehr, T., Saavedra de Miguel, L., and Fraisse, R.: GOMOS data characterisation and error estimation, *Atmos. Chem. Phys.*, 10, 9505–9519, doi:10.5194/acp-10-9505-2010, 2010. 9883

Tarasick, D. W. and Slater, R.: Ozone in the troposphere: measurements, climatology, budget, and trends, *Atmos. Ocean*, 46, 93–115, doi:10.3137/ao.460105, 2008. 9865

Terao, Y. and Logan, J. A.: Consistency of time series and trends of stratospheric ozone as seen by ozonesonde, SAGE II, HALOE, and SBUV(2), *J. Geophys. Res.*, 112, D06310, doi:10.1029/2006JD007667, 2007. 9902

Tikhonov, A. N.: On the solution of incorrectly stated problems and a method of regularization, *Dokl. Acad. Nauk SSSR*, 151, 501, 1963. 9886

Toohey, M., Hegglin, M. I., Tegtmeier, S., Anderson, J., Anel, J. A., Bourassa, A., Brohede, S., Degenstein, D., Froidevaux, L., Fuller, R., Funke, B., Gille, J., Jones, A., Kasai, Y., Kruger, K., Kyrölä, E., Neu, J. L., Rozanov, A., Smith, L., Urban, J., von Clarmann, T., Walker, K. A., and Wang, R.: Characterizing sampling biases in the trace gas climatologies of the SPARC Data Initiative, *J. Geophys. Res.*, in press, 2013. 9874, 9901

Tsou, J. J., Connor, B. J., Parrish, A., McDermid, I. S., Pierce, R. B., Boyd, I. S., Bodeker, G. E., Chu, W. P., Russell III, J. M., Swart, D. P. J., and McGee, T. J.: NDSC millimeter wave observations at Lauder, New Zealand, 1992–1998, improved methodology, validation, and variation study, *J. Geophys. Res.*, 105, 24263–24281, 2000.

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tukiainen, S., Kyrölä, E., Verronen, P. T., Fussen, D., Blanot, L., Barrot, G., Hauchecorne, A., and Lloyd, N.: Retrieval of ozone profiles from GOMOS limb scattered measurements, *Atmos. Meas. Tech.*, 4, 659–667, doi:10.5194/amt-4-659-2011, 2011. 9883
- Urban, J., Lautié, N., Jimenez, C., Le Flochmoën, E., Dupuy, E., Eriksson, P., Frisk, U., Murtagh, D., de La Noë, J., Olberg, M., and Ricaud, P.: Odin/SMR limb observations of stratospheric trace gases: level-2 processing of ClO, N₂O, O₃, and HNO₃, *J. Geophys. Res.*, 110, D14307, doi:10.1029/2004JD005741, 2005. 9890
- Urban, J., Murtagh, D., Lautié, N., Barret, B., Dupuy, E., de La Noë, J., Eriksson, P., Frisk, U., Jones, A., Le Flochmoë, E., Olberg, M., Piccolo, C., Ricaud, P., and Rösevall, J.: Odin/SMR limb observations of trace gases in the Polar lower stratosphere during 2004–2005, in: *Proc. ESA Atmospheric Science Conference, European Space Agency publications ESA-SP-628, Frascati, Italy, 8–12 May 2006, ISBN-92-9092-939-1, ISSN-1609-042X, 2006. 9890*
- van Gijsel, J. A. E., Swart, D. P. J., Baray, J.-L., Bencherif, H., Claude, H., Fehr, T., Godin-Beekmann, S., Hansen, G. H., Keckhut, P., Leblanc, T., McDermid, I. S., Meijer, Y. J., Nakane, H., Quel, E. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Tatarov, B. I., and Wolfram, E. A.: GOMOS ozone profile validation using ground-based and balloon sonde measurements, *Atmos. Chem. Phys.*, 10, 10473–10488, doi:10.5194/acp-10-10473-2010, 2010. 9882, 9883
- Van Peet, J. C. A., van der A, R. J., de Laat, A. T. J., Tuinder, O. N. E., Koenig-Langlo, G., and Wittig, J.: Height resolved ozone hole structure as observed by the Global Ozone Monitoring Experiment-2, *Geophys. Res. Lett.*, 36, L11816, doi:10.1029/2009GL038603, 2009. 9896
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., Levelt, P. F.: TROPOMI on the ESA sentinel-5 precursor: a GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, doi:10.1016/j.rse.2011.09.027, 2012. 9896
- Vigouroux, C., De Mazi Vigouroux, C., De Maziere, M., Demoulin, P., Servais, C., Hase, F., Blumenstock, T., Kramer, I., Schneider, M., Mellqvist, J., Strandberg, A., Velasco, V., Notholt, J., Sussmann, R., Stremme, W., Rockmann, A., Gardiner, T., Coleman, M., and Woods, P.: Evaluation of tropospheric and stratospheric ozone trends over Western Europe from ground-based FTIR network observations, *Atmos. Chem. Phys.*, 8, 6865–6886, doi:10.5194/acp-8-6865-2008, 2008. 9870, 9871, 9933

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Von Clarman, T., Glatthor, N., Grabowski, U., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Tsidu, G. M., Milz, M., Steck, T., Stiller, G. P., Wang, D. Y., and Fischer, H.: Retrieval of temperature and tangent altitude pointing from limb emission spectra recorded from space by the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *J. Geophys. Res.*, 108, 4736, doi:10.1029/2003JD003602, 2003. 9888
- 5 von Clarman, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H₂O, O₃, HNO₃, CH₄, N₂O, ClONO₂ and ClO from MIPAS reduced resolution nominal mode limb emission measurements, *Atmos. Meas. Tech.*, 2, 159–175, doi:10.5194/amt-2-159-2009, 2009. 9888, 9934
- 10 von Clarman, T., Stiller, G., Grabowski, U., Eckert, E., and Orphal, J.: Technical Note: Trend estimation from irregularly sampled, correlated data, *Atmos. Chem. Phys.*, 10, 6737–6747, doi:10.5194/acp-10-6737-2010, 2010. 9889
- von Savigny, C., Kaiser, J. W., Bovensmann, H., Burrows, J. P., McDermid, I. S., and Leblanc, T.: Spatial and temporal characterization of SCIAMACHY limb pointing errors during the first three years of the mission, *Atmos. Chem. Phys.*, 5, 2593–2602, doi:10.5194/acp-5-2593-2005, 2005. 9891
- 15 Wang, H. J., Cunnold, D. M., and Bao, X.: A critical analysis of stratospheric aerosol and gas experiment ozone trends, *J. Geophys. Res.*, 101, 12495–12514, 1996. 9902
- 20 Wang, H. J., Cunnold, D. M., Thomason, L. W., Zawodny, J. M., and Bodeker, G. E.: Assessment of SAGE version 6.1 ozone data quality, *J. Geophys. Res.*, 107, 4691, doi:10.1029/2002JD002418, 2002. 9902
- Waters, J. W.: Microwave limb sounding, in: *Atmospheric Remote Sensing by Microwave Radiometry*, edited by: Janssen, M. A., chap. 8, J. Wiley, New York, USA, 383–434, 1993. 9885
- 25 Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. E., Filipiak, M. J., Flower, D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo, M. S., Perun, V. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, S., Chavez, M. C., Chen, G.-S., Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M. A., Jiang, J. H., Jiang, Y., Knosp, B. W., LaBelle, R. C., Lam, J. C.,
- 30 Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, D. M., Snyder, W. V., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth Observing System

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Microwave Limb Sounder (EOS MLS) on the Aura satellite, IEEE T. Geosci. Remote, 44, 1075–1092, doi:10.1109/TGRS.2006.873771, 2006. 9886

WMO (World Meteorological Organization): Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project – Report No. 50, Geneva, Switzerland, 2007. 9861

WMO (World Meteorological Organization): Towards a Better Knowledge of Umkehr Measurements: A Detailed Study of Data from Thirteen Dobson Intercomparisons, GAW Report No. 180, Geneva, Switzerland, 2008. 9867

WMO (World Meteorological Organization): Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project – Report No. 52, Geneva, Switzerland, 2011. 9861, 9871

Worden, H. M., Logan, J., Worden, J. R., Beer, R., Bowman, K., Clough, S. A., Eldering, A., Fisher, B., Gunson, M. R., Herman, R. L., Kulawik, S. S., Lampel, M. C., Luo, M., Megretskaia, I. A., Osterman, G. B., and Shephard, M. W.: Comparisons of Tropospheric Emission Spectrometer (TES) ozone profiles to ozonesondes: methods and initial results, J. Geophys. Res., 112, D03309, doi:10.1029/2006JD007258, 2007. 9900

Yu, J. R. and She, C. Y.: Climatology of a midlatitude mesopause region observed by a lidar at Fort Collins, Colorado (40.6° N, 105° W), J. Geophys. Res., 100, 7441–7452, 1995. 9902

Zanis, P., Maillard, E., Staehelin, J., Zerefos, C., Kosmidis, E., Tourpali, K., and Wohltmann, I.: On the turnaround of stratospheric ozone trends deduced from the reevaluated Umkehr record of Arosa, Switzerland, J. Geophys. Res., 111, D22307, doi:10.1029/2005JD006886, 2006. 9868

Ziemke, J. R. and Chandra, S.: Development of a climate record of tropospheric and stratospheric column ozone from satellite remote sensing: evidence of an early recovery of global stratospheric ozone, Atmos. Chem. Phys., 12, 5737–5753, doi:10.5194/acp-12-5737-2012, 2012. 9903

SI²N: ozone profile measurements

B. Hassler et al.

Table 1a. Summary of measurement techniques and lowest and highest covered level for the five ground-based measurement systems and all described satellite measurement systems.

Name of instrument	Measurement technique	Covered vertical region: lowest level	Covered vertical region: highest level
Ground-based measurement systems			
Ozonesondes Dobson/Brewer Lidar	Electrochemical conversion of ozone Umkehr, ground-based, UV part of the spectrum Differential Absorption Lidar (DIAL), ground-based, UV laser radiation	surface tropospheric layer: surface to ≈ 250 hPa tropopause	30–35 km altitude (bursting point of balloon) to top of atmosphere 45–55 km depending on the lidar systems
FTIR	Ground-based solar absorption measurements in the infrared	Surface	40–45 km
Microwave	Measurement of the emission spectrum of a thermally excited rotational transition of ozone in the millimeter wavelength range. The ozone profile is retrieved from the pressure-broadened line shape.	Typically ≈ 20 km	Varies by spectral line frequency. Up to 72 km at night, and 66 km during the day for the transition at 110.836 GHz.
Satellite-based measurement systems			
SAGE HALOE	Solar Occultation Solar occultation instrument; 9.4 to 10.4 μm ozone channel (SPARC, 1998)	tropopause or 10 km whichever is lower cloud top or to just below the tropopause	55 km to near 0.005 hPa for temperature and ozone
ILAS POAM ACE-MAESTRO	satellite-based, solar-occultation Solar Occultation in the UV/Vis spectral range Solar occultation spectrophotometry in the Chappuis band	cloud top: ≈ 10 km 5 km (or cloud top) ≈ 5 km	to top of atmosphere 60 km ≈ 35 km
ACE-FTS GOMOS	Satellite-based, solar occultation, infrared spectrum Satellite based stellar occultation instrument in UV-VIS-NIR	cloud tops (≈ 5 km) Typically cloud top, however, lowest valid altitude ≈ 15 km	95 km 100 km
LIMS	Satellite-based instrument, limb-infrared emission experiment; 9 to 10 micrometer band for ozone	250 to 300 hPa or higher cloud top	to near 0.01 hPa for T and ozone
UARS MLS Aura MLS MIPAS	Microwave limb emission sounding (satellite) Microwave limb emission sounding (satellite) satellite-based mid-infrared limb emission	100 hPa 215 hPa It depends on the measurement mode. Most data are in Nominal observation mode: 6 km (or cloud top) for 2002 to 2004 7–12 km (following tropopause height) or cloud top for 2005 to 2012; Middle/upper atmosphere modes: above 20 or above 40 km These are nominal numbers; real numbers can vary.	0.2 hPa 0.02 hPa $\approx 68/72$ km (2002–2004/2005–2012) in nominal observation mode; up to 100 km in middle/upper atmosphere modes.
Odin-SMR SCIAMACHY Odin-OSIRIS SBUV	Microwave limb emission sounding (satellite) UV/Vis limb scatter Satellite based, UV-visible, limb scatter satellite-based nadir-view backscattered UV radiation	≈ 12 km (poles), 17–18 km (mid-latitudes) tropopause (10 km) cloud tops surface to 16 hPa (tropics);	≈ 60 km mesosphere (60 km) 55 km top of atmosphere
GOME/GOME-2	Optimal Estimation method, satellite-based instrument looking in nadir direction, UV-VIS part of the spectrum	surface to 32 hPa (mid-latitudes); surface to 40 hPa (high latitudes) surface	top of atmosphere
OMI IASI	Fourier Transform spectrometer, Thermal infrared	tropospheric layer or lower if thermal contrast is good	total columns (sensitivity up to around 45 km)
SMILES	Submillimeter-wave limb emission sounding (on-board the International Space Station)	≈ 16 km	≈ 95 km



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1b. Summary of the vertical resolution, native vertical grid and native ozone units for the five ground-based measurement systems and all described satellite measurement systems.

Name of instrument	Vertical resolution ¹	Representation grid	Native vertical grid ¹	Native ozone units ¹
Ground-based measurement systems				
Ozonesondes	100–150 m (at ascent rate of 5 ms ⁻¹ and response time of 20–30 s)	100–200 m	Flight time	Partial ozone pressure in milli Pascal (mPa)
Dobson/Brewer	≈ 10 km	≈ 10 km	Pressure levels	Dobson units (DU)
Lidar	The resolution varies as a function of altitude from 0.5 km below 20 km, 2 km around 30 km to more than 5 km above 45 km.	Depending on the systems: 0.15 or 0.3 km	Altitude levels	Ozone number density (cm ⁻³)
FTIR	≈ 8–10 km for the ground ≈ 9 km layer ≈ 8 km for the lower and middle stratospheric layer ≈ 15–20 km for the upper stratospheric layer (≈ 28–45 km)	In NDACC HDF archived files: the retrieval grid contains, depending on the station, about 41 to 47 levels (from ground to 100 km).	Altitude levels	Volume mixing ratio (ppmv) (provided also as partial columns (mol cm ⁻²))
Microwave	Typ. 8–10 km, from ≈ 20 to 40 km, then increasing to ≈ 17 km at ≈ 60 km	Varies by instrument. Typ. ≈ 2 km intervals	Pressure levels (grid on which data are provided varies by instrument)	Volume mixing ratio (ppmv)
Satellite-based measurement systems				
SAGE	1 km	1 km	Altitude levels	Ozone number density (cm ⁻³)
HALOE	Measurement: 1.8 km instantaneous field of view (IFOV); retrieval resolution is 2.3 km	2.3 km vertical resolution	Solar zenith angle or apparent altitude, sampled every 0.3 km. (provided also on pressure and associated hydrostatic altitude levels)	Transmission vs. solar zenith angle or apparent altitude (provided as volume mixing ratio (ppmv))
ILAS	ILAS-I: ≈ 1.9 km ILAS-II: ≈ 1.6 km	≈ 1 km	Tangent height (in km)	Volume mixing ratio (ppmv)
POAM	≈ 1.5 km	≈ 1 km	Altitude levels	Ozone number density (cm ⁻³)
ACE-MAESTRO	≈ 1.2 km	≈ 1.0 km	Tangent levels	Concentration (mol cm ⁻²) (provided as volume mixing ratio (ppmv))
ACE-FTS	≈ 3–4 km	1 km (interpolated to standard grid) and ≈ 3–4 km (as measured)	Altitude levels	Volume mixing ratio (ppmv)
GOMOS	Resolution of ozone measurements: 2 km below 30 km 3 km below 40 km smooth transition between 30–40 km	Vertical sampling resolution: about 0.5–1.7 km. Variable for each profile.	Altitude levels	Ozone number density (cm ⁻³)
LIMS	Measurement: 1.8 km instantaneous field of view (IFOV); Retrieval: 3.7 km	Level 2 data are provided every 0.88 km, having a vertical resolution of ≈ 3.7 km	Pressure levels (data are also provided on altitude levels)	Volume mixing ratio (ppmv)
UARS MLS	3.5–4 km in the stratosphere and 5–8 km in the mesosphere	3 km (6 levels per decade change in pressure for stratosphere)	Pressure levels	Volume mixing ratio (ppmv)

¹ specified by instrument/averaging kernel.

SI²N: ozone profile measurements

B. Hassler et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 1b. Continued.

Name of instrument	Vertical resolution ¹	Representation grid	Native vertical grid ¹	Native ozone units ¹
Aura MLS	2.5–3 km in upper troposphere to lower mesosphere, 4 to 6 km in upper mesosphere	≈ 3 km (6 levels per decade change in pressure) for pressures larger than 0.1 hPa; grid resolution degrades by a factor of two for pressures less than 0.1 hPa.	Pressure levels	Volume mixing ratio (ppmv)
MIPAS	Optimized resolution nominal mode (years 2005–2012) IMK: 2.4–3.6 km (reduced resolution nominal mode) ESA processor: 2 km at 10 km, 3.2 km at 30 km, 5 km at 70 km, mostly independent on atmospheric conditions Full resolution mode (years 2002–2004) IMK: 3.5 km at 10 km, 5 km at 42 km, 8 km at 60 km. ESA: 4 km at 10 km, 5 km at 30 km, 8 km at 70 km.	IMK: retrieval and data representation grid for reduced resolution nominal mode is: 1 km from 4–44 km, 2 km from 44–70 km; 5 km above ESA processor: grid of the measured tangent altitudes MIPAS2D: 3 km from 6 to 42 then 47, 52, 60 and 68 km	IMK: Altitude levels (ESA: provides pressure levels where the retrieved pressure corresponds to the measured tangent altitude)	Volume mixing ratio (ppmv)
Odin-SMR	2.5–3.5 km	1.5 km and 3 km	Pointing information is retrieved from pressure broadened spectral lines (provided on altitude levels: altitude-pressure-temperature information provided for interpolation between grids) Geometric height (km) Altitude levels Pressure levels	Volume mixing ratio (ppmv)
SCIAMACHY Odin-OSIRIS SBUV	3–5 km ≈ 1.5 km in UTLS and 2 km higher up ≈ 6 km at 3 hPa ≈ 15 km in troposphere	1 km ≈ 1 km 3.2 km	Geometric height (km) Altitude levels Pressure levels	Ozone number density (cm ⁻³) Ozone number density (cm ⁻³) Dobson units (provided also as volume mixing ratio (ppmv)) Dobson units (DU)
GOME/GOME-2 OMI	≈ 7–15 km	≈ 4 km	Pressure levels	Dobson units (DU)
IASI	≈ 7 km	1 km up to 40 km 2 km from 40 km to 60 km	Altitude levels	Ozone number density (cm ⁻³) (provided as volume mixing ratio (ppmv) or ozone partial columns (DU)) Volume mixing ratio (ppmv)
SMILES	2–4 km	2.5 km	Altitude levels	

¹ specified by instrument/averaging kernel



SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1c. Summary of the covered time period, the number of measurements and the time for measurements for the five ground-based measurement systems and all described satellite measurement systems.

Name of instrument	Covered time period	Average number of measurements (per day/per week/per month)	Measurement time of day
Ground-based measurement systems			
Ozonesondes	Varies from station to station, longest record from mid 1960's, early 1970's	1–2 vertical ozone soundings per week	Ozonesondes are usually launched at daytime before about noon time
Dobson/Brewer	Varies from station to station, longest record from Arosa/Switzerland: 1956–present	Average is about 15 per months (\approx 200 per year; depending on station and season)	Sunrise and sunset between 70 and 90° solar zenith angle (SZA); clear sky limitation.
Lidar	Varies from station to station, the longest records start in the late eighties (OHP, France; Hohenpeisenberg, Germany; Table Mountain, USA)	Up to 20 measurements per month (depending on weather conditions)	nighttime (only with clear sky)
FTIR	Varies from station to station, usually 1995–present.	Mean of all stations: \approx 2.5 per day (7 per week; 15 per month)	During the day.
Microwave	1992.9-present (Lauder), 1995.6-present (Mauna Loa), 2000–present (Payerne). Other records back to 1990's may be available.	4 to 48 profiles per day in suitable weather, depending on station	Measurements are made at all hours of day and night. Clear-sky conditions needed.
Satellite-based measurement systems			
SAGE	SAGE I: Feb 1979 through Nov 1981 SAGE II: Oct 1984 through Jul 2005 SAGE III: Dec 2001 through Mar 2006	Up to 15 Sunrise and 15 Sunset events per day	Sunrise/sunset
HALOE	Oct 1991 to Nov 2005.	Up to 15 Sunrise and 15 Sunset events per day	Sunrise/sunset
ILAS	ILAS-I: 30 Oct 1996 to 29 Jun 1997 ILAS-II: 2 Apr 2003 to 24 Oct 2003	ILAS-I: 28 profiles per day (14 sunrise and 14 sunset profiles) ILAS-II: 28 profiles per day (14 sunrise and 14 sunset profiles)	Sunrise/sunset Sunrise/sunset
POAM	POAM II: Sep 1993 to Nov 1996	14 measurements per day in each hemisphere	Measurement performed at local sunset in NH, but in the SH they are at local sunset in the winter and local sunrise in the summer.
ACE-MAESTRO	POAM III: Apr 1998 through Nov 2005	Up to 30 profiles per day	Sunrise/sunset
ACE-FTS	21 Feb 2004–present	Up to 30 profiles per day	Sunrise/sunset
GOMOS	Aug 2002–Apr 2012 Reduced number of measurements due to instrumental problems in year 2005 and since Oct 2011.	daily: 100–200 profiles monthly: 3000–6000 profiles	night time (around 10 p.m.)
LIMS	25 Oct 1978, to 28 May 1979.	yearly: 36 000–72 000 profiles \approx 3000 profiles per day (the LIMS observation duty cycle was of the order of 5 days on and one day off for most of its mission life)	LIMS operated in a Sun-synchronous mode, taking its tangent point scans at about 1300 and 2300 h (at the Equator), but with little change of local time with latitude until reaching high latitudes.
UARS MLS	Sep 1991 to Jul 1999	\approx 1350 profiles/day	Day and night measurements are routinely performed around the orbit, with all local times covered at any given latitude during each satellite yaw cycle (every 36 days on average)
Aura MLS	Aug 2004 to present	\approx 3500 profiles/day	Two local times are sampled every orbit at any given latitude, daytime and nighttime, with local times typically near 1–2 p.m. and near 1–2 a.m. for non-polar latitudes
MIPAS	Jun 2002–Apr 2012 with gaps in 2004	\approx 1400 profiles/day (optimized resolution phase, nominal mode) \approx 1050 profiles/day (full resolution phase, nominal mode)	10.00 a.m. and p.m. local time

SI²N: ozone profile measurements

B. Hassler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1d. Summary of the latest data version, URLs where data are available, and information about references in the SI²N special issue for the five ground-based measurement systems and all described satellite measurement systems.

Name of instrument	Latest version of data	Where data are available	Reference in SI ² N special issue
Ground-based measurement systems			
Ozonsondes	Outcome of O3S-DQA activity as part of SI ² N	WOUDC, Toronto, Canada: http://www.woudc.org/data_e.html NDACC website (selected stations): http://www.ndacc.org SHADOZ website (tropical stations): http://croc.gsfc.nasa.gov/shadoz/	(not available at this point)
Dobson/Brewer	Described in: Petropavlovskikh et al. (2005)	WOUDC, Toronto, Canada: http://www.woudc.org/data_e.html	–
Lidar	Depends on each lidar system.	NDACC website: http://www.ndacc.org	Nair et al. (2013)
FTIR	Described in: Vigouroux et al. (2008)	NDACC website: http://www.ndacc.org	(not available at this point)
Microwave	Version 6. See Boyd et al. (2013)	NDACC website: http://www.ndacc.org Special Lauder and Mauna Loa hourly processed datasets available on request from PI.	Boyd et al. (2013)
Satellite-based measurement systems			
SAGE	Version 7	Atmospheric Science Data Center, NASA Langley: http://eosweb.larc.nasa.gov/project/sage2/sage2_v7_table	Darnadeo et al. (2013)
HALOE	Version 19 ozone is described briefly in Randall et al. (2003).	NASA Goddard Earth Sciences Data & Information Services Center (GES DISC): http://daac.gsfc.nasa.gov/ HALOE Webpage: http://haloe.gats-inc.com	–
ILAS	More details: Brühl et al. (1996) and SPAPC (1998). ILAS-I: Version 6.10 ILAS-II: Version 3.0	ILAS-I: http://db.cger.nies.go.jp/ilas/ ILAS-II: http://db.cger.nies.go.jp/ilas2/	–
POAM	Latest retrieval version is 4. Version 4 ozone differs very little from version 3. The version 3 algorithm and error analysis are described in Lumpe et al. (2002). Changes for the version 4 algorithm are described in a pdf document on the POAM ftp archive.	NASA Langley Atmospheric Sciences Data Center: http://eosweb.larc.nasa.gov/project/poam3/poam3_table	–
ACE-MAESTRO	Described in: McElroy et al. (2007)	Data also available through Naval Research Laboratory: and: http://www.nrl.navy.mil/rsd/7220/poam ACE-MAESTRO: From ACE team, University of Waterloo, Waterloo, Canada (for most recent version) Publicly available data sets from: http://www.ace.uwaterloo.ca/data.html	–
ACE-FTS	Described in: Boone et al. (2005)	ACE-FTS: From ACE team, University of Waterloo, Waterloo, Canada (for v3.5 – most recent versions) (v2.2 + updates – Publicly available data sets from http://www.ace.uwaterloo.ca/data.html)	(not available yet)
GOMOS	GOMOS IPF 6.01 released in 2012	ESA EARTHNET: https://earth.esa.int/web/guest/missions/ esa-operational-eo-missions/envisat/instruments/gomos	Kyrölä et al. (2013) Vertically gridded data set: Sofieva et al. (2013)
LIMS	Described in: Remsberg et al. (2007)	NASA Goddard Earth Sciences Data & Information Services Center (GES DISC): http://daac.gsfc.nasa.gov/	–
UARS MLS	Version 5 was the last major public release of UARS MLS O ₃ data (Livesey et al., 2003)	The official public distribution location for UARS MLS data is at the NASA GES-DISC Mirador website: http://mirador.gsfc.nasa.gov	(not available yet)

SI²N: ozone profile measurements

B. Hassler et al.

Table 1d. Continued.

Name of instrument	Latest version of data	Where data are available	Reference in SI ² N special issue
Aura MLS	v2.2	The official public distribution location for Aura MLS data is at the NASA GES-DISC Mirador website: http://mirador.gsfc.nasa.gov	(not available yet)
MIPAS	IMK: V5_O3_221 (von Clarmann et al., 2009; Laeng et al., 2013) ESA: ML2PP V6 MIPAS2D – Version 2.3 (see Dinelli et al., 2010, 2012)	IMK, Germany: http://www.imk-asf.kit.edu/english/308.php Oxford University, UK: http://www.atm.ox.ac.uk/group/mipas/L2OXF/ ESA: & MIPAS2D – ISAC-CNR, Italy: www.isac.cnr.it/~rss/mipas2d.htm http://odin.rss.chalmers.se http://www.iup.uni-bremen.de/sciamachy/	Eckert et al. (2013) Raspollini et al. (2013)
Odin-SMR SCIAMACHY	Version 2.1. Version 2.5 Described in: Rahpoe et al. (2013) and Mieruch et al. (2012)	http://odin.rss.chalmers.se	– Rahpoe et al. (2013)
Odin-OSIRIS SBUV	Described in: Degenstein et al. (2009) Version 8.6	http://odin-osiris.usask.ca/?q=node/280 Goddard DISC	(not available yet) Bhartia et al. (2013); De- Land et al. (2012); Kra- marova et al. (2013a, b)
	Described in: Bhartia et al. (2013) and in McPeters et al. (2013)	http://disc.sci.gsfc.nasa.gov/daac-bin/ DataHoldingsMEASURES.pl?PROGRAM_List= RichardMcPeters GOME-2: EUMETSAT archive: http://archive.eumetsat.int/ GOME-2: TEMIS: http://www.temis.nl/profiles/ All data: In the future via Ozone_cci web portal (ESA): http://www.esa-ozone-cci.org	–
GOME/GOME-2	–	OMI: Goddard DISC http://disc.sci.gsfc.nasa.gov/	–
OMI		Upon request. Will be operationally distributed by Eumetsat in 2014	–
IASI	Described in: Hurtmans et al. (2012)	Eumetsat data archive: www.eumetsat.int	–
	Described in: Dufour et al. (2012)	ISAS/JAXA DARTS http://darts.isas.jaxa.jp/iss/smiles/ NICT https://smiles-p6.nict.go.jp/products/research_latitude-longitude. jsf	Kasai et al. (2013)
SMILES	ISAS: Version 3.0 (118-12-0602) Described in: Mitsuda et al. (2011) and Imai et al. (2013) NICT: Version 2.1.5 Described in: Baron et al. (2011); Kasai et al. (2013) and Kreyling et al. (2013)		

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.




Fig. 1. World map with locations for the long-term ground-based measurement sites: ozone sounding sites (red dots), lidar sites (green stars), microwave sites (light blue triangles), Dobson/Brewer sites with Umkehr measurements (yellow squares), and FTIR sites (dark blue circles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

SI²N: ozone profile measurements

B. Hassler et al.




Fig. 2. Temporal coverage of the described ground- and satellite-based measurement systems.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.




Fig. 3. Vertical resolution shown as function of altitude for each of the described instruments systems: **(a)** ground-based instruments, **(b)** solar/stellar occultation instruments, **(c)** nadir measuring instruments, and **(d)** limb measuring instruments.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SI²N: ozone profile measurements

B. Hassler et al.




Fig. 4. Latitudinal coverage for the solar-occultation instruments (example years: 2003 and 2005). SAGE I is similar to SAGE II (blue triangles), POAM and ILAS instruments are similar to SAGE III (turquoise triangles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)