Supplement of Atmos. Meas. Tech. Discuss., 7, 5223–5249, 2014 http://www.atmos-meas-tech-discuss.net/amtd-7-5223-2014/doi:10.5194/amtd-7-5223-2014-supplement © Author(s) 2014. CC Attribution 3.0 License.





## Supplement of

## Collection efficiency of the Soot-Particle Aerosol Mass Spectrometer (SP-AMS) for internally mixed particulate black carbon

M. D. Willis et al.

Correspondence to: M. D. Willis (megan.willis@mail.utoronto.ca) and T. B. Onasch (onasch@aerodyne.com)

## Collection Efficiency of the Soot-Particle Aerosol Mass Spectrometer (SP-AMS) for Internally Mixed Particulate Black Carbon

## **Supplementary Information**

M.D. Willis, A.K.Y. Lee, T.B. Onasch, E.C. Fortner, L.R. Williams, A.T.Lambe, D.R. Worsnop, and J.P.D. Abbatt




Figure S1: Schematic of beam width probe orientation. Left: BWP orientation for particle beam width measurements, the BWP is situated perpendicular to the laser beam and is moved horizontally. Right: BWP orientation for effective laser beam width estimation, the BWP is moved vertically.




Figure S2:  $RIE_{rBC,app}$  (top) and  $RIE_{Org,app}$  (bottom) measured as a function of BES coating thickness, demonstrating an increase in sensitivity that reaches saturation at  $R_{Org/RB} \sim 3$ . Data are plotted against SP-AMS derived  $R_{Org/RB}$  because particle mass analyzer measurements were not available during this experiment.  $RIE_{Org,app}$  values were obtained from SP-AMS data using measured  $d_{va}$  values and assuming that coated particles have a core-shell structure. Data from ARI SP-AMS (SN 215-039).




Figure S3: Top: The ratio of SP-AMS fraction of Regal Black ( $f_{RB}$ ) to the particle mass analyzer  $f_{RB}$  as a function of  $R_{\rm Org/RB}$ . Data from ARI SP-AMS (SN 215-039 (blue) and 215-130 (red)) showing an approximately 40% underestimate in  $f_{RB}$  when CE effects are not accounted for. Bottom: The fraction of Regal Black in BES coated particles ( $f_{RB}$ ), obtained from particle mass measurements, plotted against the fraction of Regal Black obtained from SP-AMS mass loadings using uncorrected RIE<sub>rBC</sub> (0.2) and RIE<sub>Org</sub> (1.4).