

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

# Assessment of performance of the inter-arrival time algorithm to identify ice shattering artifacts in cloud particle probes measurements

# A. Korolev<sup>1</sup> and P. R. Field<sup>2</sup>

<sup>1</sup>Environment Canada, Toronto, Canada <sup>2</sup>Met Office, Exeter and University of Leeds, Leeds, UK

Received: 18 August 2014 – Accepted: 11 September 2014 – Published: 8 October 2014

Correspondence to: A. Korolev (alexei.korolev@ec.gc.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

10249

## Abstract

Shattering presents a serious obstacle to current airborne in-situ methods of characterizing the microphysical properties of ice clouds. Small shattered fragments result from the impact of natural ice crystals with the forward parts of aircraft-mounted mea-

- <sup>5</sup> surement probes. The presence of these shattered fragments may result in a significant overestimation of the measured concentration of small ice crystals, contaminating the measurement of the ice particle size distribution (PSD). One method of identifying shattered particles is to use an interarrival time algorithm. This method is based on the assumption that shattered fragments form spatial clusters that have short in-
- terarrival times between particles, relative to natural particles, when they pass through the sample volume of the probe. The interarrival time algorithm is a successful technique for the classification of shattering artifacts and natural particles. This study assesses the limitations and efficiency of the interarrival time algorithm. The analysis has been performed using simultaneous measurements of 2-D optical array probes with
- the standard and antishattering "K-tips" collected during the Airborne Icing Instrumentation Experiment (AIIE). It is shown that the efficiency of the algorithm depends on ice particle size, concentration and habit. Additional numerical simulations indicate that the effectiveness of the interarrival time algorithm to eliminate shattering artifacts can be significantly restricted in some cases. Improvements to the interarrival time algorithm are discussed
- 20 are discussed.

## 1 Introduction

Ice Particle Size Distributions (PSDs) are used in atmospheric models for computing both the radiative impact of ice clouds and the microphysical process rates that control ice cloud evolution. Therefore, an inaccurate representation of the ice PSD can have adverse impacts on the accuracy of climate and weather prediction models. Measure-

adverse impacts on the accuracy of climate and weather prediction models. Measurements of ice PSDs from aircraft-based observations form the basis of size distribution representations in models. Prior to entering the instrument sample volume, an ice particle may impact the probe's upstream tips or inlet and shatter into many small fragments. These shattering products can then significantly contaminate measurements of airborne particle probes.

- Over three decades the abundance of small ice particles measured by particle probes in the tropospheric clouds remained an intriguing problem. Recent studies using simultaneous measurements of standard and modified probes (Korolev et al., 2011, 2013b) have unambiguously demonstrated that in many cases measurements made with standard cloud probes are adversely affected by shattering artifacts.
- There are two main approaches currently used within the cloud physics community to mitigate the effect of shattering. One approach is based on modifying the inlet configuration of the cloud probes to minimize the area that deflects shattered particles towards the sample volume (Korolev et al., 2013a). This method can lead to a significant reduction in the effect of shattering but is not able to completely eradicate the problem (Korolev et al., 2013b; Lawson, 2011).
- The second approach is a postprocessing methodology based on the fact that shattering products are spatially clustered. Because these closely spaced fragments will pass thought the sample volume with interarrival times much shorter than naturally occurring particles. Cooper (1977) suggested that these artifacts could be filtered out
- <sup>20</sup> by identifying the characteristically short inter-arrival times of particles associated with these shattering products. Field et al. (2002) used a Fast FSSP to measure particle spacing in ice clouds. The

Field et al. (2003) used a Fast FSSP to measure particle spacing in ice clouds. The inter-arrival time distribution in ice clouds was found to have a bimodal shape with modes at  $10^{-2}$  and  $10^{-4}$  s corresponding to approximately 1 m and 1 cm spatial sepa-

<sup>25</sup> rations. The particles from the long and short inter-arrival time modes corresponded to estimated concentrations of  $0.1-1 \text{ cm}^{-3}$  and  $\sim 100 \text{ cm}^{-3}$  respectively. No conclusions were drawn as to whether the latter localized clusters of high particle concentration were natural or artifacts. Assuming they were artifacts, their inter-arrival time algorithm

10251

suggested average and maximum concentration overestimates of a factor of 2 and 5

respectively. Field et al. (2006) applied an inter-arrival time algorithm (ITA) to filter out shattering artifacts, choosing a threshold inter-arrival time in the range  $10^{-4}$  to  $10^{-5}$  s, depend-

- ing on the instrument and the aircraft type used for the data collection, to reject the short inter-arrival mode. It was found that the OAP-2DC and CIP concentrations were reduced by up to a factor of 4 when the mass-weighted mean size exceeded 3 mm. The ice water content (IWC) estimate was reduced by up to 20–30 %, most notable in cases of narrow size distributions. It was also concluded that the effect of shattering on
  the measured size distribution reaches up to sizes of several hundred microns.
- the measured size distribution reaches up to sizes of several hundred microns. The ITA is now routinely used in 2-D probe data processing to eliminate shattering artifacts (e.g. Baker et al., 2009; Jackson et al., 2014). Moreover, it is recognised that the best practice for the operation of cloud probes in the presence of ice is provided by a combination of modified tips and the application of the interarrival time filtering
   together (e.g. Korolev et al., 2011; Baumgardner et al., 2013).
- While it has been demonstrated that the ITA reduces the effects of shattering, its accuracy and efficiency remains poorly quantified. For instance, can the ITA alone identify all shattering artefacts? Is the efficiency of the ITA dependent upon the probe specifications such as pixel resolution, response time, sample area and inlet configuration?
- Motivation for improving our understanding of the limitations and efficiency of the ITA is threefold. Firstly, there is a need to determine if the ITA can be used to successfully reanalyze the historical data collected over the past thirty years. Secondly, an improved quantitative understanding of the limitations of the ITA will provide statements of the accuracy of the measurements of the particle number concentration, ice water
- <sup>5</sup> content, extinction coefficient and other PSD derived parameters. Thirdly, knowledge of the efficiency of the ITA will aid in the design process of future cloud probes.

The objective of this paper is to evaluate the efficiency and limitation of ITA. A detailed description and main assumptions underlying the algorithm are presented in Sect. 2. Section 3 considers general limitations of the algorithm. Analysis of the results of the

**Discussion** Paper

Discussion Paper | Discussion Paper | Discussion Paper |

OAP-2DC data processing using the ITA are presented in Sect. 4. In Sect. 5 statistical simulations of ice particle shattering are used to explore the limitations of the ITA. Finally, Sect. 6 provides a summary.

## 2 Description of the inter-arrival time algorithm

#### **5 2.1 Basic assumptions**

In the following discussion the term "shattering event" will be applied to the group of shattered fragments, that (a) formed as a result of impact of a single particle with the upstream tips (or inlet) of a probe, and (b) at least one particle from the group of the shattered fragments was registered by the probe. It should be noted that if the particle rebound to the sample area without shattering, it still falls in the definition of

particle rebound to the sample area without shattering, it still falls in the definition of the "shattering event".

The successful application of the ITA is predicated on two basic assumptions: (1) the maximum inter-arrival time of the shattered fragments is shorter than the minimum inter-arrival time between intact particles, (2) shattered particles are always passing through the sample volume as a group of no less than two particles.

- The first assumption is a necessary condition for the complete separation of the interarrival time distributions  $\varphi_s(\Delta t)$  and  $\varphi_i(\Delta t)$  without overlap. Here  $\varphi_s(\Delta t)$  and  $\varphi_i(\Delta t)$ are the distribution of inter-arrival times associated with shattering events and intact particles, respectively. The absence of an overlap between  $\varphi_s(\Delta t)$  and  $\varphi_i(\Delta t)$  allows
- for the existence of a cut-off time  $\tau^*$  such that all *shattered and only shattered* particles satisfy the condition  $\Delta t < \tau^*$ , whereas all *intact particles and only intact particles* are associated with  $\Delta t > \tau^*$  (Fig. 1a). Given this assumption it is trivial to identify the shattered artifacts and intact particle based simply on a comparison of measured interarrival time  $\Delta t$  between two successive particles and the cut-of time  $\tau^*$ .

#### 10253

The second assumption forms the second necessary condition to identify shattered artifacts. A minimum of two closely spaced particles is necessary to allow artifacts to be identified.

The conceptual diagram in Fig. 2a demonstrates these two basic assumptions about particle spacing that is required for the successful segregation of shattered artifacts and intact particles using an ITA. In reality the challenges of image sampling and the statistical nature of particle spacings impose limitations on the performance of the ITA in the segregation of shattered artifacts and intact particles. As it will be shown below the first assumption cannot be satisfied due to statistical limitations, whereas the second condition is necessary but not sufficient.

#### 2.2 Inter-arrival time algorithm

Here we describe the sequence of operations composing the basic ITA. This algorithm in its basic form will be used in the present study.

1. The inter-arrival algorithm starts from the calculation of the distribution of inter-

- arrival times  $\varphi(\Delta t)$  as in Fig. 1. The calculation of  $\varphi(\Delta t)$  are performed for each averaging time intervals. Similar to Field et al. (2003, 2006) the time bins in  $\varphi(\Delta t)$ were logarithmically spaced. The width of the time bins was optimized to tradeoff accuracy of the estimation of  $\tau^*$  and the statistical significance related to the number of counts in each time bin.
- 20 2. The cut-off-time  $\tau^*$  was calculated for each averaging time interval as a minimum between two maxima associated with long and short time modes (Fig. 1). In cases when only one mode was present,  $\tau^*$  was forced to be equal to minimum interarrival time found in this averaging interval. It should be noted that the function  $\varphi(\Delta t)$  is a non-normalized distribution of counts in each time bin. Normalization using the bin width leads to a disappearance of the minimum between the short and long time modes in  $\varphi(\Delta t)$ , which hinders calculation of  $\tau^*$ .

- 3. Pairs of particles that satisfied the condition  $\Delta t < \tau^*$  were identified and marked as artifacts. It is important to note, the ITA cannot identify a single-particle shattered artifact (singleton) and that the minimum number of particles identified as artifacts is two.
- <sup>5</sup> The value of  $\tau^*$  should be calculated for each averaging time interval. As it will be shown below,  $\tau^*$  has a wide dynamic range and it depends on many microphysical, environmental and instrumental parameters. The assumption, that  $\tau^*$  remains constant, may result in large errors in identifying shattering artifacts.
- The values of  $\tau^*$  and  $\Delta t$  depend of the airplane speed. In the described algorithm it <sup>10</sup> is assumed that the aircraft speed remained approximately constant at each averaging time interval. This assumption works well for few seconds averaging intervals. However, the aircraft speed depending on the altitude may change up to two times during the flight operation. This gives another reason to recalculate  $\tau^*$  at each averaging time interval.

#### 3 Limitations of the inter-arrival time algorithm 15

This section presents a list of sampling effects that demonstrate how the assumptions underlying the inter-arrival time algorithm can be contravened. These cases impose limitations on the ability of the ITA to segregate intact particles and shattering artifacts. In the following we assume that particles are distributed randomly in space and that

<sup>20</sup> their spacing and hence interarrival time is well represented by a Poisson distribution. For the Poisson process the density function for counting one intact particle during time  $\Delta t$  is described by

$$\frac{\mathrm{d}P(\Delta t)}{\mathrm{d}t} = \frac{e^{-\Delta t/\tau}}{\tau}$$

(1)

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

(2)

where  $\tau = 1/nSu$  is the average inter-arrival time between intact particles passing through the probe's sample area S; *n* is the particle concentration; *u* is the sampling 10255

speed. Any shattered particles are deflected into the sample area of the probe after the impact with the inlet. Therefore, the shattered particles have external origin, are intermittent and their distribution can be considered as independent, but still Poissonian, with respect to the intact particles.

#### **5** 3.1 Naturally occurring particles with interarrival times shorter than the cut-off-time interval

Two closely spaced intact particles will be identified as a shattering artifact if the interarrival time  $\Delta t < \tau^*$  (Fig. 2d). Such cases break the first assumption in Sect. 2.1. The probability for coincidence of 3 or more particles falls very rapidly and has been ignored. The probability of two particles to arriving within  $\tau^*$  can be found from the Pois-10 son statistics as

$$P_2(\Delta t < \tau^*) = \frac{1}{2} \left(\frac{\tau^*}{\tau}\right)^2 e^{-\frac{\tau^*}{\tau}}$$

20

As follows from Eq. (1) the probability of such event increases with increasing particle concentration *n* (decreasing  $\tau$ ) and increases in the cut-off time  $\tau^*$ .

#### 3.2 Coincidence of a shattering event particle and an intact particle

The probability of coincidence of the arrival time of an intact particle during the shattered event can be estimated as

$$P(\Delta t < \Delta t_{\rm sh}) = \int_{0}^{\Delta t_{\rm sh}} \frac{e^{-\Delta t/\tau}}{\tau} d(\Delta t) = 1 - e^{-\Delta t_{\rm sh}/\tau}$$
(3)

Here  $\Delta t_{sh} = \sum \Delta t_{sh}(i) = L_{sh}/u$  is the duration of the shattering event registered by the probe;  $\Delta t_{sh}(i)$  is the inter-arrival time between two subsequent shattered fragments 10256

registered by the probe within the shattered event;  $L_{sh}$  is the spatial length of the shattered event along the flight direction (i.e. distance between the first and last fragments in the shattered event). Equation (2) indicates that even when  $\Delta t_{sh} \ll \tau$ , the probability of the arrival of the intact particle in the sample volume during the shattering event remains non-zero. Basically, it means that in principle it is impossible to senarate all

<sup>5</sup> remains non-zero. Basically, it means that in principle it is impossible to separate all shattering artifacts and intact particles, and the functions  $\varphi_s(\Delta t)$  and  $\varphi_i(\Delta t)$  overlap always (Fig. 1b). The relative fraction of the overlapping area of  $\varphi_s(\Delta t)$  and  $\varphi_i(\Delta t)$  characterizes frequency of misidentifying intact particles and shattering artifacts.

#### 3.3 Singletons: single particle shattering artifacts

- A significant limitation of the ITA is related to situations when only one particle from the group of the shattered fragments is registered by the probe (Fig. 2c). Such situation may occur, when most of the shattered fragments travel outside of the sample volume, but a single fragment passes through the sample volume. It may also happen when most of the shattered fragments are smaller than the probe's detecting thresh-
- <sup>15</sup> old, and only one particle exceeds the threshold and is registered by the probe. Ice particles may also rebound from the inlet without fragmentation, thus forming a single particle shattering event. Rebounding without shattering was demonstrated in Korolev et al. (2013a, Fig. 13). With respect to the ITA the single particle artifacts described above can have long inter-arrival times and are therefore indistinguishable from the
- natural population of intact particles. Due to the random nature of particle impact with the probe's arms and the direction of the trajectories of the rebound shattered fragments the probability of the single particle shattering events always remains non-zero. This imposes a significant and difficult to quantify limitation on the performance of the ITA.

10257

#### 3.4 Partially viewed ice branches

Many ice particles develop **branched** extended from few hundred micrometers up to 2–3 mm away from its center (i.e. bullet rosettes, dendrites, aggregated ice particles). Partially viewed branches of such particles could be confused with separate particles possessing short inter-arrival time (Fig. 2e), and be identified as artifacts. Rejecting

images that are in contact with the edge of the array is one way to mitigate against this problem.

#### 3.5 Diffraction fringes

Most particle imaging probes use coherent sources of light that result in the formation of diffraction fringes around the image of a particle. The binary representation of these

- of diffraction fringes around the image of a particle. The binary representation of these fringes may manifest themselves as sparse disconnected pixel images that surround the main particle image. Such optical and imaging instrumentation effects may be confused with shattered fragments and result in identifying both diffraction fringes and the intact particle producing these fringes as artifacts.
- For spherical particles such fringes become most pronounced when the dimensionless distance of the particles from the focal plane is close to  $Z_d \approx 1.9$  (Korolev, 2007, Fig. 8) where  $Z_d = \frac{4\lambda Z}{D^2}$ ;  $\lambda$  is the wavelength; Z is the distance from the object plane; D is the particle diameter. Non-circular images produce diffraction fringes over a wider range of  $Z_d$ . The probability of imaging the diffraction fringes increases with
- the increasing pixel resolution. Thus, for the probes with coarse pixel resolution 100–200 μm (e.g. HVPS, PIP, OAP-2DP), the diffraction artifacts are quite rare, whereas for the probes with 10–15 μm pixel resolution (e.g. 2D-S, CIP) the effect of the diffraction fringes may have a significant effect on misidentification of intact particles as shattering artifacts. A few examples of diffraction fringes around the CIP out-of-focus images are
- shown in Fig. 3. These images were identified by the inter-arrival time algorithms as shattering artifacts and rejected. The 2-D data processing software can be tuned to return large images to the pool of accepted images. However, setting the threshold for the

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

**Discussion** Paper

Discussion Paper | Discussion Paper

Discussion Paper

sizes of the accepted images is ambiguous, and it may result in accepting shattering artifacts and rejecting intact particles.

#### 3.6 Out-of-focus fragmented images

Out-of-focus images of particles traversing the sample area near the edges of the depth-of-field, when  $Z_d > 6$  may appear as disconnected 1–3 pixel images. Examples of such images can be found in Korolev (2007, Fig. 6). Out-of-focus images, such as images of transparent plate like ice particles, quite often appear as fragmented and will be identified by the ITA as artifacts.

#### 4 Results of measurements of inter-particle distances

- <sup>10</sup> Because shattering generates particles by a very different physical process to those that occur naturally, the mode of the Poisson distribution that describes the distribution of these particles can be very different to that associated with the natural intact particles. This difference in the distribution of the particles manifests itself through differing interarrival time populations. Therefore, the distribution  $\varphi(\Delta t)$  can be used as one of
- <sup>15</sup> the metrics for identifying shattering. The purpose of this section is to demonstrate the variety of  $\varphi(\Delta t)$  distributions and show their link to the particle size distributions. This consideration is expected to help further understanding of limitations of the ITA. In order to reduce the effect of the air speed *u* on  $\Delta t$ , the inter-particle distance

 $\Delta x = \Delta t/u$  will be used instead of  $\Delta t$ . Accordingly, the distribution  $\varphi(\Delta x)$  and the cut-

off-distance  $\chi^* = \tau^*/u$  will be used instead of  $\varphi(\Delta t)$  and  $\tau^*$ , respectively. We will also keep using the conventional term "inter-arrival time algorithm", although the term "inter-particle distance algorithm" would be more accurate.

10259

#### 4.1 Description of the data set

The data used here were collected during the Airborne lcing Instrumentation Evaluation Experiment (AIIE) flight campaign (Korolev et al., 2011, 2013b). The analysis of the inter-particle distance is focused on the measurements of two OAP-2DCs installed cide hereids in the ADP of the analysis of the second sec

- <sup>5</sup> side-by-side in the NRC Convair-580 aircraft. Both instruments have the same pixel resolution 25 μm, optics and electronics. However, one of the probes had standard configuration, whereas the second one had modified arms with the K-tips installed (Korolev et al., 2013a). While K-tips still shatter ice particles, it has been demonstrated that they significantly mitigate the effect of shattering on ice particle measurements.
- <sup>10</sup> Comparing measurements made with standard and modified OAP-2DCs before and after applying the ITA provides an opportunity to assess the efficiency of the algorithm to successfully identify and filter out shattering artifacts. The 2-D data were averaged over 5 s time intervals. For most clouds sampled during the AIIE project such averaging provided statistically significant particle numbers to estimate the function  $\varphi(\Delta x)$  and
- <sup>15</sup> cut-off-distance  $\chi^*$ .

## 4.2 Examples of the inter-particle distance distribution

The following three examples are based on the data collected during three different flights (1, 8 April 2009) and demonstrate how the presence of large particles and their concentration affect the inter-particle distance distribution  $\varphi(\Delta x)$ . The first example demonstrates a moderate level of shattering and a high concentration of ice. The sec-

ond example demonstrates a moderate level of shattening and a high concentration of ice. The second example demonstrates more intense shattering and low concentration of ice. The last example demonstrates a case where shattering has a negligible effect.

#### 4.2.1 Overlapping modes

Figure 4a and c shows the time series of inter-particle distances measured by the standard and modified OAP-2DC in an ice cloud. Each inter-particle distance in

Fig. 4a and c is represented by a dot. The red lines indicate the cut-off-distances. As seen from these two diagrams the density of points below the red line is greater for the standard probe (Fig. 4a) than that for the modified one (Fig. 4c). The concentration measured by the modified 2DC and corrected with the help of the ITA, varied from

- <sup>5</sup> 20 to 80 L<sup>-1</sup>. Whereas, the uncorrected concentration measured by the standard 2DC varied from 300 to 1600 L<sup>-1</sup>. The ice particle images measured by 2DC and 2DP are presented in Fig. 4e and f. Analysis of these images shows that the ensemble of ice particles was composed of two distinct habits: large spatial dendrites with sizes up to few millimeters and transparent plates with a characteristic size of a few hundred mineration.
- crometers. The maximum particle size  $D_{max}$  calculated for each averaging time interval remained approximately constant and did not exceed 5 mm.

The distributions of the inter-particle distances  $\varphi(\Delta x)$  calculated from the standard and modified 2DC probes data are shown in Fig. 4c and d. The inter-particle distance distribution,  $\varphi(\Delta x)$ , for the standard 2DC displays a significant overlap between the

<sup>15</sup> long and short inter-arrival modes (Fig. 4b). This may result in rejecting intact particles along with shattered artifacts when  $\Delta x < \chi^*$ , and vice versa accepting shattered fragments with intact particles for  $\Delta x > \chi^*$ .

It is possible to attempt to correct for the removal of intact particles by using Poisson statistics to estimate the fraction of intact particles rejected and then scale the remaining intact size distribution (e.g. Field et al., 2006).

The inter-particle distance distribution  $\varphi(\Delta x)$  for the modified probe has a relatively small overlap between the long and short distance modes, which suggests a better separation of shattered and intact particles. The number of particles associated with the short distance mode for the modified probe (Fig. 4d) is also reduced when com-

<sup>25</sup> pared to Fig. 4b. However, despite the larger separation between the short and long distance modes, the ITA still identifies large particles, which appear intact, as shattered fragments (e.g. Fig. 5b).

Figure 5 shows results from applying the ITA to the standard and modified OAP-2DC images sampled at the same time during one flight from the AIIE project. Images

10261

identified as shattered artifacts appear on a green background, whereas the accepted images have a white background. The number of rejected images for the standard probe (Fig. 5a) appears to be higher than that for the modified probe (Fig. 5b). Visual examination of the measurements made with the standard probe (Fig. 5a) reveals

- a number of images that we interpret as being intact (red arrows) but are rejected by the algorithm. Similar particle images can also be found in data for the modified probe (Fig. 5b), but there are far fewer images that look intact but are rejected. Some of the accepted images obtained by the standard probe we interpret as shattered artifacts (Fig. 5a, blue arrows). Shattered artifacts usually appear and elongated along the flight
- direction images due to the slower speed that they enter the sample volume. In many cases the images of the shattering artifacts also have a hole in the center, that is the result of the greater likelyhood of shatter products entering the sample volume closer to the arms and, consequently, far from the center of the depth-of-field. Further support for the assumption that the accepted images indicated by the blue arrows are shattered
  artifacts is provided by the absence of similar images recorded by the modified 2DC
  - (Fig. 5b).

The apparently erroneous acceptance of the shattered artifacts and rejection of intact images in Fig. 5a is consistent with the large overlap between the short and long distance modes indicated in Fig. 4c. This example demonstrates that the ability of the

ITA to segregate shattered artifacts and intact particles is reduced when the short and long distance modes become less separated. One of the important goals of filtering out shattering artifacts is to obtain an accurate

estimation of the PSD. Figure 6 shows the distributions of particle counts, concentration and mass calculated for all images before corrections, after corrections. Equivalent dis-

tributions are shown for the rejected images. During the image processing the distribution of counts is used as a start point for the following calculations of other distributions and bulk microphysical parameters. As seen from Fig. 6a and d for both standard and modified OAP-2DCs the number of small particles rejected by the inter-arrival algorithm is nearly two orders of magnitude greater than for the large particles. This is consistent Discussion Paper | Discussion Paper | Discussion Paper

with the concept that shattered particles are mainly composed of small fragments. Experimental studies indicate that the fragments of ice particles shattered at the aircraft speed have characteristic size from tens to hundreds of micrometers (Vidaurre and Hallett, 2009). Therefore, it is hypothesised that rejected particles larger approximately 1 mm (Fig. 6) are likely related to the cases depicted in Fig. 2b, d–f.

It is interesting to note that for the modified probe the ITA corrected and uncorrected distributions agree reasonable well for particles larger approximately 500–600  $\mu$ m (in Fig. 6d–f). However, for the standard probe the separations between ITA corrected and uncorrected distributions remain approximately constant for *D* > 600  $\mu$ m (Fig. 6a–c).

- <sup>10</sup> Despite the rejection of a large fraction of small particles (Fig. 6b), the concentration of small particles ( $D < 200 \,\mu\text{m}$ ) is only reduced after correction by a factor of 2 to 3 (Fig. 6e). At the same time the concentration of particles in the small size bins ( $D < 200 \,\mu\text{m}$ ) for the standard probe after ITA correction still remains higher than for the modified probe. This is consistent with the above conclusion that the ITA is
- <sup>15</sup> unable to filter out all shattering artifacts. The corrected concentration of small particles ( $D < 200 \,\mu$ m) measured by the modified probe is still high. It is difficult to conclude whether these particles are real or associated with shattering artifacts, for instance due to singletons, or other mis-sizing and concentration errors.

Comparisons of corrected mass distributions for standard and modified probes in Fig. 6c and f show a large difference between them for  $D < 500 \,\mu\text{m}$ . However, since large particles are the major contributors into the total mass, the discrepancy at the small size end of the PSD does not produce any significant effect on the bulk IWC. Integration of the mass distributions for standard and modified probes shows that IWC corrected standard and modified OAP-2DCs for this particular case agree to within

approximately 20%, and the mean IWC values averaged over entire time interval agree within 4%.

#### 10263

#### 4.2.2 Large particles

Figure 7 shows the changes of inter-particle distances and distributions  $\varphi(\Delta x)$  measured during a descent through precipitating aggregates of dendrites with  $D_{\text{max}}$  ranging from 0.5 to 1.5 cm. The ITA corrected concentration measured by the modified OAP-2DC varied from 0.5 to 5 L<sup>-1</sup>, whereas the uncorrected concentration measured by the

standard probe varied from 100 to 300 L<sup>-1</sup>. The appearance of the particle images measured by OAP-2DC and OAP-2DP are shown in Fig. 8e and f.

The distinctive feature of the inter-particle spatial distribution  $\varphi(\Delta x)$  calculated for the standard OAP-2DC is that it has an exceptionally high number of counts associ-

- ated with the short distance mode. This indicates that the standard probe observed mostly shattering artifacts. In contrast, for the modified probe the distribution  $\varphi(\Delta x)$ the number of counts in the short distance mode is smaller than for the long distance mode.
- The results of segregation of intact particles and shattered artifacts for standard and
  modified OAP-2DC performed by the ITA are shown in Fig. 8. The measurements made
  with the standard probe are dominated by artifacts with very few accepted images
  (Fig. 8a). The particle imagery obtained with the modified probe is largely devoid of
  the small images typically associated with shattered fragments (Fig. 8b). An absence
  of small particles in subsaturated precipitating regions is consistent with the commonly
- accepted concept of ice formation. It is worth noting that a few small accepted images (indicated by the blue arrows) still appear in the standard 2DC imagery in Fig. 8a. It is possible that these small images are related to single fragments as in Fig. 2c and were misidentified by the ITA as intact particles.

Despite the fact that most of the images in Fig. 8b appear to be intact dendrites, many of them were rejected. Visual inspection of the rejected images in Fig. 8b indicates that nearly all of them are partially viewed images. It is likely that the partially viewed branches of dendrites (e.g. Fig. 2e) has led to the ITA confusing the partially observed closely spaced arms of the dendrite with the shattered artifacts (Sect. 3.4).

The results shown in Fig. 8 demonstrate that even for the cases where there is a good separation of the short and long distance modes (Fig. 7c and d) the ITA still misidentifies shattering artifacts and intact particles.

The distributions of particle counts, concentration and mass obtained with the standard and modified probes are shown in Fig. 9. Figure 9a and d demonstrates that the

- number of counts of small particles is much greater for the standard probe than for the modified probe. This is consistent with the 2-D images shown in Fig. 8. The ITA identified most of the small image counts as artifacts (Fig. 9a). However the number of small images accepted for the standard probe is still greater than the modified probe (Fig. 9d). The same conclusion applies to the conceptorizer and mass distributions
- (Fig. 9d). The same conclusion applies to the concentration and mass distributions shown in Fig. 9b, c, and e, f.

We note that the shape of rejected distributions for the modified probe is very similar to the shape of the uncorrected distributions (Fig. 9d–f). Furthermore, the distributions of the accepted and uncorrected images approach each other only for particle sizes

 $_{15}$   $D > 1500 \,\mu\text{m}$  (Fig. 9a–c), in contrast to the case shown in Fig. 6. For both probes the number of rejected large particles remains relatively large, and it is related to misidentifying large particles as shattered artifacts.

#### 4.2.3 Small ice particles

The next example was obtained during a sampling of cirrus clouds at a temperature of  $-35^{\circ}$ C and altitude of 7500 m. The maximum size of particles varied from 200 to 400 µm, and did not exceed 500 µm. The particle concentration measured by the standard and modified OAP-2DCs agreed well and varied from 20 to  $180 L^{-1}$ .

Figure 10 shows the inter-particle spacings (Fig. 10a and b) and their distributions  $\varphi(\Delta x)$  for the standard and modified probes. As can be seen in Fig. 10c and d the

<sup>25</sup> inter-particle spacing distribution  $\varphi(\Delta x)$  is monomodal for both probes. Since the ITA used here is most efficient only for the cases with bimodal distributions, the fraction of particles rejected when  $\varphi(\Delta x)$  is monomodal is expected to be small. This is demonstrated by the small number of points that lie below the cut-off-distance line (red) in

10265

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Fig. 9a and b (for which  $\Delta x < \chi^*$ ), and that Fig. 11 shows very few images identified by the ITA as artifacts.

The distributions of particle counts, concentration and mass are shown in Fig. 12. All three distributions obtained with the standard and modified probes agree well with each other. The fraction of rejected artifacts is small and for practical purposes their

effect on the size and mass distributions has negligible effect.

These results show that for this specific case there is not much difference between the measurement obtained by the standard and modified probes. This suggests that for the cases of where the ice PSD is narrow ( $D_{max} < 400 \,\mu$ m) the effect of shattering on the

standard OAP-2DC measurements is quite small and it can be neglected. This finding also implies that there exists a threshold size below which shattering of ice crystals does not produce any significant effect on measurements. This will likely vary according to instrumental and microphysical characteristics. Of course, even though the probes agree for this case problems relating to mis-sizing and concentration errors for particles smaller than ~ 100 microns still exists (Korolev et al., 1998; Strapp et al., 2001)

#### 5 smaller than ~ 100 microns suit exists (Korolev et al., 1996, Strapp et al., 2001)

# 4.3 Statistical characteristics of inter-particle distances in shattering events

Figures 4a, b, 7a, b and 10a, b contain a plethora of information about the statistical characteristics of shattering events and their effect on ice particle measurements. This data can also be used in the development of future algorithms for the data pro-

cessing and numerical simulations of the shattering effects. In this section some of the statistical characteristics of shattering process will be investigated and compared with microphysical metrics.

Figure 13 shows distributions of the number of particles  $N_s$  within each shattering event (a1–a3), distributions of the length of spatial clusters along the flight direction  $L_s$  (b1–b3), distributions of distances between shattering events  $L_i$  (c1–c3), and the number of particles between shattering events  $N_i$  (d1–d3). These distributions were calculated for the cloud regions shown in Figs. 4, 7 and 10.

The distribution  $f(N_s)$  (Fig. 13a1–a3) for both standard and modified probes is well represented by an exponential function (Field et al., 2003). Analysis of the other cases indicates that the slope of the distribution depends on the presence of large particles in the size distribution and it is correlated with  $D_{max}$  such that increasing  $D_{max}$  leads to a shallower slope for  $f(N_s)$ : Fig. 13a2 displays the shallowest slope in  $f(N_s)$  for the

case with the greater  $D_{max}$ .

5

The distribution of  $f(N_s)$  monotonically increases with decreasing  $N_s$ , having a maximum for events with *two* shattered fragments. Extrapolating  $f(N_s)$  towards 1, i.e. shattering events with one particle, suggests that the number of singletons may be quite

high. Existence of single particle shattering events presents a principal limitation of the inter-arrival algorithm, since such particle cannot be unambiguously identified as artifacts just based on the analysis of  $\Delta x$ .

The maximum number of the shattered fragments in a shattering event measured by the standard OAP-2DC during the AIIE project reached 60. Whereas for the mod-

<sup>15</sup> ified probe the maximum number of fragments for this data set was found to be 14. Figure 13a1 and a2 also demonstrate that the modified OAP-2DC on average has a smaller number of fragment per shattering event, and therefore the antishattering tips can efficiently mitigate shattering.

The density function of the length of spatial clusters of shattered fragment  $f(L_s)$  can also be well approximated by an exponential function (Fig. 13b1 and b2). Clusters with a short length have the highest probability and they are associated with two-particles shattering events. The spatial length of the shattering clusters is presented by a cascade of scales ranging from zero to tens of centimeters. The maximum length of the shattered clusters measured by the standard and modified OAP-2DCs during the AIIE project reached 30 and 3 cm, respectively.

The density functions of distances between shattered events  $f(L_i)$  are shown in Fig. 13c1–c3. The characteristic scale of  $L_i$  is determined by the concentration of particles capable of generating shattering events. For example, the case in Fig. 13c3 has the highest total concentration of ice particles. However, only a few of them generate

10267

shattering events. As a results the  $f(L_i)$  in Fig. 13c3 has a statistically insignificant distribution. As in the previous cases  $f(L_i)$  can be well approximated by an exponential function and the characteristic value of  $L_i$  for modified OAP-2DC is larger than that for the standard probe (Fig. 13c1 and c2).

The behavior of the function  $f(N_i)$  in Fig. 13d1–d3 is very similar to that of  $f(L_i)$  (Fig. 13c1–c3). For the case depicted in Fig. 13d3 the low number of shattering events mean that most of the counts are outside of the scale.

Figure 14 shows the distributions of cut-off distances,  $f(\chi^*)$ , for the standard and modified OAP-2DCs averaged over all ice clouds sampled during the AIIE project. Be-

- <sup>10</sup> cause the value of  $\chi^*$  depends of the size distribution of ice particles and their habits, the shape of the distribution,  $f(\chi^*)$ , will be determined by a combination of cloud characteristics, aircraft and instrument properties. However, the distributions of  $f(\chi^*)$  in Fig. 14 allow a few conclusions to be made. Firstly, for OAP-2DC  $\chi^*$  can vary from tens of micrometers to approximately one meter. Secondly, the standard probe  $\chi^*_{st}$  has a mode at approximately 10 cm, whereas the modified probe  $\chi^*_{mdf}$  has a mode at approximately
  - 2 cm. And thirdly, for nearly all cases  $\chi_{st}^* > \chi_{mdf}^*$ .

#### 4.4 Effect of particle sizes

The effect of particle sizes on shattering is demonstrated in Fig. 15 that shows the maximum number of fragments per shattering event  $N_{smax}$  vs.  $D_{max}$  for standard and

<sup>20</sup> modified probes. The  $N_{smax}$  and  $D_{max}$  were calculated for each 5 s averaging interval for the entire AIIE project. Figure 15a shows that for  $D_{max} < 15 \text{ mm}$ ,  $N_{smax}$  correlates well with  $D_{max}$  and therefore the dependence  $N_{smax}(D_{max})$  can be parameterized with a linear function (Fig. 15a). However, for the modified probe the correlation coefficient between  $N_{smax}$  and  $D_{max}$  is low (0.57) and the  $N_{smax}(D_{max})$  saturates at  $N_{smax} \sim 15$ when  $D_{max} > 5 \text{ mm}$  (Fig. 15b).

Figure 16 shows the maximum length of shattering clusters  $L_{smax}$  vs.  $D_{max}$  for standard and modified probes. Similar to the case in Fig. 15  $L_{smax}$  and  $D_{max}$  were calculated for each 5 s averaging intervals. A relatively high correlation coefficient between  $L_{smax}$ 

and  $D_{\text{max}}$  (0.78) for the standard probe allows linear parameterization of  $L_{\text{smax}}(D_{\text{max}})$  (Fig. 16a). The correlation coefficient between  $L_{\text{smax}}$  and  $D_{\text{max}}$  for the modified probe is quite low (0.29), but  $L_{\text{smax}}$  never gets longer than 3 cm for all  $D_{\text{max}}$  encountered.

The nearly linear increase of  $N_{smax}$  and  $L_{smax}$  with increase of  $D_{max}$  for the standard <sup>5</sup> probe (Figs. 15a and 16a) indicates a strong dependence of particle size on particle shattering.

#### 5 Monte-Carlo simulation of inter-particle distance function

The statistical parameters obtained from the shattering analysis were used to constrain Monte-Carlo simulations of the effect of shattering on particle measurements. These simulations were used to understand the influence of various parameters on the shape of the inter-particle distance distribution,  $\varphi(\Delta x)$ .

For each simulation two fluxes of particles with the same concentration  $n_0$  are assumed to interact with the probe. The first flux represents intact particles and passes through the sample area  $S_0$  of the probe. The second flux represents the shattered

particles and pass through a "shattering" area for the probe,  $S_{sh}$ , rebound and subsequently pass through  $S_0$ . After passing though  $S_{sh}$  a particle breaks down into  $N_{sh}$  fragments.

The inter-particle distance between the intact particles was simulated assuming a Poisson distribution by combining a random number generator and an exponential

distribution with mean distance  $\overline{\chi}_0 = \frac{n_0}{S_0}$ . The inter-particle distance between the shattering events was also simulated using an exponential distribution with mean distance  $\overline{\chi}_{ev} = \frac{n_0}{S_{sh}}$ . Based on the results obtained in Sect. 4.3 the number of shattered fragments was simulated by a random number generator with exponential distribution with average  $\overline{N}_{sh}$ . The distance between the shattered fragments in each shattered event was

<sup>25</sup> also simulated by the exponential distribution with average  $\overline{\chi}_{sh} = \frac{L_s}{\overline{N}_{sh}}$ , where  $\overline{L}_s$  is the average distance of the cluster of shattered fragments. After sorting arrival times the 10269

act particles mixed

two flows were merged together to forming a time series of intact particles mixed with the shattering artifacts.

The modeling results presented below were performed for  $S_0 = 50 \text{ mm}^2$ ,  $S_{\text{sh}} = 5 \text{ mm}^2$ ,  $\overline{L}_{\text{s}} = 1 \text{ cm}$  and  $\overline{N}_{\text{sh}} = 5$ . The particle size distribution was assumed to be monodisperse.

Figure 17 shows the modeled distributions of the inter-particle distances for intact particles only  $\varphi_i(\Delta x)$  (blue), shattered particles only  $\varphi_s(\Delta x)$  (red) and all particles that passed through the sample volume  $\varphi(\Delta x)$  (black). The distribution  $\varphi_i(\Delta x)$  represents the Poisson process and has a single mode, whereas  $\varphi_s(\Delta x)$  has two modes. The long

- distance mode in  $\varphi_s(\Delta x)$  is determined by the characteristic distance between the shattered events  $\overline{\chi}_{ev}$ , whereas the short distance mode is associated with the characteristic distances between the shattered fragments  $\overline{\chi}_{sh}$ , which passed through the sample volume. The long distance mode in  $\varphi_s(\Delta x)$  also includes single particle shattering events. This is what we see for the long mode of the red line (rebounders/singletons) and
- <sup>15</sup> the blue line. Both of these are continuous processes, whereas the shorter shattering mode is intermittent conditional on a collision occurring in the shattering volume. It is important to mention that the inter-particle distribution  $\varphi(\Delta x)$  representing the flow of all particles cannot be presented as an algebraic superposition of  $\varphi_s(\Delta x)$  and  $\varphi_i(\Delta x)$ . In this regard the conceptual diagrams in Fig. 1 show an oversimplified inter-arrival time
- distributions, which help understanding, but does not reflect actual shape of  $\varphi_i(\Delta x)$ . Figure 18 show four distributions  $\varphi(\Delta x)$  calculated for the particle concentrations 1, 10, 100, and  $1000 \text{ L}^{-1}$ . Figure 18 shows that the long distance mode approaches the short distance mode, when the particle concentration  $n_0$  increases. However, the short distance mode is insensitive to the changes of the particle concentration, and it
- <sup>25</sup> remains at the same position, when  $n_0$  increases (Fig. 18a–c). The location of the cutoff-distance  $\chi^*$  has appear insensitive to the changes in  $n_0$  remaining approximately constant.

For increasing  $n_0$  the reduced separation of the long and short distance mode results in an increased overlap of the distributions associated with these modes. As described above, increasing the overlap of these distributions reduces the efficiency of the ITA to segregate intact particles and shattered artifacts. At high concentrations the short and long distance modes merge, resulting in the vanishing of the inter-modal minimum. At that stage the ITA becomes much less efficient or even disabled (e.g. Fig. 18d).

- In the above simulations the particle size distribution was assumed to be monodisperse. One of the consequences of this assumption is that all of the particles possessed the same shattering efficiency and  $\overline{N}_{sh}$  remains the same for all particles. In reality particle sizes in natural clouds are represented by broad distributions. As indicated above,  $\overline{N}_{sh}$  depends on particle sizes and that for small particles with  $D < 400 \,\mu\text{m}$
- $\overline{N}_{sh} \rightarrow 0$ . Therefore, the concentration  $n_0$  should be interpreted as a concentration of particles contribution in the effect of shattering, but not as a total concentration, which included small ice which do not affect shattering. For simplicity the effect of the particle size distribution was not included in the simulation. It should be noted that the threshold 400 µm refers OAP-2DC. Probes with different pixel resolution, response time and

<sup>15</sup> inlet configuration have different threshold sizes below which the effect of shattering becomes insignificant.

The analysis of the modeling results suggests that: (1) the number of counts in the long inter-particle distance mode is mainly controlled by the sample area  $S_0$  and particle concentration  $n_0$ , (2) the number of counts in the short distance mode depends on the area  $S_{sh}$ , which deflects particles towards the sample volume, the characteristic

- number of fragments per shattering event  $\overline{N}_{sh}$  and the particle concentration  $n_0$ . As demonstrated in Sect. 4,  $\overline{N}_{sh}$  is a function of particle size. It should also be noted that  $\overline{N}_{sh}$  also depends on  $S_{sh}$ ,  $S_0$  and spatial proximity of  $S_{sh}$  to  $S_0$ . For example as  $S_0$  decreases the probability of passing at least two shattered fragments ( $N_{sh} \ge 2$ ) through
- $S_0$  becomes diminishes too. If  $S_0$  becomes too small, most of the shattered artifacts will be associated with single particle shattered events ( $N_{sh} = 1$ ), that cannot be identified by the ITA. Therefore, it is anticipated that the inter-arrival time algorithm will be more efficient for OAP-2DS with  $S_0 \approx 50 \text{ mm}^2$ , than for the FSSP with the sample area

10271

 $S_0 \approx 3 \text{ mm}^2$ . The number of shattered fragments viewed by the probe  $N_{sh}$  also depends on spatial separation of  $S_{sh}$  to  $S_0$ . Thus, if  $S_{sh}$  is located too far from  $S_0$  then the shattered fragments may not have sufficient momentum to travel large distance across the airflow and reach  $S_0$ . Comparisons of the measurements of standard and modi-

fied probes demonstrated that  $\overline{N}_{sh}$  strongly depends upon  $S_{sh}$ . The reduced number of the shattered fragments  $N_{sh}$  for the modified probe is explained by the fact that the ant-shattering tips have a shattering area that such that  $S_{sh} \rightarrow 0$ .

#### 6 Conclusions

15

Based on the analysis of data obtained with standard and modified OAP 2DC probes in a variety of ice cloud conditions and Monte-Carlo simulations, the following conclusions have been obtained:

- 1. The interarrival time algorithm cannot segregate all shattering artifacts from the intact particles *in principle*. These limitations are imposed by the Poisson statistics of particle spatial distribution. It was demonstrated that the short inter-arrival times are not necessarily associated with shattering artifacts, and that shattered artifacts are not constrained to exhibit short inter-arrival times.
- 2. The inter-arrival time algorithm has a range of conditions when short and long interarrival modes are well separated and it can effectively segregate shattered artifacts and intact particles (e.g. low concentration).
- 3. The inter-arrival time algorithm has a number of limitations which under certain circumstances may significantly degrade its performance or even disable it. Such cases are relevant to the clouds with high particle concentration and large particles. For mixed phase clouds the inter-arrival time algorithm it may not be possible to use with probes that have a fine pixel resolution.

Discussion Paper

Discussion Paper

Discussion Paper | Discussion Paper

- It was found that in clouds with particles with D<sub>max</sub> < 400 μm the effect of shattering on measurements of the standard OAP-2DC can be neglected.
- 5. The inter-arrival distance and number of registered shattered artifacts is well represented by an exponential function. The slope of the distribution is a function of the characteristic particle size. The number of shattered fragments correlates with particle size. These findings open the door for a statistical simulation to study the effect of shattering on measurements.

This analysis of the efficiency of the inter-arrival time algorithm is based on OAP-2DC data. Nevertheless, most of the conclusions obtained in this study can be applied to other particle probes. In this regard it is worth mentioning that the effectiveness of the inter-arrival time algorithm depends on the pixel resolution, size of the sample area, the response time of the electronics and the inlet configuration.

Acknowledgements. The work of Alexei Korolev was supported by Environment Canada, Transport Canada and the US Federal Aviation Administration.

#### 15 **References**

Baker, B., Mo, Q., Lawson, R. P., O'Connor, D., and Korolev, A.: The effects of precipitation on cloud droplet measurement devices, J. Atmos. Ocean. Tech., 7, 1404–1409, 2009.

Baumgardner, D., Avallone, L., Bansemer, A., Borrmann, S., Brown, P., Bundke, U., Chuang, P. Y., Cziczo, D., Field, P., Gallagher, M., Gayet, J.-F., Heymsfield, A., Ko-

- rolev, A., Krämer, M., McFarquhar, G., Mertes, S., Möhler, O., Lance, S., Lawson, P., Petters, M. D., Pratt, K., Roberts, G., Rogers, D., Stetzer, O., Stith, J., Strapp, W., Twohy, C., and Wendisch, M.: In situ, airborne instrumentation: addressing and solving measurement problems in ice clouds, B. Am. Meteorol. Soc., 93, ES29–ES34, doi:10.1175/BAMS-D-11-00123.1, 2012.
- <sup>25</sup> Cooper, W. A.: Cloud physics investigation by the University of Wyoming in HIPLEX 1977, Rep. AS 119, Bureau of Reclamation, 321 pp., 1977.

- Field, P. R., Wood, R., Brown, P. R. A., Kaye, P. H., Hirst, E., Greenaway, R., and Smith, J. A.: Ice particle inter-arrival times measured with a Fast FSSP, J. Atmos. Ocean. Tech., 20, 249–261, 2003.
- Field, P. R., Heymsfield, A. J., and Bansemer, A.: Shattering and particle inter-arrival times measured by optical array probes in ice clouds, J. Atmos. Ocean. Tech., 23, 1357–1370, 2006.
  - Jackson, R. C., McFarquhar, G. M., Stith, J., Beals, M., Shaw, R. A., Jensen, J., Fugal, J., and Korolev, A.: An assessment of the impact of anti-shattering tips and artifact removal techniques on cloud ice size distributions measured by the 2-D Cloud Probe, J. Atmos. Ocean. Tech., accepted, 2014.
- Korolev, A. V. and Isaac, G. A.: Shattering during sampling by OAPs and HVPS. Part 1: Snow particles, J. Atmos. Ocean. Tech., 22, 528–542, 2005.
- Korolev, A. V., Strapp, J. W., and Isaac, G. A.: Evaluation of accuracy of PMS Optical Array Probes, J. Atmos. Ocean. Tech., 15, 708–720, 1998.
- Korolev, A., Emery, E. F., Strapp, J. W., Cober, S. G., Isaac, G. A., Wasey, M., and Marcotte, D.: Small ice particles in tropospheric clouds: fact or artifact? Airborne lcing Instrumentation Evaluation Experiment, B. Am. Meteorol. Soc., 92, 967–973, 2011.
  - Korolev, A., Emery, E., and Creelman, K.: Modification and tests of particle probe tips to mitigate effects of ice shattering, J. Atmos. Ocean. Tech., 30, 690–708, 2013a.
- Korolev, A. V., Emery, E. F., Strapp, J. W., Cober, S. G., and Isaac, G. A.: Quantification of the effects of shattering on airborne ice particle measurements, J. Atmos. Ocean. Tech., 30, 2527–2553, 2013b.
  - Lawson, R. P.: Effects of ice particles shattering on the 2D-S probe, Atmos. Meas. Tech., 4, 1361–1381, doi:10.5194/amt-4-1361-2011, 2011.
- Strapp, J. W., Albers, F., Reuter, A., Korolev, A. V., Maixner, U., Rashke, E., and Vukovic, Z.: Laboratory measurements of the response of a PMS OAP-2DC, J. Atmos. Ocean. Tech., 18, 1150–1170, 2001.
  - Vidaurre, G. and Hallett, J.: Particle impact and breakup in aircraft measurement, J. Atmos. Ocean. Tech., 26, 972–983, 2009.



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper







**Figure 2.** Conceptual diagram of (a) idealistic spatial sequence of intact particles and shattered artifacts passing through the sample volume. Case (c) when the inter-arrival time algorithm may confuse shattered artifact with intact particles, and (b, d, e, f) when intact particles may be confused with shattering artifacts.



**Figure 3.** Examples of diffraction fringes around out-of-focus images measured by CIP at  $15 \,\mu$ m pixel resolution. The diffraction fringes and the particle generating the fringes may be confused with shattered fragments and be rejected by the inter-arrival time algorithm.

10277





OAP-2DC standard 18:01:56.33 a	OAP-2DC modified 18:01 \$8.16 b
18.0159 13	

Figure 5. Examples of the results of the image rejection/acceptance processing with the interarrival time algorithm. The images of ice particles were simultaneously sampled by the standard (left) and modified (right) OAP-2DC during the time period shown in Fig. 4. Green backgrounds highlights the images identified by the inter-arrival time algorithm as artifacts. Images with a white background were accepted by the algorithm. As seen in (a) in some cases the standard OAP-2DC rejects large particles which appear to be intact (red arrows), at the same time it accepts particles, which have the features of shattered fragments (blue arrows).



Figure 6. Distributions of particle counts (a, d), concentration (b, e), mass (c, f) calculated for all images (grey) and accepted (blue) and rejected (red) identified by the inter-arrival time algorithm. The distributions were calculated for the standard (left) and modified (right) OAP-2DCs data collected in the cloud shown in Fig. 4.

10280

10279

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper



**Figure 7.** Comparison of inter-particle distances measured by standard (a) and modified (b) 2DC. Red lines in (a) and (b) indicate the cut-off distance  $\chi^*$ . The distribution of the interparticle distances for standard (c) and modified (d) probes. Examples of images obtained with an OAP-2DC at 25 µm pixel resolution (e) and an OAP-2DP at 200 µm pixel resolution (f). The measurements were conducted during on 8 April (1st flight) during descent through precipitating dendrites from approximately 1300 to 500 m in the Ottawa region. The temperature varied from -8 to -2°C.

#### 10281



**Figure 8.** Examples of the results of the image rejection/acceptance processing with the interarrival time algorithm. The images of ice particles were simultaneously sampled by the standard (left) and modified (right) OAP-2DC during the time period shown in Fig. 7. A green background highlights the images identified by the inter-arrival time algorithm as artifacts. Images with a white background were accepted by the algorithm. As seen in **(a)** in some cases the standard OAP-2DC rejects large particles which appear to be intact (red arrows), at the same time it accepts particles, which have the features of shattered fragments (blue arrows).



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

**Figure 9.** Distributions of particle counts **(a, d)**, concentration **(b, e)**, mass **(c, f)** calculated for all images (grey) and accepted (blue) and rejected (red) identified by the inter-arrival time algorithm. The distributions were calculated for the standard (left) and modified (right) OAP-2DCs data collected in the cloud shown in Fig. 7.



Figure 10. Same as in Fig. 7. The measurements were collected on 8 April (2nd flight) in cirrus clouds at altitude 7500 m and temperature -35 °C.

OAP-2DC standard	OAP-2DC modified	b
╕╄╶ <b>┙</b> ╡╽┍╻┍╻┍╻┍╷╷╖╼╻┍╴╎╓╺╞╎┍╺┝╓╻╒╘╻┝╻┍╺		
19:42:13.38	19:42:13.41	
19:42:14.13	19:42:14.13	
19:42:14.47	19:42:14.38	
19:42:15.13	19:42:15.13	
	ĸĸċŢĸIJŢĸIJĨĸĊŔŀĔſſŔIJĿĔĬŔ	

Figure 11. Same as in Fig. 4. The measurements of the images were obtained during the time period shown in Fig. 9.

Discussion Paper | Discussion Paper | Discussion Paper |



**Figure 12.** Distributions of particle counts **(a, d)**, concentration **(b, e)**, mass **(c, f)** calculated for all images (grey) and accepted (blue) and rejected (red) identified by the inter-arrival time algorithm. The distributions were calculated for the standard (left) and modified (right) OAP-2DCs data collected in the cloud shown in Fig. 10.



**Figure 13.** Distributions of number of particles per shattering event **(a1–a3)**, length of the clusters of shattered artifacts along the flight direction **(b1–b3)**, distance between the clusters with shattered fragments **(c1–c3)**, number of intact particles between shattered events **(d1–d3)** calculated for the cases shown in Figs. 3 (top row), 6 (middle row) and 9 (bottom row), respectively.





**Figure 14.** Distribution of cut-off distances  $\chi^*$  for standard (a) and modified (b) OAP-2DCs averaged over all ice clouds sampled during the AIIE project.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper



Figure 15. Scatter diagram of the maximum number of shattered fragments per event vs. maximum particles sizes for standard (a) and modified (b) OAP-2DCs for all ice clouds sampled during the AIIE project.

10289



Figure 16. Scatter diagram of the length of clusters of shattered fragments vs. maximum particles sizes for standard (a) and modified (b) OAP-2DCs for all ice clouds sampled during the AIIE project.







**Figure 17.** Results of Monte-Carlo simulations of distribution of inter-particle distance  $\varphi(\Delta x)$  for intact particles (blue), shattered particles (red) and all particles passed through the sample volume (black). The calculations were conducted for particle concentration  $n_0 = 100 L^{-1}$ .

10291





| Discussion Paper | Discussion Paper | Discussion Paper