

Interactive
Comment

Interactive comment on “Application of spectral analysis techniques to the intercomparison of aerosol data – Part 4: Combined maximum covariance analysis to bridge the gap between multi-sensor satellite retrievals and ground-based measurements” by J. Li et al.

A. M. Sayer (Referee)

andrew.sayer@nasa.gov

Received and published: 14 April 2014

This study builds upon previous work by the authors using principal component analysis (PCA) and related techniques to analyse spatiotemporal variations in aerosol optical depth (AOD) from a variety of satellite sensors (here, MODIS, MISR, SeaWiFS, and OMI). Ground-based AERONET data are also incorporated. The main thrust of these papers is that by decomposing datasets (singly or jointly) in terms of principle compo-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

nents/singular vectors the spatial variability, inter-sensor differences, and trends can be analysed in a new way. This work is topically relevant to AMT and, along with their prior studies, is a novel (to my knowledge) way to look at this satellite AOD data. The main development from their prior studies seems to be the presentation of the Combined Maximum Covariance Analysis (CMCA) technique, to incorporate the AERONET AOD into the analysis. After the CMCA technique is applied, the decomposed results are analysed, and some comments made about the representivity of monthly mean AOD fields from the various satellite datasets.

I have some specific comments (below), but following these (mostly minor) revisions and clarifications which the authors should address, I favour publication of this paper in AMT. Several of my comments encourage the authors to emphasise more strongly the different mechanisms which may lead to apparent disagreements between sensor monthly AOD maps (retrieval biases, and observability issues related to swath width/cloud screening). The overall clarity, quality of language, and quality of presentation are fine (aside from a few figures which I feel should be redrawn). The authors are welcome to contact me in case of any queries about my comments.

Andrew Sayer

andrew.sayer@nasa.gov

Page 3505, line 21: I suggest rephrasing to avoid starting a sentence with ‘And’.

Page 3508, line 14: rather than ‘Ångström Relationship’ I think it would be better to say Ångström power law or something similar.

Page 3508, line 16: I don’t know that I would say MODIS was ‘designed’ to retrieve aerosol properties, given that’s but one of the applications the sensor has been put to, and it is certainly far from an ideal instrument to retrieve many of the quantities we want to know about aerosols to the level of uncertainty we wish to know them at. Still, I appreciate what the authors are trying to say here, and am perhaps being a bit picky.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



I would suggest rewording to 'has been used to retrieve aerosol properties over land and ocean' or similar.

Page 3508, lines 20-21: Just FYI, in Collection 6, calibration improvements made by the MODIS Characterization Support Team (MCST) mean that Terra and Aqua should both be temporally complete for Deep Blue. But for Collection 5 data, the authors are correct that Aqua gives the more complete record.

Page 3509, lines 3-4: Again FYI, by the time that peer-review for this paper has been completed, the MODIS Aqua Collection 6 level 3 products should be available and so the authors should be able to use the 'merged' (and/or standalone DB/DT) dataset(s) as they may choose. However as this paper is in part a proof of concept of this new analysis type, I would not say that it is necessary for publication that the authors repeat the analysis using C6 (although I would encourage it, should the authors have the capability, as C6 will hopefully become one of the 'standard' datasets for the coming years).

Page 3509, lines 22-23: If I have understood this sentence correctly, the authors are talking about the narrow MISR swath meaning the revisit time may be too long to resolve small-scale temporal variability? If so, I would explicitly say 'lower spatial sampling' or similar rather than just 'lower sampling'.

Page 3510, lines 2-3: As the SeaWiFS mission ended in 2010, I think it would be better to use past tense (i.e. 'covered' rather than 'covers') here.

Page 3510, line 19: I suggest 'agreement' rather than 'agreements'.

Page 3510, line 22: I suggest rephrasing to avoid starting a sentence with 'And'. 'Additionally,' may be suitable here.

Page 3511, line 14: The Smirnov et al (2002) reference cited here is missing from the bibliography, and I think it should be Smirnov et al, RSE (2000) instead for cloud screening. The Level 2 quality assurance document is a different one, which is not

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



cited here. Both should probably be cited:

Holben, B.N., T.F. Eck, I. Slutsker, A. Smirnov, A. Sinyuk, J.S. Schafer, D.M. Giles, and O. Dubovik (2006). Aeronet's Version 2.0 quality assurance criteria Proc. SPIE, 6408(64080Q), DOI:10.1117/12.70652

Smirnov, A., Holben, B. N., Eck, T. F., Dubovik, O., and Slutsker, I.: Cloud-screening and quality control algorithms for the AERONET database, *Remote Sens. Environ.*, 73, 337–349, 2000

Page 3511, lines 26-27: It would be cleaner to write 'Gandhi College' instead of 'Gandhi_College'; AERONET site names sometimes have these extra characters so the filenames don't contain spaces, but that isn't necessary when writing text. Also, I think saying 'Singapore in Singapore City' is redundant. I'd change the end of this sentence to 'Mukdahan in Thailand, and Singapore.'

Page 3511, line 27: One of the sites listed in Table 1 is Izana, which is at around 2.4 km altitude in the Canary Islands. I think that this site is probably not useful for the analysis presented in this paper because of this high altitude, meaning that AERONET won't be sampling a lot of the total column AOD (as it will be physically located below the site). From the AERONET webpage (http://aeronet.gsfc.nasa.gov/new_web/photo_db/Izana.html): 'The observatory is situated on the top of a mountain plateau located in a pre-national park area protected by the sky law. IZO is normally above a temperature inversion layer, and so, free from local anthropogenic source influences.' So using this as a point of comparison against satellite data is not particularly representative; certainly a 2.4 km mountain can't be considered a good representation of this region (island chain) on a 1x1 degree spatial scale! The site of Santa Cruz Tenerife on the same island may be suitable, as it is low-lying and seems to have a good data volume (http://aeronet.gsfc.nasa.gov/new_web/photo_db/Santa_Cruz_Tenerife.html). If you compare typical AOD from Tenerife and Izana you can notice quite large differences,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

which supports the contention that a lot of the time the aerosol is mostly concentrated below Izana. I recommend that Izana is removed from the analysis, and Santa Cruz Tenerife considered in its place (if the authors find the data record to be suitable).

Page 3512, lines 18-19: I am not sure it is correct to say 'the interpolation performs well without introducing much uncertainty', as this isn't something which is really quantified. The gap-filled time series at Minsk looks plausible but that could just be the case because what's being used to fill in the gaps is basically a perturbation to the average seasonal cycle for the missing months. Again, I understand what the authors are trying to say here, but I don't think this assertion is supported. I think they can say that the interpolation seems reasonable, for sure. Maybe they could do a test by manually deleting and interpolating some months from the AERONET time series, and quantifying the error resulting from the interpolation there. Of course it won't be possible to know the uncertainty when there is a real data gap, but we'd be able to get some idea of typical magnitudes. Since the CMCA seems to be driven a lot by the month-to-month variability, which is in itself dominated at many locations by a seasonal cycle which is fairly repeatable between years, perhaps the reinstatement of the seasonal cycle in this interpolated data means that any errors introduced by the interpolation don't have much effect on the outcome. Either way, I think that this aspect is something which the authors should quantify and discuss a bit more.

Page 3512, line 23: I am surprised about the hole in central Australia for MODIS. Deep Blue should be providing a fairly complete coverage there, and the NDVI should be low enough that the merging process the authors are performing fills in the holes in the DT data with DB there. This has been our experience with the Collection 6 data (see e.g. Figure 19 of Levy et al., AMT, 2013), and based on browse images of monthly Collection 5 data (available online at http://modis-atmos.gsfc.nasa.gov/MYD08_M3/browse_c51.html), fairly complete Deep Blue coverage here seems to be the case for Collection 5 as well. The authors should check into this.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Page 3513, line 18: 'averaged' not 'avearged'.

Page 3514, line 8: The cross-covariance matrix is the matrix C? This is what I infer but it isn't stated explicitly. Also, shouldn't C be in bold and regular typeface rather than italic, as it is a matrix?

Page 3414, line 11: X_AERONET^T should be in bold regular typeface, as it is elsewhere, rather than italic, as it is a matrix.

Page 3514, line 17: I think it would be better to say 'in descending order of magnitude', to be explicit.

Page 3515, line 8: 'we divide V' rather than 'we divide the V matrix'

Page 3516, line 12: I think it would be clearer to say that the first 3 modes 'each' explain >10% of the variance, rather than 'all' explain it. Based on eyeballing Figure 5, it looks like together modes 1-3 explain about 70% of the total variance. If I understand things correctly, this is suggesting that 70% of the global variability of the aerosol field (on monthly time scales and spatial scales of 1 degree) can be explained in terms of three basic components. The authors mention the links to their previous work and say they won't discuss it here, but I think that it is worth mentioning this point again here briefly, as it is in my mind quite an interesting result (3 modes being so dominant).

Section 4, figure 7 general discussion: This is a figure making quite an important point about the agreement between various satellite data products. Now as the authors are looking at monthly mean fields, there are two main things which could cause these disagreements. One question is retrieval biases at high and low AODs, related mainly to aerosol model and surface reflectance assumptions respectively, which will affect the peaks and troughs of seasonal cycles. Another is related to observability, which is in part a function of swath width and aerosol temporal variability but also things like cloud cover and each algorithm's success at being able to perform a retrieval at all. See for example Figure 1 and discussion of Reid et al (Atmos. Res., 2013,

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



doi:10.1016/j.atmosres.2012.06.005) for about this in the context of south-eastern Asia (where availability of retrievals can be a challenge). And then the extent to which observability will be a problem will depend on things like the spatial and temporal scales the data are aggregated at. The authors hint at these topics a little on pages 3517-3518, but I think it would be good to mention these aspects more directly. It is my impression that the fact our validation papers against AERONET tend to dwell on level 2 comparisons (which focus mainly on retrieval-level errors rather than sampling-related representivity) means that observability issues have been given comparatively less attention when they can be important for studies such as this, which are using level 3 data. And the issue of observability is not limited only to the satellite datasets-the AERONET monthly means are (to my best knowledge) the means of daily mean data. So there are questions about how many observations does AERONET need to be representative of the aerosol burden on a given day, and how many days with observations are needed to be representative of a given month, the answers to which are probably dependent on the temporal variability of AOD at each individual site. In other words, AERONET can have this 'sampling uncertainty' on a monthly mean too.

Page 3518, line 7: I think the authors can just say 'Borneo' rather than 'Borneo Island'.

Page 3518, lines 9-15: I think it could be worth expanding on this statement a little. Specifically, as SeaWiFS lacks thermal IR channels (which are useful for cloud screening), one of the main methods of cloud detection over land is spatial variability (as clouds tend to be bright and spatially variable). Unfortunately, some aerosol plumes have a similar appearance. So there can be a tradeoff in cloud screening between errors of omission (i.e. missing a real cloud) and errors of commission (i.e. saying something is a cloud when it isn't). Our goal was to err on the side of caution and minimise the likelihood of clouds incorrectly being labelled as cloud-free. Thus, in some areas (including Borneo but also e.g. the Amazon which also often has small clouds and smoke aerosol plumes), coverage is reduced because sometimes these smoke plumes are incorrectly labelled as clouds and so the pixels discarded. And as well as

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

low-level clouds, there can often be a thin cirrus layer above the area which can result in positive biases in the retrievals if not identified. So this is also related to the observability issue I mentioned above. The cirrus can affect AERONET observations too, see e.g. Huang et al. (JGR, 2012, doi:10.1029/2012JD017757), Chew et al. (Atmos. Res., 2013, doi:10.1016/j.atmosenv.2013.06.026), and others. On the topic of future AERONET deployment the authors bring up here, in the last few years 3 AERONET sites have been acquiring data in Borneo (Kuching, Pontianak, Palangkaraya) although unfortunately yes, there was nothing available during the analysis period of this study. See Salinas et al. (Atmos Env, 2013, doi:10.1016/j.atmosenv.2013.02.016) for some first results at Kuching.

Page 3518, line 12: 'biomass' not 'bioass'.

Page 3518, line 16: Here there is a jump between looking at the full dataset to looking at the (deseasonalised) anomaly dataset). I think it would be clearer if this part became a new subsection 4.1. Then section 4.1 could be renamed 'Global analysis – full dataset' and the new section 4.2 'Global analysis – anomaly dataset' or similar.

Page 3518, line 16: 'anomaly' not 'anomly'

Page 3520, line 23: 'Gangetic' not 'Genetic'.

Page 3523, line 20: 'did not' rather than 'do not'. Also, I think it is unfair to say that the event wasn't captured by SeaWiFS or OMI. The widespread smoke from the fires is evident in the SeaWiFS and OMI records, it just isn't as prominent as in the others. I think that part of the reason is that the SeaWiFS monthly mean has some gaps in this region (I had a quick look through the source data) as the heaviest smoke was flagged as cloud (see aforementioned cloud-screening difficulties due to lack of IR bands on the sensor) meaning that some of the heaviest smoke events were not present in the SeaWiFS data. This in turn propagates into lower level 3 fields. It is possible that the same was true for OMI, although I'm not familiar enough with the algorithm/dataset to say for sure.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



References: Levy et al. (2013) is now published in final form in AMT (the present citation is to AMTD). So this should be updated to:

Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989–3034, doi:10.5194/amt-6-2989-2013, 2013.

Table 1: This table should be checked, as some sites are not where the authors are saying they are. For example, IER Cinzana is in Mali, which is in Africa (not South America). Missoula is in the USA (not Europe). IMS METU-Erdemli is in Turkey, which could be classed as Europe or Asia (but not Africa, where the authors currently place it). There may be more errors, that was from a brief look. Similarly I am not sure why an ‘others’ category is needed – couldn’t ‘North America’ be renamed ‘North/Central America’ and have these two sites listed there, or else make a new ‘Central America’ category for Mexico City and La Parguera? See also prior comment about formatting of site names (e.g. can’t FORTH_CRETE be Forth Crete?).

Figures 11, 13, 14: The presentation of these figures should be improved. The coastlines are very coarse and basic, and there is a lot of white space around the edges of the regions mapped (it looks like the clipping for the data is smaller than the clipping for the map borders).

Interactive comment on *Atmos. Meas. Tech. Discuss.*, 7, 3503, 2014.

AMTD

7, C520–C528, 2014

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

