

Interactive  
Comment

## ***Interactive comment on* “Global cloud top height retrieval using SCIAMACHY limb spectra: model studies and first results” by K.-U. Eichmann et al.**

### **Anonymous Referee #4**

Received and published: 10 September 2015

The paper presents cloud height retrievals from SCIAMACHY limb measurements over the lifetime of the instrument. It shortly presents the retrieval approach, which is a spin-off of the cloud filtering for trace gas retrievals, developed as early as 2005, as well as a cloud height retrieval sensitivity study. It provides an overview of retrieval results, including occurrence statistics, from roughly 10 years of measurements. Outstanding features showing up in the multi-year time series are discussed, and for a couple of them causes could be identified and observed patterns be explained. Also, comparisons to cloud height measurements from other ENVISAT instruments and instrument setups, namely SCIAMACHY nadir and MIPAS measurements, are shown.

For the first time presenting a complete set of SCIAMACHY limb cloud data, the paper suits AMT's publication criteria regarding scope as well as novelty. Presentation of the

C2898

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



material requires a number of improvements, though.

## 1 General Comments

**Introduction:** The authors present a fairly nice listing of existing cloud measurements. I miss an assessment of cloud height measurement quality/accuracies from the different instruments, though, as well as a review of the requirements on cloud height accuracy for current scientific problems. Are, e.g., CTHs with 1.5km or worse accuracies (as yours are by definition) valueable? Also, a more concise overview of the strenghts and weaknesses of the different instruments would be nice, leading to what SCIAMACHY limb measurements offer beyond what other sensors provide.

**Data:** State clearly, which data is used for what: SCIA limb for your original retrievals; SCIA nadir and MIPAS for comparison/validation. Which versions of the comparison data have been used? Provide references.

**Method:** The approach uses a two-wavelength (color index) ratio. I find the argumen-  
tation how the wavelengths bands are chosen insufficient as well as the explanation  
of cloud signatures in the CIR. Questions like these remain unanswered: why these  
2 bands exactly? or are any two bands ok? why 2 at all? Why is at tangent heighth  
radiance ratio at one wavelength not sufficient? what causes the different tangent  
radiance ratio behaviour in the two different bands? What CIR pattern have non-cloud  
cases, what patterns are introduced by clouds (an example of non-cloud CIR profile  
would help the understanding)? What causes the peak (is it a peak at all, i.e. is CIR  
decreasing again after a cloud? under what conditions?)? How has the threshold  
been chosen (no information at all on this is included in this paper, though i assume,  
that is is a fairly crucial choice)?

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Simulations: I find the sensitivity analysis presented too sparse to judge the quality of the method. Particularly, later in the paper the authors draw conclusions from sensitivity checks that have not been presented, i.e. which are intransparent and not reproducible by the reader.

I like to see more details on the sensitivity tests including systematic analysis of cloud height dependence of the CIR (which was obviously done and argued with later, but the reader gets presented only a selective summary, which hinders to see and understand the bigger picture), tangent height dependence, and effect of higher clouds in low tangent height measurements (clouds that might be out of FOV in the high(er) tangent height observation; that such cases can occur is mentioned in sec3 and discussed in sec4.1, but their appearance in a CIR profile and the sensitivity of the measurements to these is never analysed). Preferably also sensitivity to cloud “concentration” (extinction, geometric thickness – cloud with same  $\tau$  likely gives different signal depending on its geometric thickness. but how different is it? how much cloud needs to be within the FOV to get a detection?), and cloud microphysics (specifically, size of ice particles).

Then, I expect stringent conclusions: where are the detectability limits in dependence of cloud height and cloud thickness? Also: how sensitive are these conclusions to the threshold choice? Maybe even: how many clouds are expected to be missed at the different altitudes?

Results: For a measurement that is limited to provide one cloud detection per profile, height resolved cloud fractions seem quite a complex (and somewhat confusing) parameter to look at. On the one hand no cloud hits, i.e. no contribution to CF, will happen below a detected high-altitude cloud, which leads to a shadowing of lower clouds (lower low-cloud fractions in regions with lots of high clouds). On the other hand, high clouds that have been outside the FOV of the high tangent altitudes but are in the FOV of a

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

low tangent altitude are assigned to the low tangent altitude, both in altitude as in geographic position (displacement) and contribute to cloud fraction in the low layer. I find it hard to wrap my head around that and wonder whether the authors were successful to do so.

In the time trend discussion, different versions of the 2-band CIR are applied. However, as characteristics of different band selections are never presented or discussed, the reader hardly can follow the argumentation and conclusions, which makes the whole discussion pointless.

Validation: Most of all, I wonder, why no comparison or validation against CALIOP data is presented. CALIOP by far provides the best dataset to compare to when promising to be able to provide thin cloud CTH. Comparison to SCIA nadir and MIPAS CTH is nice, but rather underlines the issues that each of these datasets has (nadir only sensitive to thick clouds, MIPAS – as SCIA limb – just to uppermost cloud layer CTH and with cloud geolocation issues).

The modification of the SCIA nadir CTHs with their standard deviations seems fairly random. I miss a rationale for taking the standard deviation (why not just taking a constant offset? or a height dependent offset?). Also, original nadir data are never presented (neither mean nor standard deviation), so that the reader can not judge at all the effect of your – random – modifications. Nevertheless, the never presented data is discussed (*“Without the added standard deviation, nadir CTHs were systematically and globally lower. ...”*).

The authors claim a good agreement of SCIA modified nadir and limb CTH without providing sufficient evidence. From comparing Figs 7 and 12, I do not necessarily conclude a good agreement. Basic patterns are quite different (e.g. very high clouds off the western coast of middle America in nadir CTH; fairly high nadir CTH over all of South America, Southern Africa, and Australia). Any quantitative comparison is miss-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

ing, although opportunities for these two measurements are better than for any other (there should be plenty of collocations!). I like to see e.g. a scatter plot like presented for MIPAS CTH and some difference statistics (mean diffs, std of diffs e.lg. for different regions, cloud regimes, ...).

In Sec4.1. horizontal sampling issues have been disacussed in quite some depth. It appears in the discussion of the SCIA limb CTH themselves (Sec5), but is not considered anymore in the comparison to other CTH data, although I expect it to play a significant role.

Volcano event: This whole section seems a little disconnected from the rest, both in terms of target and style. I wonder, whether this would rather suit a different or separate paper.

Throughout the paper, colloquial language (or lab slang) and imprecise formulations is oasionally used (examples see below). Please reformulate properly. This could improve perceptibility of the paper a lot. References are used in odd manner at times (e.g. Reuter&Pfeifer; von Savigny et al., 2005; Hommel et al.). Use appropriately, replace with proper ones, or remove.

Generally, I have the feeling that too many discussion items and conclusions are merely stated, but not backed up sufficiently with explicitly and clearly presented evidence.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

## 2 Specific comments

**p8296, I20:** what does “*global cloud field distribution*” mean?

**p8297, I11:** “*Clouds occur in complex shapes*” – shape as in contour/outline? if not, can you find less ambiguous wording?

**p8298, I1f:** “*The balance between these effects*” – which effects? albedo enhancement and cloud emission are obviously not in balance.

**p8298, I3:** “*The light scattering ability of cloud droplets is only weakly wavelength dependent*” – This is at best imprecise. Wavelength dependence in Mie regime is weaker than in Rayleigh regime, and in VIS/NIR cloud droplets are in Mie regime, while molecules are in Rayleigh. The picture changes, however, when going to other spectral regions.

**p8298, I7:** “*This also enhances light absorption by trace gases [...] above the clouds.*” – Above? Why is that?

**p8298, I15f:** “*Due to the low temperatures [...], cirrus clouds are made of small ice crystals.*” – It’s not the low temperature, but the low absolute humidity.

**p8299, I18f:** “*Nadir viewing, passive instruments [...] were mainly designed to derive trace gas columns*” – I do not agree. Some yes. But many more were designed for weather observations (including clouds).

**p8299, I25ff:** “*Based on the [...] (ISCCP) D2 dataset, a global COD  $\tau_N = 3.9 \pm 0.3$  was found [...] showing that a lot of clouds types cannot be retrieved.*” – What type of instruments is the dataset based on? How does global mean COD show that many types (types? others than cirrus?) can not be retrieved? Or even that many cloud occurrences can not be retrieved? I’d like to see a more stringent argumentation.

**p8299, I27:** “*with a 19.6% global coverage*” – reference missing (or is it from R&S, 1999, too?).

**p8300, I3:** “*An average global cloud amount of about 68% was reported*” – Rather

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



cloud coverage? Amounts can hardly be measured as percentage (what is the reference?).

**p8300, I25ff:** Revise the paragraph. Make clear from the beginning that now, it is SCIAMACHY limb measurements and cloud retrievals that you talk about. “*SCODA was developed using the colour ratio method*” – there is not only one, well-established (aka “the”) colour ratio method.

**p8302, I3ff:** Mention overpass time. What is the horizontal spacing of two limb sequences?

**p8303, I19ff:** The whole paragraph probably fits better in Sec3. In any case, more stringent explanations are required for the reader to be able to follow you. E.g. “*Higher intensities can be seen in channel 6 when cloud scattering occurs*” – higher than what? are they really restricted limited to ch6? “*A steeper gradient between radiances from both channels*” – i.e.,  $I_{\text{ch4}}^{15.4\text{km}} / I_{\text{ch6}}^{15.4\text{km}} > I_{\text{ch4}}^{12.1\text{km}} / I_{\text{ch6}}^{12.1\text{km}}$ ? “*This means that clouds are in the field of view at tangent height 12.1 km and scatter more radiance into the line of sight.*” – This does not explain the (changing) “color” gradient. So, where does the color gradient come from? And why does a changing or high gradient imply a cloud? What are the physical processes causing that? Would be helpful to see a Fig.2 equivalent for a cloud-free case.

**p8303, I26ff:** What is the point of this paragraph? Discussing the choice of spectral retrieval windows? It is insufficient for this. And does not present any conclusions/choices.

**p8303, I26:** “*Spectral windows with molecular absorption bands*” – Reformulate, Common terminology is that (spectral) windows are the regions with low gas absorption.

**p8304, I12f:** Does tangent height knowledge refer to geometric heights, or are refraction effects included?

**p8304, I24ff:** Why/how is limb/nadir matching (and what is that?) related to platform height?

**p8306, I15:** “*The standard operational CI approach*” – what is the standard approach?”

C2904

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



provide a reference.

**Fig.3:** Wavelength bands indicated by dotted, not dashed lines (as claimed by caption).

**p8307, I1:** *“Water and ice clouds show similar reflection characteristics in this wavelength range but start to differ at wavelengths larger than 1400 nm”* – Which characteristics specifically are meant? Your analysis later on (p8312) seems to contradict this statement.

**p8307, I18:** *“where “h” denotes the high and “l” the low band.”* – high and low what? wavelengths?

**p8308, I6:** *“Figure 5a shows an example of the cloud index ratio profile”* – Fig5 seems inconsistent with Eq. (1) and Fig2. The latter indicate  $CI < 1$  ( $I_h < I_l$ ) and CI closer to 1 for lower tangent altitudes, while Fig5 shows only  $CI > 1$  and CI being large at low tangent altitudes.

**p8308, I15:** *“but we only store the highest peak”* – ‘highest’ as in highest tangent altitude or as in largest CIR?

**p8308, I27:** *“the retrieved cloud fractions”* – first occurrence of ‘cloud fraction’. not straight forward, what CF means for limb measurements, hence requires explanation. here. Besides, I find the term cloud occurrence frequency more suitable (cloud fraction, in my opinion, is rather the coverage within a sensor FOV or a modeling grid cell, i.e. describes sub-resolution issues).

**p8309, I1ff:** What area exactly do you calculate here? The area within the instrument FOV intersecting a  $\Delta z_{th}$  layer (that’s what I would expect as the correct one), or the area of a  $\Delta z_{th}$  thick plane layer intersecting a  $\Delta z_{th}$  thick shell (as Fig.2 suggests)? and to be more correct, shouldn’t it be the intersection volume? would that make a significant difference?

**p8309, I19:** *“the air mass above a cloud has less air”* – less air than what? in the cloud-free there’s as much (or little) air in this region as in the cloudy case. revise formulation.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper





**Fig.5:** “(see Figure above)” – Refer to specific, numbered figure.

**missing:** I miss a clear statement that your method restricts CTH retrieval accuracy to tangent altitude spacing, i.e., it is kind of binned data. This should also be considered in the validation discussion.

**p8309, I25f:** “*Simulating radiances in cloudy atmospheres is difficult [...] because radiative transfer models [...] treat clouds only as two-dimensional layers*” – First, cloudy simulations are complex even if one fully considers 3D properties, purely due to the complex nature and high variability of clouds themselves (as you mentioned in the Introduction). Second, please avoid making it sound like all models have this limitation (e.g., DART and MYSTIC provide 3D capabilities for scattered solar radiation). Lastly, what do you refer to by ‘two-dimensional layers’? Do you actually take into account layer inhomogeneity in the observation plane?

**p8310, I12:** “*A model accuracy of 5% was found*” – compared to what? what constitutes the truth/reference? Is the accuracy estimate valid at all with the forward model reference from 2014, the accuracy reference from 2003?

**p8310, I12:** “*Cloud top heights can best be retrieved by using the same instrument and measurement mode*” – generally? why would that be? or just if used for cloud-flagging in trace gas retrievals (if so, how is it relevant here?)? best for what? what at all is the point to discuss retrievals here again, in the forward modeling section?

**p8310, I22:** “*The SAA is the angle between the line of sight and the sun beam*” – no, that would be the single scattering angle.

**p8310, I28f:** “*The Rayleigh scattering phase function*” – aren’t you discussing clouds? which are in Mie regime?

**p8311, I7ff:** “*A spherical atmosphere [...]. The scalar discrete ordinate technique was used*” – Does it apply a true spherical or a pseudo-spherical solution for the scattering calculations?

**p8311, I13ff:** Is the spectral setup consistent with SCIAMACHY’s instrument characteristics and the spectral bands used in the cloud height retrievals? what intermediate angle steps are calculated?

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

**p8311, I17:** *“To find the limitations”* – Of what? Cloud retrievals? Trace gas retrievals? ...?

**p8311, I20:** Do you use monodisperse clouds? Monodispersions are known to exhibit oscillations in their optical properties (scattering efficiency as well as phase function), which are significantly dampened in realistic bulks. Are you sure this affects your sensitivity estimates in a negligible manner?

**p8311, I23:** What is the source of the ice crystal optical property data (assuming the droplet properties are from Mie calculations)?

**p8312, I6:** What is the rationale to use Henyey-Greenstein for aerosols? Based on what have the asymmetry parameters been chosen?

**p8312, I7ff:** *“In Fig. 6 [...]”* – Simulation setup description is redundant. However, for which tangent altitudes have the simulations been made? Do these simulations include the aerosols?

**p8312, I12f:** *“Cloud tops were in general detectable [...], because the modelled  $\Theta$  exceeds the threshold of 1.4.”* – This seems to be the wrong argumentation direction. Where at all does the threshold come from (explanation missing in Sec3)?

**p8312, I19:** *“The phase function of an ice particle is more complex and has a stronger asymmetry. Thus [...]”* – This contradicts your statement from p8307, I1.

**p8313, I26ff:** What is the effect of having the surface in the FOV on CIR? over the range of possible surface reflection properties (is sunglint an issue?). can the cloud detection threshold be reached in non-cloud cases due to surface effects? considering this, is cloud detection reliable for views with surface in FOV?

**p8314, I5ff:** *“But the cloud retrieval can be obstructed [...], when the optical thickness of cloud and aerosols has the same order of magnitude.”* – Do you say that a thin aerosol layer (say  $\tau=0.01$ ) would not obstruct a thick low water cloud (say  $\tau=5$ ), since they do not exhibit same order of magnitude  $\tau$ ? Reformulate otherwise.

**p8314, I8f:** I suggest to make it clear already in the title, that along-track, or rather along-LOS, sampling (resolution?) is discussed here (in contrast to the across-track sampling that SCIAMACHY is doing).

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

**p8314, I10ff:** That real sample path is longer than the 1-shell path  $2x(i,i)$  is NOT an effect of the  $2x(i,i)$ -shell not being resolvable. Reformulate.

**p8314, I15:** I have problem to follow. Where is tangent height going into this? A terminological distinction between true CTH (wherever along LOS) and retrieved CTH (at tangent height) could be helpful.

**p8314, I24:** “For most of the cases [...]” – Do you mean that for no ‘tomographic LOS intersections’ exist within the troposphere?

**p8314, I26ff:** “But if the LOS of a cloud free pixel [...]” – Why would the resolution be depending on (non-)cloudiness?

**p8315, I9:** “to derive a trace gas profile at the tangent height.” – ‘profile’ commonly refers to vertical dependency of a parameter (e.g. VMR(z)). How do you relate it to a tangent height, which is just 1 point in z? Or do you mean the profile at the (horizontal) geographic position of the tangent point?

**p8315, I11ff:** “the retrieved CTH will be higher in this case. These high clouds outside the along-track FOV or lowest shell can lead to a CTH overestimation” – CTH will be higher compared to what? why would a cloud outside the FOV at all lead to a wrong estimation of CTH? I do not understand how any of this could lead to a CTH OVER-estimation. The only one, I see, is an underestimation: if there is a cloud high along the LOS, which is not in the FOV for the corresponding high tangent altitude. This high cloud would cause a cloud hit for the low tangent altitude, i.e. a low CTH estimate.

**p8315, I13:** cloud fraction needs to be introduced before discussing it.

**Fig. 6:** “Simulated maximum colour index ratios” – Maximum over what? And why, maximum at all? Furthermore, what tangent altitudes are those figures for? what does ‘T\_A’ in plot titles refer to?

**p8315, I14:** Surely a matter of taste, but Sec6&7 are results, too. Hence, I’d chose a more specific Sec5 title.

**p8315, I16:** “We retrieved annual means of cloud top heights and fractions” – ‘Annual

mean cloud fraction' does not make sense. Formulate properly.

**p8315, I16ff:** Cloud fraction should be a relative number, not a pixel count. Formulate properly. Furthermore, how is the geographical position assignment done? Is the cloud assigned to the position of the tangent point?

**p8315, I21:** Any rationale for the grid size? As you have demonstrated before, the path length through a CTH shell is about 400km, i.e. roughly  $4^\circ$  big. Seems more reasonable to me to have the grid big enough such that all of the possible cloud positions within this shell fall into one cell.

**p8315, I21f:** *"All measurements [...], that exceeded the threshold"* – Here, one limb scan is one measurement? Or does it include all points within a vertical scan that exceed the threshold?

**p8316, I1:** *"A cloud image (Reuter and Pfeifer, 2011)"* – what is the Reuter&Pfeifer contribution to this image, what is special about it that it is referenced? particularly, here in text instead of in figure caption (e.g. as 'adapted from')?

**p8316, I12f:** *"which will be analysed later."* – future work? or in this paper? if the later, specifically refer to the section, where it is discussed.

**p8316, I16:** *"This is partly due to dependence of the phase function on the viewing geometry"* – what else is it/could it be due to? Phase function (as an inherent property of a particle) does not depend on viewing geometry. Single scattering angle does (and hence, 'efficient scattering coefficient' =  $\text{scat.coef} * \text{pfc}(\text{SSA})$ ).

**p8317, I4f:** *"This can be attributed to"* – Is it absolutely clear that this is the reason, or is it speculation (if so: might be attributed)? Could smearing (or displacement along LOS) effects play a role, too?

**p8317, I13:** *"Clouds are limited [...]"* – what's the point here? I don't understand what you are trying to tell the reader. Besides, are PSC not clouds (as they are limited to troposphere)? Use a better reference for the general statement on PSC altitudes (this fact has been known far before the von Savigny paper).

**p8317, I20:** *"It seems that the method is rather insensitive to low clouds"* – You never mention or discuss the "shadowing" effect of high clouds (only highest cloud in a limb

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

scan is detected, clouds further below are neglected. This will produce a low bias of low clouds in regions with frequent high(er) clouds. Might this contribute? Also, might displacement effect contribute? Particularly, displacement effects could also have a directional dependence (i expect that looking from a high-cloud region into a low-cloud region along track (or LOS) gives a different signature than looking from low- into high-cloud region).

**p8317, I24:** What sensors did Generoso et al. use?

**p8318, I1:** *“But CALIPSO [...]”* – CALIPSO (or CALIOP? as you made that distinction between platform and instrument before) should also be able to detect enhanced aerosol layers. That is, this can not be the reason for not appearing in CALIPSO. Did you check CALIPSO aerosol data? What else can explain the absence of a signature in CALIPSO? Without another explanation, I’d rather suspect it to be an artefact of your method.

**p8318, I25f:** *“aerosol layers are optically very thin and the corresponding CIR is lower than for typical tropospheric clouds.”* – is this due to low  $\tau$  or due to the aerosols as such (a stringent presentation of cloud altitude/optical depth dependency of CIR in Sec3 would help greatly)?

**p8318, I27:** *“Two other wavelength pairs were tested.”* – Which? Why and why these in particular? How were the wavelengths selected? What signature do we expect there? Is 1.6/1.5um combi always constant, or just its mean?

**p8318, I28:** *“During the longer decontamination phases”* – Does this imply some instrumental issues affecting the CIR retrievals?

**p8319, I3:** *“This indicates that the ratio is not affected by aerosols as much.”* – Which of the ratios? Not as much as what? Is there a (physical/optical/rad.transfer) explanation for that? You get the reader far too little background/explanation to follow any conclusions you draw (1.6/1.5um combi mean is so stable that I suspect it might not show any cloud signature at all).

**p8319, I3f:** *“A CIR decrease over time might reduce the sensitivity”* – Unclear. Does not make sense to me. CIR is an expression of the atmospheric state. E.g., thin clouds

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

produce smaller CIR. Still, they do not effect the sensitivity (for a given atmospheric state, the method is still sensitive to the same clouds).

**p8319, I13:** What are you referencing by Hommel et al.? You are discussing your own results here, aren't you?

**p8319, I17:** *"A considerable increase in CTH was found starting at the end of 2007"* – How long did it persist? I can see it exclusively at turn of 2007/08.

**Fig. 7:** *"The MSG picture [...] has been adapted."* – How? Why? Do you mean: 'Adapted from Reuter&Pfeifer'? Does (b) imply SCIAMACHY is looking straight south? is it like this? or is the image not geographic-north oriented?

**Fig. 8:** Is it correct that CIR are 14–22? seems very high.

**Fig. 9:** Can you clarify how height resolved cloud fractions are defined? Such that sum over all layers gives total CF?

Generally, why are the comparisons only done or presented over limited periods of time instead the full lifetime of the instruments?

**p8320, I14:** *"as in limb only the highest CTHs are detected in a grid cell."* – in my understanding nadir also detects only the "highest" CTH (or are SCIA nadir CTH multi-layer cloud heights?), just along a nadir LOS it takes longer (hence, deeper into the atmosphere) to reach the necessary path optical depth to get a radiometric effect.

**p8320, I15:** *"Only data were used for the following SACURA quality checks:"* – Sentence does not make sense. Parts missing?

**p8320, I20f:** *"Very high limb CTHs over South America and South Africa were only partly seen in nadir data."* – This is inconsistent with your Figures. CTH over South America and South Africa are higher in (modified) nadir data than in limb data!

**p8321, I5:** *"the nadir retrieval is restricted to the detection of water clouds"* – Do you say that SCIA nadir can not detect any ice clouds?

**p8321, I14f:** *"The black line and the blue area depict the SACURA nadir results and its 1- $\sigma$  scatter."* – No blue area in my Fig13.

**p8321, I26ff:** *"Better agreement can be achieved with other limb viewing instruments"* – Better agreement maybe, but other limb measurements suffer from the same

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

difficulties as SCIA limb. Hence, I'm sceptical about the independence of the two measurements and about the qualification of the MIPAS data as reference data set. Furthermore, with SCIA looking forward and MIPAS backward, one can expect the smearing/cloud displacement effects to act in different directions. Please discuss implications.

**p8322, I8f:** *“we do not see the gaps in vertical direction of the SCIAMACHY cloud top heights.”* – Reformulate properly. Of course we see the gaps in the SCIA data. Why should we see SCIA gaps in MIPAS data?

**p8322, I11f:** *“But there is [...] MIPAS top heights that clearly lie outside the vertical field of view”* – Outside of which VFOV? MIPAS'?

**p8322, I15:** *“where the lowest possible MIPAS tangent heights were at about 10 km, which can partly explain the differences.”* – How does this explain differences for individual collocated measurements? If the cloud is high enough for MIPAS to detect it (else it would not be in the scatter plot), SCIA could be able to detect it, too, would it not?

**p8322, I18ff:** *“The vertical differences are around  $-1.1$  km [...]. This difference can be mainly attributed to the different tangent height step sizes”* – I am not convinced. Fig15 shows a systematic tangent height dependency of the differences. Explain, how this is in line with the different vertical resolution you attribute it to. If it were an resolution issues, I'd expect random differences.

**p8322, I24ff:** *“Sembhi et al. (2012) compared MIPAS [...] and CALIOP [...] and found that MIPAS CTHs are in the range of up to 1 km higher for altitudes between 12 and 20 km, which is in line with our comparisons.”* – I find that a too far-fetched conclusion. Please compare SCIA limb CTH and CALIOP data directly.

**p8323, I5f:** *“Results from the lowest tangent height should be taken with caution, as MIPAS has latitude dependent tangent height cycle.”* – How does this affect MIPAS CTH? I would expect that MIPAS retrieved CTH are corrected for varying tangent height, and when comparing just collocated positive cloud hits (that is what you are doing, are you not?), one excludes cases with CTH too low to be seen by MIPAS.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Saying that, it would be interesting to see a contingency table over all (cloudy and non-cloudy) collocations.

**p8323, I6ff:** *“Higher cloud top heights [...] at the highest SCIAMACHY tangent height 18.5 km. This can partially be explained by the coarser tangent height step size of SCIAMACHY, as the 18.5 km height is generally above the tropopause.”* – How does this explain the difference? Is MIPAS restricted to tropospheric cloud detection?

**p8323, I27:** *“zonal mean CTH for clouds between 6 and 28 km”* – Why globally from 6km if MIPAS has issues measuring CTH lower 9 or 10km in the tropics?

**p8323, I27f:** *“Furthermore we have compared [...]”* – Please show results. You keep claiming good agreement without allowing the reader to judge by oneself.

**p8324, I2f:** *“the interhemispheric CTH differences in the SCIAMACHY data had disappeared.”* – Why is this here? No IH-differences discussed in this section before. How is that related to MIPAS data?

**p8324, I10f:** *“as the CIR  $\Theta$  is strongly height dependent. Because of the lower values of  $\Theta$  towards the surface the extra scattering due to aerosols have a high impact.”* – Neither of this has been shown before. Add in Sec3.

**p8324, I13:** *“But as shown in the cloud fraction distributions,”* – This has at best been speculated there, not shown (as in giving some evidence).

**p8325, I12:** *“was estimated for both volcanoes from their measurements.”* – From whose/which measurements?

**p8325, I23:** *“the lowest layer [...] is situated”* – lowest layer of what? Inconsistent with layer range discussed above.

**p8325, I26:** What is occurrence frequency? Same as/related to cloud fraction? How has it been derived? Like CF in Sections before (e.g.  $2 \times 2^\circ$  cells)?

**p8326, I12f:** *“Overall about 45% of all measurements in the lower stratosphere”* – measurement equals “cloud” hit?

**p8326, I20:** *“They detected the plume”* – Who? Doeringer et al.? It reads odd to refer to subjects that have only been introduced in parenthesis-cites before.



**p8327, I5ff:** *“Although the use of only one threshold for all atmospheric situations is simplistic in nature, it was shown that it was sufficient for the majority of cases”* – I disagree. It was not demonstrated what happens, when threshold is changed (what do we know how many cases where not or falsely detected?), i.e. it can not be rated as “shown”.

**p8327, I13f:** *“SCIATRAN model studies have shown that the method is very sensitive for a wide range of cloud optical properties and cloud top heights.”* – Not in this paper, though. At least I do not call two cloud cases with only cloud height varied “a wide range”.

**p8327, I20f:** *“SCIAMACHY CTHs were generally lower by about 1 km, which can be explained by differences in the tangent height steps.”* – Disagree. Not properly explained to readers.

**p8328, I2:** *“It was shown that the height differences can partly be attributed to the different sensitivities of the viewing geometries.”* – Was not shown, but stated.

**p8328, I6f:** *“Unfortunately, SCIAMACHY was the only instrument so far that was capable of making use of the two viewing geometries.”* – I do not see a strict requirement of having both realized within one instrument. Two simultaneously measuring nadir/downlooking and limb instruments should provide equally good data (maybe even better, since continuous data).

**p8328, I11:** *“The use other wavelength pairs in the near IR [...] enables to distinguish between clouds and aerosols.”* – Please provide a reference proving that it is indeed (and not just might be) possible or reformulate the statement.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

### 3 Technical corrections

Throughout the paper: Use consistent spelling (e.g., Nadir vs. nadir), terminology (e.g., CIR or  $\Theta$  or CIR  $\Theta$ , cloud fraction vs. cloud occurrence frequency, Figure or Fig.) and tenses (e.g., p8310 “*We have studied*” vs. “*colour index ratio will be studied*”, p8310 “*ENVISAT moves/flies*” but “*zenith angles occurred*”, p8312 “*VFOV was at*” vs. “*VFOV edges are then*”, etc.). Unit specifications for plots belong in the figure captions (or the plots themselves), not in the text.

**p8297, I25:** “*infra-red*” → infra-red

**p8303, I2:** “*The full Earth was covered after 6 days at the equator*” – Reformulate. Equator can never be full Earth.

**p8304, I22:** Either “annual cycle” or “inter-annual variabilities”.

**p8305, I4:** & **p8306, I18:** “*SACURA retrieves*” and “*The limb sounder MIPAS retrieved*” – Odd language. What is the retrieving entity, the retrieval algorithm or the instrument?

**p8305, I25ff:** “*Due to the specific spectral measurement range MIPAS [...] as a pure limb sounder recorded more limb scans [...]*” – does not makes sense. reformulate.

**p8310, I26:** “*the highest sun zenith angles*” → the largest sun zenith angles.

**p8311, I3:** “*SAA larger than 90°*” → “*SAA around than 90°*”?

**p8312, I13:** “*CIR was highest for very low SAA and lowest around 90° SAA, which is in the tropical region.*” – which of the two is (rather: occur) in the tropics? Besides, lowest CIR seems to occur for high SZA (south polar region).

**p8312, I18:** “*where the threshold was not reached (see Table 1).*” – wrong reference? Table 1 does not deal with any thresholds.

**p8312, I27:** “*aerosols are within the first 5 km*” – Reformulate properly (first of what? along LOS? from TOA? ... ?).

**p8322, I22:** “*field of view increased*” – wrong tense (at least, i expect it is always increasing).

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



**p8315, I3:** *“that the line-of-sight crosses”* – rather: that the (V)FOV crosses/intersects.

**p8317, I23:** *“High fractions were also detected at the height range (Fig. 9b).”* – Parts of sentence missing?

**p8318, I21:** *“a rather stable period from the start”* – Lab slang (start of what?). Formulate properly.

**p8320, I9:** *“low geometrical limb resolution”* → low horizontal (or along-LOS) resolution of limb measurements.

**Figure captions:** Reformulate captions with correct use of sub-figure references. For example, *“Global distribution of [...] (b) to illustrate [...]”* and *“Global annual mean [...] (b) the limb cloud fraction”* are no proper sentences and do not make sense.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)