

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

# Shape-temperature relationships of pristine ice crystals derived from polarimetric cloud radar observations during the ACCEPT campaign

A. Myagkov, P. Seifert, U. Wandinger, J. Bühl, and R. Engelmann

Leibniz Institute for Tropospheric Research (TROPOS), Permoserstr. 15,  
04318, Leipzig, Germany

Received: 24 November 2015 – Accepted: 9 December 2015 – Published: 15 January 2016

Correspondence to: A. Myagkov (myagkov@tropos.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

AMTD

doi:10.5194/amt-2015-365

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

This paper presents first quantitative estimations of ice particle shape at the top of liquid-topped clouds. Analyzed ice particles were formed in the presence of super-cooled water and in the temperature range from  $-20$  to  $-3^{\circ}\text{C}$ . The estimation is based on polarizability ratios of ice particles measured by a Ka-band cloud radar MIRA-35 with hybrid polarimetric configuration, manufactured by METEK GmbH. For this study, 22 cases observed during the ACCEPT (Analysis of the Composition of Clouds with Extended Polarization Techniques) field campaign were used. Polarizability ratios retrieved for cloud layers with the cloud top temperatures of  $\sim -5$ ,  $\sim -8$ ,  $\sim -15$ , and  $\sim -20^{\circ}\text{C}$  were 1.6, 0.9, 0.6, and 0.9, respectively. Such values correspond to prolate, quasi-isotropic, oblate, and quasi-isotropic particles, respectively. Data from a free-fall chamber were used for the comparison. A good agreement of detected shapes with well-known shape–temperature dependencies observed in laboratories was found. Polarizability ratios used for the analysis were estimated for areas located close to the cloud top where aggregation and riming processes do not strongly affect ice particles. We concluded, that in microwave scattering models ice particles detected in these areas can be assumed to have pristine shapes. It was also found that even slight variations of ambient conditions at the cloud top with temperatures warmer than  $\sim -5^{\circ}\text{C}$  can lead to rapid changes of ice crystal shape.

## 1 Introduction

Mixed-phase clouds are a crucial component of the Earth’s climate system. Their long-lasting nature impacts the radiative budget and the thermodynamic structure of the atmosphere (Sun and Shine, 1995) and microphysical processes occurring in mixed-phase clouds are the main source of precipitation (Mülmenstädt et al., 2015). Yet, current representations of mixed-phase clouds in cloud-resolving and numerical weather prediction models have significant uncertainties because numerous processes occur-

AMTD

doi:10.5194/amt-2015-365

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ring in this type of clouds are understood insufficiently (Luo et al., 2008; Klein et al., 2009; Morrison et al., 2009; Bouniol et al., 2010; Delanoë et al., 2011). The uncertainties result from an incomplete understanding of the formation, microphysical evolution, and removal processes of mixed-phase clouds. The transition from the liquid phase to the ice phase and its partitioning depends strongly on the environmental temperature and the properties of the available ice nucleating particles (DeMott et al., 2010; Hoose and Möhler, 2012; Kanitz et al., 2011; Seifert et al., 2015). Temperature, pressure, and humidity at which ice crystals are formed also define their shape and apparent density which determine the sedimentation velocity (Fukuta and Takahashi, 1999; Bailey and Hallett, 2009). The vertical and dynamical structure of mixed-phase clouds furthermore determines the type and intensity of ice multiplication processes whose occurrence are a prerequisite for the formation of intensive precipitation (Hallett, 1974; Rangno and Hobbs, 2005; Seifert and Beheng, 2006).

Ground-based remote sensing has shown a large potential for improving the understanding of the life cycle of mixed-phase clouds (Hogan et al., 2003; Ansmann et al., 2009; De Boer et al., 2009; Delanoë and Hogan, 2010; Kanitz et al., 2011; Westbrook and Illingworth, 2013). Even though microphysical retrieval techniques based on ground-based remote observations are a valuable source of information for investigation of mixed-phase clouds, further developments are required in order to increase the accuracy of these retrievals. Important but yet barely explored parameters are the shape and apparent density of an ice crystal population. Estimates of ice mass, area, or number concentration require accurate knowledge of particle shape (Westbrook and Heymsfield, 2011; Delanoë et al., 2014).

From the remote-sensing perspective, mixed-phase clouds with a single supercooled liquid layer at the top and ice virgae below are of special interest (Wang et al., 2004; Smith et al., 2009). Further, we denote such clouds as single-layer clouds. Single-layer clouds have less complex microphysical and dynamical properties (Fleishauer et al., 2002; Ansmann et al., 2009; Zhang et al., 2012) compared to convective cloud systems where more than 25 different transfer processes may take place (Seifert and Beheng,



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



approximated scattering properties of columnar-shaped and plate-wise ice particles by prolate and oblate spheroids, respectively. Dielectric properties of ice particles depend on apparent ice density (Oguchi, 1983), which characterizes the ratio of air and ice within the approximating spheroidal particle. Observational weather radars are often operating nearly horizontally (at low elevation angles of the antenna). The authors claimed that under such conditions prolate and oblate particles can have similar polarimetric signatures and can not be reliably distinguished. As the result the proposed method is only applicable for data with high values of  $Z_{DR}$  which can be only induced by strongly oblate particles and which, therefore, can be undoubtedly separated from prolate particles. In contrast to weather radars, cloud radars are often operated in modes with scanning in azimuth and elevation directions (Kollias et al., 2014; Lamer et al., 2014; Ewald et al., 2015). Matrosov et al. (2012) showed observational evidence that elevation dependencies of polarimetric variables can be efficiently used to discriminate between different types of ice particles.

In a previous study, Myagkov et al. (2015a) combined the two mentioned approaches into an algorithm for a quantitative characterization of shapes and orientations of ice particles based on polarimetric observations from a newly developed 35 GHz cloud radar with hybrid polarimetric configuration. This mode is widely used in polarimetric weather radars but is rarely implemented in cloud radars. In hybrid mode radars measure the differential reflectivity  $Z_{DR}$  and the correlation coefficient  $\rho_{HV}$ . Ice particles of different types have distinct polarimetric scattering signatures that are utilized in the retrieval. For instance, differences in modeled polarimetric variables for oblate, prolate, and spherical solid ice particles are shown in Fig. 1. The algorithm contains two steps. First, using elevation dependencies of  $Z_{DR}$  and  $\rho_{HV}$  ice particles are classified as either oblate or prolate. Second, at a certain altitude polarizability ratio  $\rho_e$  and degree of orientation  $\rho_a$  are estimated from  $Z_{DR}$  and  $\rho_{HV}$  for every elevation angle in range from 30 to 60°. Note, that 90° elevation corresponds to the zenith direction. For every altitude mean and standard deviation of  $\rho_e$  and  $\rho_a$  are calculated. The polarizability ratio  $\rho_e$  depends on shapes and dielectric properties of scatterers, while the degree of orienta-

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion  $\rho_a$  characterizes the width of the canting angle distribution. It is assumed that the mean orientation of ice particles is horizontal. The algorithm was applied to a complex cloud system observed during the ACCEPT (Analysis of the Composition of Clouds with Extended Polarization Techniques) field campaign in Cabauw, the Netherlands.

5 Even though the results of the retrieval do not contradict other studies, the additional validation of the retrieved parameters would be of benefit. Note, that in this paper we only analyze polarizability ratios  $\rho_e$ .

During the ACCEPT campaign the performance of the hybrid-mode cloud radar was evaluated, but in-situ observations were not available. Nevertheless a number of active and passive remote sensors that were deployed during the campaign can be used for the determination of ambient conditions under which ice crystals were formed. Within this study, we use such information to indirectly validate the retrieval presented in Myagkov et al. (2015a) by comparing the obtained microphysical properties of ice crystals with those grown in a wind tunnel under laboratory conditions, and present an analysis of 22 case studies from the ACCEPT campaign.

The paper is organized as follows. Section 2 describes the instruments used for this study and the data set. Five case studies are presented in detail in Sect. 3. The results of the analysis of the 22 cases and their comparison with laboratory studies are presented in Sect. 4. Summary and conclusion are given in Sect. 5.

## 2 Instrumentation and data set

The ACCEPT measurement campaign was initiated by the Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany, the Technical University of Delft, the Netherlands, and METEK GmbH, Elmshorn, Germany within the ITaRS (Initial Training for atmospheric Remote Sensing) project. The major goal of the campaign was to evaluate the polarimetric capabilities of a newly developed polarimetric cloud radar of type MIRA-35 and to estimate the potential of its implementation into existing observational stations. The cloud radar had the hybrid polarimetric configuration. A detailed descrip-

tion can be found in Myagkov et al. (2015a). In the following, we denote this radar as the hybrid mode MIRA-35.

The campaign took place from 7 October to 17 November 2014 at the Cabauw Experimental Site for Atmospheric Research (CESAR), located in the Netherlands (51.971° N, 4.927° E). The CESAR observatory operated by Royal Netherlands Meteorological Institute (KNMI) is well equipped with a broad range of atmospheric remote-sensing instruments. In addition to the instruments available at CESAR and the hybrid mode MIRA-35, which was rented from METEK GmbH, the main instruments of Leipzig Aerosol and Cloud Remote Observations System (LACROS; Bühl et al., 2013) were brought to Cabauw. In Table 1 the instruments that were used for this particular study are presented.

As mentioned before, the shape retrieval of Myagkov et al. (2015a) requires information about the elevation dependencies of the polarimetric variables differential reflectivity  $Z_{DR}$  and correlation coefficient  $\rho_{HV}$ . Therefore, we installed the hybrid mode MIRA-35 into the scanning unit of the LACROS container. An implemented scanning cycle took 15 min and included two elevation (range-height-indicator, RHI) scans from 30 to 150°. Note, that 90° elevation corresponds to the zenith direction. The elevation scans were performed perpendicular in azimuthal direction with  $0.5^\circ \text{ s}^{-1}$  angular velocity. For an accurate determination of the polarimetric variables it is required to correct for antenna coupling effects. Within every scanning cycle the radar was pointed vertically for several minutes. This allowed us to use occasional light rain events for calibration of polarimetric variables as explained in Myagkov et al. (2015a, b).

During the campaign a second cloud radar MIRA-35 was operated as well. The radar is owned by TROPOS and has the conventional linear depolarization configuration. Further this radar is denoted as LDR MIRA-35. The radar was unmounted from the scanning unit of the LACROS container and installed into a trailer without scanning unit and, thus, was pointed vertically. In order to avoid interference between the two cloud radars, their operation frequencies were set to differ by approximately 300 MHz (35.17 GHz vs 35.5 GHz for hybrid-mode and LDR-mode MIRA-35, respectively). Ad-

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ditionally, the trailer was placed about 30 m away from the LACROS container to avoid any near-field interference. In this study, we use data from LDR MIRA-35 to estimate the temporal and spatial dimensions of the observed cloud systems.

When both liquid water droplets and ice crystals are present in a volume, cloud radar alone can hardly detect the liquid-water signatures. It is, however, well known that polarimetric lidars are powerful tools for the detection of supercooled liquid particles within mixed-phase clouds (Schotland et al., 1971; Seifert et al., 2010). We employed the multiwavelength Raman lidar Polly<sup>XT</sup> for this purpose. The lidar was set up near the Cabauw meteorological tower which is located about 300 m north from the measurement site where most of the other instruments were operated. In order to avoid specular reflection from horizontally aligned planar surfaces of ice crystals the laser beam of Polly<sup>XT</sup> was pointed to 5° off-zenith (85° elevation). The container with the lidar was oriented in such a way that the beam was above the radar site at about 4 km height.

Temperature is the main parameter controlling the efficiency of heterogeneous nucleation of ice crystals (DeMott et al., 2015) and it is lowest at cloud top. Therefore, in this study cloud top temperature is used as the reference parameter when crystal properties are investigated. In order to retrieve the temperature at the cloud tops, we used, in order of priority, either locally launched radiosondes, the microwave radiometer HATPRO, or assimilated meteorological datasets. Radiosondes of type Vaisala RS-92 were occasionally launched on-site. If no local radiosonde information was available, radiosonde data from the 00:00 UTC launch at De Bilt (WMO code: 06260; 20 km north-east of CESAR site) was used in case the cloud was observed around 00:00 UTC. If no recent radiosonde ascent was available, temperature data of HATPRO was used in non-precipitating conditions. Finally, for precipitating cases, when the operation of the microwave radiometer is hampered, we used temperature profiles from the GDAS1 dataset (available at: <http://ready.arl.noaa.gov/gdas1.php>) provided by the Global Data Assimilation System (GDAS, Kanamitsu, 1989) operated by the U.S. National Center for Environmental Prediction (NCEP).

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion









## 3.2 Case 2

An analysis of a mid-level mixed-phase cloud observed on 18 October 2014, 01:00–02:00 UTC is given in Fig. 3. The time period chosen for the retrieval is 01:23–01:27 UTC. The cloud top estimated from equivalent radar reflectivity measured by LDR MIRA-35 (Fig. 3a) was at around 5 km height. The thickness of the cloud layer exceeded 1.2 km. Reflectivity values reached values as high as 10 dBZ that indicate the presence of large ice particles. LDR values registered by LDR MIRA-35 for the analyzed period were mostly low even though areas with increased LDR (up to  $-17$  dB) can be seen occasionally at 4–4.5 km height. At the top of the cloud a liquid layer characterized by high attenuated backscatter coefficients and low volume depolarization ratios (Fig. 3c and d, respectively) is visible. It should be noted, that in the second half of the analyzed period the lidar also detected an internal liquid layer at 4.5 km height. In contrast to case 1, where low volume depolarization ratios were observed with Polly<sup>XT</sup>, the ice virgae observed in the case 2 produced volume depolarization ratios exceeding 0.5.

Elevation scans of the differential reflectivity and the correlation coefficient depicted in Fig. 3e and f, respectively, show that the cloud was spatially inhomogeneous. Within the right-half scan only slight angular changes in  $Z_{DR}$  and  $\rho_{HV}$  are visible. In the left-half scan high  $Z_{DR}$  and low  $\rho_{HV}$  values were observed at lower elevations at 4.5 km height at which the lidar detected the internal liquid layer. For the correction of the polarization coupling we used vertical measurements in a short precipitation event on 18 October 2014, 2:30–2:50 UTC. The results of the shape retrieval are shown in Fig. 3g and h. It can be seen that due to the spatial inhomogeneity the calculated polarizability ratios are available only for a limited number of range bins. Nevertheless, the results show that in the left-half scan prolate particles characterized by the polarizability ratio of  $1.52 \pm 0.2$  were detected. In the right-half scan the polarizability ratios closest to the cloud top were  $1.07 \pm 0.1$ . Such values correspond to particles with quasi-spherical shapes and/or low apparent ice density. Further we denote such particles as quasi-isotropic as they

# AMTD

doi:10.5194/amt-2015-365

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





trieved polarizability ratios are slightly higher in the right half-scan which is caused by the observed increased values of  $Z_{DR}$ . The polarizability ratios closest to the cloud top are  $1.5 \pm 0.16$ . The distance from the cloud top is about 240 m. A temperature profile retrieved from the microwave radiometer HATPRO indicated that the temperature at the cloud top was  $-6.1^\circ\text{C}$  (see Fig. 4i).

### 3.4 Case 4

A complex mixed-phase cloud system observed on 10 November 2014, 02:00–03:00 UTC is presented in Fig. 5. A cloud layer with the cloud top at around 5 km height in the time period 02:53–02:57 UTC is considered for the analysis. The cloud layer was more than 2 km thick. The equivalent radar reflectivity at the cloud top did not exceed  $-10$  dBZ while it reached values up to 10 dBZ towards the cloud bottom (Fig. 5a). LDR values measured by LDR MIRA-35 were about  $-30$  dB (Fig. 5b). In Fig. 5c it can be seen that the laser beam often could not penetrate through the whole cloud layer because of strong attenuation. Nevertheless, some indications of liquid water at the cloud top are present. For example, two areas at 5.4 km height characterized by the low volume depolarization ratio can be identified. There is also a thick internal liquid layer at about 4 km height specified by the increased attenuated backscatter coefficient and the low volume depolarization ratio.

Figure 5e and f shows almost no angular dependencies of  $Z_{DR}$  and  $\rho_{HV}$ . Some slight changes are visible at 4 km height where the lidar detected the internal liquid layer. For the correction of the antenna coupling we used observations during a short precipitation event on 1 November 2014 17:00–18:00 UTC. The retrieved polarizability ratios were close to 1 characterizing quasi-isotropic particles. The retrieved profiles indicated rapid changes of the polarizability ratio from 0.92 to 1.05, e.g., from 4.5 to 4.8 km height in the left-half scan. Such changes result from misclassification of prolate and oblate particles which is caused by a variability in  $Z_{DR}$  and  $\rho_{HV}$  due to measurement noise and/or differences in scattering properties of ice populations. Biases in polarimetric variables caused by the polarization coupling, also lead to inaccurate classification.

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Without the polarization coupling correction given by (Myagkov et al., 2015b) the misclassification for the used radar would occur in the range of polarizability ratios from 0.8 to 1.2. For the following analysis we chose the polarizability ratio of  $0.92 \pm 0.08$  observed 450 m below the cloud top. The temperature measured at 5.5 km height by the radiosonde launched at midnight from De Bilt was about  $-20^\circ\text{C}$ .

### 3.5 Case 5

Figure 6 depicts a precipitating cloud system with the cloud top located at around 2.3 km height which was observed on 7 November 2014, 20:00–21:00 UTC. The SNR in this case was not high enough to retrieve polarizability ratios, although elevation dependencies of the measured polarimetric variables allow us to classify the general shape of the observed ice particles. We analyzed two scans performed by the hybrid MIRA-35 which correspond to the time periods shown in Fig. 6a enframed by black rectangles. The time gap between these periods is 11 min. It can be seen that values of  $Z_e$  in the first period do not exceed  $-15$  dBZ. Ice formation was not intensive enough to produce precipitation reaching the ground. Corresponding ice particles caused low depolarization which is indicated by low LDR values of around  $-30$  dB shown in Fig. 6b. In contrast, ice formation in the second time period was much more intensive.  $Z_e$  values close to the top of the cloud were around  $-20$  dBZ, while those observed 1 km below the cloud top exceeded 10 dBZ. Ice particles were large enough to produce liquid precipitation at the ground with 10 dBZ equivalent radar reflectivity. Ice particles in the second period were characterized by LDR values of up to  $-15$  dB. The melting layer with LDR of  $-12$  dB is clearly seen at around 1.4 km height.

In Fig. 6c and d range-height cross sections of SNR for the first and second time period are given, respectively. In both cases SNR are of the same order of magnitude. Elevation scans of differential reflectivity shown in Fig. 6e and f yield  $Z_{DR}$  values close to 0 dB in the zenith-pointing direction, while at lower elevations  $Z_{DR}$  reached 4 and 2 dB, respectively. For both cases  $Z_{DR}$  had less pronounced angular dependence at 1.5 km height. This effect can be caused by aggregation as particles become more spherical

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and/or less dense. Angular dependencies of  $\rho_{HV}$  at the cloud tops show a different behavior. In Fig. 6g  $\rho_{HV}$  has the highest value in the zenith-pointing direction and slightly decreases at lower elevations. For the second time period  $\rho_{HV}$  has its minimum value of about 0.93 at vertical pointing direction and increases up to 0.98 at lower elevations.

5 Observed elevation dependencies at the cloud tops indicate the presence of oblate and prolate particles for the first and the second time periods, respectively. Unfortunately, we do not have continuous temperature profiles from the microwave radiometer for this case because of precipitation. Temporal resolution of the GDAS model is 3 h and can therefore not be used to capture temperature variations within 15 min. The temperature profile given in Fig. 6i shows the cloud top temperature of  $-4^{\circ}\text{C}$  at 2.3 km height  
10 at 21:00 UTC. In Fig. 6a it can be seen that cloud top altitudes for the analyzed time periods differ by about 200 m. Also vertical variations of LDR (Fig. 6b) indicate changes of the  $0^{\circ}\text{C}$  isotherm. Thus, temperatures at the top could be different by few degrees which can cause crucial differences in the ice particle shape. We want to emphasize  
15 that such strong indications of the presence of oblate particles at such warm temperatures were observed twice during the whole field campaign. In both cases the existence of these particles was registered for not longer than about 5 min.

#### 4 Comparison with laboratory studies

The analysis shown in Sect. 3 was applied to 22 cases. The number of cases corresponding to cloud top temperature ranges from  $-7$  to  $-3$ ,  $-13$  to  $-7$ ,  $-17$  to  $-13$ ,  
20 and  $-25$  to  $-17^{\circ}\text{C}$  are 9, 2, 9, and 2, respectively. Most of the analyzed clouds had a liquid layer at the top which was identified using the polarimetric measurements from the lidar. In the case of precipitating clouds we rely on the conclusion of Westbrook and Illingworth (2011) and De Boer et al. (2011) that 95 % of ice crystals at temperatures warmer than  $-25^{\circ}\text{C}$  are formed in presence of liquid water. For every case we  
25 chose the polarizability ratio closest to the cloud top detected by the LDR MIRA-35. The distance from the cloud top mostly does not exceed 500 m.



It is known that the dielectric constant of ice is almost linearly dependent on apparent ice density (Oguchi, 1983). In this study, we assume the following dependence:

$$\epsilon = 2.36\rho_a + 1, \quad (3)$$

where  $\rho_a$  is in  $\text{g cm}^{-3}$ . We used the analytical spheroidal scattering model and information about geometrical axis ratios and dielectric constants of ice crystals grown in the fall chamber to obtain their polarizability ratios  $\rho_e$  (see Eqs. 39–44 in Myagkov et al., 2015a). For the polarimetric scattering model we assume that spheroids have the same volume and geometrical axis ratio as the hexagonal prisms used for the calculation of the apparent ice density in Eq. (2). In Fig. 8 values of  $\rho_e$  retrieved from the laboratory studies are shown as blue filled circles. The highest  $\rho_e$  values of about 1.6–1.7 were observed in the temperature range from  $-7$  to  $-6^\circ\text{C}$ . Even though prolate particles with the highest geometrical axis ratio were formed at  $-5^\circ\text{C}$ , they had low apparent ice density and therefore their  $\rho_e$  did not exceed 1.4. Polarizability ratios of 0.4 correspond to ice crystals that formed at temperatures of  $-12$  and  $-16^\circ\text{C}$ , where ice crystals were found to cause apparent ice densities exceeding  $0.8 \text{ g cm}^{-3}$ . Dendrites that form at  $-15^\circ\text{C}$  had low  $\rho_a$  that led to values of  $\rho_e$  of 0.6. In Fig. 8 we also show polarizability ratios retrieved from the polarimetric observations of the hybrid MIRA-35 during the ACCEPT campaign (red filled dots with error bars). A good agreement between findings from the free-fall chamber and remote observations can be seen. In the temperature range of  $-6$  to  $-4^\circ\text{C}$  values of  $\rho_e$  retrieved close to cloud tops varied from 1.4–1.8. At temperatures of  $-9$  to  $-7^\circ\text{C}$  isometric particles were detected with  $\rho_e$  of 0.8–1.2. In the temperature range of  $-17$  to  $-13^\circ\text{C}$  observed oblate particles mostly had  $\rho_e$  of 0.4–0.6. At temperatures from  $-25$  to  $-20^\circ\text{C}$  ice crystals had  $\rho_e$  of 0.8–1. At the same time differences in measured and calculated polarizability ratios can be seen in Fig. 8 at temperatures of  $-4$  to  $-3^\circ\text{C}$ . Such differences can be caused by uncertainties in temperature values from the GDAS1 data set. Even though the number of cases available from 6 weeks of measurements is quite low, we show that ice particles formed close to the top of mid-level mixed-phase clouds at temperatures warmer than

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



~ -25°C in general have a similar dependence of shape and apparent ice density on ambient temperature as the ones grown in the free-fall chamber.

## 5 Summary and outlook

Polarimetric cloud radars have a great potential for the provision of valuable information about ice crystal microphysical properties based on remote sensing. Within this study it was demonstrated, that the scanning capabilities of modern cloud radars allow for the classification and quantitative characterization of ice particle shape. For this purpose a 35 GHz cloud radar MIRA-35 with the hybrid polarimetric configuration was deployed during the ACCEPT measurement campaign in Cabauw, the Netherlands in October and November of 2014. The radar provided elevation dependencies of the differential reflectivity and the correlation coefficient which were used to estimate the polarizability ratios of ice crystals approximated by spheroids. The polarizability ratio thus depends on the geometrical axis ratio and the apparent ice density. The radar was collocated with a vast number of active and passive remote sensors providing continuous information about cloud geometry, ambient conditions, and the presence of supercooled liquid layers. The combined analysis of the available data allowed us to derive temperature-dependent polarizability ratios of ice crystals in the cloud-top region of mixed-phase cloud layers. During the ACCEPT campaign in-situ observations were not available. Therefore, we used data about ice crystals grown in a free-fall cloud chamber to validate our retrieval indirectly. The measurements available from laboratory studies include accurate information about axis lengths and mass of ice crystals grown at water saturation conditions. We utilized these parameters to calculate polarizability ratios from the laboratory measurements. A comparison of polarizability ratios of ice crystals investigated in the cloud chamber and the ones observed close to the cloud tops showed a good agreement. At temperatures in the range of -6 to -4°C columnar-shaped particles with  $\rho_e$  of 1.2–1.7 and 1.4–1.8 were found in laboratory studies and remote observations, respectively. Isometric particles occurred at temper-

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





gramme (FP7/2007-2013) under grant agreement no. 262254. Authors want to acknowledge METEK GmbH, particularly Matthias Bauer-Pfundstein and Alexander Partus for their assistance in maintenance of cloud radars during the ACCEPT campaign. We also highly appreciate T. Takahashi, Hokkaido University of Education, Sapporo, Japan, for providing the data from the free-fall cloud chamber. The authors also would like to acknowledge the logistic support received from KNMI and TU Delft.

## References

- Althausen, D., Engelmann, R., Baars, H., Heese, B., Ansmann, A., Müller, D., and Komppula, M.: Portable Raman lidar PollyXT for automated profiling of aerosol backscatter, extinction, and depolarization, *J. Atmos. Ocean. Tech.*, 26, 2366, doi:10.1175/2009JTECHA1304.1, 2009. 28
- Ansmann, A., Tesche, M., Seifert, P., Althausen, D., Engelmann, R., Fruntke, J., Wandinger, U., Mattis, I., and Müller, D.: Evolution of the ice phase in tropical altocumulus: SAMUM lidar observations over Cape Verde, *J. Geophys. Res.*, 114, D17208, doi:10.1029/2008JD011659, 2009. 3
- Bailey, M. and Hallett, J.: Growth rates and habits of ice crystals between  $-20^{\circ}\text{C}$  and  $-70^{\circ}\text{C}$ , *J. Atmos. Sci.*, 61, 514–544, doi:10.1175/1520-0469(2004)061<0514:GRAHOI>2.0.CO;2, 2004. 9
- Bailey, M. P. and Hallett, J.: A comprehensive habit diagram for atmospheric ice crystals: confirmation from the laboratory, AIRS II, and other field studies, *J. Atmos. Sci.*, 66, 2888, doi:10.1175/2009JAS2883.1, 2009. 3, 9
- Bouniol, D., Protat, A., Delanoë, J., Pelon, J., Piriou, J.-M., Bouyssel, F., Tompkins, A. M., Wilson, D. R., Morille, Y., Haeffelin, M., O'Connor, E. J., Hogan, R. J., Illingworth, A. J., Donovan, D. P., and Baltink, H.-K.: Using continuous ground-based radar and lidar measurements for evaluating the representation of clouds in four operational models, *J. Appl. Meteorol. Clim.*, 49, 1971–1991, doi:10.1175/2010JAMC2333.1, 2010. 3
- Bühl, J., Seifert, P., Wandinger, U., Baars, H., Kanitz, T., Schmidt, J., Myagkov, A., Engelmann, R., Skupin, A., Heese, B., Klepel, A., Althausen, D. and Ansmann, A.: LACROS: the Leipzig Aerosol and Cloud Remote Observations System, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 8890, 889002, doi:10.1117/12.2030911, 2013. 7

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bühl, J., Seifert, P., Myagkov, A., and Ansmann, A.: Closing the ice-nucleus life cycle with active remote sensing: analysis of mixed-phase cloud layers in a Cloudnet dataset, to be submitted to AMTD, 2016. 4, 20

De Boer, G., Eloranta, E. W., and Shupe, M. D.: Arctic mixed-phase stratiform cloud properties from multiple years of surface-based measurements at two high-latitude locations, *J. Atmos. Sci.*, 66, 2874, doi:10.1175/2009JAS3029.1, 2009. 3

De Boer, G., Morrison, H., Shupe, M. D., and Hildner, R.: Evidence of liquid dependent ice nucleation in high-latitude stratiform clouds from surface remote sensors, *Geophys. Res. Lett.*, 38, L01803, doi:10.1029/2010GL046016, 2011. 4, 16

Delanoë, J. and Hogan, R. J.: Combined CloudSat-CALIPSO-MODIS retrievals of the properties of ice clouds, *J. Geophys. Res.*, 115, D00H29, doi:10.1029/2009JD012346, 2010. 3

Delanoë, J., Hogan, R. J., Forbes, R. M., Bodas-Salcedo, A., and Stein, T. H. M.: Evaluation of ice cloud representation in the ECMWF and UK Met Office models using CloudSat and CALIPSO data, *Q. J. Roy. Meteor. Soc.*, 137, 2064–2078, doi:10.1002/qj.882, 2011. 3

Delanoë, J. M. E., Heymsfield, A. J., Protat, A., Bansemer, A., and Hogan, R. J.: Normalized particle size distribution for remote sensing application, *J. Geophys. Res.-Atmos.*, 119, 4204–4227, doi:10.1002/2013JD020700, 2014. 3

DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T. and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *P. Natl. Acad. Sci. USA*, 107, 11217–11222, doi:10.1073/pnas.0910818107, 2010. 3

DeMott, P. J., Prenni, A. J., McMeeking, G. R., Sullivan, R. C., Petters, M. D., Tobo, Y., Niemand, M., Möhler, O., Snider, J. R., Wang, Z., and Kreidenweis, S. M.: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles, *Atmos. Chem. Phys.*, 15, 393–409, doi:10.5194/acp-15-393-2015, 2015. 8

Ewald, F., Winkler, C., and Zinner, T.: Reconstruction of cloud geometry using a scanning cloud radar, *Atmos. Meas. Tech.*, 8, 2491–2508, doi:10.5194/amt-8-2491-2015, 2015. 5

Fleishauer, R. P., Larson, V. E., and Vonder Haar, T. H.: Observed microphysical structure of midlevel, mixed-phase clouds, *J. Atmos. Sci.*, 59, 1779–1804, doi:10.1175/1520-0469(2002)059<1779:OMSOMM>2.0.CO;2, 2002. 3

Fukuta, N.: Experimental studies on the growth of small ice crystals, *J. Atmos. Sci.*, 26, 522–531, doi:10.1175/1520-0469(1969)026<0522:ESOTGO>2.0.CO;2, 1969. 4

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fukuta, N. and Takahashi, T.: The growth of atmospheric ice crystals: a summary of findings in vertical supercooled cloud tunnel studies, *J. Atmos. Sci.*, 56, 1963–1979, doi:10.1175/1520-0469(1999)056<1963:TGOAIC>2.0.CO;2, 1999. 3, 4
- Görsdorf, U., Lehmann, V., Bauer-Pfundstein, M., Peters, G., Vavriv, D., Vinogradov, V., and Volkov, V.: A 35 GHz polarimetric Doppler radar for long term observations of cloud parameters – description of system and data processing, *J. Atmos. Ocean. Tech.*, 32, 675–690, doi:10.1175/JTECH-D-14-00066.1, 2015. 28
- Hallett, J.: Production of secondary ice particles during the riming process, *Nature*, 249, 26–28, doi:10.1038/249026a0, 1974. 3
- Hogan, R. J., Francis, P. N., Flentje, H., Illingworth, A. J., Quante, M., and Pelon, J.: Characteristics of mixed-phase clouds. I: Lidar, radar and aircraft observations from CLARE'98, *Q. J. Roy. Meteor. Soc.*, 129, 2089–2116, doi:10.1256/rj.01.208, 2003. 3
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817–9854, doi:10.5194/acp-12-9817-2012, 2012. 3
- Kampe, H. J., Weickmann, H. K., and Kelly, J. J.: The influence of temperature on the shape of ice crystals growing at water saturation, *J. Atmos. Sci.*, 8, 168–174, doi:10.1175/1520-0469(1951)008<0168:TLOTOT>2.0.CO;2, 1951. 17
- Kanamitsu, M.: Description of the NMC global data assimilation and forecast system, *Weather Forecast.*, 4, 335–342, doi:10.1175/1520-0434(1989)004<0335:DOTNGD>2.0.CO;2, 1989. 8
- Kanitz, T., Seifert, P., Ansmann, A., Engelmann, R., Althausen, D., Casiccia, C., and Rohwer, E. G.: Contrasting the impact of aerosols at northern and southern midlatitudes on heterogeneous ice formation, *Geophys. Res. Lett.*, 38, L17802, doi:10.1029/2011GL048532, 2011. 3
- Klein, S. A., McCoy, R. B., Morrison, H., Ackerman, A. S., Avramov, A., Boer, G. D., Chen, M., Cole, J. N. S., Del Genio, A. D., Falk, M., Foster, M. J., Fridlind, A., Golaz, J.-C., Hashino, T., Harrington, J. Y., Hoose, C., Khairoutdinov, M. F., Larson, V. E., Liu, X., Luo, Y., McFarquhar, G. M., Menon, S., Neggers, R. A. J., Park, S., Poellot, M. R., Schmidt, J. M., Sednev, I., Shipway, B. J., Shupe, M. D., Spangenberg, D. A., Sud, Y. C., Turner, D. D., Veron, D. E., Salzen, K. V., Walker, G. K., Wang, Z., Wolf, A. B., Xie, S., Xu, K.-M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- mixed-phase arctic cloud experiment. I: single-layer cloud, Q. J. Roy. Meteor. Soc., 135, 979–1002, doi:10.1002/qj.416, 2009. 3
- Kollias, P., Bharadwaj, N., Widener, K., Jo, I., and Johnson, K.: Scanning ARM cloud radars. Part I: Operational sampling strategies, J. Atmos. Ocean. Tech., 31, 569–582, doi:10.1175/JTECH-D-13-00044.1, 2014. 5
- Lamer, K., Tatarevic, A., Jo, I., and Kollias, P.: Evaluation of gridded scanning ARM cloud radar reflectivity observations and vertical doppler velocity retrievals, Atmos. Meas. Tech., 7, 1089–1103, doi:10.5194/amt-7-1089-2014, 2014. 5
- Luo, Y., Xu, K.-M., Morrison, H., McFarquhar, G. M., Wang, Z., and Zhang, G.: Multi-layer arctic mixed-phase clouds simulated by a cloud-resolving model: comparison with ARM observations and sensitivity experiments, J. Geophys. Res., 113, D12208, doi:10.1029/2007JD009563, 2008. 3
- Matrosov, S. Y., Reinking, R. F., Kropfli, R. A., Martner, B. E., and Bartram, B. W.: On the use of radar depolarization ratios for estimating shapes of ice hydrometeors in winter clouds, J. Appl. Meteorol., 40, 479–490, doi:10.1175/1520-0450(2001)040<0479:OTUORD>2.0.CO;2, 2001. 4
- Matrosov, S. Y., Mace, G. G., Marchand, R., Shupe, M. D., Hallar, A. G., and McCubbin, I. B.: Observations of ice crystal habits with a scanning polarimetric W-band radar at slant linear depolarization ratio mode, J. Atmos. Ocean. Tech., 29, 989–1008, doi:10.1175/JTECH-D-11-00131.1, 2012. 5
- Melnikov, V. and Straka, J. M.: Axis ratios and flutter angles of cloud ice particles: retrievals from radar data, J. Atmos. Ocean. Tech., 30, 1691–1703, doi:10.1175/JTECH-D-12-00212.1, 2013. 4
- Morrison, H., McCoy, R. B., Klein, S. A., Xie, S., Luo, Y., Avramov, A., Chen, M., Cole, J. N. S., Falk, M., Foster, M. J., Del Genio, A. D., Harrington, J. Y., Hoose, C., Khairoutdinov, M. F., Larson, V. E., Liu, X., McFarquhar, G. M., Poellot, M. R., von Salzen, K., Shipway, B. J., Shupe, M. D., Sud, Y. C., Turner, D. D., Veron, D. E., Walker, G. K., Wang, Z., Wolf, A. B., Xu, K.-M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM mixed-phase arctic cloud experiment. II: Multilayer cloud, Q. J. Roy. Meteor. Soc., 135, 1003–1019, doi:10.1002/qj.415, 2009. 3
- Mülmenstädt, J., Sourdeval, O., Delanoë, J., and Quaas, J.: Frequency of occurrence of rain from liquid-, mixed-, and ice-phase clouds derived from A-Train satellite retrievals, Geophys. Res. Lett., 42, 6502–6509, doi:10.1002/2015GL064604, 2015. 2

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Myagkov, A., Seifert, P., Bauer-Pfundstein, M., and Wandinger, U.: Cloud radar with hybrid mode towards estimation of shape and orientation of ice crystals, *Atmos. Meas. Tech. Discuss.*, 8, 9105–9163, doi:10.5194/amt-d-8-9105-2015, 2015a. 5, 6, 7, 9, 10, 13, 18, 28

Myagkov, A., Seifert, P., Wandinger, U., Bauer-Pfundstein, M., and Matrosov, S. Y.: Effects of antenna patterns on cloud radar polarimetric measurements, *J. Atmos. Ocean. Tech.*, 32, 1813–1828, doi:10.1175/JTECH-D-15-0045.1, 2015b. 7, 15, 29

Oguchi, T.: Electromagnetic wave propagation and scattering in rain and other hydrometeors, *IEEE Proceedings*, 71, 1029–1078, 1983. 5, 18

Oue, M., Kumjian, M. R., Lu, Y., Verlinde, J., Aydin, K., and Clothiaux, E. E.: Linear depolarization ratios of columnar ice crystals in a deep precipitating system over the Arctic observed by zenith-pointing Ka-band Doppler radar, *J. Appl. Meteorol. Clim.*, 54, 1060–1068, doi:10.1175/JAMC-D-15-0012.1, 2015. 4, 13, 20

Rangno, A. L. and Hobbs, P. V.: Microstructures and precipitation development in cumulus and small cumulonimbus clouds over the warm pool of the tropical Pacific Ocean, *Q. J. Roy. Meteor. Soc.*, 131, 639–673, doi:10.1256/qj.04.13, 2005. 3

Rose, T., Crewell, S., Löhnert, U., and Simmer, C.: A network suitable microwave radiometer for operational monitoring of the cloudy atmosphere, *Atmos. Res.*, 75, 183–200, doi:10.1016/j.atmosres.2004.12.005, 2005. 28

Schotland, R. M., Sassen, K., and Stone, R.: Observations by lidar of linear depolarization ratios for hydrometeors, *J. Appl. Meteorol.*, 10, 1011–1017, doi:10.1175/1520-0450(1971)010<1011:OBLOLD>2.0.CO;2, 1971. 8

Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteorol. Atmos. Phys.*, 92, 45–66, doi:10.1007/s00703-005-0112-4, 2006. 3

Seifert, P., Ansmann, A., Mattis, I., Wandinger, U., Tesche, M., Engelmann, R., Müller, D., Pérez, C., and Haustein, K.: Saharan dust and heterogeneous ice formation: eleven years of cloud observations at a central European EARLINET site, *J. Geophys. Res.*, 115, D20201, doi:10.1029/2009JD013222, 2010. 8

Seifert, P., Kunz, C., Baars, H., Ansmann, A., Bühl, J., Senf, F., Engelmann, R., Althausen, D., and Artaxo, P.: Seasonal variability of heterogeneous ice formation in stratiform clouds over the Amazon Basin, *Geophys. Res. Lett.*, 42, 5587–5593, doi:10.1002/2015GL064068, 2015.

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Smith, A. J., Larson, V. E., Niu, J., Kankiewicz, J. A., and Carey, L. D.: Processes that generate and deplete liquid water and snow in thin midlevel mixed-phase clouds, *J. Geophys. Res.*, 114, D12203, doi:10.1029/2008JD011531, 2009. 3

Straka, J. M., Zrni, D. S., and Ryzhkov, A. V.: Bulk hydrometeor classification and quantification using polarimetric radar data: synthesis of relations, *J. Appl. Meteorol.*, 39, 1341–1372, doi:10.1175/1520-0450(2000)039<1341:BHCAQU>2.0.CO;2, 2000. 4

Sun, Z. and Shine, K. P.: Parameterization of ice cloud radiative properties and its application to the potential climatic importance of mixed-phase clouds, *J. Climate*, 8, 1874–1888, doi:10.1175/1520-0442(1995)008<1874:POICRP>2.0.CO;2, 1995. 2

Suortti, T. M., Kats, A., Kivi, R., Kämpfer, N., Leiterer, U., Miloshevich, L. M., Neuber, R., Paukkunen, A., Ruppert, P., Vömel, H., and Yushkov, V.: Tropospheric comparisons of vaisala radiosondes and balloon-borne frost-point and Lyman- $\alpha$  hygrometers during the LAUTLOS-WAVVAP experiment, *J. Atmos. Ocean. Tech.*, 25, 149, doi:10.1175/2007JTECHA887.1, 2008. 28

Takahashi, T.: Influence of liquid water content and temperature on the form and growth of branched planar snow crystals in a cloud, *J. Atmos. Sci.*, 71, 4127–4142, doi:10.1175/JAS-D-14-0043.1, 2014. 4, 17

Takahashi, T. and Fukuta, N.: Supercooled cloud tunnel studies on the growth of snow crystals between  $-4^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$ , *J. Meteorol. Soc. Jpn.*, 66, 841–855, 1988. 17

Takahashi, T., Endoh, T., Wakahama, G., and Fukuta, N.: Vapor diffusional growth of free-falling snow crystals between  $-3$  and  $-23^{\circ}\text{C}$ , *J. Meteorol. Soc. Jpn.*, 69, 15–30, 1991. 4, 17, 36, 37

Tao, W.-K. and Moncrieff, M. W.: Multiscale cloud system modeling, *Rev. Geophys.*, 47, RG4002, doi:10.1029/2008RG000276, 2009. 4

Wang, Z., Sassen, K., Whiteman, D. N., and Demoz, B. B.: Studying altocumulus with ice virga using ground-based active and passive remote sensors, *J. Appl. Meteorol.*, 43, 449–460, doi:10.1175/1520-0450(2004)043<0449:SAWIVU>2.0.CO;2, 2004. 3

Westbrook, C. D. and Heymsfield, A. J.: Ice crystals growing from vapor in supercooled clouds between  $-2.5^{\circ}\text{C}$  and  $-22^{\circ}\text{C}$ : testing current parameterization methods using laboratory data, *J. Atmos. Sci.*, 68, 2416–2429, doi:10.1175/JAS-D-11-017.1, 2011. 3

Westbrook, C. D. and Illingworth, A. J.: Evidence that ice forms primarily in supercooled liquid clouds at temperatures  $-27^{\circ}\text{C}$ , *Geophys. Res. Lett.*, 38, L14808, doi:10.1029/2011GL048021, 2011. 4, 16

- Westbrook, C. D. and Illingworth, A. J.: The formation of ice in a long-lived supercooled layer cloud, Q. J. Roy. Meteor. Soc., 139, 2209–2221, doi:10.1002/qj.2096, 2013. 3
- Zhang, D., Wang, Z., Heymsfield, A., Fan, J., Liu, D., and Zhao, M.: Quantifying the impact of dust on heterogeneous ice generation in midlevel supercooled stratiform clouds, Geophys. Res. Lett., 39, L18805, doi:10.1029/2012GL052831, 2012. 3

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Shape-temperature relationship of pristine ice crystals


A. Myagkov et al.



**Figure 1.** Elevation dependencies of modeled (a)  $Z_{DR}$  and (b)  $\rho_{HV}$  for strongly oblate (axis ratio  $\ll 1$ ), strongly prolate (axis ratio  $\gg 1$ ), and spherical (axis ratio of 1) solid ice particles. Particles are assumed to be oriented near horizontally with the degree of orientation of 0.99. The figure is based on look-up tables given in Myagkov et al. (2015b).

## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.



Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 2.** Case study of ice crystal shapes observed with hybrid-mode MIRA-35 on 12 October 2014 at Cabauw, the Netherlands. Shown are time-height cross sections of the following parameters: **(a)** equivalent radar reflectivity  $Z_e$  from LDR MIRA-35; **(b)** radar linear depolarization ratio from LDR MIRA-35; **(c)** attenuated backscatter coefficient at 1064 nm from Polly-XT; and **(d)** volume linear depolarization ratio at 532 nm from Polly-XT, range-height cross sections of **(e)** differential reflectivity  $Z_{DR}$  and **(f)** correlation coefficient  $\rho_{HV}$  measured by hybrid MIRA-35, **(g, h)** vertical profiles of polarizability ratio for the left- and right-half scans, respectively, **(i)** vertical temperature profile from the microwave radiometer HATPRO, **(j)** photo taken by a web-camera installed at the CESAR site. The red rectangle shows the analyzed cloud layer and the time period corresponding to a full elevation scan of hybrid MIRA-35.  $Z_e$  and LDR were calculated from total powers in the co- and cross-channels.  $Z_{DR}$  and  $\rho_{HV}$  were obtained for the spectral peaks. Note, that  $Z_{DR}$  and  $\rho_{HV}$  are not corrected for the antenna coupling here to make figures more illustrative. Red horizontal lines in **(h, i, j)** represents the cloud top. Vertical profiles in **(h)** and **(i)** indicated by red line correspond to mean values of polarizability ratio. Horizontal blue bars in **(h)** and **(i)** denote 2 standard deviations of polarizability ratio.

## Shape-temperature relationship of pristine ice crystals


A. Myagkov et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.




**Figure 3.** Same as in Fig. 2, but for 18 October 2014 and without a photograph.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.




**Figure 4.** Same as in Fig. 2, but for 20 October 2014 and without a photograph.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.



**Figure 5.** Same as in Fig. 2, but for 10 November 2014 and without a photograph.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Shape-temperature  
relationship of  
pristine ice crystals

A. Myagkov et al.



**Figure 6.** Case study of strong short-term variability of ice particle shapes observed at Cabauw, the Netherlands, on 7 November 2014. Shown are time-height cross sections of **(a)** equivalent radar reflectivity  $Z_e$  from LDR MIRA-35 and **(b)** radar linear depolarization ratio from LDR MIRA-35; range-height cross sections of **(c)** signal-to-noise ratio, **(e)** differential reflectivity  $Z_{DR}$  and **(g)** correlation coefficient  $\rho_{HV}$  measured by hybrid MIRA-35 from 20:31 to 20:35 UTC; **(d)** signal-to-noise ratio, **(f)** differential reflectivity  $Z_{DR}$  and **(h)** correlation coefficient  $\rho_{HV}$  measured by hybrid MIRA-35 from 20:46 to 20:50 UTC, **(i)** vertical temperature profile from the GDAS1 model. Two black rectangles in **(a)** show the analyzed cloud layer and the time periods corresponding to full elevation scans of hybrid MIRA-35.  $Z_e$  and LDR were calculated from total powers in the co- and cross-channels. SNR,  $Z_{DR}$  and  $\rho_{HV}$  were obtained for the spectral peaks. Note, that  $Z_{DR}$  and  $\rho_{HV}$  are not corrected for the antenna coupling here to make figures more illustrative. Red horizontal lines in **(i)** represents the cloud top at 20:45 UTC.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Shape-temperature relationship of pristine ice crystals

A. Myagkov et al.



**Figure 8.** Temperature dependence of polarizability ratios for ice crystals grown in the free-fall chamber (blue filled circles) and for ones located close to tops of mixed-phase clouds, retrieved from hybrid MIRA-35 (red filled circles). Note, that  $\rho_e > 1$  corresponds to prolate particles,  $\rho_e < 1$  corresponds to oblate particles. Vertical red bars represent  $\pm 1$  standard deviation of observed polarizability ratios. Data from the free-fall chamber (Takahashi et al., 1991) were provided by Prof. Takahashi, Hokkaido University of Education, Sapporo, Japan.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)