Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2016-139-RC2, 2016 © Author(s) 2016. CC-BY 3.0 License.



AMTD

Interactive comment

Interactive comment on "Trends of tropical tropospheric ozone from twenty years of European satellite measurements and perspectives for Sentinel-5 Precursor" by Klaus-Peter Heue et al.

Anonymous Referee #2

Received and published: 8 July 2016

Overall this is a good paper on tropospheric ozone trends but it can be cleaned up a bit regarding clarity of sections with editing/typo fixes, etc. I suggest acceptance of the paper with some needed changes but generally minor as listed below:

The most important result from Beig and Singh [2007] is that they were able to detect decadal increases in tropospheric ozone over southern Asia for the TOMS record (1979-2005). In their analysis they were able to evaluate CCD measurements for 30S to 30N. For greater impact is it possible to extend your product to include higher northern latitudes and also evaluate Asia, but now for later time period? Perhaps averaging your measurements annually or seasonally each year when stratospheric ozone variability is small?

Printer-friendly version

Discussion paper



Your trend results are very interesting, in particular Figures 8-9 showing the regional changes. Comparisons with other products show differences with these results, reflecting both differing time periods and also that all these decadal changes are pretty small and difficult to detect considering signal to noise. The increases you find in TTOC in the tropics through 2015 are not inconsistent to large extent with increases in global TTOC that have been reported the last few years in the BAMS State of the Climate Reports.

In the Introduction for Feng and Kobayashi [2009] you have numbers stated of 5% to 20% crop loss due to tropospheric ozone which is very large and seems too much of a broad generalization by them given how the assessments are made under many years of controlled environment experiments with wide range of plant species. Wheat, corn, soybeans, etc. will have different ozone sensitivities and very different than many other plant species which may not have any negative reaction to ozone at all (maybe true for some crops also). It would be more convincing for readers to include some other reference(s), perhaps even more recent if possible, that claims similar numbers for crop loss from ozone.

The discussion of the important role of tropospheric ozone is very short in the Introduction. You might mention more directly regarding emphasizing the importance of tropospheric ozone that it is the third most important radiative greenhouse gas [IPCC, 2014], but also has the good property as being main source of air purifier/oxidizer OH. I don't know how much of the photochemistry discussion with details is really needed in the Introduction, maybe okay, but it seems to jump away then from this detailed photochemistry discussion into retrieval methods and measurements (which is the emphasis of the paper).

The harmonization of the datasets was a bit hard to follow but I am sure all okay in fundamental approach. The time series regression fits that you show in Figure 6 are for tropical averages – but does a similar plot for 12.5 N (or 12.5S) have any indication of instrument offset(s) despite following the multi-instrument harmonization? This might

AMTD

Interactive comment

Printer-friendly version

Discussion paper



in part explain the problems in winter latitudes with high standard deviations in Figure 2?

In discussion of Figure 1 the small < 2 DU column for tropopause to effective cloud pressure is probably generally typical over the remote tropical Pacific over thick clouds. The Ziemke et al. [2009] Figure 8 used MLS ozone profiles to get the SOC and compared 1-1 with ACCO from OMI. Away from the Pacific the ACCO and SOC difference was much greater than 2 DU – hence the need for applying near zonally invariant SOC to get gridded TTOC everywhere. That Figure 8 seemed to summarize why the simple CCD method works as well as it does including assumption of near-zero zonal variability of SOC (from MLS in the figure).

Most of the figures are difficult to make out, especially in terms of text readability. Please increase sizes.

There are several typos and some sentences that can use some re-editing. I found several in the reading, but didn't list them all here (you may have already found them and corrected them):

Page 3, line 6: "...increase (Wang et al., 2009)..." (or reference at end of sentence) Page 14, line 10: "...quasi-biennial..." Page 17, line 7: "increase"

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2016-139, 2016.

AMTD

Interactive comment

Printer-friendly version

Discussion paper

