

Interactive comment on “Intercomparison of total column ozone data from the Pandora spectrophotometer with Dobson, Brewer, and OMI measurements over Seoul, Korea” by Jiyoung Kim et al.

J. Kim

jigang615@yonsei.ac.kr

Received and published: 8 September 2016

General comments: The manuscript presents the results of a two-year comparison of total ozone measurements from Pandora, Brewer and Dobson spectrophotometers and satellite estimates (OMI, two algorithms). Results are given in terms of a least-squares analysis and some statistical indicators are provided (max, min, average, standard deviation, relative differences, Pearson’s correlation index, R2) together with the analysis of variance (ANOVA). The manuscript addresses relevant questions and could potentially provide interesting results, since a complete set of instrument operates at the

Printer-friendly version

Discussion paper



measuring station of Seoul.

ANSWER: We really appreciate the reviewer's all comments and suggestions about this manuscript. They are really helpful, so we can improve the details and quality of our manuscript based on these comments. The following content is the answers for the reviewer's comments and the revised manuscript is attached in supplement

Specific comments: 1. No mention is made in the manuscript of the processing/algorithms of the measurements. Do they differ for different instruments? For example, how much can the used spectroscopic sets of cross-sections (and their relative dependence on effective ozone C2 AMTD Interactive comment Printer-friendly version Discussion paper temperature) impact on the comparison?

ANSWER: General manuscripts of the processing/algorithms of four measurements are mentioned in Section 1. And the specific mention for each instrument in Seoul is summarized in Section 2.

2. Table 2: do the authors have an idea why some instruments have a negative intercept compared to Pandora and others a positive intercept? How do the authors cope with straylight, which affects all data from the presented ground-based instruments?

ANSWER: The intercept is associated with the slope. Actually, the intercept is relatively small when the slope value is large. Moreover, Pandora measurements are relatively small compared to other measurements (underestimation). Therefore, it is possible to find the negative value of the intercept if the difference is somewhat large at the small TCO. However in Table 2, all absolute values of t-ratio are in 0.27~1.14, so it is hard to say that the intercept is significantly negative. As for stray light, it affects all data especially in UV band, and its valid wavelength range is mentioned in Sabburg et al. (2002). In this study, Pandora, Brewer and Dobson data didn't use this wavelength range. Moreover, Dobson measures TCO using 2 pair wavelengths, so the instrument effect is small.

Printer-friendly version

Discussion paper



3. Lines 179-180 explain that measurements from Pandora at high SZAs are removed from the analysis, but what about the other instruments? The comparison is performed in terms of monthly and daily averages. Then, a section (3.3) explains that large inconsistencies may originate due to the ozone variability when daily averages are performed on different datasets (because of different sampling frequencies). Could the authors additionally perform a comparison of nearly simultaneous measurements? It is stated that the Pandora temporal resolution is about 2 minutes, therefore those measurements could be interpolated to the nearest Brewer/Dobson/OMI estimate, thus avoiding the bias illustrated in Sect. 3.3;

ANSWER: Basically, Dobson measures TCO under condition of $\mu < 2.5$ as mentioned in Section 2.1 (noon/close to noon) and Brewer instrument is also set up to $\mu < 3.0$. As mentioned in Section 3.3, large inconsistencies may generate due to the ozone variability, but the main purpose of this study is focused on the comparison of practical Pandora data (daily TCO) with other measurements rather than detailed validation as stated in line 371-374. Section 3.3 is the additional part in order to explain diurnal variation of TCO considered to be the main cause of bias and Pandora data clearly shows this variation due to its high temporal resolution as mentioned in line 321. Validation of real-time data from each instrument leading to minimize bias is not the main purpose.

4. Line 196: if the only condition for the comparison is that the number of daily observations must be greater than 10 days, it is likely that some differences are due to the sampled subset. Since the results orient the following comparison of the instruments, I think that the authors should revise their criteria or state the uncertainty of their results due to the day-to-day variability and different days sampled for each month in the datasets (as done in Sect. 3.3 for the daily averages);

ANSWER: As mentioned above, the main purpose of this study is to compare practical Pandora data. So in spite of some differences due to sampling issue, daily Pandora TCO was calculated and the comparison result actually shows good agreement with Brewer, Dobson and OMI measurements, respectively. Also, monthly TCO values are

[Printer-friendly version](#)[Discussion paper](#)

relatively less affected by daily variation and they are calculated when all data sets are available. The number of daily data is considerably small especially in summer season due to lack of clear days. Considering this, the comparison condition (daily observations > 10 per month) is set and monthly TCO values from all instruments are in good agreement with each other. The result of small mean relative difference stands for small day-to-day variability.

5. The dependence on solar zenith angle is listed as an important factor impacting the comparison (line 365). Could the authors present some plots of the differences between instruments as a function of the solar zenith angle or the airmass? What is the expected magnitude of the SO₂ effect in Seoul (line 365)?

ANSWER: Two graphs in Figure 1 are generated from the comparison of Brewer and Dobson measurements in Seoul, Korea. The left figure shows the comparison of Brewer data and Dobson data after SO₂ correction from 1999-2005 except for year 2001 in Seoul and the right figure shows the same comparison but from January 2007 to March 2007. From both figures, the agreements between the Dobson and Brewer dataset are reasonably good after SO₂ correction with slopes of 0.996 (left) and 1.004 (right) respectively and R² values of 0.986 (left) and 0.985 (right) respectively. Although there are some differences in data sampling, it can be known that slopes and R² values in these figures are closer than those in Figure 4(e).

Technical corrections: 1. Line 5 (and 209): it should be clarified why Pandora is taken as a reference for the comparison. Is it because of its high temporal resolution? In this case, this should be C3 AMTD Interactive comment Printer-friendly version Discussion paper explicitly stated;

ANSWER: The reason Pandora data is taken as a reference for the comparison is that the operation of Pandora instrument was recently started compared to Dobson and Brewer in Seoul. So this study shows the reliability of Pandora data roughly through the result of inter-comparison with other instruments.

[Printer-friendly version](#)[Discussion paper](#)

2. Line 10: reporting the slope and R2 for the comparison with the Brewer is redundant, since these data are already provided few lines above;

ANSWER: The redundant phrase is deleted.

3. Line 12-13: are both instruments affected by these factors in a similar way?

ANSWER: Line 12-13 states the difference between the Pandora and Dobson data is affected by these factors. The limited amount of data and SO2 effect generate the bias in Dobson and SZA dependence affects Pandora measurements. And the temperature and humidity affect both Dobson and Pandora measurements.

4. Line 13: does "temperature" mean effective ozone temperature or instrumental temperature? Or both?

ANSWER: It means Effective ozone temperature.

5. Line 25-27: please, add some bibliographic references;

ANSWER: Bibliographic references are added in manuscript.

6. Line 31: the recovery of the "ozone hole" is still an open question (e.g., Solomon et al., Science, 2016), and the cited articles (1997-2003) do not pretend to report the recovery of the ozone hole, contrary to what the authors state. Please, notice that slow-down of depletion does not necessarily mean recovery. Moreover, a quite confusing explanation of the "ozone hole" is offered to the reader (without even specifying where it occurs) and no mention to the decrease of ozone at midlatitudes is made. Please, rewrite this part;

ANSWER: This part is rewritten.

7. Line 94: why "An" OMI?

ANSWER: It is corrected to OMI

8. Line 216: how can the authors state that Fig. 3 shows a "generally gaussian dis-

Printer-friendly version

Discussion paper



tribution” based on only 6 bins? Can they support their sentence on the basis of a normality test?

ANSWER: The sentence is rewritten.

9. Line 221: "Dobson unit" or "Dobson instrument"?

ANSWER: It indicates Dobson instrument

10. Line 224: are you comparing Pandora, Brewer and OMI from 2012-2014 to the Dobson in the period 1985-2000? Please rewrite this sentence, since it is very confusing. Furthermore, can you assess an increase/decrease by comparing datasets from different instruments?

ANSWER: Line 224 is rewritten to clarify the sentence in manuscript.

11. Lines 312: "fixed temperature". Do you mean effective ozone temperature? Does humidity refer to instrumental humidity or atmospheric humidity?

12. Line 365: "temperature". Instrumental or ozone effective?

ANSWER: It refers to effective TCO retrieval temperature.

13. Table 1: please, write more clearly the year (2012-2013) and the day;

ANSWER: The sentence in Table 1 is revised to be more clearly..

14. Fig. 4: define the used acronyms (OMT, OMD, DBS, etc.). Also, Fig. 4d has no dashed lines. Explain in the caption what the dashed lines indicate;

ANSWER: Definition of acronyms and corrected line are shown in fig. 4.

15. Fig. 5, caption: "blue lines". There are no blue lines in the figures;

ANSWER: It changes to dashed lines in Fig. 5

16. Figs. 4-5: why $n=115$ for all plots? Only days with all instruments measuring were chosen? Is it necessary to report the same number in all plots?

Printer-friendly version

Discussion paper



ANSWER: In this comparison, only days with all instruments measuring were chosen to ensure high reliability of intercomparison results as mentioned in line 229-230.

Please also note the supplement to this comment:

<http://www.atmos-meas-tech-discuss.net/amt-2016-146/amt-2016-146-SC2-supplement.pdf>

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2016-146, 2016.

Printer-friendly version

Discussion paper






Figure 1. The comparison of Brewer and Dobson measurements in Seoul, Korea. The left figure shows the comparison of Brewer data and Dobson data after SO₂ correction from 1999-2005 except for year 2001 in Seoul. And the right figure shows the same comparison but from January 2007 to March 2007.

Fig. 1.

Printer-friendly version

Discussion paper

