Dust Opacities inside Dust Devil Column in the Taklimakan Desert

Zhaopeng Luan[1,2,3], Yongxiang Han[1,2], Feng Liu[1,2], Tianliang Zhao[1,2] , Chong Liu[1,2], Mark J. Rood[4], Xinghua Yang[5], Qing He[5], Huichao Lu[3]

[1] Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science and Technology, Nanjing, 210044, China

[2] Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science and Technology, Nanjing, 210044, China

[3] Tai'an Meteorological Brueau, Tai'an, 271000, China

[4] Department of Civil & Environmental Engineering, University of Illinois at Urbana-Champaign, Illinois ,61820, USA

[5] Institute of Desert Meteorology, China Meteorological Administration, Urumqi ,830002, China

Correspondence Author: Tianliang Zhao (tlzhao@nuist.edu.cn)

Abstract. The distribution of particulate matter (dust) in dust devils (DDs), which are well-defined vortexes of wind that range from 1 m to 1,000 m tall, is quantitatively quantified here based on light transmission. We applied the Digital Optical Method (DOM) with digital still cameras to quantify the opacity of the DDs in the Taklimakan Desert, China. This study presents the following unique and important results and interpretations: 1) the distinct horizontal distributions of opacity indirectly indicated the possible existence of DDs' zone of weak winds in the center of the swirling vortex, similar like the eye of tropical cyclone, which is difficult to be observed directly; 2) The opacity of the DDs decreases with increasing height, however, the dust does appears to settle out, and the relatively calm eye leads to a minimum in dust opacity at

the eye; 3) The horizontal distribution of opacity is quasi symmetric with a bimodal across the eye of the DDs; and 4) A new robust method is developed for the representation of three-dimensional structure of opacity based on the observed two-dimensional structures provided by DOM. This research not only proposes a highly reliable, low cost and efficient methodology to characterize the inside structure of DDs, but also provides the information on estimation of dust emissions caused by DDs.

Keywords: dust devil; digital optical; opacity; dust devil eye; 3-D structure

1 Introduction

Well-defined vortexes of wind that range from 1 m to 1,000 m tall, also known as dust devils (DDs) are the most common small scale (< 50 m diameter at ground level) dust transmitting system in the atmosphere(Kanak, 2005;Kanak et al., 2000;Leovy, 2003). It is a special case of columnar, ground-based convective vortex occurring in the lower atmospheric boundary layer (Gu et al., 2008;Koch and Renno, 2005). Occurrences of wind devils are associated with weak wind, sunny weather, and local surface pressure fluctuations which are dominated by heterogeneous solar radiation due to uneven heating of the ground, that leads to a rising convection vortex rotation containing particulate matter (PM, dust) under certain conditions of angular momentum (Duan et al., 2013).The theory and numerical model results suggest that DDs have significant potential for high dust loadings. DDs contribute to one third of the total natural particulate mass emitted to the atmosphere annually (Koch and Renno,

2005). Emissions of total primary particulate mass by DDs to the atmosphere is as high as 65% in the United States (Gillette and Sinclair, 1990). About 77%-87% of the primary particulate mass emitted to the atmosphere in Chinese deserts is caused by DDs (Han et al., 2008;Deng et al., 2011;Zhang et al., 1994), while DDs generate 30% of the primary particulate mass emitted to the atmosphere in the Sahara Desert (Marsham et al., 2008). These results imply that DDs may play an equal or more important role than sandstorms when considering the total emissions of PM into the atmosphere. DDs affect not only air quality of arid areas but also impact global or regional climate through the “umbrella” (Wei et al., 1998) and “condensation nucleus” effects (Wang and Zhang, 2001) in the atmosphere as well as the “iron fertilizer effect” in open oceans (Han et al., 2006) that occur because of the remote entrainment, transmission and then deposition of dust from the atmosphere.

However, physical processes and mechanisms of dust entrainment by DDs are poorly understood. It is extremely difficult to observe DDs because of their sporadic and unpredictable mobile paths. So the basic structure and characteristic parameters have long been based on human observations by Ives (1947) and Sinclair (1964). DDs’ central core structure has been remarkably observed with mobile Doppler radar (Bluestein et al., 2004) and their center pressure has been characterized by pressure logger (Lorenz, 2012;Lorenz, 2013). However, data obtained from the mobile Doppler radar and pressure logger depend on locating the instruments along the estimated path of the DDs where they may appear, and the measurements are mainly focused on meteorological parameters. The high-cost measurement implementation

requires extensive and expensive resources. In addition, LES（Large Eddy Simulation）have also been applied to DD studies by Toigo (2003), Ito et al.(2010) and Kanak et al. (2000; 2005). The condition of weaker wind and stronger surface heat flux favorable for the formation of dust devil is confirmed according to their studies. Gu et al. (2003; 2010) even provided the meteorological characteristics inside of DDs by numerical simulation. However, field observations of DDs to compare to and validate the modeled results including vortex dynamics are critical and necessary (Leovy, 2003).

The Digital Optical Method (DOM) was developed to measure opacities of plumes in the atmosphere emitted from stationary point sources (Du et al., 2007; Du et al., 2009). DOM uses a digital still camera and software for processing the digital pictures to determine plume opacity. This method was also used to quantify plume opacities for fugitive dust plumes in the atmosphere (Du et al., 2013). DOM was developed to quantify plume opacity at low cost, easy implementation, with improved accuracy compared to human observations of plume opacity, and to provide digital record.

We conducted a DD field campaign in the Taklimakan Desert, the largest desert in China and the world's second largest liquidity desert during July 2014. This is the first time to use DOM to observe DDs. Two dimensional (2D) structures of dust devils are archived with the digital images of the DD and processing with DOM software. This effort provides horizontal and vertical distributions of opacity values in DDs, and presents the first three-dimensional (3D) opacity structure of DDs. It is important to quantify the structure which is helpful to estimate dust emissions of DDs. The study thus services to provide insightful information for DD numerical simulation and

validation.

2 Methods

2.1 Image acquisition and processing

The observational investigation site is located in the Xiaotang region (40°50′N，84°10′E，altitude 943.9m), in the hinterland location of the Taklimakan Desert (see Fig.1). Xiaotang is a typical desert-Gobi transitional zone where DDs occur frequently according to local meteorological observations. During the observations from 2 to 14 July, 2014 digital images were obtained with two digital still cameras (Sony Cyber-shot Model DSC-P100). In order to obtain the appropriate background for taking the pictures, the camera is back to the sun within a 140° sector (ASTM D-7520). All pictures archived as JPEG files. Most observed DDs show inverted circular cone and are quasi-symmetric shape across the DD's eye with a height of several tens to hundreds meters. None of the DDs have absolute symmetry due to the heterogeneous distribution of dust particles in the DD caused by the time-dependent dynamics.

A digital image of a typical DD is illustrated in Fig.1 to describe how to determine the DD opacity values. The vertical curved lines were added to the image to represent auxiliary lines describing vertical grids for estimation of opacity values. The sky background of the DD is relatively uniform as shown in Fig.1. The upright electric pole right next to the DD with a height of 4.5 m was a reference to measure the height and diameter of the DDs. The DD with a base radius of 6.4 m and a height of 23.0 m

has a maximum diameter of 15.8 m at the height of 16.0 m. In order to examine the spatial distribution of dust concentration within the DD's dust cone, we horizontally set the cone into 0.4 m grids with 15 auxiliary upward lines parallel to the DD's conical surface. The centerline (line 0) is a vertical line from the bottom of the DD's eye. For the sake of clarity, we count lines on the left of the centerline from 1 to 7, while denoting lines to the right of the center as A to H (Fig.1b).

2.2 Calculation of opacity

DOM's transmission model was used in this study due to the uniform background sky conditions (e.g., uniform clear or overcast sky) for the DDs. And the challenge to install contrasting backgrounds behind and next to the DDs as needed with DOM's contrast model (Du et al., 2007). The transmission model determines the plume's opacity based on the radiance from the plume and the radiance from the plume's background. Part of the radiance (N_0) from the sky is lost as it passes through the plume due to light scattering and/or absorption. N denotes the radiances received by the charge-coupled device (CCD) of the digital camera that correspond to the sky background, in terms of pixel values:

$$N = N_0 T_0 + N^*$$

T_0 denotes the transmittance of the plume-free atmosphere along the path between the camera and the sky background. N^* is the path radiance of the atmosphere along the same path , which is from direct ,diffuse, and reflected radiances scattered into the sight path by ambient air and aerosols (Du et al., 2013), and can be estimated with an equilibrium radiance model for uniform illumination and negligible absorption:

$$N^* = N_0(1-T_0)$$

From the two equation, getting the results of $N=N_0$.

Therefore, when the radiance reaches the camera (N_p), it is caused by the attenuated radiance value from the plume (N_{T1}), diffusive radiance value (N_{T2}), and attenuation of the radiance caused by the surrounding atmosphere (which is negligible compared to the DD). The attenuated radiance value (N_{T1}) results from N_0 after the light is scattered and/or absorbed by the plume and the diffusive radiance value (N_{T2}) is caused by other sources of light than the uniform sky background (Fig.2). According to the definition of opacity:

$$Opacity = 1 - \frac{N_{t1}}{N_0} = 1 - \frac{N_p - N_{t2}}{N}$$

Because the camera did not directly measure $N_{t2,}$ the proportionality coefficient, K, is defined by N_{t2} = K*N*Opacity (Du et al., 2007), and then the plume opacity could be determined by the transmission model as described by:

$$Opacity = \frac{1 - \frac{N_p}{N}}{1-K}$$

N_p is the equivalent radiance value recorded by the camera, in terms of pixel values, caused by radiance from the plume (N_T) and path radiance of the atmosphere. N is the equivalent radiance value recorded by the camera, in terms of pixel values, after N_0 passes through the DD-free atmosphere. K value of 1.4 in the transmission model is used.

3 Results and discussion

3.1 The vertical profile of opacity

Fig. 3a, b, and c demonstrate the opacity profiles of the vertical grid lines, as shown in Fig.1. It can be seen that: 1) The opacity profiles are non-uniform distributions in the vertical direction, and the vertical variation of opacity becomes more significant from the inner to the outer portions of the conical dust plume (See from the both Fig.3a and b), 2) The opacity profiles from center to both sides show quasi-symmetry but none of them are absolutely symmetrical; and 3) The opacity values generally decrease with height. However, the opacity decreases more rapidly at lower levels than upper ones. The averaged lapse rate (percent change in opacity values with increasing height) for the profile is 4.1% per meter below 10 m height over which the lapse rate drops to 0.6% per meter (Fig.3c).

3.2 The horizontal variation of opacity

From the horizontal variation of the opacity at different levels, it is observed that: 1) the horizontal opacity values increase first and then decrease from the center part to both sides at the base of the DD, implying that the DD is quasi-symmetric; 2) The opacity caused by dust aerosols and the corresponding dust concentrations are decreasing with height within the DD, and the opacity is a monotonically decreasing function as height increase. The maximum opacity is observed at the bottom of the DD (Fig. 4a); and 3) on the right side of the DD, the magnitude of opacity is greater than the left side (Fig.4). Therefore DOM method is able to capture all important well-known natures of DD including non-absolute symmetry and internal inhomogeneity.

3.3 Two-dimensional distribution of the quantized DD's opacity

2D grid boxes shown in Fig. 5 describe the more detailed 2D spatial distributions of the opacity values. Each grid box with a unique averaged opacity value corresponds to a specific dust concentration. The 2D distribution image reflects the basic features of inhomogeneity and quasi-symmetry inside the DD. The grids used to quantify spatial distribution of opacity values are able to present the similar spatial variability like grid boxes defined in DD numerical models to simulate spatial distribution of physical properties (Mason et al., 2013;Gu et al., 2006).

The opacity is related to the PM's concentration, composition, and size distribution in DDs (Metzger et al., 1999). The value of the opacity is assumed to decrease with increasing height due to gravity that prevents larger particles from traveling to the upper parts of the DD. Therefore, it is expected that smaller particles could be transmitted to a higher altitude (Gu et al., 2003;Gu et al., 2007). A distinct vertical opacity gradient develops with small values at the center portion of the DD. It suggests that airflow inside DD is relatively stable and vertical mixing is weak. At the same time larger particles are transported into the outer spiral dust bands where the mixing is much stronger because of entrainment. In addition, the large number of sand vortexes at lower portions of spiral bands explains the coexistence of descending larger particles and ascending fine particles. And these results are consistent with those from numerical simulations by Gu (2007) and Gierasch (1973; 1974).

3.4 The DD eye and the three dimensional distribution of opacity

Previous research have clarified the principal characteristics of DDs that the centers of DDs have low pressure, weak airflow, and almost zero tangential velocity (Fiedler

and Kanak, 2001;Kanak, 2006). Dust concentration in this inner core region is much lower compared to other portion of DD (Balme and Greeley, 2006;Gu et al., 2003). It is similar to the eye of hurricanes or typhoons characterized by light wind, clear skies. Though the DD's eye is difficult to observe directly, the variation of opacity with a bimodal distribution at the same height（Fig.4）indicates that the DD's eye does exist. Fig. 6 illustrates a conceptual mode of the horizontal cross section of a DD. The DD DD's eye diameters are measured by 2R and 2r, respectively. The line segment FG is the distance between two peak opacity points (F and G) away from the eye, H with minimum opacity is the center of the eye. The opacity value at point A in Fig. 6, for instance, is the accumulated opacity through distance from B to C. The changes in opacity from B to C, however, are unknown. So, we assume the horizontal opacity values are constant between B and C.

A more detailed example illustrated by Fig. 4 describes the DD having a diameter of 12 m and a height of 4 m. The horizontal variation of opacity values shows an obvious bimodal distribution. The peak opacity value is located at 3 m away from the DD's centerline. Therefore, it can be determined that the outer radius of DD is 6 m and the inner radius of DD's eye is 3 m at the same height. Every segment (e.g., B to C) is divided into four equal sub-segments, and the horizontal distribution can be derived accordingly. With this method, we can obtain a 3D structure of the DD by calculating the horizontal distribution at different heights as shown in Fig. 7 (i.e., at 3m, 6 m, 9 m, and 12 m). The opacity values increase first and then decreases from the center to both sides of the DD that make up the bimodal distribution. The 3D

structure of DD as described in Fig. 7 represents the fundamental features of inhomogeneity and quasi-symmetric opacity distribution.

The results demonstrate that the methodology to derive the 3D opacity structure in a DD is feasible and practicable. The future effort to improve this methodology is to identify and quantify discrepancies between the assumed opacity values and observed ones from B to C. It can be realized using more than one camera to provide digital images of the same DD from different angles. The more precise and accurate 3D structure of DD will be obtained with the improved measurements.

4 Summary and Conclusions

This study is the first to apply Digital Optical Method (DOM) to quantify the opacity of dust devils (DDs) in the Taklimakan Desert. The two dimensional (2D) distribution of opacity values inside the DD was obtained by DOM and correspondent observational data analysis. Analysis results show that the opacity values of DDs decrease monotonically with height; the opacity decreases more significantly at lower levels than upper ones. Also, the averaged lapse rate for the profile is 4.1% below 10 m height over which the lapse rate drops to 0.6%. The horizontal opacity values from the DD's eye to both edges of the DD increases, reaches a peak value, and then diminishes rapidly. The quasi-asymmetry distribution of dust particles inside DDs can be characterized by the bimodal distribution from observed opacity values from the DD's eye to the outward horizontal directions. The distinct horizontal distribution of opacity values indicates the possible existence of the DD's eye which is difficult to be

observed directly. A typical 3D structure of opacities is developed based on the assumption of constant opacity through horizontal cross-sections of a DD. Such results are encouraging and support the use of DOM as alternative to other instrumentally intensive, generally expensive methodologies for DD observations.

5 Prospect

In order to improve the quality of DD observations using DOM, the following key issues will be addressed: 1) the proportionality coefficient (K) is not available so far for desert dust devils. The K value we used in our opacity calculations is for white plume (light scattering plume) whose optical properties is definitely different form DD (desert dust particles), the estimate of K for desert dust is identified as a urgent need for our future study; 2) a scaling model of DD size from DD images is required if there is no reference to determine DD scale; 3) the results documented in this study are far from generalized characteristics of DD opacity. In order to realize statistical analysis of large sized samples, more observations including carefully-designed specific field campaigns are needed for characterizing DD opacities; 4) the validation of calculated opacities inside DD column is also a challenge for us. It is essential to collect other independent observation data of DD for this purpose.

A new method to convert 2D structure of opacity using multiple-camera digital optical method (MDOM) to 3D will be developed. The results from further improved MDOM will be related to the estimate of vertical gradient of dust concentration. In addition, the wind shear and Reynolds stress will be considered in the improved system in order to realize the parameterization of vertical dust flux caused by DDs.

The opacity of dust devils may also be affected by meteorology like ambient wind and temperature difference between surface and air, which may influence the strength and size of dust devils. High-strength dust devils can roll up more dust particles, affecting the opacity of dust devils. So in the following study, more observation sites will be set to measure meteorology such as ambient wind, temperature difference between surface and air, wind of dust devils, temperature and pressure difference between core of dust devils and surrounding environment, etc.

Acknowledgments

This research was supported the National Natural Science Foundation of China (41375158; 41175093; 41405013) and the Research Starting Project of NUIST (20110304) of China. Any opinions, findings, and conclusions or recommendations expressed in this paper are those of the authors and do not necessarily reflect the view of the Nanjing University of Information Science and Technology, University of Illinois at Urban and Champaign, or any organizations and government agencies.

References

American Society of Testing Materials (2013) Standard Test Method for Determining the Opacity of a Plume in the Outdoor Ambient Atmosphere (ASTM D-7520), http://www.astm.org/Standards/D7520.htm.

Balme, M., and Greeley, R.: Dust devils on Earth and Mars, Reviews of Geophysics, 44, 2006, doi: 10.1029/2005RG000188.

Bluestein, H. B., Weiss, C. C., and Pazmany, A. L.: Doppler radar observations of dust devils in Texas, Monthly weather review, 132, 209-224, 2004,

doi: http://dx.doi.org/10.1175/1520-0493(2004)132<0209:DROODD>2.0.CO;2.

Duan, J.P., Han, Y. X., Zhao, T. L., and Song, H. D.: The contribution of dust devils to atmospheric dust aerosols and its relation with solar radiation. China Environmental Science, 33**,** 43-48,2013. (In Chinese with English abstract)

Deng, Z.Q., Han, Y.X., Bai, Z.H.,and Zhao, T.L.:Relationship between dust aerosol and solar radiation in gebi desert in North China.China Environment Science, 31(11):1761-1767,2011.(In Chinese with English abstract)

Du, K., Rood, M. J., Kim, B. J., Kemme, M. R., Franek, B., and Mattison, K.: Quantification of plume opacity by digital photography, Environmental science & technology, 41, 928-935, 2007, doi: 10.1021/es061277n.

Du K., Rood, M. J., Kim, B.J., Kemme, M. R., Franek, B., and Mattison, K.: Evaluation of digital optical method to determine plume opacity during nighttime. Environmental science & technology, 43**,** 783-789,2009, doi**:** 10.1021/es800483x.

Du, K., Shi, P., Rood, M. J., Wang, K., Wang, Y., and Varma, R. M.: Digital Optical Method to quantify the visual opacity of fugitive plumes, Atmospheric Environment, 77, 983-989, 2013, doi:10.1016/j.atmosenv.2013.06.017.

Fiedler, B. H., and Kanak, K. M.: Rayleigh- Bénard convection as a tool for studying dust devils, Atmospheric Science Letters, 2, 104-113, 2001,doi : 10.1006/asle.2001.0046.

Gillette, D. A., and Sinclair, P. C.: Estimation of suspension of alkaline material by dust devils in the United States, Atmospheric Environment. Part A. General Topics, 24, 1135-1142, 1990, doi:10.1016/0960-1686(90)90078-2.

Gu, Z., Zhao, Y., Yu, Y., and Feng, X.: Numerical study of the formation evolution and structure of dust devil. Acta Meteorologica Sinica, 61**,** 751-760,2003. (In Chinese with English abstract)

Gu, Z. L., Wei, W., and Zhao, Y.Z.: An overview of surface conditions in numerical simulations of dust devils and the consequent near-surface air flow fields. Aerosol Air Qual. Res, 10**,** 272-281,2010, doi: 10.4209/aaqr.2009.12.0077.

Gu, Z. L., Qiu, J., Lu, L.Y., and Zhao, Y.Z.: Advances in Study of Dust Devils, Journal of Desert Research, 5, 022, 2007. (In Chinese with English abstract)

Gu, Z., Zhao, Y., Li, Y., Yu, Y., and Feng, X.: Numerical simulation of dust lifting within dust devils-simulation of an intense vortex, Journal of the atmospheric sciences, 63, 2630-2641, 2006, doi: http://dx.doi.org/10.1175/JAS3748.1.

Gu, Z., Qiu, J., Zhao, Y., and Hou, X.: Analysis on dust devil containing loess dusts of different sizes, Aerosol Air Qual. Res, 8, 65-77, 2008,doi:10.4209/aaqr.2007.03.0026.

Gierasch, P., and Goody, R.: A model of a Martian great dust storm. Journal of the Atmospheric Sciences, 30**,** 169-179,1973,doi: http://dx.doi.org/10.1175/1520-0469(1973)030<0169:AMOAMG>2.0.CO;2.

Gierasch, P. J.: Martian dust storms. Reviews of Geophysics, 12**,** 730-734,1974, doi**:** 10.1029/RG012i004p00730.

Han, Y., Fang, X., Xi, X., Song, L., and Yang, S.: Dust storm in Asia continent and its bio-environmental effects in the North Pacific: A case study of the strongest dust event in April, 2001 in central Asia, Chinese Science Bulletin, 51, 723-730, 2006,doi: 10.1007/s11434-006-0723-2.

Han, Y., Dai, X., Fang, X., Chen, Y., and Kang, F.: Dust aerosols: a possible accelerant for an increasingly arid climate in North China, Journal of Arid Environments, 72, 1476-1489, 2008, doi:10.1016/j.jaridenv.2008.02.017.

Ives, R. L.: Behavior of dust devils. Bull. Amer. Meteor. Soc, 28**,** 168-174,1947.

Ito, J., Tanaka, R. and Niino, H.: Large eddy simulation of dust devils in a diurnally-evolving convective mixed layer. Journal of the Meteorological Society of Japan, 88,63-77,2010,doi:10.2151/jmsj.2010-105.

Kanak, K. M., Lilly, D. K., and Snow, J. T.: The formation of vertical vortices in the convective boundary layer, Quarterly Journal of the Royal Meteorological Society, 126, 2789-2810, 2000,doi: 10.1002/qj.49712656910.

Kanak, K. M.: Numerical simulation of dust devil-scale vortices, Quarterly Journal of the Royal Meteorological Society, 131, 1271-1292, 2005,doi: 10.1256/qj.03.172.

Kanak, K. M.: On the numerical simulation of dust devil–like vortices in terrestrial and Martian convective boundary layers, Geophysical research letters, 33, 2006, doi: 10.1029/2006GL026207.

Koch, J., and Renno, N. O.: The role of convective plumes and vortices on the global aerosol budget, Geophysical research letters, 32, 2005, doi**:** 10.1029/2005GL023420.

Leovy, C. B.: Mars: The devil is in the dust, Nature, 424, 1008-1009, 2003, doi:10.1038/4241008a.

Lorenz, R.: Observing desert dust devils with a pressure logger, Geoscientific Instrumentation, Methods and Data Systems, 1, 209-220, 2012,doi:10.5194/gi-1-209-2012.

Lorenz, R. D.: Irregular dust devil pressure drops on Earth and Mars: Effect of cycloidal tracks, Planetary and Space Science, 76, 96-103, 2013, doi:10.1016/j.pss.2013.01.001.

Marsham, J., Parker, D., Grams, C., Johnson, B., Grey, W., and Ross, A.: Observations of mesoscale and boundary-layer scale circulations affecting dust transport and uplift over the Sahara, Atmospheric Chemistry and Physics, 8, 6979-6993, 2008,doi:10.5194/acp-8-6979-2008.

Mason, J. P., Patel, M. R., and Lewis, S. R.: Radiative transfer modelling of dust devils, Icarus, 223, 1-10, 2013, doi:10.1016/j.icarus.2012.11.018.

Metzger, S. M., Carr, J. R., Johnson, J. R., Parker, T. J., andLemmon, M. T.: Dust devil vortices seen by the Mars Pathfinder camera. Geophysical research letters, 26**,** 2781-2784, 1999, doi**:** 10.1029/1999GL008341.

Sinclair, P. C.: Some preliminary dust devil measurements. Monthly Weather Review, 92,1964,available at: http://citeseerx.ist.psu.edu/viewdoc/download?doi= 10.1.1.395.3338&rep=rep1&type=pdf.

Toigo, A. D., M. I. Richardson, S. P. Ewald, and P. J. Gierasch: Numerical simulation of Martian dust devils. Journal of Geophysical Research: Planets (1991–2012), 108,2003, doi**:** 10.1029/2002JE002002.

Wei L. and Shen,Z.B.:The radiative characteristics of atmospheric dust observed from satellite.J.Plateau Meteorology, 17(4):347-355,1998.(In Chinese with English abstract)

Wang, M., and Zhang, R.: Frontier of atmospheric aerosols researches, Climatic and Environmental Research, 6, 119-124, 2001.(In Chinese with English abstract)

Zhang X.Y.,An, Z.S., Zhang, G.Y., Chen, T., Liu, D.S., Arimoto, R.:Transport, deposition and reflected climate change of Chinese atmospheric particulate-I. Modern Atmospheric Aerosols, J. Science in China, 24(12):1314-1322,1994. (In Chinese with English abstract)

Fig.1. A typical DD observed on July 2, 2014 with (a) the original photo, and（b）the image marked with black lines, letters and numbers for the analysis of DD vertical structures

Fig.2. Schematic describing the transmission model to determine plume opacity

Fig.3. Vertical profiles of opacity of dust plume (a and b are the line-specific vertical profiles of opacity of dust plume, c is averaged vertical profile and vertical standard deviations of opacity of dust plume, the averaged lapse rate for the profile below 10m height is 4.1% over which it is 0.6%)

Fig.4. Horizontal variation of the DD's opacity(%) at different levels

Fig.5. Distribution of the DD's opacity in a vertical cross section

Fig.6. DD's horizontal section model and the corresponding variation of opacity

Fig.7. 3D structure of the DD's opacity