

Answer to Report #1

Thank you very much for your valuable comments. I answered them in the following and changed the manuscript accordingly especially at the end of section 4 where the reasons for the oscillation in the mean squared residuals are discussed.

The revised version of the Wüst et al. manuscript is strongly improved compared to the original version. The authors clarified the scope of their method and adapted the description according to both reviewers' comments. I am happy that the scope is much clearer now. Using the spline method for the description of the background and taking the residuals as wave disturbance is a widely used approach in data analyses. Therefore, it has been useful, e.g., to change the term "approximation error" into "sum of squared residuals". Nevertheless I am still concerned about the structure of the manuscript. Furthermore, a few answers to my previous review stimulated some new comments.

General comments:

1. The authors made clear that the description of (gravity) waves from the residuals of the spline fit is the major topic of this paper. In detail, the presumably artificial oscillation in averaged SABER GW activity is the main motivation to develop improved methods of spline fitting. On the other hand they only shortly speculate about the true origin of this oscillation. Instead, they state in the Discussion that the analysis "is beyond the scope of this manuscript". I suggest to at least try to examine the true reason for this oscillation. Is it visible in every profile or in monthly averages? Is it only a result of averaging? Does it also appear if some random GW are added to a CIRA profile? Does it also appear if the sampling point distance is set to, e.g., 15 km? See also below my addition to the old comment #9.

I included two more figures (9 and 10, one of them figure 1 of the former answer to the reviewer comments) and discussed them at the end of section 4 in order to make clearer where the mentioned oscillations probably come from.

2. From my point of view it should be made clearer that the difference between normal and repeating spline fitting is largest if the sampling point distance is close to the half wavelength.

I added this information in the last paragraph of section 4.

In case of the constant background the differences between both methods vanish for distances larger than 2 km or smaller than about 1.3 km. In case of the CIRA background, differences are always only about 20% of the sum of squared residuals. Here it would be very interesting to see, which of both methods compares better with the true background profile. This comparison can only be done with CIRA because here the background is exactly known.

I included this comparison in figure 7 (d)

3. From my point of view, the essential information from this paper is twofold:
 - i) The spline method may produce some artefacts in separating background and waves if the data set contains waves with wavelengths of about double the sampling point distance. This can at least partly be overcome with the repeating spline.
 - ii) The description of the background by a too coarse spline may produce artificial oscillations in the residuals which should not be confused with GW. This interesting result from the answer to Reviewer 1 should also be presented in the paper, because it demonstrates the limitations of (non-repeating) splines in general, if the background is unknown (as usual).

I added this information in the summary.

Further concerns about the structure of the manuscript are described in the specific comments below.

Specific Comments to the Author Response:

1. Old Comment 7: In the Summary (p. 9, l. 8) the “ability of a spline to approximate oscillations” is mentioned, not the ability to reproduce a background state (and wave-induced residuals).

Answer: We hope that we have clarified all confusing points mentioned above and leave the formulation in the summary as it was since this meets the purpose of the manuscript.

My comment: At different positions of the manuscript you write that the motivation is to remove the oscillations in the wave activity (Fig. 1). Obviously this can only be done by improved description of the background by the spline, not by optimal approximation of the oscillations.

Changed, see answer to specific comment no. 8

2. Old Comment 9. (i) P. 8, l. 17-18: i) It is still surprising that the annual mean residuals (~500 profiles) show such a pronounced oscillation. Please comment on this.

Answer: See page 10, first paragraph for possible explanations. The answer to the major comment, number 3 of the other reviewer helps to better understand the information which we added to this paragraph.

My comment: As written above, the new figure provided in the answer to Reviewer 1 would also improve the manuscript. It shows limitations of the standard spline approximation by an impressive example. Furthermore, it gives some hints on the question why oscillations with a wavelength similar to the sampling point distance appear in the average data of Fig. 1, but may not be visible in a single GW profile.

See answer to general comment no. 1

Specific Comments to the manuscript (line numbers refer to the manuscript without tracked changes):

1. p.1, l. 14: Maybe I am too strict on this, but I suggest writing “approximation of the background state”

Done

2. Section 3, case study with constant background: Obviously in this section it is intended to show a method for a good approximation of the oscillations (disturbed profile), see, e.g., p. 5, l. 27, “sinusoidal approximation is better adapted”. This is somewhat outside of the main motivation. Therefore it should be made clear that, e.g., this section helps to understand the general behavior of splines if the data set contains waves with a wavelength of double the sampling point distance – which may happen in the general case of an unknown mixture of waves.

We provide the information about the purpose of this section now right at the section beginning. Furthermore, we deleted the sentence “sinusoidal approximation is better adapted” and re-formulated the sentence “the oscillation is adapted very well between 20 km and 40 km for a phase of $\pi/2$ and $3\pi/2$.” (p. 6, ll. 2–3, version with changes marked)

3. Section 3, case study with CIRA: Now it seems that a good approximation of the original CIRA profile is intended. Therefore, it would be very helpful to show also the original profile in Fig. 7. This helps the reader to estimate whether the conventional spline or the repeating spline removes the waves better. Additionally, also the residuals between both retrieved profiles and the original CIRA should be calculated.

I included the original CIRA-profile in figure 7 a & b. I also plotted the difference between the both retrieved profiles and the original CIRA-background (see figure 7d) and changed the manuscript accordingly (p6. ll.19–22, version with changes marked).

4. p. 7, l. 1: Again, I thought that the motivation is not to reproduce oscillations by the spline, but to reproduce the background and to calculate the GW from the residuals.

I changed the sentence into “In section 3, we showed that the quality of spline to approximate the background and also its ability to filter for a specific part of the wave spectrum varies” to point out that the purpose is the background approximation including from case to case also large-scale gravity wave signature for example.

5. p. 8, l. 8-13: The calculation is straightforward. I suggest shortening this part and giving the numbers in dB and K^2 without further explanation.
Done
6. p. 8, l. 14-16: I think it is hard to say from the comparison with Shuai et al., which method is better. Their numbers are comparable to both, Fig. 1 and Fig. 8, especially if potential longitudinal variations are taken into account. I suggest adding an appropriate statement.
I changed “These values agree very well with the ones provided here” to “These values agree very well with the ones provided here but it cannot be decided whether they match better with the ones based on the repeating or non-repeating approach (fig. 1 or 8 (a))”.
Unfortunately I do not have access to the Shuai et al. paper. If they show profiles, it would be helpful to learn whether these are also showing the “surprising” oscillations as Fig. 1. Shuai et al. provide exclusively colour-coded means (figure 2) or profiles of the maximum of monthly averaged GW activity over all years (figure 3). The latter shows fluctuations but they can also be due to the fact that they look for the maximum at each height. Due to the scaling of the colour bar and the use of dB (which includes a logarithm and makes fluctuations appearing smaller) it is from my point not possible to provide a serious statement about possible fluctuations.
7. p. 9, l. 3: From my understanding, the motivation of this paper is to remove these oscillations (see Introduction and other places). Therefore, the analysis should not be “beyond the scope”. See above.
See answer above.
8. p. 10, l. 8: I suggest (again) rephrasing “ability of the spline to approximate oscillations”. See above.
I changed “oscillations” to the “background state (and large-scale wave-induced fluctuations)”.

Typos:

1. p. 9, l. 1: “not very much” should read “not vary much”
Done
2. Fig. 6: “reisiduals” should read “residuals”
Done

Smoothing data series by means of cubic splines:
quality of approximation and introduction of a repeating spline
approach

Sabine Wüst ¹, Verena Wendt ^{1,2}, Ricarda Linz ¹, Michael Bittner ^{1,3}

¹Deutsches Fernerkundungsdatenzentrum, Deutsches Zentrum für Luft- und Raumfahrt, 82234 Oberpfaffenhofen, Germany

²Umweltforschungsstation Schneefernerhaus, Zugspitze, Germany, now at: Institut für industrielle Informationstechnik, Hochschule Ostwestfalen-Lippe, Ostwestfalen-Lippe, Germany

³Institut für Physik, Universität Augsburg, 86159 Augsburg, Germany

Correspondence to: Sabine Wüst (sabine.wuest@dlr.de)

Abstract. Cubic splines with equidistant spline sampling points are a common method in atmospheric science for the approximation of background conditions by means of filtering superimposed fluctuations from a data series. What is defined as background or superimposed fluctuation depends on the specific research question. The latter also determines whether the spline or the residuals—the subtraction of the spline from the original time series—are further analysed.

Based on test data sets, we show that the quality of approximation of the background state ~~approximation~~ is not increasing continuously with increasing number of spline sampling points / decreasing distance between two spline sampling points. Splines can generate considerable artificial oscillations in the background and the residuals.

We introduce a repeating spline approach which is able to significantly reduce this phenomenon. We apply it not only to the test data but also to TIMED-SABER temperature data and choose the distance between two spline sampling points in a way that we are sensitive for a large spectrum of gravity waves.

1 Introduction

It is essential for the analysis of atmospheric wave signatures like gravity waves that these fluctuations are properly separated from the background. Therefore, particular attention must be attributed to this step during data analysis. Splines are a common method in atmospheric science for the approximation of atmospheric background conditions. The shortest wavelength or period which can be resolved by the spline is twice the sampling point distance according to the Nyquist theorem. Depending on the field of interest, either the smoothed data series or the residuals—the subtraction of a spline from the original time series—are further analysed (see for example the work of Kramer et al., 2016; Baumgarten et al., 2015; Zhang et al., 2012; Wüst and Bittner, 2011; Wüst and Bittner, 2008; Young et al., 1997; Eckermann et al., 1995).

Algorithms for the calculation of splines are implemented in many programming languages and in various code packages making them easy to use. Nevertheless, spline approximations need sometimes to be handled with care when it comes to physical interpretation.

Figure 1 explains our motivation for the work presented below. It shows the squared temperature residuals averaged over one year for the years 2010–2014 versus height between 44° N and 48° N and 5° E and 15° E (approximately 500 profiles per year). This region includes the Alps where gravity waves are supposed to be generated. The vertical temperature profiles are derived from the SABER (Sounding of the Atmosphere using Broadband Emission Radiometry) instrument on board the satellite TIMED (Thermosphere Ionosphere Mesosphere Energetics Dynamics), data version 2.0 (details about this data version can be found in Wüst et al. (2016) and references therein, for example). For the calculation of the residuals, we applied a cubic spline routine with equidistant sampling points. As mentioned above, the shortest wavelength which can be resolved by the spline is two times the distance between two consecutive spline sampling points. At the same time, this wavelength is the largest resolvable one in the residuals. The number of spline sampling points (and the length of the data series) therefore determines the sensitivity of the spline to specific wavelengths. The distance of 10 km between two spline sampling points makes sure that we are sensitive for a large spectrum of gravity waves. Therefore, we take the squared temperature residuals as a simple proxy for gravity wave activity with vertical wavelengths up to 20 km. Obviously, the mean squared residuals do not only reveal a strong and continuous increase with height (note the logarithmic x-axis) as it is expected since gravity wave amplitudes should increase due to the exponentially decreasing atmospheric background pressure with altitude. Superimposed on this general increase of gravity wave activity are well-pronounced oscillations with a wavelength of ca. 10 km, which is nearly equal to the distance between two spline sampling points.

Since we are not aware of any physical reason for this oscillation, we formulate the hypothesis that this is an artefact of the analysis. In order to avoid or at least reduce such problems we here propose an repeating variation of the cubic spline approach, which we explain in section 2. In section 3, we apply the original and the repeating approach to test data sets. The results are discussed in section 4. A brief summary is given in section 5.

2 Methods and algorithms

The approach we investigate here relies on cubic splines with equidistant sampling points. Since spline theory is well-elaborated, we will therefore not go into much detail here. The algorithm we use is based on Lawson and Hanson (1974).

The first step for the adaption of a spline function to a data series on an interval [a, b] is the choice of the number of spline sampling points (also called knots). These points divide the interval for which the spline is calculated into sub-intervals of equal length. For each sub-interval a third-order polynomial needs to be defined, that means the coefficients have to be determined. At the spline sampling points, not only the function value, but also the first and second derivatives of the two adjacent polynomials need to be equal. The optimal set of coefficients is calculated according to a least square approach where the sum of the squared differences between the data series and the spline is minimized.

As mentioned above, the number of spline sampling points (and the length of the data series) determines the sensitivity of the spline to specific wavelengths. Since the length of the data series must be an integer number of the distance between two spline sample points only certain distances between two consecutive spline sampling points can be chosen if the whole data series is approximated. We would like to operate the spline algorithm in the way that we provide the shortest wavelength which shall be resolved by the spline. That means that we have to cut the upper part of the profile from case to case. This is only possible for data sets of sufficient length as the SABER temperature profiles we used for our purpose. In detail, our spline algorithm works as follows. The scheme includes the repeating as well as the non-repeating algorithm.

- *Step 1: Provision of shortest wavelength*

We provide the algorithm the shortest wavelength which shall be resolved by the spline (in the following denoted as \lim). It is equal to the doubled distance between two spline sampling points; therefore the distance between two spline sampling points is equal to $\lim/2$.

- *Step 2: Determination of x-values of the spline sampling points*

The minimal x-value of the data series is subtracted from the maximal x-value, the difference is divided by $\lim/2$. If the result is a whole-number, 1 is added. If this is not the case, the closest integer less than the result is calculated and 1 is added. This is the number of spline sampling points used for the next step. It is denoted as n:

$$n = \left(\frac{x_{\max} - x_{\min}}{\lim/2} - \frac{x_{\max} - x_{\min}}{\lim/2} \bmod 1 \right) + 1 \quad (1)$$

Knowing $\lim/2$ and the minimal x-value, the x-values of the further spline sampling points can be calculated.

- *Step 3: Calculation of spline approximation*

The spline approximation is calculated based on Lawson and Hanson (1974). If the length of the data series is not equal to an integer multiple of $\lim/2$, the surplus part at the end of the data series is not subject of this step. For the non-repeating approach, the spline algorithm stops here.

- *Step 4 (only in the case of the repeating approach): Iteration of starting point*

The first point of the data series is removed and step 2 and 3 are repeated. If the starting point is equal to the original minimal x-value plus $\lim/2$, the algorithm proceeds with step 5.

5

- *Step 5 (only in the case of the repeating approach): Calculation of the final spline*

The mean of all splines derived before is calculated. That is the final (repeating) spline.

For the repeating approach, the length of the data series is not the same in each iteration since data at the beginning and the end of the data series are not necessarily part of each iteration: at the beginning of the data series, this holds for all x-values between the minimal x-value and the minimal x-value plus $\lim/2$ (see step 4), at the end of the data series, this is the case for all values between the maximal x-value and the maximal x-value minus $\lim/2$ (see step 3).

For the non-repeating approach, data are cut only at the end of the data series if the length of the data series is not equal to an integer multiple of $\lim/2$.

3 Case studies

The purpose of this section is to help understanding the general behaviour of splines if the data set contains waves with a wavelength of double the sampling point distance—which may happen in the general case of an unknown mixture of waves.

We generate a basic example using an artificial sine with a vertical wavelength of 3 km, a phase of zero and an amplitude of one. The function is sampled every 375 m (that means at its zero-crossings, at its extrema and once in between the zero-crossing and the next extremum / the extremum and the next zero-crossing).

The values for the sampling rate and the vertical wavelength are set arbitrarily. However, the spatial resolution of 375 m is motivated through the spatial resolution of TIMED-SABER, an instrument which is commonly used for the investigation of gravity waves (e.g. Zhang et al., 2012; Ern et al., 2011; Wright et al., 2011; Krebsbach and Preusse, 2007) and which delivered also the temperature profiles we used in fig. 1.

Fig. 2 (a) shows the test data series (dotted line) between 15 km and 100 km height. This great height range is chosen since it facilitates the demonstration of our results. A (non-repeating) spline with a distance of 1.5 km between two spline sampling points is fitted (solid line). According to the Nyquist theorem, the chosen distance between two spline sampling points is small enough for resolving the oscillation in our test data. In part (b) and (c) of fig. 2, a spline with a distance of 1.6 km and 1.4 km between two spline sampling points is calculated. Part (d) to (f) of fig. 2 focus on the height range of 15 km to 50 km of fig. 2 (a) to (c): here, the height-coordinates of the spline sampling points are plotted additionally (dashed-dotted lines). The asterisks mark the sampling points of the original sine. The spline adaption in fig. 2 (a) / (d) differs significantly from the spline adaption in fig. 2 (b) / (e) and 2 (c) / (f): apart from a slight oscillation at the beginning / end of the height interval, the spline is equal to zero in fig. 2 (a) / (d). The spline approximation plotted in fig. 2 (b) and (c) shows a beat-like structure in the whole height range.

In order to give an overview concerning the quality of adaption not only for some chosen examples as they were shown in fig. 2, the test data set is approximated by a cubic spline with varying numbers of spline sampling points. The squared differences between the spline and the test data are summed up between 20 km and 40 km (this height interval is chosen in order to be consistent with fig. 7 later). We call this value the sum of squared residuals which is equal to the approximation error in this case. It is not decreasing continuously with increasing number of spline sampling points / decreasing distance between two spline sampling points but it is characterized through a superimposed oscillation which reaches its maximum for a distance of ca. 1.5 km between two spline sampling points (fig. 3, solid line). When changing the phase of the test data set to $\pi/2$ (instead of zero), the sum of squared residuals for a distance of ca. 1.5 km between two spline sampling points is much lower (fig. 3, dashed line), that means the sinusoidal oscillation is better adapted in this case. This makes clear that the sum of squared residuals depends on the phase of the oscillation (one can also say on the exact position of the spline sampling points).

The analysis described above is repeated, but the phase of the oscillation is varied between 0 and 2π . The sum of squared residuals (between 20 km and 40 km) is calculated for three different distances between two spline sampling points: 1.5 km

(fig. 4, solid line), 1.4 km (fig. 4, short dashes) and 1.6 km (fig. 4, long dashes). The dependence on the phase is most pronounced for a distance of 1.5 km: ~~the sum of squared residuals the oscillation is adapted very well between 20 km and 40 km is minimal~~ for a phase of $\pi/2$ and $3\pi/2$. For a phase of 0 and π , the contrary holds.

This example directly motivates the application of the repeating spline approach on the same test data set (see fig 5 (a)–(f) which can be directly compared to fig 2 (a)–(f): the black line represents the final spline approximation and the different colours refer to the spline approximations during the different iteration steps). In this case, the sum of squared residuals depends much less on the distance between two spline sampling points (fig. 6 (a)) and on the phase of the test data set (fig. 6 (b)). Only for a distance of 1.6 km between two spline sampling points, a slight phase dependence is still visible (fig. 6 (b)).

Until now, we showed only test data which are not superimposed on a larger-scale variation like the atmospheric temperature background. Now, three sinusoids with vertical wavelengths of 3, 5, and 13 km, phase 0, $\pi/3$, and $\pi/5$, and amplitude 2.0 (growing amplitude with height neglected for simplicity reasons) are superimposed on a realistic vertical temperature background (fig. 7 (a)). The background is based on CIRA-86 (COSPAR International Reference Atmosphere, Committee on space Research and NASA National Space Science Data Center, 2006) temperature data for 45° N for January, which is brought on a regular grid using a cubic spline with a distance of 3 km between two spline sampling points. It was checked that no additional signatures are caused thereby. The sum of squared residuals shows three steps but no superimposed oscillations (fig. 7 (b)): the first step at ca. 6 km to 7 km (distance between two spline sampling points), the second one at ca. 2 km to 3 km and the last one at 1 km to 2 km. Following Nyquist's sampling theorem, this observation can be explained through the ability of the spline to adapt the original wavelengths. Even if the sum of squared residuals decreases much smoother for the repeating approach than for the non-repeating one with decreasing distance between two spline sampling points, the approximation of the CIRA-background is in both cases not optimal (fig. 7 (d) and dashed lines in fig. 7 (a) and (b)). The realistic background makes clear why we restrict the calculation of the sum of squared residuals to the height range between 20 km and 40 km: this height interval is chosen in order to exclude especially the stratopause since the fast changing temperature gradient can cause additional problems for the spline approximation. Furthermore, the choice of this interval makes sure that the data used for fig. 7 (b) are part of each iteration step (which is not the case for the data at the beginning and the end of the data series, see section 2).

4 Discussion

In section 3, we showed that the ~~ability-quality of spline to approximate the background of a spline to approximate oscillations and therefore also its ability~~ to filter for a specific part of the wave spectrum varies

- a) with the number of spline sampling points, and
- b) with the exact position (height coordinate) of the spline sampling points.

While the first statement can be explained Nyquist's sampling theorem, the second one is not well-known.

When the distance between two spline sampling points matches exactly half the wavelength of the test data, the approximation is worst for a phase of 0 and π . In this case, the spline sampling points are located exactly between the extrema of the test data. If the height coordinates of the spline sampling points agree with the height coordinates of the extrema of the test data, the contrary holds (in appendix A, we provide a mathematical explanation for this observation). The dependence of the quality of approximation on the phase of the test data decreases with greater / smaller distances between two spline sampling points (fig. 3). These findings directly motivate the use of the presented repeating spline approach which is characterized by varying positions (height coordinate) of the spline sampling points.

Furthermore, we showed that if the distance between two spline sampling points is only slightly larger or smaller than half the wavelength present in the data series and if enough wave trains are present (which might not be the case in reality), the (non-repeating) spline reminds of a beat (see fig. 2 (b) and (c), an explanation is given in appendix B). The subtraction of such a "beat" will lead to an artificial oscillation in the residuals with a periodically increasing and decreasing amplitude reaching ca. 70–80% of the original amplitude at maximum (fig. 2 (e) and (f)). This oscillation must not be interpreted as a gravity wave of varying amplitude, for example, and the described effect has to be taken into account when analysing wavelengths similar to the doubled distance between two spline sampling points.

For our case studies, we used a constant and a realistic CIRA-based temperature background profile. For both background profiles, we showed the sum of squared residuals decreases much smoother with increasing number of spline sampling points for the repeating approach compared to the non-repeating one (compare fig. 3 to fig. 6 (a)) and the amplitude of the "beat"-like structure is reduced.

However, the motivation for this work was—as already mentioned—the results shown in fig. 1 which are characterized by a strong superimposed oscillation with a wavelength of approximately 10 km for which we do not have a physical explanation.

Figure 8 (a) now depicts the mean squared residuals after the application of the repeating spline to the same data set, fig. 8 (b) focuses on the year 2014 (the dashed line is based on the application of the repeating spline, the solid line refers to the non-repeating spline). This year is chosen arbitrarily and allows the direct comparison between the repeating and non-repeating approach. The amplitude of the superimposed oscillation is reduced significantly but the oscillation can still be

observed. This supports our hypothesis that the strong superimposed oscillation described in fig. 1 is an artefact of the non-repeating spline de-trending procedure. Furthermore, it becomes obvious now that gravity wave activity increases less with altitude between approximately 45 km and 60 km height compared to the height range below and above. This is in accordance with literature (e.g. Mzé et al., 2014; Offermann et al., 2009). For most heights, the mean squared residuals are smaller for the repeating approach than for the non-repeating one. At 38 km height for example, the difference reaches ca. 2.5 K², which is approximately 32 % (referring to the mean value of both approaches). Shuai et al. (2014) use an earlier version of TIMED-SABER temperature data (1.07) and a different de-trending procedure as we do in order to derive monthly averages of the squared temperature fluctuations for the years 2002–2010. They provide this parameter in dB ($10 \cdot \log_{10}(T_{GW}^{\prime 2})$) with the squared temperature fluctuation $T_{GW}^{\prime 2}$ ~~which makes a short calculation necessary.~~

For 100 km (25 km) height, we extract a yearly mean of ca. 21 dB (4 dB) for 50°N from their figure 2 that means a squared temperature fluctuation of ca. 126 K² (2.5 K²).

$$10 \cdot \log_{10}(T_{GW}^{\prime 2}) = 21 \Leftrightarrow \log_{10}(T_{GW}^{\prime 2}) = 2.1 \Leftrightarrow T_{GW}^{\prime 2} = 10^{2.1} \sim 126$$

For 25 km height, we read a mean value of ca. 4 dB:

$$10 \cdot \log_{10}(T_{GW}^{\prime 2}) = 4 \Leftrightarrow \log_{10}(T_{GW}^{\prime 2}) = 0.4 \Leftrightarrow T_{GW}^{\prime 2} = 10^{0.4} \sim 2.5$$

These values agree very well with the ones provided here but it cannot be decided whether they match better with the ones based on the repeating or non-repeating approach (fig. 1 or 8 (a)). ~~Furthermore~~ However, the overall structure which is characterized by a slow increasing or even a nearly constant gravity wave activity in the upper stratosphere can be observed in their fig. 2 (and in parts also fig. 3(a)) and our fig. 8 (a).

In order to give a comprehensive comparison of the repeating and the non-repeating spline algorithm, we calculate also the mean (non-squared) residuals. In this case, the results look very similarly; in both cases, they again show an oscillation with a vertical wavelength of 10–20 km (fig. 8 (c) for the non-repeating approach, fig. 8 (d) for the repeating spline approach). We can explain this in the following way: when calculating the mean (non-squared) residuals and the mean squared residuals at a specific height, one refers to two different parameters of the distribution of residuals at that specific height. While the mean (non-squared) residuals estimate the mean of the distribution, the mean squared residuals refer to the variance of the distribution. We conclude: at a defined height, the repeating approach changes the mean of the distribution of the residuals only slightly, but it reduces its spread significantly. For individual profiles, the approximation through the repeating approach is therefore less variable in average and can be recommended. The repeating approach can also be recommended if squared residuals are needed for further analysis (e.g. for the calculation of the wave potential energy). If non-squared residuals will be analysed, it does not make a difference in average which approach is applied, for the individual profile, however, this does not necessarily hold. In this case, only waves with amplitudes larger than 0.5 K in the stratosphere and 1.0 K in the mesosphere (fig. 8 (c) and (d)) should be taken seriously.

It is known that the tropo-, strato-, and mesopause where the temperature gradient becomes zero and changes are challenging for approximation methods. The same holds for the beginning and the end of a data series. This becomes evident when a smooth profile like a CIRA-temperature profile is de-trended with different numbers of spline sampling points (see fig. 9 (a)–(d)). In these cases, the residuals show oscillations for both approaches, the repeating and the non-repeating spline which become smaller with decreasing distance between two spline sampling points. Position and strength of the oscillations change especially above 40 km height only slightly compared to fig. 7 (d) which shows the difference between the approximated background and the real one in the presence of typical gravity wave signatures (fig. 10 (a)). That means the non-optimal approximation of the smooth temperature background, i.e., without any gravity waves, is the most-likely reason for the oscillations observed in the mean SABER residuals for both approaches (see fig. 8 (c) and (d)).

However, for the non-repeating approach strength and position of the oscillation in the residuals (de-trended CIRA background) change when another starting height is chosen while the oscillation is only slightly shifted in the vertical for the repeating approach (fig. 10 (b) and (c)). For fig. 1, the starting height varied mostly in the range of ca. 2 km. A comparison of fig. 1 and fig. 8 (c) reveals that the height coordinates of the local extrema of the mean residuals corresponds approximately to the ones of the mean squared residuals. The less pronounced dependence of the repeating spline approach on the starting height (fig. 10 (b)) is therefore the most likely reason for the lower variance of the residuals (as described above).

Since the beginning and the end of the data series do not vary much (we analysed the SABER data between ca. 12–14 km and 110 km height), possible side effects might not cancel out. We speculate that these are at least two of the reasons for the superimposed oscillation in fig. 8 (c) and (d). However, analysing this hypothesis is beyond the scope of this manuscript.

There exist many methods to approximate / de-trend / filter time series (see e.g. Baumgarten et al. (2015), and references therein) and we do not claim that the presented repeating cubic spline is the best method for every purpose and every data series. It is just one possible algorithm which reduces disadvantages-artefacts of the non-repeating cubic spline routine as it was proposed by Lawson and Hanson (1974) if the data set contains waves with wavelengths of about double the sampling point distance which is mostly not known in advance-like the dependence of approximation on the exact position (height coordinate) of the spline sampling points. Furthermore, it reduces its dependence on the starting height. However, it comes along with enhanced computational effort which is of special importance when analysing large data sets.

5 Summary

It is essential for the analysis of atmospheric wave signatures like gravity waves that these fluctuations are properly separated from the background. Therefore, particular attention must be attributed to this step.

- 5 Cubic splines with equidistant sampling points are a common method in atmospheric science for the approximation of superimposed, large-scale structures in data series. The subtraction of the spline from the original time series allows the investigation of the residuals by means of different spectral analysis techniques. However, splines can generate artificial oscillations in the residuals—especially if the background is described by a coarse spline or if the data set contains waves with wavelengths of about double the sampling point distance—which must not to be interpreted in terms of gravity waves.
- 10 The ability of a spline to approximate ~~oscillations~~ the background state (and large-scale wave-induced fluctuations) varies not only with the number of spline sampling points, but also with their exact position.
When the distance between two spline sampling points equals exactly or approximately half the wavelength of the waves present in the data, the last mentioned effect is most pronounced.
- 15 Since knowledge about the wavelengths present in the data set is normally not available in advance, this directly motivates the use of a ~~repeating~~ spline which is based on changing starting points. It comes along with enhanced computational effort but it can be recommended for the approximation / de-trending of individual profiles and if squared residuals are needed for further analysis (e.g. for the calculation of the wave potential energy).

20

Acknowledgement

We would like to thank the TIMED-SABER team for their great work in providing an excellent data set.

We also thank the Bavarian Ministry for Environment and Consumer Protection for financially supporting our work: V.

- 25 Wendt was paid by the Bavarian project BHEA (Project number TLK01U-49580, 2010–2013). The work of S. Wüst was subsidised in parts by this project.

At least, we thank Julian Schmoeckel, formerly University of Augsburg, for helping to produce the test data sets and the figures.

30

Appendix A

Between two spline sampling points, a spline is equal to a cubic polynomial of the form

$$f(z) = az^3 + bz^2 + cz + d \quad \text{with} \quad a, b, c, d \in \mathbb{R}. \quad (2)$$

Its derivatives are

$$5 \quad f^{(1)}(z) = 3az^2 + 2bz + c, \quad (3)$$

$$f^{(2)}(z) = 6az + 2b, \quad (4)$$

$$f^{(3)}(z) = 6a. \quad (5)$$

Between two spline sampling points, the second derivative of a spline depends linearly on the height coordinate z . That means the curvature of the spline can change from negative to positive or vice versa between two spline sampling points but it can only increase or decrease linearly or it can stay constant. At the spline sampling points, all derivatives of the two adjacent polynomials must agree. For example, a spline cannot form two parabolas with different signs in two adjacent intervals in order to approximate a sine / cosine since the second derivative (curvature) would be positive constant in one interval and negative constant in the other. If the spline sampling points are not distributed in a way such that the curvature of the original function increases or decreases linearly between two spline sampling points, the spline cannot approximate the original function properly.

Therefore, the ability of the spline to reproduce a sine / cosine does not only depend on the number of spline sampling points, it varies also with their position.

Appendix B

The optimal spline parameters are determined through a least-square approach: depending on the spline parameters, the squared differences between the spline and the original data set are minimized. The maximum wavelength which a spline can approximate in principle is equal to two times the distance between two spline sampling points.

Let us denote the oscillation which has to be approximated with $f_1(z)$ and the spline with $f_2(z)$.

If those two oscillations which will be subtracted from each other are characterized by very similar wave numbers k_1 and k_2 , then a beat with the following wave numbers will occur.

$$25 \quad f_1(z) - f_2(z) = \sin k_1 z - \sin k_2 z \stackrel{(*)}{=} 2 \cos\left(\frac{k_1+k_2}{2} z\right) \sin\left(\frac{k_1-k_2}{2} z\right) \quad (6)$$

where

$\frac{k_1+k_2}{2}$ is the wave number of the beat, which is very similar to the original wave number, and


$\frac{k_1-k_2}{2}$ is the wave number of the envelope.

(*) : application of an addition theorem

References

- Baumgarten, G., Fiedler, J., Hildebrand, J., and Lübken, F.-J.: Inertia gravity wave in the stratosphere and mesosphere observed by Doppler wind and temperature lidar, *Geophys. Res. Lett.*, 42, 10,929–10,936, doi: 10.1002/2015GL066991, 2015.
- 5 Committee on Space Research; NASA National Space Science Data Center: COSPAR International Reference Atmosphere (CIRA-86): Global Climatology of Atmospheric Parameters. NCAS British Atmospheric Data Centre. <http://catalogue.ceda.ac.uk/uuid/4996e5b2f53ce0b1f2072adadaeda262>, 2006.
- Eckermann, S. D., Hirota, I., and Hocking, W. K.: Gravity wave and equatorial wave morphology of the stratosphere derived from long-term rocket soundings. *Q.J.R. Meteorol. Soc.*, 121, 149–186, doi: 10.1002/qj.49712152108, 1995.
- 10 Ern, M., Preusse, P., Gille, J. C., Hepplewhite, C. L., Mlynczak, M. G., Russell III, J. M., and Riese, M.: Implications for atmospheric dynamics derived from global observations of gravity wave momentum flux in stratosphere and mesosphere. *J. Geophys. Res.*, 116, doi: 10.1029/2011JD015821, 2011.
- Kramer, R., Wüst, S., and Bittner, M.: Climatology of convectively generated gravity waves at Prague based on operational radiosonde data from 13 years (1997–2009). *J. Atmos. Sol.-Terr. Phys.*, 140, 23–33, doi: 10.1016/j.jastp.2016.01.014, 15 2016.
- Krebsbach, M., and Preusse, P.: Spectral analysis of gravity wave activity in SABER temperature data, *Geophys. Res. Lett.*, 34, L03814, doi: 10.1029/2006GL028040, 2007.
- Lawson, C. L., and Hanson, R. J.: Solving least squares problems, Prentice-Hall, Inc., Englewood Cliffs, New Jersey, U.S.A. 1974.
- 20 Mzé, N., Hauchecorne, A., Keckhut, P., and Théti, M.: Vertical distribution of gravity wave potential energy from long-term Rayleigh lidar data at a northern middle-latitude site, *J. Geophys. Res. Atmos.*, 119, 12,069–12,083, doi: 10.1002/2014JD022035, 2014.
- Offermann, D., Gusev, O., Donner, M., Forbes, J. M., Hagan, M., Mlynczak, M. G., Oberheide, J., Preusse, P., Schmidt, H. and Russell III, J. M.: Relative intensities of middle atmosphere waves, *J. Geophys. Res.*, 114, D06110, doi: 10.1029/2008JD010662, 2009.
- 25 Shuai, J., Zhang, S., Huang, C., Yi, F., Huang, K., Gan, Q., and Gong, Y.: Climatology of global gravity wave activity and dissipation revealed by SABER/TIMED temperature observations. *Science China Technological Sciences*, 57, 998–1009, doi: 10.1007/s11431-014-5527-z, 2014.
- Wright, C. J., Rivas, M. B., and Gille, J. C.: Intercomparisons of HIRDLS, COSMIC and SABER for the detection of 30 stratospheric gravity waves, *Atmos. Meas. Tech.*, 4, 1581–1591, doi: 10.5194/amt-4-1581-2011, 2011.
- Wüst, S., and Bittner, M.: Gravity wave reflection: case study based on rocket data. *J. Atmos. Sol.-Terr. Phys.*, 70, 742–755, doi:10.1016/j.jastp.2007.10.010, 2008.

- Wüst, S., and Bittner, M.: Resonant interaction between two planetary waves with zonal wave number two? *J. Atmos. Sol.-Terr. Phys.*, 73, 771–778, doi:10.1016/j.jastp.2011.01.004, 2011.
- Wüst, S., Wendt, V., Schmidt, C., Lichtenstern, S., Bittner, M., Yee, J.-H., Mlynczak, M. G., and Russell III, J. M.: Derivation of gravity wave potential energy density from NDMC measurements. *J. Atmos. Sol.-Terr. Phys.*, 138–139, 32–46, doi:10.1016/j.jastp.2015.12.003, 2016.
- 5 Young, L. A., Yelle, R. V., Young, R., Seiff, A., and Kirk, D. B.: Gravity Waves in Jupiter's Thermosphere, *Science*, 276, 108–111, doi:10.1126/science.276.5309.108, 1997.
- Zhang, Y., Xiong, J., Liu, L., and Wan, W.: A global morphology of gravity wave activity in the stratosphere revealed by the 8-year SABER/TIMED data, *J. Geophys. Res.*, 117, D21101, doi:10.1029/2012JD017676, 2012.



5 Figure 1: Mean squared temperature residuals for the years 2010 to 2014 (colour-coded): they are derived from TIMED-SABER, data version 2.0 by using a cubic spline routine with equidistant sampling points for de-trending. The distance between two spline sampling points is 10 km. All vertical SABER temperature profiles which were retrieved between 44° N and 48° N and 5° O and 15° O are used (that means approximately 30-50 profiles per month and approximately 500 profiles per year).










Figure 2: This figure shows the approximation of a cubic spline using different numbers of spline sampling points:

- (a) A spline with a distance of 1.5 km between two spline sampling points is fitted (solid line) to the test data (dotted line).
- (b) Same as (a) but the distance between two spline sampling points is 1.6 km.
- (c) Same as (a) but the distance between two spline sampling points is 1.4 km.
- 5 (d) Same as (a) but restricted to the height range between 15 km and 50 km. The dashed-dotted lines refer to the height-coordinate of the spline sampling points. The asterisks show the sampling points of to the original sine.
- (e) Same as (b) but restricted to the height range between 15 km and 50 km.
- (f) Same as (c) but restricted to the height range between 15 km and 50 km.



5 Figure 3: This figure shows the differences between the spline and the approximated test data (solid line: phase of 0, dashed line: phase of $\pi/2$) which are summed up between 20 km and 40 km. They are plotted versus the distance between the number of spline sampling points / distance between spline sampling points. The number of spline sampling points / distance between spline sampling points refers to the whole height range between 15 km and 100 km.




Figure 4: Dependence of the sum of squared residuals on the phase of the wave with a wavelength of 3.0 km and a distance of 1.4 km (short dashes), 1.5 km (solid line) and 1.6 km (long dashes) between two spline sampling points.











Figure 5: Here, the results based on the repeating spline approach are shown. The different colours refer to the different spline approximations (to keep it as clear as possible, we only show the first four iterations, a fifth one exists for case (b) and (e), see step 4 of the algorithm). The black line represents the final spline approximation. The distance between two spline sampling points in part (a) to (f) agrees with the respective values in figure 2 part (a) to (f). While part (a) to (c) show the height range between 15 km and 100 km, part (d) to (f) focus on the height range between 15 km to 50 km. The asterisks have the same meaning as in figure 2




Formatiert: Englisch (Großbritannien)



(b)

Formatiert: Schriftart: 9 Pt., Fett




5


Figure 6: Part (a) is equivalent to figure 3, part (b) is equivalent to figure 4, but here the repeating spline approach is used.

10

(a)





Formatiert: Schriftart: Fett




(b)

Formatiert: Schriftart: Fett




(c)



Formatiert: Schriftart: Fett

(d)



Formatiert: Schriftart: Fett

Formatiert: Standard

Formatiert: Standard, Einzug: Erste
Zeile: 0,5 cm

Figure 7: a) The solid oscillating (non-oscillating) line depicts three sinusoids with vertical wavelengths of 3, 5, and 13 km, phase 0, $\pi/3$, and $\pi/5$, and amplitude 2.0 (the CIRA background) and. The dashed line shows the spline approximations for the repeating (black) and non-repeating (grey) spline approach (dashed line, sampling point distance 10 km). Part b) as part a) but focussing on the height interval between 20 and 40 km for which the sum of squared residuals is calculated in part c). The range of the x- and y-axis differs from the ones used in fig. 3 and 6(a). Part d) shows the difference between the spline (i.e., the approximated background, grey: non-repeating approach, black: repeating approach) and the CIRA-background (i.e. the original background). The range of the x- and y-axis differs from the ones used in fig. 3 and 6(a).






Figure 8: a) As figure 1, for de-trending the repeating cubic spline routine with equidistant sampling points as it is described in section 2 is used. b) Mean squared residuals for the repeating (dashed line) and the non-repeating (solid line) approach for the year 2014. c) Mean (non-squared) residuals for the *non-repeating* approach for the years 2010 to 2014, and d) mean (non-squared) residuals for the *repeating* approach.



Formatiert: Schriftart: Fett



Formatiert: Schriftart: 10 Pt., Nicht Fett

Formatiert: Schriftart: Fett

Figure 9: (a) Original CIRA-temperature profile (which was de-trended between ca. 14 and 110 km as the SABER data), (b)–(d) residuals for a sampling point distance of 2.5 km, 5 km, and 10 km between 20 and 100 km height (grey: non-repeating spline, black: repeating spline).

Formatiert: Zeilenabstand: einfach

Formatiert: Schriftart: 9 Pt.

Formatiert: Schriftart: 9 Pt.

Formatiert: Schriftart: Fett

Formatiert: Schriftart: 9 Pt.

Formatiert: Schriftart: Fett

Formatiert: Schriftart: 9 Pt.

Formatiert: Schriftart: 9 Pt.




Figure 10: (a) Superposition of fig. 9 (d) (dashed lines, residuals after de-trending the CIRA temperature background without superimposed gravity waves) and fig. 7 (d) (solid lines, difference between the original and the approximated background in the presence of gravity waves), (b) and (c): residuals of the repeating (black) and non-repeating (grey) spline approximation (max. wavelength: 20 km) of the CIRA temperature profile de-trended between varying starting heights (15–20 km) and 115 km.