

We thank the reviewer for carefully reading the revised manuscript and for providing further suggestions. Please find below our answers and responses to the follow-up questions and suggestions (printed in *italic*). Corresponding adjustments made on the text in the manuscript are shown in green.

1. *In my opinion, Figure 1 has been over simplified and no longer fulfills its purpose of aiding the reader to piece the different bits of the retrieval together. I suggest using the flowchart to emphasize the key assumptions and intricacies of the retrieval (e.g. smoothing LWC at cloud base, choosing which assumptions are applied). As the retrieval is quite complex, a good flowchart will go a long way to guiding the reader through the method section.*

Following the comments of the second reviewer in the first round, we simplified the flowchart in an attempt to make it more understandable and to emphasize the overall structure of the retrieval. We have now revised Fig. 1 to show more details as currently requested by the reviewer, while maintaining as few distractions as possible from the overall flow. In the new Fig. 1 in the manuscript, we have added the following: the process of determining cloud boundaries, the procedure to determine drizzle reflectivities, and the branching into case I and case II. For each of these, we put the reference to the relevant section of the text to guide the reader. The text describing the flowchart on page 3 has been adjusted accordingly.

2. *Performance of the retrieval in Case I, where drizzle is detected only above cloud base. I am worried that there is not enough information to constrain both drizzle and cloud properties. How sensitive are the forward models to the first guess of drizzle scale factor? For example if the drizzle LWC is doubled or halved in CASE I of Fig 6., and all other settings of the retrieval remain unchanged, does the retrieval give satisfactory results?*

We recognize the reviewer’s concern about the information content in Case I. We mentioned in the manuscript that our retrieval strategy for Case I is adversely limited due to this (page 9 line 14). We tried to exploit the signal but the little amount of information prompted us to adopt a simple retrieval scheme for the drizzle, i.e. using the cloud LWC parametrization with a scaling factor. Here, our goal is to get an order-of-magnitude estimate of the drizzle quantities, rather than to achieve a high accuracy.

We have modified the synthetic signals in Case I of Fig. 6 as requested, i.e. by varying the drizzle LWC. Changing drizzle LWC consequently changed its r_e , N and α while drizzle Z was kept the same. We also kept the retrieval setting identical to the case I retrieval shown in Fig. 6 (e.g. same random seed that determines the retrieval starting point or ‘initial guess’). Fig. A below shows the retrieval results for the two requested LWC profiles, i.e. half and twice the original LWC. Fig. B and C expand on this and display the retrieval results when drizzle LWC is decreased or increased by a factor of 3 and 4, respectively. The layout and legend of those figures are the same as in Fig. 6.

In Fig. A to C, the lidar backscatter and MWR signals (both truth and retrieval) show no discernible change due to the modifications in LWC_{dzt} . The cloud properties are robustly recovered in all cases, within the expected level of uncertainties. Fig. A to C also show that changing the drizzle true properties changes the retrieval results as well, despite identical retrieval starting points, cloud properties and reflectivities. This means that the modifications in drizzle LWC can be tracked by the retrieval procedure through the lidar and MWR signals. Similarly to what is shown in Fig. 6, the drizzle reflectivities in Fig. A-C are better retrieved in the lower half than in the upper half of the cloud. Drizzle LWC, extinction coefficient and number concentrations are all retrieved within one order of magnitude from the truth. The drizzle r_e is admittedly poorly recovered in the 0.25x and 0.33x LWC_{dzt} runs, where the retrieved r_e spans almost the entire radius range allowed in the retrieval, but it gets visibly better constrained for cases with large LWC. This is because the true extinction coefficient gets larger as the true LWC is increased, leaving stronger traces of drizzle in the lidar signal that help constrain the drizzle retrieval. For this reason, drizzle properties are better retrieved for larger LWC_{dzt} than for smaller LWC_{dzt} .

To give a more complete picture of how the starting point of the state vector could influence the retrieval, we performed three retrievals using different random seeds. These runs are based on the

Case I retrieval shown in Fig. 6. The input and settings of the runs are all identical except for the random seed used by the differential evolution routine. The results are shown in Fig. D, which is essentially a copy of the left part of Fig. 6 (case I) with three additional profiles resulting from the different random seeds/starting points (shown in magenta, green and yellow). They all confirm the results from Fig. A-C. The retrieved clouds match the truth with an accuracy consistent with what is expected from the exercise in section 3.1. The retrieved drizzle LWC, extinction coefficient and number concentration give a good order-of-magnitude estimate of the truth with uncertainties of about 50% for the drizzle r_e retrieval.

We have added a few lines in section 3.2 under 'Case I' in the manuscript to note the limited information content and our expectations for the case I retrieval. If the reviewer would like us to elaborate on it, we can show the discussion and the figures here either in section 3.2 or as a separate appendix in the manuscript.

3. *What causes the kink in retrieved drizzle properties at 0.5 km in CASE II of Fig. 6?*

We mentioned the 'kink' as a 'dip' in the manuscript (page 18 line 9). The kink at 530 m originates from the drizzle Z . Since the drizzle r_e is parametrized, the kink does not appear here. Instead, the smoothness of r_e profile propagates the kink to the drizzle LWC, α and N profiles.

This dip in drizzle Z comes about because the cloud reflectivity between 500 and 600m is not well recovered (the effect is most visible in panels IIe and IIg in Fig. 6). More specifically, the retrieved cloud reflectivities at these heights are slightly larger than the true ('observed') Z , resulting in $Z_{\text{excess}} = 0$ (eq. 19). The smoothing that is applied afterwards to produce Z_{drl} (see text below eq. 19) replaces the zeros with non-zero values that are smaller than the neighbouring range gates, creating the kink.

We have added this additional explanation in section 3.2 for completeness.

4. *How sensitive are the results of drizzle below cloud base to the lidar backscatter? In CASE II of Fig. 6, the lidar backscatter below cloud base is identical or at least very similar to the clear air backscatter (CASE I Fig. 6). Perhaps a greater LWC of drizzle would allow the synthetic drizzle to be detectable above the molecular backscatter.*

The drizzle results below the cloud base are very sensitive to the lidar backscatter. The drizzle retrieval scheme below the cloud base relies on inferring the magnitude of drizzle extinction coefficient from the lidar backscatter. Noticeable enhancement of backscatter values (on top of the clear air/molecular backscatter) below the cloud base provides a strong indication that the drizzle extinction coefficient is comparable to or larger than the air extinction coefficient. Conversely, when the lidar backscatter appears identical or very similar to the molecular backscatter, one can be sure that the drizzle extinction is significantly smaller than the molecular extinction (drizzle extinction can never be zero as long as Z is detected below the cloud base). In Case II of Fig. 6, the drizzle extinction coefficient amounts to 5% of the molecular extinction coefficient and contributes very little to the lidar signal.

Larger amounts of drizzle would certainly lead to a more detectable drizzle backscatter. To show this, we modified the synthetic drizzle from Case II, i.e. we increased LWC by approximately a factor of 10 and decreased r_e by almost a factor of 2. This results in drizzle extinction coefficient being almost 20 times larger. One can see the effect on the lidar backscatter signal between 200 m and the cloud base. Fig. E shows the retrieval results for the modified signals. Discrepancies between the retrieval and the truth are largest close to the cloud top where the drizzle is least constrained. The cloud retrievals differ slightly from case II Fig. 6, which leads to the differences in the drizzle retrievals above the cloud base. The drizzle properties below the cloud base are more accurately retrieved now, as expected, due to the stronger drizzle signal in the lidar backscatter.

We have added some text and reorganized the paragraphs under 'Case II' in section 3.2 to address this point.

5. *There is too much discussion of the retrieval in cloud only mode (Sect. 3.1). I suggest removing Figure 3,4 and 5, replacing them with a Case 0: cloud only set of panel plots, similar to those in Fig 6. I think this will make the paper more concise and focused on the problem at hand.*

Section 3.1 is dedicated to test and verify the forward models and the cloud LWC model. Here, we use simulated signals where drizzle is absent in the LES scene and is therefore not considered in the retrieval. We view the discussion in section 3.1 as a necessary element and as a basis for discussing the drizzle retrieval that follows. Making sure that the cloud retrieval is robust is crucial especially because the drizzle retrieval depends on and derives from that.

Fig. 3, 4 and 5 allow the reader to visually assess the 'time' evolution and stability of the retrieval. This is an important aspect considering that the retrieval is performed on a column-by-column basis. Replacing them with a set of panel plots similar to Fig. 6 would defeat that purpose. Furthermore, the comparison of the true and retrieved cloud properties for a single profile is already shown in Fig. 6. Since that comparison is representative of the results in section 3.1 as well, it would be redundant to repeat such plots.

We think that categorizing the exercise in section 3.1 as 'Case 0' would be misleading. The actual retrieval scheme as depicted in Fig. 1 does not offer the possibility to ignore/dismiss drizzle prior to the retrieval (as done in Section 3.1), i.e. drizzle is always assumed to be present at the beginning until the optimization process decides otherwise.

For the reasons above, we prefer to keep section 3.1 as it is. However, if the reviewer still feels strongly about this particular point, we could proceed to place some of the text and the figures in the appendix.

6. *Pg 9, ln 32 The extinction coefficient, effective radius, k_1 and k_2 are all dependent on each other, so the choice of which to include in the state vector is arbitrary. I suggest removing this sentence.*

The sentence has been removed.

7. *Pg 17 ln 13 is too informal*

The sentence has been improved to include a more quantitative evaluation.

8. *Pg 18 ln 12 Surely a thorough error analysis is exactly within the scope of the paper? I suggest removing this sentence.*

The sentence has been removed.




Figure A: Case I retrievals for the modified drizzle LWC. Panels Ia-II show the results where the true drizzle LWC is halved while panels IIa-II correspond to the run where the true drizzle LWC is doubled. The red dash-dotted lines mark the cloud base height $z_{cb,opt}$. Filled circles: synthetic signals, squares: the cloud truth, diamonds: drizzle truth; red solid lines: retrieved signals, blue dashed lines: retrieved cloud properties and blue dotted lines: retrieved drizzle properties.




Figure B: Case I retrievals for the modified drizzle LWC. Panels Ia-II show the results where the true drizzle LWC is multiplied by one third while panels IIa-III correspond to the run where the true drizzle LWC is tripled. Legend: see the caption of Figure A.




Figure C: Case I retrievals for the modified drizzle LWC. Panels Ia-II show the results where the true drizzle LWC is multiplied by one fourth while panels IIa-III correspond to the run where the true drizzle LWC is increased by a factor of four. Legend: see the caption of Figure A.




Figure D: Case I retrievals with different random seeds. Panels Ia-Ij are a reproduction of the same panels in Fig. 6 with the additional green, magenta and yellow lines displaying the retrieval results using different random seeds. Legend: see the caption of Figure A.




Figure E: Case II retrievals. Panels Ia-II are a copy of the same panels in Fig. 6, repeated here for comparison. Panels IIa-III show the retrieval results where the drizzle extinction coefficient is increased by a factor of ~ 20 . Legend: see the caption of Figure A.

Simultaneous and synergistic profiling of cloud and drizzle properties using ground-based observations

Stephanie P. Rusli^{1,2}, David P. Donovan², and Herman W. J. Russchenberg¹

¹Department of Geoscience and Remote Sensing, Faculty of Civil Engineering and Geosciences, TU Delft, Delft, The Netherlands

²Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

Correspondence to: S. P. Rusli (s.rusli-1@tudelft.nl)

Abstract. Despite the importance of radar reflectivity (Z) measurements in the retrieval of liquid water cloud properties, it remains non-trivial to interpret Z due to the possible presence of drizzle droplets within the clouds. So far, there has been no published work that utilizes Z to identify the presence of drizzle above the cloud base in an optimized and a physically-consistent manner. In this work, we develop a retrieval technique that exploits the synergy of different remote sensing systems to carry out this task and to subsequently profile the microphysical properties of the cloud and drizzle in a unified framework. This is accomplished by using ground-based measurements of Z , lidar attenuated backscatter below as well as above the cloud base, and microwave brightness temperatures. Fast physical forward models coupled to cloud and drizzle structure parametrization are used in an optimal estimation type framework in order to retrieve the best-estimate for the cloud and drizzle property profiles. The cloud retrieval is first evaluated using synthetic signals generated from large-eddy simulation output to verify the forward models used in the retrieval procedure and the vertical parametrization of the liquid water content. From this exercise it is found that, on average, the cloud properties can be retrieved within 5% of the mean truth. The full cloud-drizzle retrieval method is then applied to a selected ACCEPT campaign dataset collected in Cabauw, The Netherlands. An assessment of the retrieval products is performed using three independent methods from the literature, each was specifically developed to retrieve only the cloud properties, the drizzle properties below the cloud base, or the drizzle fraction within the cloud, respectively. One-to-one comparisons, taking into account the uncertainties or limitations of each retrieval, show that our results are consistent with what is derived using the three independent methods.

1 Introduction

Low-level liquid water clouds are known to have a large areal extent (Hartmann et al., 1992) and consequently a strong impact on the Earth's energy balance (Ramanathan et al., 1989; Slingo, 1990). Observations of these clouds to characterize the microphysical and radiative processes are therefore needed for climate studies. One important aspect of such observations is the presence of drizzle, which is found to be a common occurrence in stratocumulus clouds (Fox and Illingworth, 1997). Drizzle alters the cloud droplet spectra and thus the microphysical structure and radiative properties of the clouds (Feingold et al., 1997; vanZanten et al., 2005). Most notably, drizzle is thought to play a significant role in determining the cloud lifetime (Albrecht, 1989). Additionally, drizzle within the cloud complicates matters by dominating the radar reflectivity signal. Accurately

separating the drizzle contribution from the cloud contribution to the received radar signal is necessary to properly derive the cloud and drizzle properties.

Since liquid water clouds tend to **form** at relatively low altitudes in the atmosphere, it is easier to observe them from the surface than from space. Ground based remote sensing systems have the potential to deliver high-resolution time-series data to
5 evaluate and monitor cloud and drizzle properties on a regional scale. A synergistic way in utilizing different remote sensors is a powerful approach that has been widely used to provide a more complete and comprehensive view of these clouds. Active sensors operating in different frequency windows such as radar and lidar provide complementary information on the clouds vertical structure since they 'see' different parts of the cloud (Donovan and van Lammeren, 2001). Microwave radiometers that measure the accumulated radiation along a column provide a particularly accurate way to derive the liquid water path of clouds
10 (Westwater 1978; Peter and Kämpfer 1992).

Various methods that exploit sensor synergy to profile microphysical properties of the liquid water cloud have been developed (Frisch et al. 1995a; Austin and Stephens 2001; McFarlane et al. 2002; Löhnert et al. 2001; Brandau et al. 2010; Martucci and O'Dowd 2011). However, these methods either avoid, ignore or do not fully capture the presence of drizzle. Other techniques that focus on drizzle retrieval are limited in their application to the region below the cloud base (O'Connor et al., 2005;
15 Westbrook et al., 2010). Retrieving the properties of drizzle that is interspersed within the cloud is indeed more difficult. A few hundred meters into the cloud, lidar backscatter signal no longer carries useful information due to the strong attenuation by cloud droplets. While radar has the capability to penetrate deeper into the cloud, the radar reflectivity is known to be sensitive to drizzle droplets that are larger in size as compared to the cloud droplets. Since the observed reflectivity contains contributions from both cloud and drizzle droplets, its interpretation is not straightforward.

Fielding et al. (2015) set a precedence by jointly retrieving cloud and drizzle properties using ground-based radar, lidar and sun photometer observations. **Their** retrieval departs from the assumption that drizzle is present only when the maximum observed reflectivity in a given column exceeds a single threshold value. While the existence of such a reflectivity threshold is supported by many observational studies, the empirically-determined value differs among these studies and can span quite a wide range. Sauvageot and Omar (1987); Frisch et al. (1995b); Mace and Sassen (2000) suggest different Z thresholds in the
25 range of -20 and -15 dBZ. Baedi et al. (2002) showed that the reflectivity due to a non-drizzle component of the cloud reaches a maximum at about -20 dBZ while that of the drizzle component does not get lower than about -10 dBZ, leaving on average a ~ 10 **dB** reflectivity gap between the drizzle-contaminated and drizzle-free droplet spectrum. Furthermore, Wang and Geerts (2003) demonstrate that the value of this threshold varies with altitude within the cloud layer and it can increase from around -25 dBZ near the cloud base to about -12 dBZ close to the cloud top. A theoretical approach by Liu et al. (2008) reveals a
30 dependence of the threshold value on the droplet number concentration, a finding that compares favorably with observations. In remote sensing applications, where droplet concentration is one of the unknown variables to retrieve, setting a single Z threshold value in advance may lead to an unaccounted bias in the retrieval.

In this work we develop a retrieval technique that combines ground-based radar, lidar and microwave radiometer (MWR) measurements to simultaneously profile the cloud and drizzle properties without placing apriori constraints on the presence
35 of drizzle droplets within the cloud. There is no predefined reflectivity threshold and so drizzle is always assumed to be

present until that possibility is excluded by the best-fit to the data. The MWR brightness temperature and the lidar attenuated backscatter up to a few hundred meters above the cloud base provide much of the constraint on the cloud component, whereas the radar reflectivity is used to then infer the drizzle contribution. Drizzle droplets are set apart from the cloud droplets through the use of a critical effective radius threshold in the algorithm. This choice of threshold is motivated by the recognition that a characteristic or critical droplet radius exists, above which intense droplet coalescence triggers rapid drizzle formation. This radius is found to be 12-14 microns as shown by satellite and ground-based observations (Suzuki et al., 2010; Rosenfeld, 2000; Rosenfeld and Gutman, 1994), aircraft measurements (Gerber, 1996; Boers et al., 1998; Freud and Rosenfeld, 2012) and numerical simulations (Magaritz et al., 2009; Pinsky and Khain, 2002; Benmoshe et al., 2012). This retrieval technique allows us to retrieve not only drizzle microphysical properties below the cloud base but also within the cloud at the same time. We apply this algorithm to synthetic signals for a test case, as well as to observational data collected in the Fall of 2014 as part of the ACCEPT field campaign in Cabauw, The Netherlands. The retrieved cloud and drizzle products from the ACCEPT dataset are evaluated against the results of three independent retrieval methods that use the lidar depolarization signal (Donovan et al., 2015), the lidar attenuated backscatter and radar Doppler spectral moments (O'Connor et al., 2005), and the radar Doppler spectra (Kollias et al., 2011a, b; Luke and Kollias, 2013) as their main tools.

The remainder of this paper is organized as follows. Section 2 describes the retrieval procedure in detail, including the theoretical assumptions and the forward models. The test application of this technique to synthetic data based on the large-eddy simulation output is presented in Section 3. In Section 4, we perform the cloud and drizzle retrieval on a ground-based dataset. The retrieval products are then evaluated through comparisons with results from three independent retrieval techniques in Section 5. To conclude the paper, a summary is provided in Section 6.

2 Retrieval technique

The target group for this retrieval technique is single-layered liquid water clouds. Retrieval is not performed when rain is detected on the ground. The retrieval addresses an inverse problem of deriving the cloud and drizzle profiles that give rise to the observed radar reflectivity Z , lidar attenuated backscatter β and microwave brightness temperatures T_B . The end products include the optical extinction coefficient, liquid water content, number concentration and the effective radius of both the cloud and the drizzle components separately, as a function of height.

The overall structure of the retrieval method is depicted in a flowchart in Fig. 1. This flowchart shows the main components of the algorithm, accompanied by references to the relevant sections. Here we give a general overview of how the algorithm works. The state vector refers to the collection of control parameters that we aim to optimize in order that our forward models match the observations. These control parameters are used to construct cloud and drizzle profiles through the parametrization of the droplet size distributions and their vertical structures outlined in section 2.1. The algorithm starts with determining the cloud boundaries and calculating the cloud microphysical properties (section 2.2.1). It then continues to compute drizzle reflectivities and it proceeds to derive the microphysical properties of drizzle by following one of the two available schemes, depending on whether drizzle is detected below the cloud base (section

2.2.2). If the cloud and drizzle properties satisfy the physical constraints set out in section 2.2.3, the forward models are computed for each instrument (sections 2.2.4-2.2.6). These forward models map the theoretical construction of the cloud and drizzle to the observables, i.e. create simulated observations. The cost function quantifies the difference between these predicted observations and measured signals. The optimization process to find the best fit solution (section 2.3) to the inverse problem seeks to minimize the cost function. It is worth noting here that this optimization process also accounts for important factors such as lidar calibration uncertainty and the changes in the cloud base height within the measurement temporal resolution.

2.1 Theoretical basis and parametrization

2.1.1 Cloud and drizzle droplet size distribution

We treat the cloud and the drizzle droplets as two separate entities and they are assigned independent and unimodal droplet size distribution (DSD) functions, the combination of which results in a bimodal distribution. Here we assume that the number density of the cloud or drizzle droplets as a function of their size can be described by the a generalized gamma distribution (Walko et al., 1995):

$$n(r) = \frac{N}{r_n \Gamma(\nu)} \left(\frac{r}{r_n} \right)^{\nu-1} \exp\left(\frac{-r}{r_n}\right), \quad (1)$$

where $N = \int_0^\infty n(r) dr$ is the total number concentration, r_n is the droplet characteristic radius and ν is the shape parameter.

The moments of this DSD,

$$\langle r^k \rangle = \frac{\int_0^\infty r^k n(r) dr}{\int_0^\infty n(r) dr},$$

are central in defining and deriving the physical properties of the cloud and the drizzle as listed below.

- Effective radius r_e :

$$r_e = \frac{\langle r^3 \rangle}{\langle r^2 \rangle} = r_n(\nu + 2). \quad (2)$$

- Extinction coefficient α :

$$\alpha = \int_0^\infty Q_{\text{ext},\lambda}(r) \pi r^2 n(r) dr \approx 2\pi N \langle r^2 \rangle, \quad (3)$$

where the extinction efficiency $Q_{\text{ext},\lambda}(r) \approx 2$ assuming that the droplets are much larger than the wavelength λ of the incident light.

- Liquid water content LWC:

$$\text{LWC} = \frac{4}{3} \pi \rho_w N \langle r^3 \rangle. \quad (4)$$

In addition, the sixth moment of the distribution function delivers the radar reflectivity factor Z by virtue of Rayleigh approximation, which is valid in the case of scattering of particles whose size is small compared to the radar wavelength. The exact expression is:

$$Z = 2^6 \int_0^\infty r^6 n(r) dr = 64N \langle r^6 \rangle. \quad (5)$$

5 Moreover, the moments of the DSD are assumed to be related to each other such that:

$$\langle r^a \rangle = k_{ab} \langle r^b \rangle^{a/b},$$

in which k_{ab} is a function of shape parameter ν . Using the property of the gamma function we derive, for instance,

$$k_{23}^3 = \frac{\nu(\nu+1)}{(\nu+2)^2}, \quad (6)$$

$$k_{36}^2 = \frac{\nu(\nu+1)(\nu+2)}{(\nu+3)(\nu+4)(\nu+5)}, \quad (7)$$

10 that allow one to relate LWC to α (k_{23}) or Z (k_{36}).

2.1.2 Cloud structure

To profile the cloud, we adopt an approximation for LWC vertical profile introduced in Boers et al. (2006). Here we repeat what is necessary and adjust some of the notation.

Near the cloud base, LWC is assumed to vary linearly with height (i.e. constant lapse rate). Deeper into the cloud, entrainment
15 leads to a decrease in the LWC lapse rate. **The LWC at a given height above cloud base \tilde{z} is related to its adiabatic value through a subadiabatic fraction $f(\tilde{z})$ such that:**

$$\text{LWC}(\tilde{z}) = f(\tilde{z}) \text{LWC}_{\text{ad}}(\tilde{z}) = f(\tilde{z}) \rho_a A_{\text{ad}} \tilde{z}. \quad (8)$$

ρ_w is the density of water and LWC_{ad} the adiabatic LWC. ρ_a and A_{ad} are the density of air and the adiabatic lapse rate of the liquid water content mixing ratio, respectively; both are a function of the temperature and pressure at the cloud base. The
20 subadiabatic fraction changes as a function of height and is governed by two variables W and H :

$$f(\tilde{z}) = \left[1 - \exp(-W\hat{h}) \right] \left[1 - \frac{\exp(-\hat{h}(1-\hat{z}))}{1 - \exp(-\hat{h})} + \frac{\exp(-\hat{h})}{1 - \exp(-\hat{h})} \right], \quad (9)$$

with $\hat{h} = x/H$ and $\hat{z} = \tilde{z}/x$. **x is cloud depth**, W represents the vertical weight of the liquid water distribution and the relaxation length scale H indicates how much the liquid water content departs from adiabaticity. The smaller W is, the more liquid water there is close to the cloud top. The smaller H is, the closer the actual LWC becomes to the adiabatic profile.

25 Boers et al. (2006) consider two mixing scenarios to describe the vertical variation in $f(\tilde{z})$, namely inhomogeneous and homogeneous mixing. Both scenarios are implemented in the retrieval algorithm so the user can decide which one to use. In

the first one, the variation of $f(\tilde{z})$ is attributed to the vertical change in N :

$$N(\tilde{z}) = f(\tilde{z})N_{\text{ad}} \quad (10)$$

where N_{ad} is the adiabatic value of N . In the homogeneous mixing case, evaporation causes the droplet sizes to decrease while preserving the total number of droplets:

$$5 \quad N(\tilde{z}) = N_{\text{ad}} \quad (11)$$

For the retrieval in this paper, we assume for simplicity that N is constant with height. This assumption corresponds to the homogeneous mixing case (eq. 11).

2.1.3 Drizzle structure

The drizzle signature is strongly imprinted in radar reflectivity measurements, making Z indispensable for drizzle retrieval. Owing to the proportionality between the moments of the DSD, the observed reflectivity is related to the drizzle microphysical properties and the vertical shape of Z can be used to profile drizzle. This is especially true below the cloud base where drizzle is isolated from the cloud and Z is related to drizzle alone.

At the very early stages of drizzle formation, when drizzle is still contained within the cloud and there are no detected drizzle droplets falling from the cloud, the analogy of eq. 9 is used to also describe how drizzle LWC varies with height. Using equations 4, 5 and 7, r_e can then be written in terms of ν , Z and LWC:

$$r_e^3 = \frac{\pi \rho_z Z}{48 \text{LWC}} \frac{(\nu + 2)^3}{(\nu + 3)(\nu + 4)(\nu + 5)} \quad (12)$$

from which N and α can be computed.

As drizzle starts to grow and leave the cloud, we use a different drizzle parametrization. The vertical profile of the drizzle effective radius above the cloud base is parametrized as a function of height z via an exponential function:

$$20 \quad r_e(z) = r_{e,\text{cb}} \exp \left(\frac{k_1 (z - z_{\text{cb}})}{z_{\text{dt}} - z_{\text{cb}}} \right)^{-0.5}. \quad (13)$$

$r_{e,\text{cb}}$ is the value of drizzle effective radius at cloud base, k_1 describes the rate at which r_e decreases towards cloud top. The subscript 'dt' denotes drizzle top and 'cb' cloud base. The choice of such an exponential function is motivated by the results of in-situ drizzle measurements showing that within the cloud, the drizzle effective radius displays an exponential-like increase towards the cloud base (Wood, 2005a; Lu et al., 2009).

Below the cloud base, drizzle droplets are not expected to keep growing. Instead, they are assumed to evaporate and shrink. In this region, the parametrization of r_e is based on a simple power-law:

$$r_e(z) = r_{e,\text{cb}} \left(\frac{z - z_{\text{db}}}{z_{\text{cb}} - z_{\text{db}}} \right)^{k_2}, \quad (14)$$

with k_2 describing the rate at which r_e decreases from the cloud base to the drizzle base (denoted by the subscript 'db').

In the retrieval, the two parameters k_1 and k_2 are positive and are solved using values of r_e at three different heights: below, at and above the cloud base (see Section 2.2). The droplet size information at the cloud base is crucial since it acts as a scaling factor and is the point where these two functions meet. Once the vertical profiles of ν , Z and r_e are specified, one can derive LWC, α and N as a function of height.

5 2.2 Retrieval scenario

Following the retrieval flowchart in Fig. 1, we address each of the main steps here. There is a total of 12 elements in the state vector: 7 for the cloud component (sec 2.2.1), 4 for the drizzle component (sec 2.2.3), and one element to compensate for a possible offset in the lidar signal due to imperfect calibration (sec 2.2.6). These state vector elements are used throughout the algorithm and their roles are explained below; we mark these 12 parameters with an asterisk(*) to help the reader distinguish

10 them from other variables.

2.2.1 Cloud profiles

The profile of cloud LWC is constructed according to eq. 8, using the subadiabatic fraction $f(\tilde{z})$. $f(\tilde{z})$, as formulated in eq. 9, is a function of relative height and depends on three variables: the vertical weight, the relaxation length scale and the cloud depth. Thus, specifying the cloud LWC profile requires us to determine four parameters: W_{cld}^* , \hat{h}_{cld}^* , cloud base height $z_{\text{cb,opt}}$ and cloud top height $z_{\text{ct,opt}}$. We use lidar and radar observations to get estimates of the cloud base and top altitudes and let the exact locations be optimized in the algorithm. This way, the limited range resolution of the radar and lidar is taken into account in the model. Variables ft_{cb}^* and ft_{ct}^* are employed to serve this purpose, that is to allow the model cloud boundaries to be located at any spatial point within a given range.

ft_{ct}^* is used to optimize the cloud top height $z_{\text{ct,opt}}$ based on radar measurements, such that:

$$20 \quad z_{\text{ct,opt}} = z_{\text{ct}} + ft_{\text{ct}}^* \Delta z_r. \quad (15)$$

Δz_r is the range resolution of the radar and z_{ct} is the height of the radar range gate above the last radar detection. The value of ft_{ct}^* is restricted to be in the range [-1,0].

The cloud base location is estimated from the lidar attenuated backscatter (β) profile. There is an added complication that the measured lidar signals appear to have suffered from spatial broadening around the cloud base: the attenuated backscatter increases towards the peak value more mildly than theoretically expected for clouds with sharp, well defined boundary, suggesting a somewhat gradual increase in extinction coefficient. This is also true for cases where there is no radar detection below the cloud base. Possible causes for this may include turbulence, entrainment and changes in cloud base heights within the measurement temporal resolution that blur the cloud boundary, creating a transitional region seen in the lidar signal. This effect is comparatively small in magnitude with respect to the maximum backscatter (see Fig. 2) but would certainly affect the determination of cloud base height and drizzle quantification. We account for this effect by smoothing the cloud LWC around the cloud base, as described in the following paragraphs. Here we first show that this smoothing helps to achieve a better fit. Fig. 2 compares the profiles of the forward-modelled β with and without LWC smoothing against observational data

(circles). Without smoothing, the best-fit (forward-modelled) lidar backscatter is amplified strongly within 2 range gates (30 m) before it gets attenuated (dashed line). In the height range between 1.20 and 1.26 km, the relative error at the signal peak is the smallest, leading to the good fit at this range gate at the expense of the much worse fit at the earlier range gates. The cloud base height inferred from the model β appears to be an overestimate and at this height β is underestimated by more than an order of magnitude. When LWC is smoothed around $z_{\text{cb,opt}}$, the fit improves significantly as shown by the solid line.

To optimize the cloud base height $z_{\text{cb,opt}}$, we first find the lidar estimate of cloud base height (z_{cb}) as the first-order approximation. The lidar attenuated backscatter profile is searched to find the height where β reaches its maximum value: z_{peak} . We then identify as z_{cb} the lowest range gate where β rises by more than 50% to the next range gate and at the same time shows a continuous increase from there on up to z_{peak} . $z_{\text{cb,opt}}$ is located somewhere between z_{cb} and z_{peak} , such that:

$$z_{\text{cb,opt}} = z_{\text{cb}} + ft_{\text{cb}}^*(z_{\text{peak}} - z_{\text{cb}}), \quad (16)$$

where ft_{cb} is constrained to the range (0,1). In practice, when the broadening effect is clearly larger than the lidar range resolution then one can set the possible range of ft_{cb}^* to [0,1] and write:

$$z_{\text{cb,opt}} = z_{\text{min}} + ft_{\text{cb}}^*(z_{\text{max}} - z_{\text{min}}). \quad (17)$$

z_{min} and z_{max} are $z_{\text{cb}} + \Delta z_l$ and $z_{\text{peak}} - \Delta z_l$, respectively, with Δz_l denoting the lidar range resolution.

Once the cloud LWC profile is set up, the smoothing is applied to the region around $z_{\text{cb,opt}}$ via the centered moving average scheme. The width of the smoothing window is $2n+1$ where n is the number of lidar range gates between z_{cb} and $z_{\text{cb,opt}}$. The LWC values within the smoothing window are weighted as $\exp(-p_{\text{cb}}^*d)$. p_{cb}^* acts as a coefficient of the exponential weight and is part of the state vector, whereas d is the distance in the unit of range gates, such that $d=0$ for the central value, $d=1$ for the values next to it, and so on. The smoothing is performed only up to $n+1$ gates above $z_{\text{cb,opt}}$. Above this height, the impact of the smoothing is insignificant: as LWC_{cld} increases up to the peak value in an approximately linear fashion, the effect of the smoothing quickly diminishes with height.

After the smoothed LWC profile is available, N_{ad}^* and ν_{cld}^* (both are assumed to be constant with height) given in the state vector can be used to derive profiles of the other cloud properties, i.e. the cloud droplet number concentration (eq. 10 or 11), the extinction coefficient (eq. 3) and the effective radius (eq. 2). In total, seven variables in the state vector are used to construct the profiles of cloud properties: ν_{cld}^* , W_{cld}^* , \hat{h}_{cld}^* , N_{ad}^* , ft_{cb}^* , ft_{ct}^* , p_{cb}^* .

2.2.2 Drizzle profile

From the properties derived in the previous section, the radar reflectivity of the cloud component Z_{cld} can be computed (eq. 5). The difference between Z_{cld} and the observed reflectivity Z_{obs} is recorded as Z_{excess} , such that:

$$Z_{\text{excess}} = Z_{\text{obs}} - Z_{\text{cld}} \quad \text{for } Z_{\text{obs}} > Z_{\text{cld}} \quad (18)$$

$$= 0.0 \quad \text{for } Z_{\text{obs}} \leq Z_{\text{cld}}. \quad (19)$$

To enforce some level of spatial continuity for the drizzle above z_{cb} , the resulting Z_{excess} is smoothed. For simplicity, we apply a simple moving average to Z_{excess} values within cloud boundaries using three range gates as the smoothing width to result in drizzle reflectivity Z_{dzl} . Below the cloud base, $Z_{\text{dzl}} = Z_{\text{excess}} = Z_{\text{obs}}$.

At this point, we can distinguish two types of profiles: with and without radar detection below the cloud base. While radar signal below the cloud base is an obvious sign of the presence of drizzle, its absence, however, does not imply that there are no drizzle droplets within the cloud layers. We classify a profile as completely non drizzling when Z_{dzl} is zero below the cloud base and $Z_{\text{dzl}} > 0$ at less than 3 radar range gates above the cloud base (positive Z_{dzl} at only one or two gates does not constitute a drizzle profile and is likely dominated by noise). Consequently, the drizzle properties are all set to zero at all heights and the algorithm proceeds to compute the forward models and the cost function considering only contributions from the cloud.

We categorize the drizzling profiles into two cases with different retrieval mechanisms:

- Case I: drizzle is detected only above the cloud base

This case represents very early stages of drizzle formation when the drizzle droplets have not reached below the cloud base and its vertical extent is not known in advance of the retrieval. The absence of a clear, isolated drizzle signature adversely limits the retrieval strategy so we resort to adopting the same vertical model as for the cloud (eq. 9). Here the function $f(\tilde{z}_d)$ is defined by W_{dzl}^* , \hat{h}_{dzl}^* , with the cloud depth replaced by drizzle depth. The drizzle base and top heights are set to the closest radar range gate beyond the first and last detected Z_{dzl} , respectively, where $Z_{\text{dzl}} = 0$. The drizzle LWC profile is thus derived using:

$$\text{LWC}_{\text{dzl}}(\tilde{z}_d) = q_{\text{dzl}}^* f(\tilde{z}_d) \text{LWC}_{\text{ad}} = q_{\text{dzl}}^* f(\tilde{z}_d) \rho_a A_{\text{ad}} \tilde{z}_d. \quad (20)$$

\tilde{z}_d is height above the drizzle base while LWC_{ad} is identical to that for the cloud, computed using the temperature and pressure at the model cloud base. q_{dzl}^* is a drizzle scaling factor to account for the fact that the adiabatic and drizzle LWC can be a few orders of magnitude apart, a range that is not covered by drizzle $f(\tilde{z}_d)$ alone.

Given Z_{dzl} , LWC_{dzl} and ν_{dzl}^* (assumed independent of height), the profile of drizzle effective radius can be derived using eq. (12). The drizzle droplet number concentration and extinction coefficient profiles follow from eq. (3) and (4). To sum up, the elements of the state vector used to build the drizzle profile in case I are: ν_{dzl}^* , W_{dzl}^* , \hat{h}_{dzl}^* , q_{dzl}^* .

- Case II: drizzle is detected below and above cloud base

The drizzle retrieval here is based on the vertical structure of r_e above (eq. 13) and below (eq. 14) the cloud base. The r_e profile is split into the two functional forms to account for the different expected behaviours of drizzle droplets across heights. As in case I, drizzle base and top are set to the closest radar range gate beyond the first and last Z_{dzl} , respectively. This means, to solve eq. 13 and 14, we need to determine 3 parameters: drizzle effective radius at the cloud base, k_1 and k_2 . In the algorithm, we choose to express these parameters in terms of drizzle extinction coefficients at three height levels: within the cloud, at the cloud base and below the cloud base. The drizzle effective radius and extinction coefficient are related to each other through Z_{dzl} and ν_{dzl}^* . By combining equations 3, 5 and 7, it follows that:

$$r_{e,\text{dzl}}^4 = \frac{\pi Z_{\text{dzl}}}{32\alpha_{\text{dzl}}} \frac{(\nu_{\text{dzl}} + 2)^3}{(\nu_{\text{dzl}} + 3)(\nu_{\text{dzl}} + 4)(\nu_{\text{dzl}} + 5)}. \quad (21)$$

The three extinction coefficients are α_{id}^* , α_{cb}^* and α_{ic}^* . The first two refer to drizzle extinction coefficients at the first radar gate (the lowest range gate with radar detection) and at the cloud base. These two variables are used to construct the power-law profile (eq. 14). At these two heights, LWC_{cl} and α_{cl} are zero, allowing for an unambiguous drizzle retrieval. To specify the drizzle profile above the cloud base, α_{ic}^* is needed. It denotes drizzle extinction coefficient at a certain height within the cloud. Together with α_{cb}^* , α_{ic}^* solves eq. 13. The height choice for α_{ic}^* considers the following. The strong attenuation of the lidar signal within the cloud means that useful constraints are available only in the region between cloud base and ~ 200 m above it. Drizzle top is usually found at about 200 m into the cloud or higher. To obtain a meaningful solution for eq. 13, the height choice for α_{ic}^* should be well above the cloud base. Taking those into account, we opt to retrieve α_{cb}^* at 150 m above the cloud base.

Having constructed profiles for Z_{drl} and the drizzle effective radius, one can easily derive the other microphysical properties with the knowledge of ν_{drl}^* . As in case I, ν_{drl}^* is held constant with height. Finally, the four state vector parameters used in case II are ν_{drl}^* , α_{id}^* , α_{cb}^* , α_{ic}^* .

2.2.3 Physical constraints

- Following the above scheme, both the cloud and drizzle properties can be computed for a given state vector. There is, however, no guarantee that these properties are a sensible representation of the system in question. To mitigate this problem we impose several physical constraints that act as filters for the state vector:
- We apply a droplet size threshold to separate the cloud and drizzle regime. It has been shown that there exists a critical effective radius between 12-14 microns, above which coalescence increases and drizzle forms very rapidly (Rosenfeld et al. (2012) and references therein). Based on this, we adopt 13 microns as the separation threshold, which means that at any altitude, the cloud r_e has to be smaller than 13 microns and when drizzle is present, its r_e cannot be less than that. For profiles without radar signal below the cloud base, this threshold plays an important role in categorizing the profile as drizzling or non-drizzling.
 - The radar reflectivity due to drizzle must not be higher than the cloud reflectivity near the cloud top. The cloud top region is critical for the cloud-drizzle separation because this region is where the drizzle starts to form and where the difference between cloud and drizzle droplet size is minimal. Since the cloud LWC is highest near the cloud top, it is likely that the cloud number concentration and therefore cloud reflectivity will be dominant here. From this it also follows that the location of the maximum radar reflectivity near the cloud top is an indicator of the location of the LWC_{cl} peak.
 - For case II, it is important that the drizzle effective radius achieves its maximum value at the cloud base. This follows from the scenario that drizzle droplets grow as they fall through the cloud layers via accretion and then evaporate after they leave the cloud, thereby reducing their size. For case I, the increase of the drizzle effective radius towards the drizzle base is preferred through the use of a penalty function (see section 2.3), but not forced.

- Drizzle effective radius must not be larger than 250 microns due to the use of Rayleigh approximation on which the radar forward model is based. For a 35-GHz radar, the validity of the approximation sets an upper limit of droplet radius at about 280 microns. The 250 micron upper limit is imposed as a safeguard and is more of a technical limitation than a physical one. For the selection of drizzling clouds in our study here, this is not a concern.

5 2.2.4 Radar forward model

Equation (5) relates the cloud/drizzle microphysical properties to the radar reflectivity. The equation assumes the validity of Rayleigh approximation. For comparison with the observed reflectivity, the contribution from the cloud and drizzle must be added and attenuation effects have to be incorporated. The observed reflectivity that we use here is taken from the Cloudnet categorization product (Illingworth et al., 2007) and has been corrected for two-way attenuation due to atmospheric gases and in some cases, liquid water. Since the liquid water attenuation is dependent on the availability or the reliability of liquid water path measurements, it is not consistently applied to radar pixels containing cloud/drizzle droplets in the Cloudnet algorithm. For this reason, we recover the measured reflectivity before the liquid, but after the gas, attenuation correction Z_{obs} using the information provided in the same Cloudnet product. The liquid attenuation is then incorporated in the forward model to compute

$$15 \quad Z_{\text{fm}} = (Z_{\text{cld}} + Z_{\text{dzl}}) \exp(-2\tau). \quad (22)$$

Hereinafter, the subscripts 'cld' and 'dzl' refer to the cloud and the drizzle components, respectively. The optical depth τ is calculated from cloud and drizzle LWC using the approximation for attenuation coefficient given in Liebe et al. (1989). Z_{fm} is compared to Z_{obs} during the fitting in the retrieval.

2.2.5 MWR forward model

20 To simulate microwave brightness temperatures T_B , gaseous absorption by water vapor and oxygen is computed according to Rosenkranz (1998) and the absorption due to liquid water according to Liebe et al. (1993). The forward radiative transfer calculation is then performed by integrating the radiation intensity along the vertical path up to an altitude of 30 km, neglecting the variation in optical depth due to scattering. As such, T_B measurements provide constraints on the liquid water path (LWP) of a given column. **This forward model also requires additional information on the pressure, temperature and humidity profiles up to 30 km, which can be obtained from a numerical forecast model or radiosonde data.**

2.2.6 Lidar forward model

A publicly available code¹ for the calculation of lidar signals including multiple scattering is used to simulate the lidar attenuated backscatter. In treating the multiple scattering, the code allows an explicit computation of higher scattering orders following an approach by Eloranta (1998), and a fast calculation using the photon variance-covariance method (Hogan, 2006,

¹<http://www.met.reading.ac.uk/clouds/multiscatter/>

2008). Once the relevant parameters are available, e.g. lidar set-up, extinction coefficient and droplet size profiles, lidar attenuated backscatter from below and within the cloud can be calculated.

In the retrieval, we attempt to reconstruct the attenuated backscatter profile in the cloudy, drizzling regions as well as at drizzle-free altitudes between the ground and the cloud. The latter is necessary to estimate a possible offset in lidar calibration.

- 5 At these altitudes, scattering due to air molecules and aerosol particles are expected to prevail. While it is straightforward to approximate the extinction coefficient due to air molecules α_m from the temperature and pressure profile, the aerosol extinction coefficient α_a is largely unknown. Since multiple scattering does not play an important role in this region below the cloud/drizzle base, we use Klett inversion for a two-component atmosphere (Klett, 1981; Kovalev, 1995) to infer α_a from the observed β profile, such that for $z < z_0$:

$$10 \quad \alpha'(z) = \left[\frac{\left(\frac{P'(z)z^2}{P'(z_0)z_0^2} \right)}{\frac{1}{\alpha_0'^*} + 2 \int_z^{z_0} \frac{P'(z')z'^2}{P'(z_0)z_0^2} dz'} \right] \quad (23)$$

where

$$\alpha'(z) = \alpha_a(z) + S_a \beta_m(z), \quad (24)$$

$$P'(z) = S_a P(z) \exp \left(2 \int_0^z (\alpha_m(z') + S_a \beta_m(z')) dz' \right). \quad (25)$$

- The zero subscript refers to the Klett reference point with z_0 is set to $\min(z_{cb}, z_{db})$ and $\alpha'(z_0)$ is equivalent to $\alpha_0'^*$, one of the state vector elements. S_a is the extinction-to-backscatter ratio for aerosol. Given that the lidar operates at 355 nm, we adopt $S_a = 50$ sr, a representative value for aerosol particles. $\beta_m(z) = \alpha_m(z)/S_m$ is the attenuated backscatter due to molecular scattering and is calculated assuming $S_m = 8\pi/3$. $P(z)$ is the attenuated backscatter power as a function of height and is defined as:

$$P(z) = \frac{C_{ldr}(\beta_z(z) + \beta_m(z))}{z^2 \exp \left(2 \int_0^z (\alpha_a(z') + \alpha_m(z')) dz' \right)}. \quad (26)$$

- 20 Using the equations above, the $\alpha_0'^*$ value in the state vector, and the fact that lidar calibration factor C_{ldr} cancels out in eq. 23, the α_a vertical profile below the reference point can be derived. The $\alpha_a(z)$, the drizzle and cloud extinction coefficient profiles (as derived in sections 2.2.1 and 2.2.2) are then stitched together and used as input for the multiple scattering code to construct the complete attenuated backscatter profile below and within the cloud/drizzle. The lidar calibration factor C_{ldr} can now be computed at each range gate in the cloud- and drizzle-free region:

$$25 \quad C_{ldr}(z) = \frac{S_a \beta_{obs}(z) \exp(2 \int_0^z (\alpha_a(z') + \alpha_m(z')) dz')}{\alpha'(z)}. \quad (27)$$

Since C_{ldr} serves to compensate for a systematic offset due to inaccurate calibration, its values are expected to be approximately constant with heights. We have confirmed that the values are very similar across heights to within about 2%. Finally, we multiply the forward-modelled β profile by the median value of $C_{ldr}(z)$ for comparison with the observed β .

2.3 Finding the optimal solution

This retrieval procedure attempts to solve the inverse problem of deriving cloud and drizzle properties from observations by minimizing the cost function

$$cf = [y - F(x)]^T S_y^{-1} [y - F(x)] \quad (28)$$

5 to arrive at the optimal solution for the state vector x . y is the measurement vector defined as:

$$y = [T_{B,\text{obs},1}, \dots, T_{B,\text{obs},nf}, \beta_{\text{obs},1}, \dots, \beta_{\text{obs},nl}, Z_{\text{obs},1}, \dots, Z_{\text{obs},nr}], \quad (29)$$

with nf, nl, nr represents the number of MWR frequency channels (14), the number of lidar range gates, and the number of radar range gates with detection, respectively. $F(x)$ is the vector of forward-modelled observables, with the same composition as y . For a T_B measurement at a frequency i with an uncertainty $\sigma_{T_B,i}$, the diagonal element (i, i) of the measurement covariance matrix S_y is $\sigma_{T_B,i}^2$. All non-diagonal elements (cross-channel or cross-instrument elements) are set to zero assuming no correlation. The elements (m, n) of matrix S_y corresponding to radar and lidar data are calculated according to

$$S_{y,m,n} = E([y_m - E(y_m)][y_n - E(y_n)]), \quad (30)$$

which results in:

$$S_{y,m,n} = \sigma_C^2 y_m y_n \quad \text{for } m \neq n \quad (31)$$

$$15 \quad S_{y,m,n} = \sigma_C^2 y_m^2 + \sigma_{y_m}^2 \quad \text{for } m = n \quad (32)$$

where σ_{y_m} and σ_C are the random uncertainties of the measured signal and the instrument calibration, respectively. σ_C is set to be small (comparable to the desired fit accuracy for Z and β).

For case I, a penalty term is added to the cost function to bias $r_{e,\text{dzt}}$ toward the desired profile. Along the drizzle profile, it is checked whether $r_{e,\text{dzt}}$ is larger than the one directly below it. Since the determining factor of drizzle is the radar reflectivity, the penalty is applied to the radar part of the cost function such that every single violation would add the radar term to the original cost function.

The cost function (combined with the penalty function for case I) is minimized using Differential Evolution (DE), a global stochastic optimization technique similar to population-based optimization routines (Storn and Price, 1997). DE does not require gradient information, which is an advantage given the complexity and the non-linearity of the cost function. It is designed to deliver robust results and a fast convergence while maintaining a small number of control variables. This minimization algorithm begins with a population of state vectors that constitute a generation. The vector values are chosen to cover the allowed parameter space and the population is then updated with each generation. For each member vector, a new vector is created through mutation and parameter mixing/crossover to replace the old one if it results in a smaller cost function value. Otherwise, the old member vector is retained to be part of the subsequent generation population. This mutation-crossover scheme, along with the strategy to start with a set of vectors, instead of a single initial vector, make it less likely for the algorithm to

get trapped in local minima. To use the algorithm, initial guesses for the state vector values are not needed, but the lower and upper limits for each state vector parameter are required.

DE comes in several variants, which differ in the way the mutation and crossover are done. Here we choose the DE/best/1/bin variant with a population size $NP=10$, a mutation factor F that randomly changes between 0 and 1.9 on a generation-by-generation basis, and a crossover constant $CR=0.8$ (Storn and Price, 1997). For the retrieval, we use the numerical implementation of DE provided within the Python-based environment for scientific computing SciPy², where the stop conditions are specified by the tolerance (0.01) and the maximum number of generations (150).

The minimization of the cost function is performed over bounded state vector values, from which a physically-sensible solution should be found. Unless stated otherwise, the lower and upper limits of state vector values that we use in this work are listed in Table 1. The shape parameter for the cloud DSD is expected to vary between 2 and 10 (Miles et al., 2000; Gonçalves et al., 2008), depending on for example airmass and location (marine or continental). For drizzle DSD, an exponential fit ($\nu = 1$) is found to be a good approximation (Wood, 2005b). Here we allow ν to vary within a wide range. From our investigation (Section 3) it appears that constraining ν to a fixed value when the radar calibration accuracy is unknown can potentially create a significant bias in the retrieval products. The limits for \hat{h} and W cover the subadiabatic range of LWP that is viewed to be common (Boers et al., 2006). Since small drizzle droplets present minimal effects on β , the extinction coefficient of drizzle at the cloud base α_{cb} is constrained to be comparable to the air extinction coefficient. The value of α_{id} is expressed relative to the drizzle extinction coefficient at the cloud base and α_{ic} is determined relative to the value of the cloud extinction coefficient value at the same height. α'_0 is given a large range because it is rather sensitive to a small change in the retrieved lidar offset.

The uncertainties for the optimal solution are computed using Monte Carlo realizations that were generated by perturbing the observations. Each random realization of the observations is drawn from a Gaussian distribution centered on the measurements with the dispersion taken from the measurement (random) errors. The retrieval procedure is then performed on all realizations resulting in a set of solutions. The RMS difference with the optimal solution is calculated to represent the uncertainties of the retrieval. For each column observation, we create ten realizations which should provide a conservative estimate of the random uncertainty. Systematic uncertainties due to inaccurate radar calibration are not included in the Monte-Carlo error estimate. Assuming that the calibration offset can be under- or overestimated by up to a factor of two (3 dB), the resulting systematic errors on the retrieval products are found to be larger than the random uncertainties (see Section 3).

3 Test using synthetic data

3.1 Cloud retrieval

Before applying the technique to real observational data we test it on a set of synthetic signals generated from large-eddy simulation (LES) results. Similar to the work described in Donovan et al. (2015), the simulation set-up is based on output from the Dutch Atmospheric LES model (DALES) (Heus et al., 2010) for conditions corresponding to the FIRE campaign

²<http://www.scipy.org>

(Duynerkerke et al., 2004; de Roode and Los, 2008). Given the LWC from the LES, the DSD is assumed to be a monomodal gamma distribution, i.e. drizzle droplets are not present. The shape parameter and the number concentration along the vertical column are externally imposed and they are largely constant. ECSIM (Voors et al., 2007; Donovan et al., 2015) was used to generate the radar, lidar and MWR signals. Applying the algorithm to these signals serves primarily as a sanity check for the retrieval code, to verify the forward models and the assumption on the vertical shape of LWC.

The synthetic signals are simulated for a zenith-pointing 32 GHz radar, a lidar operating at 353 nm, and a MWR with 14 frequency channels between 20-60 GHz to mimic the instruments used in the ACCEPT campaign (see Section 4). The radar and lidar signals are sampled at a fine spatial resolution: 2.5 m vertically and 25 m horizontally. To mimic real observations we degrade the vertical resolution of both the radar and lidar to 22.5 m. Along the horizontal axis (corresponding to the time axis), we lower the resolution to 150m by averaging radar reflectivity and lidar attenuated backscatter data at each range gate, and averaging the brightness temperatures at each frequency channel. The standard deviation of the mean serves as the uncertainty. The atmosphere below the cloud is rather static, making the standard deviation of β in this region unrealistically low. This condition virtually assigns a lot of weight to the part below the cloud in the fitting process, which leads to inaccurate retrieval. For this reason, we set the noise floor for the β profile to 1% below the cloud base and 5% above the cloud base. Similarly for Z and T_B , the minimum relative error is set to 0.03 and 0.01, respectively. It is these simulated measurements with adjusted resolution, together with the uncertainties, that are fed to the retrieval code.

Fig. 3 shows the input synthetic signals as compared to the signals recovered by the retrieval. Apart from the lower edge of the cloud at horizontal distances < 7.5 , Z is generally well reproduced. The lidar signal is also recovered despite the noise. It is fitted up to 300 m into the cloud, after which the noise prevails. The histograms of the reflectivity and the attenuated backscatter residuals (truth - retrieval) are displayed in panels 3e and 3f. Most of the residuals are relatively small, the peaks of the histograms are centered at zero and the distributions are quite symmetric with no particularly strong tendency towards positive or negative values. T_B , averaged over distance, at each frequency channel coincides well with the data. Average T_B fluctuations over time are small: less than 4% as shown in Fig. 3f. The Root-Mean-Square Deviation (RMSD) between the data and the retrieval is also very low, i.e. less than 1%, suggesting a good match between the two. The maximum rms is found at 31.4 GHz, where the extinction due to liquid water dominates the microwave signal.

Fig. 4 displays the true microphysical and optical properties in comparison with the retrieved ones. The structures in the LWC, r_e and α are mostly reproduced. Since the retrieval is performed on a column-by-column basis the retrieval is not entirely smooth along the horizontal axis and this effect is particularly visible in N . The mild vertical structure in the true droplet concentration is not reproduced due to the model assumption of constant N . From the histograms, it can be seen that generally the LWC and the extinction coefficient are retrieved more accurately than the effective radius and the number concentration. The retrieved r_e and N tend to be higher and lower than the truth, respectively, by a few percents (see also Fig. 5 and Table 2). The distribution of ΔN appears less Gaussian than those for the other microphysical properties due to the column gradient in the true N that is not matched by the retrieval assumption.

Fig. 5 displays the vertically collapsed version of Fig. 4. LWC and α are integrated into LWP and optical depth, respectively. N and r_e are vertically averaged, with the latter weighted by α . The error bars represent the random measurement error from

the Monte Carlo realizations; there is no systematic error due to the radar calibration. The fluctuations of the variables along the horizontal axis are easily reproduced with very little bias, which is mostly seen in r_e and N , as shown before by the histograms in Fig. 4. The mean values of the LWP, r_e , optical depth and the number concentration, averaged over the horizontal axis, and the deviation from the truth are given in Table 2.

- 5 The true shape parameter is not strictly constant along the vertical direction; it is mostly close to 6, and decreasing to around 2 at the cloud base or cloud top. The retrieval is performed with $\nu = 5.5$ and with the radar calibration factor fixed to 1 to match the true values. The lidar calibration factor is retrieved on average with a 5% accuracy. For comparison purposes, we also include in the last column of Table 2 the run where the shape parameter ν is free within a fixed range, i.e. between 2 and 10. The result is that the noise of the retrieved products becomes higher but there is very little systematic offset. The optimized
- 10 ν is found to have a mean of 5.98 (RMSD=2.01), very close to the true ν . By comparing the last two columns in Table 2, it is apparent that when ν is not fixed, the RMSD increases significantly due to the large column-to-column fluctuation but the mean values are hardly affected. The extinction coefficient is found to be relatively stable against the variation in ν , possibly because its retrieval is largely dependent on the β profile.

- We also investigate the effect of under- or overestimating the radar calibration offset. For this purpose, we apply a shift of
- 15 ± 3 dB (a factor of two) to the forward-modelled Z and perform the retrieval with ν bounded between 2 and 10. When the offset is underestimated (forward modelled Z is multiplied by 0.5, or $C_r = 0.5$), LWC and r_e are overestimated by 10-15% while α generally becomes lower by a few percent, and vice versa when $C_r = 2$. The retrieved number concentration tends to fluctuate and is on average 15% higher than the mean truth for $C_r = 2$. The relatively mild systematic impact of doubling or halving the radar calibration offset is possibly due to the fact that the shape parameter is allowed to vary within a certain
- 20 range; the true shape parameter is not recovered in both cases of C_r . The magnitude of the systematic difference between the retrieval products and the truth increases when ν is fixed to the true value, especially for the number concentration where the mean retrieved N becomes 36% lower than the truth.

- What is demonstrated with this exercise is that the forward models are able to reproduce the radar, lidar and MWR signals and that the LWC parametrization that we use for the cloud indeed provides a realistic description of LWC vertical structure.
- 25 Given accurate instrument calibration, the systematic mismatch between the retrieval and the truth is found to be very small for this test case, both when ν is fixed to approximately the true value or when it is optimized. From all four retrieval products, the largest mean offset from the truth is found for the number concentration N at less than 5%.

3.2 Cloud and drizzle retrieval

- In this section we present two examples of the cloud and drizzle retrieval using synthetic data to illustrate the drizzle retrieval
- 30 scheme described in section 2.2.2. The synthetic data is produced as follows. Cloud LWC is provided by the LES results as in section 3.1. We select one LWC profile and derive the effective radius and extinction coefficient profiles by imposing a monomodal gamma DSD and a value for the number concentration. The DSD shape parameter and the number concentration are set to be independent of height. Two idealized drizzle profiles for both case I and case II are then constructed following the parametrizations introduced in section 2.1.3. These cloud and drizzle profiles serve as the truth, to which the retrieval products

are later compared against. The two drizzle truths are combined with the cloud truth to result in two example profiles, which are then forward modelled to produce synthetic radar, lidar and MWR signals. The truths and the retrieval results for the cloud and drizzle components representing the two retrieval cases are displayed in Fig. 6, which is discussed below.

– Case I

This is the case where drizzle presence is restricted to the cloudy region. There is only a limited amount of information that we can use to retrieve the cloud and drizzle properties in such a situation. Since drizzle contributes only to a small fraction of the total signals, the task of retrieving its properties becomes even more challenging. For the drizzle properties, our goal here is to get an order-of-magnitude estimate, rather than to achieve a high accuracy.

In the example here, drizzle is added to all gates within the cloud. Panels 1e and 1i show the LWC models for the cloud and drizzle, parametrized using eqs. 8 and 20. The retrieval scheme is as follows. LWC_{cld} and N_{cld} are first constructed to derive Z_{cld} . Z_{dzl} follows from the difference between the synthetic Z and Z_{cld} . Based on the vertical extent of Z_{dzl} , LWC_{dzl} is constructed. The effective radius of drizzle is derived using eq. 12 and the other properties follow from eqs. 2-7.

The optimized total Z (panel Ia), the attenuated backscatter (I_c) and T_B (Id) match the synthetic signals very well. The decomposition of the retrieved Z into a cloud part and a drizzle part is shown in panel Ib. The retrieved Z_{cld} provides a good match to the truth. **The true cloud LWC (Ie), effective radius (If), and extinction coefficient (Ig) are recovered to a high accuracy: the maximum difference between the truth and the retrieved values at any range gate amounts to no more than 5%.**

As for the drizzle, the retrieved Z_{dzl} (panel Ib) is not a perfect match to the truth in the upper half of the cloud, where the lidar signal is weak. The drizzle properties are small in magnitude, especially in the upper part of the cloud, making its retrieval very sensitive to noise. Additionally, drizzle retrieval is based on excess Z , which means that any discrepancy between the retrieved and the true Z_{cld} has to be compensated by the retrieved Z_{dzl} . Since Z_{dzl} is typically much lower than Z_{cld} at the top part of the cloud, this compensation mechanism impacts the Z_{dzl} retrieval quite strongly. In some observational cases (see section 4), as also reflected in this example, Z_{dzl} close to the cloud top can be very low, e.g. about 1% of the Z_{cld} , making it comparable to the noise/uncertainty level. In this sense, the droplet size threshold that we apply (section 2.2) helps to distinguish weak drizzle signal from noise, but the retrieval accuracy would still be limited. Panels (Ii)-(Ii) show the drizzle properties as derived from the knowledge of drizzle LWC and Z . While the vertical shape is not exactly recovered, the retrieved values are within one order of magnitude from the truth.

– Case II

This is the case where isolated drizzle can be found below the cloud base, so here we can be more certain about the drizzle spatial extent. As in case I, the retrieval starts with building LWC_{cld} and N_{cld} to derive Z_{cld} . Z_{dzl} is computed from the difference between the synthetic Z and Z_{cld} above the cloud base; below the cloud base, Z_{dzl} is equal to the

synthetic Z . Then, the profile of drizzle effective radius is constructed using the two parametrizations given in eqs. 13-14. The other drizzle properties follow from eqs. 2-7.

The optimized total Z , the attenuated backscatter and T_B are shown in panels (IIa),(IIc) and (IId). The decomposition of Z (panel IIb) in the upper half of the cloud is plagued by the same problem as in case I. Just below the cloud top, the separation between the cloud and the drizzle contribution deviates from the truth, and this deviation is propagated to the retrieved properties (panels IIe - IIh for cloud and Ili-III for drizzle). The retrieved cloud LWC and cloud extinction coefficient are too large by $\sim 20\text{-}25\%$ at the peak close to the cloud top. The vertical shape of the drizzle r_e follows an exponential function within the cloud. **The retrieved effective radius here is about 6 microns smaller than the truth.** The smooth exponential profile of the drizzle r_e causes a dip in the Z_{drl} value at 530-m to also appear in the α_{drl} and N_{drl} profiles. **This dip in drizzle Z comes about because the cloud reflectivity between 500 and 600 m is not well recovered (the effect is most visible in panels IIe and IIg in Fig. 6). More specifically, the retrieved cloud Z at these heights are slightly larger than the true ('observed') Z , resulting in $Z_{\text{excess}} = 0$ (eq. 19). The smoothing that is applied afterwards to produce Z_{drl} replaces the zeros with non-zero values that are smaller than the neighbouring range gates, creating the dip.**

Below the cloud base, the drizzle r_e is restricted to not decrease towards the drizzle base. Here we see that it is roughly constant with height. **The retrieval scheme in this region relies on inferring the magnitude of drizzle extinction coefficient from the lidar backscatter. The three control points (i.e. the drizzle extinction coefficients at three height levels) are retrieved with limited accuracies. The differences between the retrieval and the truth are on average about 38%, 14%, 46% and 52% of the true values for drizzle LWC, r_e , α and number concentration, respectively. Note that in this example, the small amount of drizzle extinction hardly changes the lidar backscatter below the cloud base. The drizzle extinction coefficient here amounts to about 5% of the molecular extinction. Other simulations were conducted where the amount of drizzle was varied. It was found that the drizzle retrieval accuracy below the cloud base increased as drizzle amounts increased. For example, when α_{drl} was increased by a factor of ~ 20 , the retrieval accuracies improved: drizzle LWC, r_e , and α below the cloud were retrieved within 8%, while the number concentration within 25% from the truth (not shown).**

In general, the cloud retrieval is more robust than the drizzle retrieval. Although the retrieved cloud properties for an individual cloud profile at a given height can be off by $\sim 25\%$, collective retrieval of a sample of profiles and column-averaged properties are expected to have better accuracies, as shown in section 3.1. Drizzle retrieval is very sensitive to non-idealized cloud structure and also to the uncertainties of the observed signals. From the examples using idealized profiles of drizzle, it is shown that the retrieval error at a given range gate can be large (starting at 30% level) but overall, profiles of the retrieved drizzle properties are reasonably close to the truth.

4 Application to ground-based observations

The observational data was collected during the ACCEPT campaign that took place in October and November 2014 in Cabauw, The Netherlands (see Myagkov et al. (2016) and Pfizenmaier et al. (2017) for more information about the measurement campaign). We use the data acquired from three co-located instruments:

- 5 – A zenith-pointing MIRA-35 radar
 It is a Ka-band cloud radar with Doppler capabilities. The signal was recorded with a spatial resolution of about 30 m.
- A UV lidar (Leosphere ALS-450) operating at 355 nm
 The attenuated backscatter measurements are available every 30 sec with a vertical resolution of 15 m.
- A microwave radiometer (MWR) HATPRO
- 10 The brightness temperature was measured at 14 frequency channels: the first seven between 20-35 GHz and the other seven between 50-60 GHz. The temporal resolution is 1 sec with regular gaps due to automatic calibration periods.

For the inversion procedure, we use the calibrated radar reflectivity factor, as well as model forecast of temperature and humidity delivered in the Cloudnet categorization product (Illingworth et al., 2007). The calibrated reflectivity here is already corrected for gas attenuation and has the same temporal resolution as the lidar although their time stamps do not exactly
15 coincide.

The retrieval is performed on a column-by-column basis with a time interval of 30 sec. For each 1D column, a set of radar, lidar and MWR data was created by first finding the lidar and radar profiles that are less than 15 sec apart. The corresponding T_B profile was computed by averaging T_B measurements within 15 sec of the average time stamp of the radar and lidar. The standard deviation of the mean was then adopted as the measurement error. Since the full overlap distance of the lidar is
20 expected to be around 100-200m, column profiles with radar detections down to < 200 m were not retrieved. There are gaps in the observations where a complete data set for the three instruments is not available, e.g. breaks in the MWR data stream during instrument calibration periods.

We selected two periods with a total time of approximately 4 hours on October 25 and 26, when one layer of liquid water cloud is present. The scene includes clouds with clear drizzle/precipitation events and also clouds without obvious signature
25 of drizzle below the cloud base, suitable for the dual retrieval mode (case I and case II). The cloud top is located between 1400 and 1500 m, with the cloud thickness varying between 200-400 m. The cloud base height (as determined in the retrieval) fluctuates between 1050 and 1250m during the two periods. Despite low reflectivity values, virga/drizzle is observed below the cloud base for the majority of the time with its maximum occurring on Oct 26. The extent of the drizzle below the cloud is variable, with a depth of up to 600 m.

30 The observed signals and their recovery in the retrieval is shown in Fig. 7 for each instrument. In general, the reflectivity within the cloud increases with height indicating particle condensational growth. For the most part, the radar reflectivity is not higher than -28 dBZ. On Oct 26 at around 3.8hr, Z is maximum at -12 dBZ. In the retrieval, cloud and drizzle contributions to the total reflectivities are separated and these are shown in panels (c) and (d). Below the cloud base, Z belongs only

to the drizzle. Above the cloud base, the reflectivity of the cloud increases with height and peaks close to the cloud top. Conversely, drizzle reflectivity increases downwards from the top and reaches maximum in the cloud base region before decreasing again towards the drizzle base. It follows that within the cloud, the cloud reflectivity dominates towards the cloud top while drizzle dominates near the bottom. In almost all profiles where no virga is visible below the cloud base, the retrieval
5 algorithm finds drizzle to be present within the cloud although with small reflectivities. This is usually caused by the significant excess reflectivity near the cloud base that cannot be attributed to the cloud component.

Panels (e) and (f) compare the observed and retrieved lidar attenuated backscatter. The fitting of the β profile starts from an altitude between 200 m (from the ground) and the drizzle base, and continues up to 200 m above the cloud base. Several lidar β profiles show double backscatter peaks that we deem unsuitable for the algorithm, in which case the retrieval is not performed
10 resulting in white gaps in the time-height map. We found 25 such column profiles, corresponding to 5% of the available data. It can be seen that the drizzle below the cloud base remains largely transparent and undetected by the lidar. The histograms of the residual signals are similar to those obtained for the LES exercise (Fig. 3e and 3f): centered at zero and largely symmetric. Compared to Fig. 3e, the ΔZ distribution here is narrower because a part of the Z residual is attributed to drizzle. The mean brightness temperature at each frequency channel is shown in Fig. 7i. The observations show little variation over time (less
15 than 8%). On average, the fit to the observed T_B is reasonably good with a RMSD of 5% or less, as indicated by the red line. The largest variation or difference is seen around 30 GHz, where the cloud contribution to the microwave extinction spectrum is significant.

The retrieved microphysical and optical properties for both cloud and drizzle are shown in Fig. 8. Cloud and drizzle LWC in panels (a) and (b) show a similar time-height distribution to the respective reflectivity field. Cloud LWC increases with height
20 until the peak is reached close to the cloud top, while most water in drizzle is found at a lower altitude. The highest LWC is found at the time of maximum observed reflectivity. Drizzle water content is generally 2 orders of magnitude smaller than the cloud LWC. The average of LWC_{cld} and LWC_{drl} maximum values are 1.9×10^{-1} and 1.2×10^{-3} g/m³, respectively. The temporal variations of the cloud and drizzle LWP are positively correlated in time, as was also found by Fielding et al. (2015).

The effective radius of the cloud droplets (panel c) is found to be well below the threshold values of 13 microns. As the cloud
25 droplets grow via condensation, their size increases with height to a peak value of 5.1 microns, on average. Drizzle effective radius (panel d), on the other hand, increases towards the cloud base as imposed by the parametrization. At the cloud base, the mean drizzle effective radius is found to be ~ 22 microns. During the intense drizzle period between 3.8-4.0 hr, it can be as high as 60 microns.

The extinction coefficients of the cloud and drizzle (panels e and f) are mainly determined from the observed lidar attenuated
30 backscatter. The extinction coefficient of drizzle is smallest below the cloud base, in accordance with the relatively low observed β in this region. It increases with height and peaks within the cloud but it is still orders of magnitude smaller than the cloud. The mean maximum α_{cld} and α_{drl} are 0.06 and 1.2×10^{-4} m⁻³, respectively. The number concentration of the cloud (panel g) droplets shows a somewhat high and rapid fluctuation. From our LES exercise (Section 3), we learned that setting ν_{cld} as a free parameter can indeed cause this, but we also expect that the fluctuation averages out to a minimally biased mean value. The

cloud number concentration averages to about 549 cm^{-3} . The drizzle droplet concentration (panel h) has a mean of $\sim 0.06 \text{ cm}^{-3}$, consistent with the in-situ measurements of marine-stratocumulus in the MASE-II experiment (Lu et al., 2009).

In Fig. 9 we show the mean vertical profiles of the radar reflectivity and the derived microphysical properties of the cloud and drizzle. These profiles are constructed from averaging the retrieved profiles between 3.8 and 4.0 hour UTC when there is significant drizzle. In all but the lowermost parts of the cloud, drizzle gives a very small contribution to the total water content. Close to the cloud top, the effective radius of the drizzle droplets are found to be around 30 microns for this particular time segment. At around the same height, the cloud droplets reach their maximum size of ~ 6 microns with the cloud liquid water content and reflectivity dominating over the drizzle. As the drizzle droplets grow exponentially towards the cloud base via coalescence, its reflectivity increases, matching the cloud reflectivity about halfway through the cloud. Near the cloud base, drizzle reflectivity becomes dominant due to the much larger size of the drizzle droplets compared to the cloud droplets. Inside the cloud layers, this behaviour of cloud and drizzle reflectivities is found to be quite typical over the observation period. From the cloud base towards the drizzle base, Z_{drl} decreases monotonically, and so does the LWC_{drl} . Once drizzle drops escape the cloud, they are expected to evaporate below cloud base and shrink as they fall through the air, in accordance with the gradient seen in the $r_{e,\text{drl}}$ profile. The jagged feature in the $r_{e,\text{drl}}$ profile below the cloud base is an artifact of the profile averaging caused by the variable drizzle base height during the 12-minute period.

Lastly, the number concentration of the drizzle is 3-4 orders of magnitude smaller than the cloud at all heights above the cloud base. Unlike the cloud number concentration, N_{drl} shows variation in height. The highest density of drizzle is found approximately where its LWC is highest, that is within the cloud. From this point towards the cloud base, the drizzle number density keeps decreasing as $r_{e,\text{drl}}$ rises sharply, which could be due to the accretion of smaller drizzle droplets by the bigger ones to form even larger droplets. Below the cloud base, some droplets experience complete evaporation, depleting the number density of drizzle as it approaches the ground.

5 Comparisons with Other Retrieval Methods

To assess the results presented in Section 4, we perform comparisons with three independent retrieval methods, applied to the ACCEPT dataset within the same period. The three retrieval techniques offer a tool to evaluate our retrieval below and above the cloud base separately. Below the cloud base, the drizzle comparison is made with the readily-available results from the method of O'Connor et al. (2005) as part of the Cloudnet algorithm package. Above the cloud base, the cloud properties are retrieved using the depolarization lidar technique developed by Donovan et al. (2015). The amount of drizzle within the cloud is derived from radar doppler spectra analysis, as described in Kollias et al. (2011a, b). In Appendix A, we present the application of this technique to the ACCEPT data, following the implementation in Luke and Kollias (2013); the comparison with our retrieval is discussed in Section 5.2.

5.1 Cloud properties above the cloud base

A depolarization lidar-based (DL) method (Donovan et al., 2015) was applied to the lidar dataset used in our retrieval to derive the cloud properties. While our retrieval method utilises only the total attenuated backscatter, the DL method exploits the parallel and perpendicular polarization components of the received signal to infer the cloud extinction coefficient and droplet size.

Fig. 10 displays the time series of the cloud properties at a specific, arbitrary height (chosen to be 100 m) above the cloud base, as derived from the DL (blue) and our (black, red) retrieval methods. The DL method determines the cloud base height that is then used as a height reference for the subsequent retrieval of the extinction coefficient and effective radius, from which the LWC and N are then derived. The retrieval is performed with a temporal resolution of 180 sec. For a fair comparison, we interpolate our retrieved profiles to 100 m above the cloud base as well. Since our retrieval has a higher temporal resolution (30 sec), we use our own cloud base height estimate (z_{cb}) as a reference. It is therefore imperative to first make sure that the cloud base height estimates from the two methods match. The bottom panel of Fig. 10 shows that this is indeed the case.

We average our results over the 180-sec time interval to match the time stamps and the temporal resolution of the DL method. The time-averaged products (red line in Fig. 10) are then compared with the results of the DL method. The extinction coefficients retrieved by both methods are very similar; we find that the mean difference and the RMSD amount to -5% and 11%, respectively. The fractional quantities here and in the following are produced using the mean of the DL results as a reference. The effective radii do not compare as well, with a mean difference of -10% and a RMSD of 21%. Our effective radius is highest at around 3.88 hr, coinciding with a high radar reflectivity situated in the middle of an intense drizzle episode. This is not picked out by the DL retrieval. The difference in r_e is mostly visible between 17.0-17.4 hr and 4.8-5.4hr. The mismatch does not necessarily correlate with the drizzle quantity and can amount to up to 1.5 microns but is still within the expected DL uncertainty range.

Our retrieved LWC is on average lower by about 15% with a RMSD of 26%. The DL number concentration is derived assuming that the DSD follows a single-mode gamma distribution (eq. 1) with a shape parameter $\nu = 6$. Uncertainties of N are propagated from the errors of r_e and α , taking into account a range of shape parameter values between 4 and 10 which is typical for liquid water clouds. Our number concentration fluctuates rather strongly with time, contributing to the large RMSD (45%). The shape parameter in the retrieval procedure is set as a free parameter to minimize biases and as discussed in Sect 3, this can lead to a strong fluctuation in the number density. In comparison to the DL method, our mean N is higher by about 15%. Given that the DL uncertainties of α are between 15-20%, and that the large fraction of the DL-retrieved r_e , LWC and N have $\sim 50\%$ errors, the differences between the two methods are well within the range of uncertainties.

5.2 Drizzle reflectivity above the cloud base

The qualitative comparison between our and the Doppler retrieval results is shown in Fig. 11 for the drizzle (a), cloud (b) and the total (c) reflectivities. Each circle represents a time-height pixel for which both methods are applicable; the number of time-height pixels that can be retrieved using only one of the methods, and hence are not used in the comparison, is less than

8%. In general, the distribution of the circles is consistent with the one-to-one line. The spread is higher for lower Z , indicating the higher fraction of noise for low- Z retrieval. The larger values of our ('retrieved') total Z in (c) compared to the Doppler counterpart can be attributed to the absence of attenuation correction in the Doppler spectra. On average, there is a difference of 0.9 dB between the two sets of total Z . This difference has very little effect on the trend that we see in the scatter plots (a) and (b), in which the circles are color coded according to their relative heights within the cloud. We divide the cloud into three horizontal parts: (i) one quarter into the cloud (cloud base region: red), (ii) the top quarter (cloud top region: green), and (iii) the middle region in between (blue). The insets show the same scatter plot without the blue points to highlight the division between the cloud base and cloud top regions.

The correlation between our and Doppler Z varies across altitudes within the cloud. Close to the cloud base there is a clear tendency that our retrieved drizzle reflectivities are larger than those from the Doppler analysis. The cluster of red circles in (a) is almost exclusively located to the left of the one-to-one line. Consequently, our cloud reflectivities become smaller than the Doppler Z_{cd} , as seen in (b). The primary cause for this stems from the assumptions used in both methods. In our retrieval, the cloud LWC, effective radius and thus reflectivity near the cloud base is assumed to decrease downwards until it reaches zero at the cloud base. This way, drizzle can gradually maximize its share of the total reflectivity, guaranteeing the continuity of the drizzle reflectivity when crossing the cloud base. Such a restriction in the vertical profile is not in place for the Doppler retrieval and the applicability of the method is limited to situations where drizzle is not dominant.

It should be noted that the accuracy of the Doppler retrieval is crucially determined by the shape of the spectra. The choice of the time interval during which the individual spectra are averaged is known to play a role in determining the shape of the composite spectra (Giangrande et al., 2001). Here we fix the time window to 30 seconds to allow for a one-to-one comparison with our retrieval results. High turbulence is shown to cause an underestimation of drizzle reflectivity, as derived through spectral decomposition, by up to 10-15 dB (Luke and Kollias, 2013). This means that even if drizzle indeed dominates the cloud, turbulence can smear the spectrum in such a way that it appears cloud-dominated. A compensation for this effect is thus critical in determining the correct drizzle fraction. In the scatter plots above, the presented reflectivity values have been turbulence-corrected according to Luke and Kollias (2013) who estimate the correction factor as a function of spectral broadening, determined from their extensive dataset of marine stratocumulus clouds. Ideally, turbulence-corrected drizzle reflectivities just below and just above the cloud base should have similar values, which was indeed the case in Luke and Kollias (2013). After applying their procedure to the ACCEPT dataset, however, we find that the drizzle reflectivities above the cloud base is lower by several dB (the mean is 4 dB lower) than those just below, as shown in Fig. 12. This could be the result of the uncertainties in the estimate of the correction factor, or an artifact of the spectrum averaging process. We do not attempt to formulate the correction factor necessary to eliminate this reflectivity gap due to the insufficient amount of data to provide a statistically significant estimate.

Towards the cloud top there is a mild trend that our retrieved drizzle reflectivity is smaller than that derived from the spectra, thus opposite to what is seen in the cloud base region. In our retrieval, the drizzle reflectivity is derived from the excess reflectivity that is not claimed by the cloud component and it typically increases towards the cloud base. Such a pattern is not enforced and is also hardly visible in the Doppler retrieval, leading to discrepancies in the results.

The best agreement between the two methods is found for the middle part of the clouds. If a shift of a few dB is introduced to Z_{drl} above the cloud base to correct for the reflectivity gap indicated in Fig. 12, then the agreement would improve in the cloud base and mid-cloud regions, but would worsen in the cloud top region. Overall, the correlation coefficient assessed by including all points is 0.38 for drizzle and 0.75 for the cloud. When we examine only the mid-cloud reflectivity distribution, the correlation coefficient improves to 0.54 and 0.87 for drizzle and cloud, respectively. The cloud boundaries are problematic areas for the comparisons due to their transitional nature, lower reflectivities and therefore high uncertainties. However, despite the distinct retrieval procedures, different sources of information and different assumptions, the two methods show reasonable agreement in quantifying the amount of drizzle within the cloud.

5.3 Drizzle LWP below the cloud base

Drizzle parameters below the cloud base are retrieved as one of the level 2a Cloudnet products using the algorithm introduced in O'Connor et al. (2005). The retrieval makes use of lidar backscatter and the first three moments of the Doppler radar spectra. The radar data comes from MIRA radar, the same as the one used for our retrieval. The lidar backscatter information is obtained from an independent observation with a different instrument, i.e. a CHM15X ceilometer.

The drizzle property that we can directly compare is the liquid water path of the drizzle below the cloud base, displayed in Fig. 13. The graph demonstrates a strong correlation between the two sets of LWP values, spanning a few orders of magnitude. The points align well with the one-to-one line and the correlation coefficient is found to be 0.99. The mean values of $\log_{10}(\text{LWP})$ are -2.15 (ours) and -1.87 (O'Connor et al. (2005) method) with an RMSD of 0.29.

6 Summary

We developed a method to simultaneously profile cloud and drizzle properties by exploiting the synergy of ground-based radar, lidar and microwave radiometer measurements. This method has the advantage that the (non-)presence of drizzle is inferred from the best-fit to the data, rather than being imposed prior to the retrieval. The lidar forward model simulates the attenuated backscatter not only below the cloud base but also a few hundred meters into the cloud, taking multiple scattering into account. The cloud and drizzle components are distinguished using a droplet size threshold of 13 microns, a choice that is empirically motivated by the results of numerous observational and numerical studies. The combined droplet size distribution of cloud and drizzle follows a bimodal gamma distribution function. To aid the retrieval and to ensure some level of smoothness, the general shape of cloud and drizzle vertical profiles are parametrized based on empirical findings in the literature. The retrieval products include a full set of microphysical parameters (LWC, droplet effective radius and number density) and the optical extinction coefficient for the cloud and drizzle components.

The cloud retrieval was tested using synthetic signals generated from LES output. The vertical (along the height) and horizontal (along the time axis) variations in the true microphysical properties were largely reproduced, thereby verifying forward models and the cloud LWC model. On average, the column-averaged effective radius and the column-integrated quantities (LWP and optical depth) were retrieved within 1% of the mean truth, while the number density was retrieved within 5%. This

indicates that in the absence of radar calibration error, the retrieval method can potentially achieve a high accuracy. From this LES exercise, it appears that when the radar calibration is inaccurate, assuming a single (incorrect) value of the shape parameter ν in the retrieval can introduce a significant bias. When ν was set as a free parameter in the fitting, temporal noise increased but the mean bias in the retrieved cloud properties decreased. A radar systematic error of 3 dB led to a mean bias of $\sim 15\%$ in LWP, r_e , and N and this value became higher when ν was fixed to the true value. Examples of the full cloud and drizzle retrieval using synthetic signals were given for the two drizzle cases implemented in the algorithm. The reflectivity decomposition into the cloud and drizzle components were most problematic in the upper half of the cloud. The retrieval of the cloud component appeared to be more robust than the drizzle.

The retrieval algorithm was applied to a dataset collected during the ACCEPT campaign in Cabauw, The Netherlands. Single layer liquid water clouds with a varying amount of virga were selected. The clouds were between 200-400 m thick with a LWP that varied mostly between 10 and 100 g/m². The cloud and drizzle LWP were found to be positively correlated, with drizzle LWP about two orders of magnitude smaller. The effective radius of the cloud droplets was found to be less than 5 microns on average, far lower than the threshold value of 13 microns. The mean drizzle effective radius at the cloud base where it is expected to be the largest amounted to ~ 22 microns, but increased to as high as ~ 60 microns in intense drizzle periods. The cloud number concentration averaged to about 549 cm⁻³, around four orders of magnitude larger than the drizzle number density. Such a ratio of cloud to drizzle number density is comparable to what has been measured for marine-stratocumulus clouds in the MASE-II experiment (Lu et al., 2009).

Different elements of the ACCEPT retrieval products were assessed through comparisons with the results of three independent retrieval methods. The first method relied on the lidar depolarization signal to retrieve cloud properties at 100 m above the cloud base. The second one used radar Doppler spectra to quantify drizzle reflectivity within the cloud boundaries. Lastly, the third technique derived drizzle LWP below the cloud base using information from both radar and lidar. Overall, the cloud and drizzle properties retrieved using the three retrieval methods and the method described in this paper show a high degree of consistency within the expected uncertainties. Considering the different approaches and the limitation of each retrieval technique, we find the agreement in the results encouraging.

In closing, the application examples of the retrieval algorithm presented here show promising results. Application to datasets with larger size and variety is necessary to establish and improve the validity of the method. Retrieval evaluations using radiances measured from space or on the ground, and comparison with CCN (cloud condensation nuclei) measurements could be part of the future development. From the computational point of view, there is room for optimization that would lead to a faster implementation of the algorithm and make it more suitable for a large-scale application (currently, it takes roughly 2.5 hours on a dual core, 3 GHz i7 MacBook Pro with 16 GB of RAM to process the 1.2-hour ACCEPT data on Oct 25, 2014 shown in Figures 7 and 8).

Appendix A: Drizzle retrieval using radar Doppler spectra

Kollias et al. (2011a, b) introduce a method to improve drizzle retrieval within stratiform clouds by analysing the higher-order moments of the radar Doppler spectrum. Luke and Kollias (2013) implement this method and show that it works successfully for almost 50% of the spectra close to the cloud top, decreasing to about 15% towards the cloud base. Here we apply the procedure in Luke and Kollias (2013) to the ACCEPT campaign dataset from the chosen time period in Section 4. The aim is to compare the drizzle reflectivity as derived from this spectral analysis with Z_{dzi} retrieved using the technique presented in this paper.

The Doppler spectra were acquired using the same cloud radar, so instrument calibration is not an issue. The spectra were recorded every second with a velocity resolution of 0.0825 m/s and a vertical resolution of 30 m. For comparison purposes, we used the time-height coordinate in Fig. 7 such that one composite spectrum corresponds to one time-height pixel. This means applying a running window of 30 sec at a particular height above the cloud base. The spectra were shifted such that their spectral peaks coincided and were then averaged per velocity bin. The average velocity of the individual peaks was assigned to be the velocity location of the composite spectrum peak. From each composite spectrum, we computed the skewness as an indicator of drizzle presence. Negative skewness suggests that drizzle is the dominant component of the spectrum. Since the spectral decomposition technique is valid only for spectra with positive skewness, we continued to process only the composite spectra having skewness equal to or larger than 0.1. Here, positive velocities correspond to downward motions (approaching the zenith-pointing radar).

Each spectrum was decomposed by assuming that the portion to the left of the maximum power was entirely due to the cloud –such that it represented the left half of cloud spectrum– and that the drizzle contribution remained to the right side of the spectral peak. The full cloud spectrum was constructed by assuming that the right half followed a Gaussian shape with a dispersion estimated from the known left portion. Having constructed a complete cloud spectrum, the drizzle spectrum was then produced from the difference between the the cloud and the composite spectra.

With the cloud and drizzle spectra at hand, the reflectivity of each can be simply calculated from the 0th moment. At this stage, we computed the correction factor to compensate for the turbulence as a function of the spectral broadening σ_t (Fig. A2 in Luke and Kollias (2013)). For spectra with σ_t larger than 0.1 m/s, this correction was applied to the drizzle reflectivity. To preserve the total power, the cloud reflectivity was corrected (reduced) by the same amount.

Fig. A1 presents the results of the procedure above. Panel (a) shows the observed (total) reflectivity computed from the 0th moment of the composite spectra. This is comparable to the reflectivities shown in Fig. 7. Note that the reflectivities from the spectra are not corrected for gas attenuation, while those from the Cloudnet product are. The cloud base location is marked by the black line as determined from the lidar attenuated backscatter profile. The skewness displayed in (b) is strongly negative around 4 and 5.6 hr coinciding with high reflectivities and a high amount of drizzle below cloud base. Spectral decomposition is not performed for pixels/areas with skewness less than 0.1 and these show up as white gaps in panels (d)-(h).

The mean Doppler velocity shown in (c) demonstrates primarily updraft motion above the cloud base with a few downdraft streaks, most notably around 17.2 hr. The mean velocity below cloud base is strikingly higher than above cloud base. This is

- indicative of falling drizzle drops that evaporate, cool the air and cause downdraft motions. The reflectivity-weighted mean velocity for drizzle in panel (d) includes air motion, to allow for a consistent comparison between velocities below and above the cloud base. Below the cloud base there is no retrieval so the velocity fields in (c) and (d) are identical. The cloud reflectivity shows similarities to the observed one, suggesting that cloud is the dominant component. In panel (f), the drizzle reflectivities below the cloud base are identical to the observed reflectivities in (a). Visually, there is a continuous transition between the reflectivities above and below the cloud base. In Fig. 12, we show that there is a mean difference of 4 dB between the drizzle reflectivities immediately above or below the cloud base (see Section 5.2). Finally, the air motion in panel (g) is determined from the average velocity of the spectral peaks within the 30-sec time interval during the construction of composite spectra, and the width of the composite spectra is presented in panel (h).
- 10 *Acknowledgements.* This research is supported by the Netherlands Organisation for Scientific Research (NWO) through the User Support Programme Space Research project ALW-GO/13-20. The ACCEPT campaign was partly funded by the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 262254 and was carried out in cooperation with the Leibniz Institute for Tropospheric Research (TROPOS), LMU Munich, and METEK GmbH. The authors thank Lukas Pfizenmaier (TU Delft) for his assistance with the ACCEPT dataset, and Alexander Myagkov and Patric Seifert (TROPOS) for providing the clean radar Doppler spectra. We are also
- 15 grateful to Ulrich Löhnert for the provision of the original microwave radiative transfer code upon which the microwave forward model used in this work was based, and to Christine Unal for the fruitful scientific discussions.

References

- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, doi:10.1126/science.245.4923.1227, 1989.
- Austin, R. T. and Stephens, G. L.: Retrieval of stratus cloud microphysical parameters using millimeter-wave radar and visible optical depth
5 in preparation for CloudSat: 1. Algorithm formulation, *Journal of Geophysical Research*, 106, 28, doi:10.1029/2000JD000293, 2001.
- Baedi, R., Boers, R., and Russchenberg, H.: Detection of Boundary Layer Water Clouds by Spaceborne Cloud Radar, *Journal of Atmospheric and Oceanic Technology*, 19, 1915, doi:10.1175/1520-0426(2002)019<1915:DOBLWC>2.0.CO;2, 2002.
- Benmoshe, N., Pinsky, M., Pokrovsky, A., and Khain, A.: Turbulent effects on the microphysics and initiation of warm rain in deep convective clouds: 2-D simulations by a spectral mixed-phase microphysics cloud model, *Journal of Geophysical Research (Atmospheres)*, 117,
10 D06220, doi:10.1029/2011JD016603, 2012.
- Boers, R., Jensen, J. B., and Krummel, P. B.: Microphysical and short-wave radiative structure of stratocumulus clouds over the Southern Ocean: Summer results and seasonal differences, *Quarterly Journal of the Royal Meteorological Society*, 124, 151–168, doi:10.1002/qj.49712454507, 1998.
- Boers, R., Acarreta, J. R., and Gras, J. L.: Satellite monitoring of the first indirect aerosol effect: Retrieval of the droplet concentration of
15 water clouds, *Journal of Geophysical Research (Atmospheres)*, 111, D22208, doi:10.1029/2005JD006838, 2006.
- Brandau, C. L., Russchenberg, H. W. J., and Knap, W. H.: Evaluation of ground-based remotely sensed liquid water cloud properties using shortwave radiation measurements, *Atmospheric Research*, 96, 366–377, doi:10.1016/j.atmosres.2010.01.009, 2010.
- de Roode, S. R. and Los, A.: The effect of temperature and humidity fluctuations on the liquid water path of non-precipitating closed-cell stratocumulus clouds, *Quarterly Journal of the Royal Meteorological Society*, 134, 403–416, doi:10.1002/qj.222, 2008.
- 20 Donovan, D. P. and van Lammeren, A. C. A. P.: Cloud effective particle size and water content profile retrievals using combined lidar and radar observations, 1, Theory and examples, *Journal of Geophysical Research: Atmospheres*, 106, 27, doi:10.1029/2001JD900243, 2001.
- Donovan, D. P., Klein Baltink, H., Henzing, J. S., de Roode, S. R., and Siebesma, A. P.: A depolarisation lidar-based method for the determination of liquid-cloud microphysical properties, *Atmospheric Measurement Techniques*, 8, 237–266, doi:10.5194/amt-8-237-2015, 2015.
- 25 Duynerkerke, G. P., de Roode, R. S., van Zanten Margreet, C., Calvo, J., Cuxart, J., Cheinet, S., Chlond, A., Grenier, H., Jonker, P. J., Köhler, M., Lenderink, G., Lewellen, D., Lappen, C.-L., Lock, P. A., Moeng, C.-H., Müller, F., Olmeda, D., Piriou, J.-M., Sánchez, E., and Sednev, I.: Observations and numerical simulations of the diurnal cycle of the EUROCS stratocumulus case, *Quarterly Journal of the Royal Meteorological Society*, 130, 3269–3296, doi:10.1256/qj.03.139, 2004.
- Eloranta, E. W.: Practical Model for the Calculation of Multiply Scattered Lidar Returns, *Applied Optics*, 37, 2464–2472,
30 doi:10.1364/AO.37.002464, 1998.
- Feingold, G., Boers, R., Stevens, B., and Cotton, W. R.: A modeling study of the effect of drizzle on cloud optical depth and susceptibility, *Journal of Geophysical Research*, 102, 13, doi:10.1029/97JD00963, 1997.
- Fielding, M. D., Chiu, J. C., Hogan, R. J., Feingold, G., Eloranta, E., O’Connor, E. J., and Cadeddu, M. P.: Joint retrievals of cloud and drizzle in marine boundary layer clouds using ground-based radar, lidar and zenith radiances, *Atmospheric Measurement Techniques*, 8,
35 2663–2683, doi:10.5194/amt-8-2663-2015, 2015.
- Fox, N. I. and Illingworth, A. J.: The Retrieval of Stratocumulus Cloud Properties by Ground-Based Cloud Radar., *Journal of Applied Meteorology*, 36, 485–492, doi:10.1175/1520-0450(1997)036<0485:TROSCP>2.0.CO;2, 1997.

- Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, *Journal of Geophysical Research (Atmospheres)*, 117, D02207, doi:10.1029/2011JD016457, 2012.
- Frisch, A. S., Fairall, C. W., and Snider, J. B.: Measurement of Stratus Cloud and Drizzle Parameters in ASTEX with a K_{α} -Band Doppler Radar and a Microwave Radiometer., *Journal of Atmospheric Sciences*, 52, 2788–2799, doi:10.1175/1520-0469(1995)052<2788:MOSCAD>2.0.CO;2, 1995a.
- Frisch, A. S., Lenschow, D. H., Fairall, C. W., Schubert, W. H., and Gibson, J. S.: Doppler Radar Measurements of Turbulence in Marine Stratiform Cloud during ASTEX., *Journal of Atmospheric Sciences*, 52, 2800–2808, doi:10.1175/1520-0469(1995)052<2800:DRMOTI>2.0.CO;2, 1995b.
- Gerber, H.: Microphysics of Marine Stratocumulus Clouds with Two Drizzle Modes., *Journal of Atmospheric Sciences*, 53, 1649–1662, doi:10.1175/1520-0469(1996)053<1649:MOMSCW>2.0.CO;2, 1996.
- Giangrande, S. E., Babb, D. M., and Verlinde, J.: Processing Millimeter Wave Profiler Radar Spectra, *Journal of Atmospheric and Oceanic Technology*, 18, 1577–1583, doi:10.1175/1520-0426(2001)018<1577:PMWPRS>2.0.CO;2, 2001.
- Gonçalves, F. L. T., Martins, J. A., and Silva Dias, M. A.: Shape parameter analysis using cloud spectra and gamma functions in the numerical modeling RAMS during LBA Project at Amazonian region, Brazil, *Atmospheric Research*, 89, 1–11, doi:10.1016/j.atmosres.2007.12.005, 2008.
- Hartmann, D. L., Ockert-Bell, M. E., and Michelsen, M. L.: The Effect of Cloud Type on Earth’s Energy Balance: Global Analysis., *Journal of Climate*, 5, 1281–1304, doi:10.1175/1520-0442(1992)005<1281:TEOCTO>2.0.CO;2, 1992.
- Heus, T., van Heerwaarden, C. C., Jonker, H. J. J., Pier Siebesma, A., Axelsen, S., van den Dries, K., Geoffroy, O., Moene, A. F., Pino, D., de Roode, S. R., and Vilöguerau de Arellano, J.: Formulation of the Dutch Atmospheric Large-Eddy Simulation (DALES) and overview of its applications, *Geoscientific Model Development*, 3, 415–444, doi:10.5194/gmd-3-415-2010, 2010.
- Hogan, R. J.: Fast approximate calculation of multiply scattered lidar returns, *Applied Optics*, 45, 5984–5992, doi:10.1364/AO.45.005984, 2006.
- Hogan, R. J.: Fast Lidar and Radar Multiple-Scattering Models. Part I: Small-Angle Scattering Using the Photon Variance-Covariance Method, *Journal of Atmospheric Sciences*, 65, 3621, doi:10.1175/2008JAS2642.1, 2008.
- Illingworth, A. J., Hogan, R. J., O’Connor, E. J., Bouniol, D., Brooks, M. E., Delanoë, J., Donovan, D. P., Eastment, J. D., Gaussiat, N., Goddard, J. W. F., Haeffelin, M., Baltink, H. K., Krasnov, O. A., Pelon, J., Piriou, J.-M., Protat, A., Russchenberg, H. W. J., Seifert, A., Tompkins, A. M., van Zadelhoff, G.-J., Vinit, F., Willén, U., Wilson, D. R., and Wrench, C. L.: Cloudnet, *Bulletin of the American Meteorological Society*, 88, 883, doi:10.1175/BAMS-88-6-883, 2007.
- Klett, J. D.: Stable analytical inversion solution for processing lidar returns, *Applied Optics*, 20, 211–220, doi:10.1364/AO.20.000211, 1981.
- Kollias, P., RéMillard, J., Luke, E., and Szyrmer, W.: Cloud radar Doppler spectra in drizzling stratiform clouds: 1. Forward modeling and remote sensing applications, *Journal of Geophysical Research (Atmospheres)*, 116, D13201, doi:10.1029/2010JD015237, 2011a.
- Kollias, P., Szyrmer, W., RéMillard, J., and Luke, E.: Cloud radar Doppler spectra in drizzling stratiform clouds: 2. Observations and microphysical modeling of drizzle evolution, *Journal of Geophysical Research (Atmospheres)*, 116, D13203, doi:10.1029/2010JD015238, 2011b.
- Kovalev, V. A.: Sensitivity of the lidar solution to errors of the aerosol backscatter-to-extinction ratio: influence of a monotonic change in the aerosol extinction coefficient, *Applied Optics*, 34, 3457, doi:10.1364/AO.34.003457, 1995.
- Liebe, H. J., Hufford, G. A., and Manabe, T.: Millimeter-wave attenuation and delay rates due to fog/cloud conditions, *IEEE Transactions on Antennas and Propagation*, 37, 1617–1623, doi:10.1109/8.45106, 1989.

- Liebe, H. J., Hufford, G. A., and Cotton, M. G., eds.: Propagation modeling of moist air and suspended water/ice particles at frequencies below 1000 GHz, 1993.
- Liu, Y., Geerts, B., Miller, M., Daum, P., and McGraw, R.: Threshold radar reflectivity for drizzling clouds, *Geophysical Research Letters*, 35, L03807, doi:10.1029/2007GL031201, 2008.
- 5 Löhnert, U., Crewell, S., Simmer, C., and Macke, A.: Profiling Cloud Liquid Water by Combining Active and Passive Microwave Measurements with Cloud Model Statistics, *Journal of Atmospheric and Oceanic Technology*, 18, 1354–1366, doi:10.1175/1520-0426(2001)018<1354:PCLWBC>2.0.CO;2, 2001.
- Lu, M.-L., Sorooshian, A., Jonsson, H. H., Feingold, G., Flagan, R. C., and Seinfeld, J. H.: Marine stratocumulus aerosol-cloud relationships in the MASE-II experiment: Precipitation susceptibility in eastern Pacific marine stratocumulus, *Journal of Geophysical Research* (Atmospheres), 114, D24203, doi:10.1029/2009JD012774, 2009.
- 10 Luke, E. P. and Kollias, P.: Separating Cloud and Drizzle Radar Moments during Precipitation Onset Using Doppler Spectra, *Journal of Atmospheric and Oceanic Technology*, 30, 1656–1671, doi:10.1175/JTECH-D-11-00195.1, 2013.
- Mace, G. G. and Sassen, K.: A constrained algorithm for retrieval of stratocumulus cloud properties using solar radiation, microwave radiometer, and millimeter cloud radar data, *Journal of Geophysical Research*, 105, 29, doi:10.1029/2000JD900403, 2000.
- 15 Magaritz, L., Pinsky, M., Krasnov, O., and Khain, A.: Investigation of Droplet Size Distributions and Drizzle Formation Using A New Trajectory Ensemble Model. Part II: Lucky Parcels, *Journal of Atmospheric Sciences*, 66, 781, doi:10.1175/2008JAS2789.1, 2009.
- Martucci, G. and O'Dowd, C. D.: Ground-based retrieval of continental and marine warm cloud microphysics, *Atmospheric Measurement Techniques*, 4, 2749–2765, doi:10.5194/amt-4-2749-2011, 2011.
- McFarlane, S. A., Evans, K. F., and Ackerman, A. S.: A Bayesian algorithm for the retrieval of liquid water cloud properties from microwave radiometer and millimeter radar data, *Journal of Geophysical Research (Atmospheres)*, 107, 4317, doi:10.1029/2001JD001011, 2002.
- 20 Miles, N. L., Verlinde, J., and Clothiaux, E. E.: Cloud Droplet Size Distributions in Low-Level Stratiform Clouds., *Journal of Atmospheric Sciences*, 57, 295–311, doi:10.1175/1520-0469(2000)057<0295:CDSIL>2.0.CO;2, 2000.
- Myagkov, A., Seifert, P., Wandinger, U., Bühl, J., and Engelmann, R.: Relationship between temperature and apparent shape of pristine ice crystals derived from polarimetric cloud radar observations during the ACCEPT campaign, *Atmospheric Measurement Techniques*, 9, 3739–3754, doi:10.5194/amt-9-3739-2016, 2016.
- 25 O'Connor, E. J., Hogan, R. J., and Illingworth, A. J.: Retrieving Stratocumulus Drizzle Parameters Using Doppler Radar and Lidar., *Journal of Applied Meteorology*, 44, 14–27, doi:10.1175/JAM-2181.1, 2005.
- Peter, R. and Kämpfer, N.: Radiometric determination of water vapor and liquid water and its validation with other techniques, *Journal of Geophysical Research*, 97, 18, doi:10.1029/92JD01717, 1992.
- 30 Pfitzenmaier, L., Dufournet, Y., Unal, C. M. H., and Russchenberg, H. W. J.: Retrieving Fall Streaks within Cloud Systems Using Doppler Radar, *Journal of Atmospheric and Oceanic Technology*, 34, 905–920, doi:10.1175/JTECH-D-16-0117.1, 2017.
- Pinsky, M. B. and Khain, A. P.: Effects of in-cloud nucleation and turbulence on droplet spectrum formation in cumulus clouds, *Quarterly Journal of the Royal Meteorological Society*, 128, 501–533, doi:10.1256/003590002321042072, 2002.
- Ramanathan, V., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., and Hartmann, D.: Cloud-Radiative Forcing and Climate: Results from the Earth Radiation Budget Experiment, *Science*, 243, 57–63, doi:10.1126/science.243.4887.57, 1989.
- 35 Rosenfeld, D.: Suppression of Rain and Snow by Urban and Industrial Air Pollution, *Science*, 287, 1793–1796, doi:10.1126/science.287.5459.1793, 2000.

- Rosenfeld, D. and Gutman, G.: Retrieving microphysical properties near the tops of potential rain clouds by multispectral analysis of AVHRR data, *Atmospheric Research*, 34, 259–283, doi:10.1016/0169-8095(94)90096-5, 1994.
- Rosenfeld, D., Wang, H., and Rasch, P. J.: The roles of cloud drop effective radius and LWP in determining rain properties in marine stratocumulus, *Geophysical Research Letters*, 39, L13801, doi:10.1029/2012GL052028, 2012.
- 5 Rosenkranz, P. W.: Water vapor microwave continuum absorption: A comparison of measurements and models, *Radio Science*, 33, 919–928, doi:10.1029/98RS01182, 1998.
- Sauvageot, H. and Omar, J.: Radar Reflectivity of Cumulus Clouds, *Journal of Atmospheric and Oceanic Technology*, 4, 264, doi:10.1175/1520-0426(1987)004<0264:RROCC>2.0.CO;2, 1987.
- Slingo, A.: Sensitivity of the Earth’s radiation budget to changes in low clouds, *Nature*, 343, 49–51, doi:10.1038/343049a0, 1990.
- 10 Storn, R. and Price, K.: Differential Evolution - A Simple and Efficient Heuristic for Global Optimization over Continuous Spaces, *Journal of Global Optimization*, 11, 341–359, doi:10.1023/A:1008202821328, 1997.
- Suzuki, K., Nakajima, T. Y., and Stephens, G. L.: Particle Growth and Drop Collection Efficiency of Warm Clouds as Inferred from Joint-CloudSat and MODIS Observations, *Journal of Atmospheric Sciences*, 67, 3019–3032, doi:10.1175/2010JAS3463.1, 2010.
- vanZanten, M. C., Stevens, B., Vali, G., and Lenschow, D. H.: Observations of Drizzle in Nocturnal Marine Stratocumulus., *Journal of Atmospheric Sciences*, 62, 88–106, doi:10.1175/JAS-3355.1, 2005.
- 15 Voors, R., Donovan, D., Acarreta, J., Eisinger, M., Franco, R., Lajas, D., Moyano, R., Pirondini, F., Ramos, J., and Wehr, T.: ECSIM: the simulator framework for EarthCARE, in: *Sensors, Systems, and Next-Generation Satellites XI*, vol. 6744 of *SPIE Proceedings*, p. 67441Y, doi:10.1117/12.737738, 2007.
- Walko, R. L., Cotton, W. R., Meyers, M. P., and Harrington, J. Y.: New RAMS cloud microphysics parameterization part I: the single-moment scheme, *Atmospheric Research*, 38, 29–62, doi:10.1016/0169-8095(94)00087-T, 1995.
- 20 Wang, J. and Geerts, B.: Identifying drizzle within marine stratus with W-band radar reflectivity, *Atmospheric Research*, 69, 1–27, doi:10.1016/j.atmosres.2003.08.001, 2003.
- Westbrook, C. D., Hogan, R. J., O’Connor, E. J., and Illingworth, A. J.: Estimating drizzle drop size and precipitation rate using two-colour lidar measurements, *Atmospheric Measurement Techniques*, 3, 671–681, 2010.
- 25 Westwater, E. R.: The accuracy of water vapor and cloud liquid determination by dual-frequency ground-based microwave radiometry, *Radio Science*, 13, 677–685, doi:10.1029/RS013i004p00677, 1978.
- Wood, R.: Drizzle in Stratiform Boundary Layer Clouds. Part I: Vertical and Horizontal Structure., *Journal of Atmospheric Sciences*, 62, 3011–3033, doi:10.1175/JAS3529.1, 2005a.
- Wood, R.: Drizzle in Stratiform Boundary Layer Clouds. Part II: Microphysical Aspects., *Journal of Atmospheric Sciences*, 62, 3034–3050, doi:10.1175/JAS3530.1, 2005b.
- 30

Table 1. Lower and upper bounds for the state vector values used in the optimization. The units, when relevant, are given in the square brackets.

State vector element	Lower limit	Upper limit
ν_{cld}	2	20
\hat{h} (cloud and drizzle)	0.001	35
W (cloud and drizzle)	0.001	1
N_{ad} (cloud; [m^{-3}])	10^7	5×10^9
ft_{cb}	0	1
ft_{ct}	-1	0
ν_{dzl}	1	10
q	0.001	0.03
α_{ic} [α_{cld}]	10^{-5}	10^{-2}
α_{cb} [m^{-1}]	10^{-6}	10^{-4}
α_{id} [α_{cb}]	0.001	1
α'_0 [m^{-1}]	10^{-10}	10^{-3}
p_{cb}	1	3

Table 2. Values of the LWP, r_e , optical depth and the number concentration (as shown in Fig. 5), averaged over the horizontal distance. The RMSD between the truth and the retrieved values is given as the error of the retrievals. The last column is given here for completeness - see discussion in the text.

	Truth	Retrieval with $\nu = 5.5$	Retrieval with optimized ν
LWP [g/m^2]	171.68	171.96 ± 5.31	171.64 ± 9.60
Effective radius [μm]	20.26	20.39 ± 0.23	20.44 ± 0.84
Optical depth	12.68	12.62 ± 0.46	12.56 ± 0.45
Number concentration [cm^{-3}]	21.26	20.30 ± 1.47	20.83 ± 5.14




Figure 1. The retrieval flowchart. For each main step, a reference to the section that provides the details is given.




Figure 2. Lidar attenuated backscatter profiles as a function of height. The circles outline an example of a measured β profile, taken during the ACCEPT campaign on October 26, 2014 at 5.04 hour UTC. The dashed black line shows the forward-modelled β that best fits the measurements when no smoothing is applied to LWC. The solid red line represents the forward-modelled β when LWC is smoothed.




Figure 3. Synthetic signals generated using ECSIM based on LES. The top and middle panels compare radar reflectivity Z and lidar attenuated backscatter β , respectively, between the synthetic signals that are fed to the retrieval as input (left) and best-fit produced by the retrieval (right). Panels (e) and (f): histograms of the differences between the true and the retrieved signals (truth-retrieval). The spread of the distribution is indicated by the blue vertical lines that mark the interval within which 95% of the total occurrences are found. The red curves show the cumulative distributions. Panel (g) shows brightness temperatures T_B averaged over distance at each frequency channel: black circles and line show the synthetic measurements (data mean) while the dashed red line shows the retrieval mean. In (h) we plot the standard deviation of the data mean (black line in (g)) divided by the data mean itself (black) and the RMSD between the retrieval and the data, divided by the data mean (red).




Figure 4. True (left panels) and retrieved (right panels) microphysical and optical properties corresponding to the synthetic signal shown in Fig. 3 as a function of vertical and horizontal distance. The first four rows from top to bottom: liquid water content, effective radius, optical extinction coefficient and number concentration. The last row: histograms of the differences between the true and the retrieved cloud properties (truth - retrieved). The spread of the distribution is indicated by the blue vertical lines that mark the interval within which 95% of the total occurrences are found. The right y-axes of the four histograms are all identical and correspond to the cumulative distributions shown by the red curves.




Figure 5. Microphysical and optical properties collapsed along the vertical axis: N is column-averaged, LWC and α are integrated into LWP and optical depth, respectively, and r_e are vertically averaged with the corresponding α as the weights. The black line represents the truth and the circles are the retrieved values. The error bars denotes the random errors obtained from the Monte Carlo realizations.




Figure 6. Examples of cloud and drizzle profiling for the two drizzle cases described in section 2.2.2. Panels Ia-II show the results for case I and panels IIa-III for case II. The red dash-dotted lines mark the cloud base height $z_{cb,opt}$. Filled circles: synthetic signals, squares: the cloud truth, diamonds: drizzle truth; red solid lines: retrieved signals, blue dashed lines: retrieved cloud properties and blue dotted lines: retrieved drizzle properties.




Figure 7. Measured and retrieved signals of the selected cases from the ACCEPT campaign. The breaks along the horizontal axes in (a)-(f) mark the change of date from October 25 to 26, 2014. Panels (a)-(d) show the radar reflectivity as observed and as retrieved, along with the decomposition into drizzle and cloud reflectivities. The black line delineates the cloud base determined in the retrieval. Panels (e)-(f) display the observed and the retrieved attenuated lidar backscatter β . Panels (g) and (h): histograms of the differences between the observed and the retrieved (total) signals (observed - retrieved). The red curves show the cumulative distributions. The spread of the distribution is indicated by the blue vertical lines that mark the interval within which 95% of the total occurrences are found. Panel (i) shows the brightness temperatures T_B averaged over distance at each frequency channel: black circles and line show the observations while the dashed red line shows the retrieval. In (j) we plot the standard deviation of the observation mean divided by the mean itself (black) and the RMSD between the retrieval and the observation, divided by the observation mean (red).




Figure 8. Optical and microphysical parameters of the cloud (left) and the drizzle (right) as obtained from the retrieval, as a function of time and height. From top to bottom: liquid water content, effective radius, extinction coefficient and number concentration.




Figure 9. Mean vertical profiles of the cloud (solid lines) and drizzle (dashed lines) properties from the observation on October 26, 2014 between 3.8 and 4.0 hr UTC. Clockwise from the top left: radar reflectivity factor in dBZ, liquid water content in g/m^3 , number density per cm^3 and effective radius in microns. Note that the number density of the drizzle has been multiplied by a factor of 1000 for illustration purposes.




Figure 10. Time series of the optical and microphysical properties of the cloud at 100 m above the cloud base. The break along the horizontal axis marks the change of day from Oct 25 to Oct 26. The data around 5.6 hr is deemed unsuitable for retrieval using the depolarization lidar method. The cloud properties and their respective uncertainties derived using the depolarization method are shown by the blue lines and the shaded area. The results of our retrieval are represented by the black lines (temporal resolution: 30 sec) and red lines (time average, to match the time stamps and the 180-sec resolution of the depolarization results). Top to bottom: liquid water content, effective radius, extinction coefficient, number concentration and the altitude at which the cloud properties are evaluated.




Figure 11. Radar reflectivity from our retrieval ('retrieved Z') plotted against those derived from the Doppler spectral decomposition ('doppler Z'). Drizzle, cloud and total reflectivities are shown separately in (a), (b), (c), respectively. The points are color-coded according to their location in the cloud. Red points are from the cloud base area, blue points from the middle of the cloud and green points are from the cloud top region (see the text for details). The insets in (a) and (b) show the same plot with the blue points excluded. The diagonal line is the one-to-one line.




Figure 12. Comparison between the drizzle reflectivities at one range gate above and one range gate below the cloud base. The reflectivities above the cloud base are obtained from the Doppler spectral decomposition. The ones below the cloud base are simply the observed reflectivity. The diagonal line marks the one-to-one correspondence.




Figure 13. Comparison with the drizzle liquid water path below the cloud base as retrieved using the technique of O'Connor et al. (2005), available as one of the Cloudnet products. The diagonal line marks the one-to-one correspondence.




Figure A1. Results of the retrieval using Doppler spectra data on Oct 25 and 26, 2014: (a) zeroth moment of the Doppler composite spectra, (b) skewness of the composite spectra above the cloud base, (c)-(d) first moments of the composite spectra and the drizzle spectra, (e)-(f) 0th moments of the cloud and the drizzle spectra, respectively, as obtained from the spectral decomposition, (g) air velocity and (h) spectral broadening due to turbulence. The breaks along the time axis mark the change of day. The black line delineates the cloud base.