



Observations of VOC emissions and photochemical products over US oil- and gas-producing regions using high-resolution H_3O^+ CIMS (PTR-ToF-MS)

5 Abigail Koss^{1,2,3}, Bin Yuan^{1,2†}, Carsten Warneke^{1,2}, Jessica B. Gilman¹, Brian M. Lerner^{1,2*}, Patrick R. Veres^{1,2}, Jeff Peischl^{1,2}, Scott Eilerman^{1,2}, Rob Wild^{1,2}, Steven S. Brown^{1,3}, Chelsea R. Thompson^{1,2}, Thomas Ryerson¹, Thomas Hanisco⁴, Glenn M. Wolfe^{4,5}, Jason M. St. Clair^{4,5}, Mitchell Thayer⁶, Frank N. Keutsch^{6,7}, Shane Murphy⁸, Joost de Gouw^{2,3}

- 10 1. NOAA Earth System Research Laboratory (ESRL), Chemical Sciences Division, Boulder, CO, USA
2. Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA
3. Department of Chemistry and Biochemistry, University of Colorado Boulder, CO, USA
4. NASA Goddard Space Flight Center, Greenbelt, MD, USA
5. Joint Center for Earth Systems Technology, University of Maryland Baltimore County, Baltimore, MD, USA
15 6. University of Wisconsin Madison, Madison, WI, USA
7. Paulson School of Engineering and Applied Sciences and Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA
8. University of Wyoming, Laramie, WY, USA
† now at Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen, Switzerland
20 * now at Aerodyne Research, Inc., Billerica, MA, USA

Correspondence to: Carsten Warneke (Carsten.warneke@noaa.gov)

Abstract. VOCs related to oil and gas extraction operations in the United States were measured by H_3O^+ chemical ionization time-of-flight mass spectrometry (H_3O^+ ToF-CIMS / PTR-ToF-MS)
25 from aircraft during the SONGNEX campaign in March-April 2015. This work presents an overview of major VOC species measured in nine oil and gas producing regions, and a more detailed analysis of H_3O^+ ToF-CIMS measurements in the Permian Basin within Texas and New Mexico. Mass spectra are dominated by small photochemically produced oxygenates, and compounds typically found in crude oil: aromatics, cyclic alkanes, and alkanes. Mixing ratios of
30 aromatics were frequently as high as those measured downwind of large urban areas. In the Permian, the H_3O^+ ToF-CIMS measured a number of underexplored or previously unreported species, including aromatic and cycloalkane oxidation products, nitrogen heterocycles including pyrrole ($\text{C}_4\text{H}_5\text{N}$) and pyrroline ($\text{C}_4\text{H}_7\text{N}$), H_2S , and a diamondoid (adamantane) or unusual monoterpene. We additionally assess the specificity of a number of ion masses resulting from H_3O^+
35 ion chemistry previously reported in the literature, including several new or alternate interpretations.



1. Introduction

Recent advances in fossil fuel extraction technology, especially horizontal drilling and hydraulic fracturing, have enabled a surge in crude oil and natural gas production in several regions across the United States over the past decade (US Energy Information Administration). A particular environmental concern is the release of air pollutants. Emissions can affect global climate, by the release of greenhouse gases (Miller et al., 2013; Brandt et al., 2014); regional air quality, by contributing ozone and particulate precursor species (Kemball-Cook et al., 2010; Edwards et al., 2014; McDuffie et al., 2016); and local air quality, by releasing air toxics harmful to human health (McKenzie et al., 2012; Adgate et al., 2014).

Detailed measurements of volatile organic compound (VOC) emissions, and their atmospheric reaction products, are needed to understand and mitigate these air quality concerns. Several studies have used gas chromatography (GC) techniques to characterize oil- and gas- related VOCs in relatively high chemical detail (Simpson et al., 2010; Gilman et al., 2013; Swarthout et al., 2013). These studies have demonstrated that comprehensive VOC characterization is invaluable for source identification and air quality modeling in these regions. To date, there are few such studies, and they have been limited in measurement of secondary species and in time resolution. The Permian Basin, located in west Texas and eastern New Mexico, is the physically largest and most productive oil field in the United States, but non-methane VOC measurements from this region have rarely been reported.

Proton-transfer-reaction mass spectrometry (PTR-MS) is a well-established chemical ionization technique used to measure VOCs, especially polar and aromatic species, in ambient air. This technique uses H_3O^+ reagent ions and can have time resolution of 1 Hz or better. The recent development of PTR-MS instruments that use high-resolution time-of-flight mass analyzers (PTR-ToF-MS) has greatly expanded the number of measureable species, enhanced the technique's suitability to mobile platforms, and improved our ability to speciate specific ion masses (Jordan et al., 2009; Graus et al., 2010; Cappellin et al., 2011; Yuan et al., 2016a). For example, the resulting complex PTR-ToF-MS mass spectra in forested environments (Kim et al., 2010) and biomass burning (Brilli et al., 2014; Stockwell et al., 2015) have been reported. A recent study comparing PTR-ToF-MS and PTR-quadrupole MS instruments in an oil and gas producing



70 region in Utah pointed to the scientific advances possible with the application of PTR-ToF-MS, especially the separate measurement of hydrocarbon masses from isobaric oxygenates, and the measurement of previously undetectable photo-oxidation products (Warneke et al., 2015). In this work PTR-MS refers to the proton-transfer technique, and PTR-ToF-MS refers to PTR-MS instruments with a high-resolution time-of-flight mass analyzer. The instrument discussed in this work is called “H₃O⁺ ToF-CIMS”, an instrument similar to the PTR-ToF-MS but developed at NOAA.

75 This work comprises a detailed analysis of PTR-ToF-MS mass spectra obtained from measurements in oil- and gas-producing regions, supported by a comprehensive suite of other chemical instrumentation. We outline an interpretation of PTR-ToF-MS measurements in these regions, and report observed mixing ratios of commonly measured PTR-MS species in nine oil and gas producing regions. We present detailed VOC observations for the Permian Basin. This work provides detailed information about the VOC chemistry of these regions, will aid in the interpretation of PTR-MS (especially PTR-ToF-MS) measurements in oil and gas producing regions, and will support future analysis of ambient measurements in these regions.

2. Methods

85 2.1 Measurement location and context

Measurements were made from the NOAA WP-3D Orion research aircraft during the Shale Oil and Natural Gas Nexus (SONGNEX) campaign in March and April 2015. The SONGNEX campaign surveyed nine large oil and natural gas production regions in the central United States, several smaller producing regions, and locations with associated infrastructure. All research flights 90 took place during daytime (late morning to mid-afternoon). The aircraft was equipped with a suite of chemical and meteorological instrumentation, which is described in section 2.2. This work reports measurements in nine regions: Bakken (ND), Upper Green River (WY), Uintah (UT), Denver-Julesburg (CO), San Juan (CO, NM), Permian (NM, TX), Barnett (TX), Eagle Ford (TX), and Haynesville (TX, LA) (Fig. 1a). SONGNEX measurements coincided with a peak in fossil 95 fuel production in many of these regions, but were just after a downturn in the drilling of new wells due to a drop in the price of crude oil (Fig. 1b, 1c). Analysis is restricted to data collected in the



fossil fuel producing areas of the basins, and within the PBL (typically $\leq 600\text{m}$ AGL). Selection of data is shown in Fig. SI 3.

100 The Permian oil and gas field, located in western Texas and eastern New Mexico, was surveyed on three flights, on April 6, April 9, and April 23, 2015. Detailed interpretation of the H_3O^+ ToF-CIMS data focuses on measurements taken during the Permian flight on April 23. This flight featured high signal on many VOC masses, providing the best overall signal-to-noise ratio of any SONGNEX flight and allowing the observation of many VOCs that may have been below
105 detection limit on other flights. Additionally, there are few non-oil and gas emission sources in this region, which simplifies the interpretation of VOC measurements. During the April 23 flight, the average boundary layer temperature was $19\text{ }^\circ\text{C}$, the relative humidity ranged from 20-80% (average 34%), and wind speeds were typically between 2 and 10 m s^{-1} , averaging 5.4 m s^{-1} . The maximum concentration of ozone measured was 62 ppb, and the average 49 ppb.

110

The Permian Basin is an approximately $200\,000\text{ km}^2$ area encompassing a number of geologically distinct fossil fuel producing reservoirs, including several shale oil formations. The region is characterized by significant and intense oil production activity (accounting for nearly 20% of 2013 US domestic oil production), currently largely driven by recent development of shale oil
115 formations (Budzik and Perrin, 2014). As of September 2015, there were approximately 124 000 actively producing oil wells and 10 000 gas wells in this region, and approximately 1300 new wells were drilled in April 2015 (Railroad Commission of Texas, 2017; State of New Mexico Oil Conservation Division, 2017). Point sources reported in the NEI 2011 inventory are largely associated with oil and gas production, including, for example, refineries, processing facilities,
120 compressor stations, and tank batteries. The region has a population of 1 million, of whom about one-fifth live in the largest urban area, Midland-Odessa (U.S. Census). The climate is arid and the land cover consists mainly of desert and grassland. The April 23 SONGNEX flight track; locations of oil and gas wells, urban areas, and NEI 2011 point sources, and the spatial distributions of a few VOCs of interest are shown in Figure 2.

125

To support the interpretation of SONGNEX H_3O^+ ToF-CIMS measurements, this work also refers to measurements taken by an Ionicon PTR-ToF-MS instrument in the Uintah Basin, Utah, oil and



gas field during winter of 2013. The Ionicon PTR-ToF-MS instrument relies on the same measurement principle as the SONGNEX H_3O^+ ToF-CIMS and has similar mass resolution. The PTR-ToF-MS Uintah Basin measurements have been previously described by Warneke et al. (2015), and a comparison between the H_3O^+ ToF-CIMS and PTR-ToF-MS instruments is given by Yuan et al. (2016a). Finally, we include H_3O^+ ToF-CIMS headspace measurements (at 25 °C) of a crude oil sample. The sample was purchased from ONTA (Geology) Goal and Petroleum Inc. and is a blend of oil from several reservoirs in West Texas. The sample was bottled at a central collection facility after transport in tanker trucks and before being sent to a refinery. It is possible that some VOCs were removed prior to transport: sulfur and nitrogen-containing organics are often removed at processing facilities, although we do not know how this sample was treated at the collection facility. Some of the most highly-volatile VOCs may have been depleted during transport.

2.2 Instrumentation

2.2.1 Description of H_3O^+ ToF-CIMS instrument

The H_3O^+ ToF-CIMS and its operation during the SONGNEX campaign have been previously reported (Yuan et al., 2016a). The basic operational principle is the same as other PTR-MS instruments. H_3O^+ ions are generated from water vapor in a hollow cathode discharge ion source. The H_3O^+ reagent ions are then mixed with ambient air, containing VOCs, in a drift tube section. The proton from H_3O^+ is transferred to VOCs with sufficiently high proton affinity, and the resulting ionized VOCs are transferred to a mass analyzer (in the H_3O^+ ToF-CIMS, VOC ions are guided through a set of RF-only segmented quadrupoles to a time-of-flight unit). The hollow cathode, drift tube, pressure controlled inlet, background, and calibration components are custom built; the ion transmission and time-of-flight analyzer were produced by Aerodyne Research Inc./Tofwerk. The H_3O^+ ToF-CIMS has a drift tube E/N (electric field strength divided by number density) of about 120×10^{-17} Td (V cm^2). Data are presented in this work at 1s time resolution unless otherwise noted.

During field operation, the instrument background was determined for 90 seconds every 20-40 minutes by flowing ambient air through a catalyst, and a 10-component gas standard was added



for 90 seconds every 1-2 hours (single-point calibration) to record instrument stability and sensitivity. In addition to the in-flight single-point calibrations, multi-point calibrations were performed at the beginning and end of each flight. A small amount of trichlorobenzene ($C_6H_3Cl_3$) was continuously introduced into the instrument as a mass calibrant. Data were analyzed using Tofware high resolution peak-fitting software (Aerodyne Research Inc./Tofwerk). Data were then corrected for humidity-dependent sensitivities and background-subtracted. In addition to the in-flight calibrations, laboratory calibrations of a larger number of species were performed using standard cylinders and permeation tubes. More details on the data quality assurance procedures related to instrument operation, background subtraction, humidity dependence, and calibration are included in Yuan et al. (2016a).

The H_3O^+ ToF-CIMS instrument has a mass resolution of approximately 3900-5900 $m/\Delta m$ over the mass range discussed in this work (m/z 12-200), which is sufficient to determine the elemental formulas of most detected ion masses. Evaluation of data quality related to high-resolution peak-fitting are discussed in Supplementary Information (Section SI 1). Also included in the Supplementary Information are an extensive set of high-resolution mass spectra showing isobaric contributions to nominal masses, which will be useful to operators of both unit-mass- and high-resolution PTR-MS instruments (Fig. SI 4).

In this work we report signal intensity using normalized counts-per-second (ncps), and VOC mixing ratio (ppbv) for calibrated species. Normalized counts-per-second (ncps) is the instrument signal relative to $10^6 H_3O^+$ ion counts, corrected for humidity effects, and background subtracted. VOC sensitivities were determined by (1) direct calibration, where available; (2) calculated using proton-transfer rate constants, either known or calculated based on polarizability and dipole moment (Sekimoto et al., 2017); or (3) an average sensitivity determined from the calibrated and calculated sensitivities. Accuracy is within 15% for directly calibrated compounds and generally within a factor of two for calculated sensitivities. Some ion masses, such as m/z 45.992 NO_2^+ (Section 3.4.4) and m/z 81.070 $C_6H_8H^+$ (Section 3.4.3) have an ambiguous interpretation and these are discussed in terms of instrument signal (ncps) and not mixing ratio.



Measured 1-s detection limits (for a signal-to-noise ratio of 3) range from approximately 40 pptv (aromatics) to 400 pptv (methanol). Some mixing ratios reported in this work are smaller than 40 pptv. In a few cases, the signal-to-noise ratio is less than three, but variability is still discernable, and these species are discussed mostly for the *absence* of significant enhancement (e.g. styrene, cresol). Other species are presented as an average over a longer period of time: for example, average boundary layer enhancements presented in Figure 5, and time series in Figure 13. As averaging time increases, the limit of detection decreases: a typical aromatic compound with 1s detection limit of 40 pptv (sensitivity of 500 count ppbv⁻¹ s⁻¹, background of 5 counts per second) has a 10s detection limit of about 9 pptv (a calculation is included in the supplementary information, Section SI 2).

2.2.2 PTR-MS application to oil and gas emissions: strengths and limitations

In the WP-3D SONGNEX payload, the primary strengths of the H₃O⁺ ToF-CIMS include measurement of small acids and carbonyls not detected by other instruments, and a much higher measurement rate of aromatics and cyclic alkanes, which were also measured by the whole air sampler, which typically collected sample for five seconds once every three minutes (Lerner et al., 2017). Aromatics and compounds with heteroatoms are generally detected sensitively (aromatics: measured average 180 ncps/ppb, polar compounds: 200 ncps/ppb) with a few exceptions of compounds that can dissociate by dehydration, such as small alcohols and aldehydes (de Gouw and Warneke, 2007). Cyclopentane and cyclohexane cannot be detected, but alkyl-substituted cyclic alkanes are detected, at approximately 5% of the sensitivity of aromatics. Alkenes containing four or more carbon atoms are detected sensitively (estimated average 300 ncps/ppb), although larger alkenes can fragment at the high E/N conditions used in our instrument (Gueneron et al., 2015).

PTR-MS is notably limited in its measurement of alkanes: saturated alkanes smaller than hexane have too low proton affinity to be detected, and C₆ and larger branched-and straight-chain alkanes are detected with very low sensitivity, reacting with H₃O⁺ at a rate one or two orders of magnitude slower than aromatics (Arnold et al., 1998). Additionally, the larger alkanes fragment extensively (Gueneron et al., 2015). This behavior makes alkanes difficult to measure sensitively and selectively. This is particularly limiting in measurements of emissions from oil and gas operations,



where alkanes are typically dominant in terms of mixing ratio. NO_2^+ cannot be converted to a mixing ratio in a meaningful way and was excluded from the concentration comparison. For hydrocarbons, an average sensitivity was applied (Section 2.2.1).

220

During SONGNEX, fast measurement of methane and ethane were provided by cavity ring-down and direct-absorption spectroscopy (respectively), and speciated C2-C8 alkanes were measured by whole-air-sampling/GC-MS. Data from a number of other chemical instruments that were deployed during the SONGNEX mission are used in this analysis. Descriptions of these
225 instruments are given in Table 1. Further information can be found at <https://esrl.noaa.gov/csd/groups/csd7/measurements/2015songnex/P3/datainfo.html>.

3. Results and Discussion

3.1 Overview of H_3O^+ ToF-CIMS measurements

In this section, we provide a comparison of the mixing ratios of several volatile species between
230 all flights made in the San Juan, Uintah, Upper Green River, Denver-Julesburg, Barnett, Permian, Haynesville, Bakken, and Eagle Ford regions. Species measured by H_3O^+ ToF-CIMS that are included in this comparison are the sum of C6-C10 aromatics, methanol, H_2S , and acetone. Figure 3 shows an overview of the mixing ratios of these compounds in the boundary layer during each flight. Additionally, monoterpenes (measured by iWAS/GC-MS), NO_x (measured by the
235 chemiluminescence instrument), CH_4 (measured by cavity ring-down spectroscopy), and average wind speed within the boundary layer are shown in Figure 3 for chemical and meteorological context.

BTEX species (benzene, toluene, and C8 aromatics), and higher weight aromatics, are important
240 air toxics and ozone and aerosol precursors that are commonly reported in VOC measurements of oil and gas-producing regions. PTR-MS measures these compounds with high specificity and sensitivity (de Gouw and Warneke, 2007). Figure 3a shows that aromatics concentrations in oil and gas basins were typically high – comparable to concentrations downwind of large urban areas.



245 The H_3O^+ ToF-CIMS measurement of benzene, toluene, and C8 aromatics agrees well with
measurements made by whole-air-sampling followed by GCMS analysis (iWAS/GC-MS) (Yuan
et al., 2016a; Lerner et al., 2017). There were large differences in observed mixing ratios between
basins. On all flights, sharp, concentrated plumes of aromatics were encountered (resulting in
maximum and average mixing ratio much higher than median). Large differences in mixing ratio
250 between different flights in the same basin (e.g. Uintah, Denver-Julesburg) are partially the result
of differences in meteorological conditions while in other basins (Permian, Haynesville, Upper
Green River, Eagle Ford, Bakken), average mixing ratios were more consistent between flights.
Benzene was not measured by the H_3O^+ ToF-CIMS during the March 24 San Juan flight due to an
instrument issue. For comparison, Figure 3 also shows boundary layer statistics from two flights
255 during other recent P3 aircraft campaigns: SENEX (June 16 2013 flight, Atlanta GA metropolitan
area) and CALNEX (May 05 2010 flight, Los Angeles CA metropolitan area). SENEX and
CALNEX data are from a PTR-quadrupole MS instrument. The maximum mixing ratios of
aromatics during every SONGNEX flight were considerably higher than those measured during
SENEX and CALNEX (note the log scale); averages within several basins (especially Uintah,
260 Denver-Julesburg, and Permian) were comparable to or higher than aromatics measured over the
Los Angeles metropolitan area (population of 18 million).

H_2S is an air toxic that can be emitted from oil and gas sources, and can seriously harm human
health (Tarver and Dasgupta, 1997; Li et al., 2014). Significant enhancements of H_2S were seen
265 only in the Permian and Haynesville regions (Fig. 3c). All three Permian flights saw broad
enhancements in H_2S likely associated with oil and gas production. Emissions from a paper mill
were captured during the Haynesville flights and a mixing ratio of 27.7 ppbv was measured 20.5
km downwind of the point source, which was the highest H_2S mixing ratio measured during
SONGNEX. The Eagle Ford flights had a higher limit-of-detection due to an unknown instrument
270 issue and the maximum mixing ratio is not statistically significant.

Oxygenated compounds provide insight into photochemical aging, and comprised the majority of
 H_3O^+ ToF-CIMS product ion signal. The most abundant oxygenated compounds were methanol
and small (C1- C4) carbonyls and acids. These compounds include m/z 45.034 $\text{C}_2\text{H}_4\text{OH}^+$
275 (acetaldehyde), m/z 59.049 $\text{C}_3\text{H}_6\text{OH}^+$ (acetone), m/z 73.065 $\text{C}_4\text{H}_8\text{OH}^+$ (2-butanone), m/z 61.028



$C_2H_4O_2H^+$ (acetic acid), and m/z 75.044 $C_3H_6O_2H^+$ (propionic acid). Support for our interpretation of these ion masses is provided in Section 3.4.1.

Mixing ratios of acetone were similar in each basin, with average mixing ratios within a range of
280 about 1.5 ppb (Fig. 3b). Mixing ratios of acetaldehyde, MEK, and acetic acid for each flight are in
Figure SI 5. The relative abundances of oxygenates were generally similar between basins (Fig.
4), although there was higher variability in organic acids and formaldehyde. The highest relative
abundances of organic acids were observed in the Denver-Julesburg Basin and are likely due to
primary emissions from concentrated animal feeding operations and not photochemistry (Eilerman
285 et al., 2016; Yuan et al., 2017).

Figure 4 compares the distribution of oxygenates to that measured during the UBWOS 2013
campaign (Utah Division of Air Quality, 2014), the SENEX and CALNEX flights, and the
CALNEX ground site in Los Angeles (Veres et al., 2011; Warneke et al., 2013). The mix of VOC
290 precursors from oil and gas fields (dominated by small alkanes) is very different from urban and
biogenically influenced air. However, the mix of products was similar to that measured during
CALNEX. The similarity between the SONGNEX and CALNEX profiles is somewhat surprising
since it has been shown that the oxidation mechanisms in urban and oil and gas producing regions
are quite different (Yuan et al., 2015). The distribution of carbonyls was similar to that measured
295 during the SENEX flight, but had significantly lower formaldehyde and formic acid, likely due to
the much lower concentration of their precursor, isoprene.

Methanol (detected at m/z 33.034 CH_4OH^+) was the single most abundant VOC detected by the
 H_3O^+ ToF-CIMS. There is agreement to within the stated uncertainties between the H_3O^+ ToF-
300 CIMS measurement and the iWAS/GC-MS measurement ($R^2 = 0.9$, slope = 1.23), despite the
difficulty in retrieving methanol from the whole-air-sampling system (Lerner et al., 2017). There
was high variability in methanol mixing ratios between basins, between flights within the same
basin, and within individual flights (Fig. 3d). Also shown, in Figure 3f, is the iWAS/GC-MS
measurement of monoterpenes, as a proxy for biogenic emissions.

305



The two flights with the highest methanol concentrations were the April 25 Haynesville flight, and the April 23 Permian flight, both of which had mixing ratios comparable to those observed during SENEX, a summertime campaign over a biogenically productive region. All three Permian flights had high methanol mixing ratios.

310

To summarize, oil and gas-producing regions, even those in rural areas such as the Permian and Uintah basins, can have VOC mixing ratios comparable to those measured in urban areas. The concentrations and relative distribution of photochemically produced species were relatively similar between basins. There are significant differences between the basins in the mixing ratios and composition of primary compounds such as aromatics and H₂S. Forthcoming work, including measurements from iWAS/GC-MS, will investigate these differences and their origins in greater detail.

315

3.2 Overview of measurements in the Permian Basin

320 The highest overall mixing ratios of VOCs were detected during the April 23 flight over the Permian Basin. In the following sections we provide a detailed interpretation of PTR-ToF-MS mass spectra in oil and gas producing regions, based on observations in the Permian Basin.

325 An averaged mass spectrum from the boundary layer in the Permian Basin flight on April 23 is shown in Figure 5a, where all peaks shown are well above the detection limit. Enhancements are relative to the average mixing ratio in a ten-minute free-troposphere measurement immediately prior to descent into the basin. (The time period and altitude of this measurement are shown in Figure SI 3).

330 Compounds containing one or two oxygen atoms dominate the product ions measured by the H₃O⁺ ToF-CIMS. These oxygenates are comprised mainly of methanol (m/z 33.034 CH₄OH⁺) and small photochemical products. Other important ions include masses typically attributed to aromatics and cyclic alkanes; m/z 34.995 H₂S·H⁺ (hydrogen sulfide); and m/z 45.992 NO₂⁺. Figure 5b compares the average boundary layer concentrations of species detected by H₃O⁺ ToF-CIMS. NO₂⁺ cannot
335 be converted to mixing ratio in a meaningful way and was excluded from the concentration



comparison. For fragmentary hydrocarbons, an average sensitivity was applied (Section 2.2.1). It can be seen that methanol and H₂S, while comprising a relatively small portion of the overall signal, are actually quite important in terms of actual abundance.

340 The relationship between major H₃O⁺ ToF-CIMS product ion signals measured over the Permian and the West Texas crude oil headspace sample is shown in Figure 6. Photochemical products like the small oxygenates, and methanol, are greatly enhanced compared to hydrocarbon masses, which are generally similar to the composition of crude oil. Alkane and alkene masses are somewhat more abundant relative to aromatics in the SONGNEX measurements, compared to the crude oil.

345 This could be due to compositional differences between crude oil and VOC emission sources in the Permian, additional signal on alkane masses from photochemical products (Section 3.2.4), or photochemical removal of aromatics (Section 3.3.2). A few species, particularly H₂S, were enhanced in the Permian but not detected in the crude oil. There are large differences in the H₂S content of various Permian basin reservoirs (Railroad Commission of Texas) and it is possible that

350 our crude oil sample was derived from low-sulfur-content reservoirs, or that sulfur species were removed prior to bottling.

During the April 23 flight, several compositionally distinct air masses were sampled (Fig. 2). One air mass, in the southwestern part of the flight path, was enriched in aromatics, methane, ethane,

355 and other primary compounds (less aged); another, to the east, was relatively more enriched in oxygenates that are typically photochemically produced, such as acetone, acetaldehyde, and PAN species (more aged). HYSPLIT back-trajectory modeling indicates that over the 24 hours prior to sampling, the more aged air mass circulated over the southwestern part of the oil field. The highest concentrations of acetone and acetaldehyde occur over the topographically lower areas of the

360 flight. A reasonable explanation for the more aged air mass is that it consists of emissions and chemical products from the previous day pooled in low-lying areas. Most VOCs are enhanced in the less-aged spatial distribution (e.g. toluene), the more-aged spatial distribution (e.g. acetaldehyde), or a combination of the two (e.g. m/z 83.086 C₆H₁₀H⁺). Figure 7 compares the correlation of the measured VOCs with toluene, a primary emission, and acetaldehyde, a secondary

365 product. Alkanes, aromatics and cycloalkanes have a higher correlation with toluene, while oxygenates have a higher correlation with acetaldehyde. A small number of species, such as H₂S



and m/z 71.049 $C_4H_6OH^+$, have different distributions. The physical separation between less and more aged emissions during this flight was used to help identify VOC ions and to interpret their source.

370

The three transects in the western part of the flight path contain similar VOC composition, but the concentrations of toluene, C8 aromatics, and larger aromatics decrease relative to benzene from south to north. The relative decreases of the more reactive aromatics is likely the result of longer photochemical processing, which is consistent with the southerly wind direction during the flight, generally lower concentrations from south to north, and the time of each transect (northernmost transect latest in the day). We used the ratios of toluene, C8 aromatics, and C9 aromatics to benzene to calculate a relative OH exposure for each transect, in order to explore photochemical aging of other VOCs. This method has been used extensively in atmospheric chemistry (de Gouw et al., 2005;Warneke et al., 2013) and details are shown in the supplementary information (SI Section 3).

375

Not all species had significant enhancement in this area of the flight (e.g. H_2S had a different spatial distribution), so we were not able to use this method to investigate photochemical aging these species.

380

In the following discussion, we examine several groups of compounds measured during the April 23 Permian flight in greater detail: hydrocarbons (aromatic, cyclic alkane, alkene, and alkane masses), secondary compounds, and compounds with heteroatoms. Table 2 lists ion masses discussed in this work and our interpretation of that measurement. Table 2 also highlights (in bold text) measurements that provide new understanding of atmospheric composition and chemistry: VOCs that have been previously unreported or underexplored, and VOC ion masses where our assessment is a new interpretation of a mass previously reported in the literature.

385

390 **3.3 Hydrocarbon masses**

3.3.1 Aromatics

Alkyl-substituted aromatic species are commonly measured by PTR-MS and the interpretation of these ion masses is well established (de Gouw and Warneke, 2007;Blake et al., 2009). Aromatics can be important to the photochemical production of ozone in oil and gas producing regions, and can have high yields of secondary organic aerosol (Henze et al., 2008;Edwards et al., 2014).

395



Quantifying the distribution of these species emitted from oil and gas sources is important to modeling work and to distinguishing between oil and gas and urban sources.

Figure 8 summarizes the distribution of C7 C10 alkyl-substituted aromatics relative to benzene measured during SONGNEX and compares the findings with other sources. This figure compares (1) measurements from all flights, (2) the average aromatics enhancement over the Permian during the April 23 flight, (3) fresh emissions sampled in a plume from a point source (a natural gas processing plant, 32.49N 101.35W) on the April 6 and April 9 Permian flights, (4) aromatics in the headspace of West Texas crude oil, (5) a literature survey of oil and gas sources, and (6) a literature survey of urban sources. The profiles of the average for all SONGNEX flights, and of aromatics measured in the Permian, are similar to the profile typically measured in oil and gas producing regions. The Permian profile is very similar to the composition of West Texas crude oil, and there is significant diversity in aromatics profiles between different oil and gas regions and sources. The average oil and gas profile is clearly differentiated from urban emissions (vehicular sources) by roughly equal enhancements of toluene and benzene, followed by a steady decrease in enhancement with increasing carbon number. The average oil and gas, and urban composition shown in Figure 8d were calculated by averaging all oil and gas sources (panels a and b) and all literature profiles shown in panel 8c for the urban profile.

The H_3O^+ ToF-CIMS also allowed for the investigation of more highly alkyl-substituted aromatics with 4 double bond equivalents; and less saturated aromatics, including PAHs and styrenes. During the April 23 Permian flight, we found that benzene and C7-C9 alkyl substituted aromatics were by far the most abundant aromatic species, accounting for 95% of total aromatic signal. These species were observed in both broad enhancements and in localized plumes from point sources. No PAHs with significant enhancement (above estimated 1s detection limit of 30-40 pptv) were observed. Over all flights, styrene (m/z 105.070 $\text{C}_8\text{H}_8\text{H}^+$) was consistently enhanced by up to 1.1 ppbv in plumes from point sources, and up to 60 pptv during one leg of the April 6 flight, where it was not correlated with other aromatics.



3.3.2 Cycloalkanes

425 Cycloalkanes are an important component of crude oil (Smith, 1968; National Research Council, 1985; Drozd et al., 2015), and have been detected by GC in significant concentrations in the atmosphere over oil and gas producing regions (Simpson et al., 2010; Gilman et al., 2013; Edwards et al., 2014). Although cycloalkanes are measured by PTR-MS with lower sensitivity than
430 not specific for cycloalkanes, but may still be useful for characterizing the VOC composition of air masses, and may be specific for cycloalkanes in relatively unaged air masses.

Several previous laboratory and field experiments have explored the PTR-MS response to cycloalkanes. PTR-MS is somewhat less sensitive to cyclic alkanes than to aromatics and
435 oxygenates (Midey et al., 2003; Gueron et al., 2015). At the E/N conditions in our instrument, cycloalkanes experience significant fragmentation, creating important product ions at m/z 69.070 $C_5H_9^+$, 83.086 $C_6H_{11}^+$, 85.101 $C_6H_{13}^+$, 97.101 $C_7H_{13}^+$, 111.117 $C_8H_{15}^+$, and 125.132 $C_9H_{17}^+$ (Midey et al., 2003; Warneke et al., 2003; Yuan et al., 2014; Gueron et al., 2015). Cyclic alkanes have been measured with PTR-MS in crude oil using a GC interface (Yuan et al., 2014), and in ambient
440 air in the Uintah basin (Warneke et al., 2014); based on these measurements we expect that m/z 69.070 is produced generally by C6-C9 cycloalkanes; m/z 83.086 mainly from methylcyclopentane, cyclohexane, and methylcyclohexane; m/z 85.101 mainly by methylcyclopentane; m/z 97.101 mainly by methylcyclohexane; and m/z 111.117 and 125.132 by C9 and C10 cycloalkanes, respectively. These masses were clearly enhanced in H_3O^+ ToF-CIMS
445 measurements over the Permian and other basins. Cumulatively, they account for 50% of the hydrocarbon concentration (ppbv) measured by H_3O^+ ToF-CIMS.

Chemically specific measurements of methylcyclohexane and cyclohexane were made by iWAS/GC-MS. The comparison between the H_3O^+ ToF-CIMS and iWAS/GC-MS
450 methylcyclohexane measurements shows that the H_3O^+ ToF-CIMS measurement is on average three times higher and the correlation coefficient R^2 is 0.71. The ratio of m/z 97.101 to iWAS methylcyclohexane is higher in the more aged air mass (Fig. 9a). This suggests that m/z 97.101 includes a significant contribution from a secondary VOC that fragments to produce $C_7H_{13}^+$. Some



possibilities include an aldehyde or alcohol with formula m/z 115.112 $C_7H_{14}OH^+$ (e.g. heptanal or
455 methylcyclohexanol) or a larger molecule (e.g. m/z 129.127 $C_8H_{16}OH^+$ losing $-CH_3OH$).

Comparisons of methylcyclohexane between other GC and PTR-MS instruments previously
deployed in the Uintah Basin, Utah, indicate that this behavior is not an H_3O^+ ToF-CIMS specific
issue and is not unique to the Permian Basin. In the winter of 2012, when photochemistry was
460 low, quadrupole PTR-MS m/z 97 measurements agreed with the methylcyclohexane measurement
from a GC-MS instrument within 23% (Warneke et al., 2014). However, in the winter of 2013,
when photochemistry was much higher (Edwards et al., 2014), the relationship between PTR-ToF-
MS m/z 97.101 $C_7H_{13}^+$ and GC-FID methylcyclohexane was dependent on photochemical
exposure (Fig. 9b).

465 The relatively high correlations of other cycloalkane indicator masses (m/z 83.086, m/z 69.070,
 m/z 111.117, etc.) with acetaldehyde suggest that they too experience interference from similar
secondary products (Fig. 7). The secondary compounds could be cycloalkane oxidation products,
or other alcohols or aldehydes, which fragment by losing a water molecule -- a common PTR-MS
470 fragmentation mechanism (Yuan et al., 2016a).

3.3.3 Interpretation of hydrocarbon ion masses in oil and gas regions

The H_3O^+ ToF-CIMS measurements during SONGNEX offer new insights into some VOC ion
masses commonly measured by PTR-MS.

475 It is clear from the comparison to iWAS/GC-MS measurement of isoprene (Fig. 10) that m/z
69.070 $C_5H_9^+$ is not isoprene, which is the dominant contributor in many air masses (de Gouw and
Warneke, 2007; Blake et al., 2009). We interpret it as the sum of a cycloalkane fragment and a
secondary compound from the oxidation of oil and gas precursor emissions; several other studies
have also suggested non-biogenic interpretations of this mass (Yuan et al., 2014; Gueneron et al.,
480 2015). Similarly, m/z 137.132 $C_{10}H_{16}H^+$, usually interpreted as monoterpenes, was enhanced far
above the monoterpene species (α - and β -pinene) detected by iWAS/GC-MS. m/z 137.132
 $C_{10}H_{16}H^+$ was also observed using a PTR-ToF-MS instrument in the Uintah Basin, UT in winter
of 2013 (Fig. SI 6). The Uintah observations indicate that this unknown species is emitted in



several oilfields and is likely not an instrument artifact. Possible alternative sources of this VOC
485 ion mass are (1) another isomer of $C_{10}H_{16}$, but not a monoterpene; (2) a monoterpene species not
detected by iWAS/GC-MS; or (3) another species, such as an alcohol or aldehyde, which
fragments to m/z 137.132. Each possibility is discussed in more detail below.

(1) *Other $C_{10}H_{16}$ isomer.* Other, non-terpene isomers of $C_{10}H_{16}$ are found in fossil fuels. In
490 particular, the presence of “diamondoids”, characterized by cage-like structures, is well known
(Dahl et al., 1999; Araujo et al., 2012). The smallest diamondoid molecule, adamantane
(tricyclo[3,3,1,1(3,7)]decane, $C_{10}H_{16}$), has a tetrahedral structure formed by three cyclohexane
rings (Fig. 10). Adamantane can have concentrations from <1% to 100% of typical benzene
495 atmospheric measurement, m/z 137.132 $C_{10}H_{16}H^+$ was up to 20% the concentration of benzene
(assuming similar sensitivity as monoterpenes). Alkyl-substituted adamantanes are relatively
abundant (Stout and Douglas, 2004; Wang et al., 2006). We detected enhancement of m/z 151.148
 $C_{11}H_{18}H^+$, but larger molecules were below the detection limit of the H_3O^+ ToF-CIMS during this
flight. The measured headspace of West Texas Crude showed relative enhancements of $n(CH_2)$ -
500 substituted $C_{10}H_{16}$ that are more consistent with the adamantane series than with terpenes, where
we would expect to see enhancements at every $n(C_5H_8)$, but the evidence is not conclusive (Fig.
SI 7). Adamantane has a longer retention time than the GC elution time used for the iWAS samples,
so iWAS/GC-MS was not able to confirm or exclude the presence of this molecule.

505 (2) *Other monoterpene.* Terpene-derived (“isoprenoid”) saturated compounds are known to be in
crude oil, and some larger molecules such as phytane and pristane are used as geochemical
biomarkers (Tissot and Welte, 1984). There is less information available about the presence of
unmodified monoterpenes, but the presence of an unusual monoterpene (such as limonene) derived
from crude oil or industry solvents is certainly possible. Data from the iWAS/GC-MS was again
510 inconclusive, but suggested that it was not limonene.

(3) *Fragment of another species.* M/z 137.132 $C_{10}H_{16}H^+$ is not likely to be a fragment of another
species. We did not detect any oxygenated species that could easily fragment to produce m/z 137.
A larger hydrocarbon could fragment to this mass. However, in crude fuels, hydrocarbon



515 concentrations in a particular homologous series generally decrease with carbon number. An
anomalously intense larger mass would require its own explanation.

We do not currently have enough information to state conclusively if the observed m/z 137.132
 $C_{10}H_{16}H^+$ ion is from adamantane, a monoterpene, or another isomer. Collection of more evidence
520 is outside the scope of this manuscript and is currently under separate investigation. If the observed
 $C_{10}H_{16}$ molecule is indeed adamantane, it represents a previously undetected class of atmospheric
VOCs. Regardless of identity, this observation highlights that there may be significant emissions
of higher-mass hydrocarbons from fossil fuels that are rarely measured. These compounds could
be used to identify emission sources, and could potentially be SOA precursors (de Gouw et al.,
525 2011).

3.3.4 Other hydrocarbon masses

The H_3O^+ ToF-CIMS also measured other hydrocarbon signals that could not be tied to a single
VOC, and these hydrocarbon masses comprise a significant fraction of the total hydrocarbon signal
(Fig. 5). These other important hydrocarbon ion masses include m/z 41.039 $C_3H_4H^+$, m/z 43.054
530 $C_3H_6H^+$, m/z 57.070 $C_4H_6H^+$, and m/z 71.086 $C_5H_{10}H^+$. Unit-mass-resolution PTR-MS has not
been able to investigate these ions because of strong interference from isobaric masses, such as
 $C_2H_2OH^+$ at m/z 43.018. Several studies have reported that these masses are non-specific and are
produced by alkenes, and fragmentation of alkanes, alkenes, aldehydes, and alcohols (Buhr et al.,
2002; Jobson et al., 2005; Gueneron et al., 2015). Laboratory tests with the H_3O^+ ToF-CIMS
535 confirmed that multiple alkane and alkene VOCs present during the April 23 flight (as measured
by iWAS/GC-MS) fragment to these masses. These ion masses had significant intensity in both
the southwestern (less aged) and eastern (photochemically enhanced) areas of the flight, suggesting
contributions from fragments of both primary and secondary species.

3.4 Hydrocarbon oxidation chemistry

540 3.4.1 Major secondary compounds

Our interpretation of the most abundant photochemically produced ion masses is consistent with
interpretations often presented in the literature: m/z 45.034 $C_2H_4OH^+$ is specific for acetaldehyde,



m/z 59.049 $C_3H_6OH^+$ for acetone, m/z 73.065 for MEK, m/z 61.028 $C_2H_4O_2H^+$ for acetic acid, and m/z 75.044 $C_3H_6O_2H^+$ for propionic acid.

545 The carbonyl species with more than three carbon atoms have both ketone and aldehyde isomers, and both isomers could be products of alkane oxidation. For example, oxidation of propane in the presence of NO_x is expected to yield approximately 26% propanal and 74% acetone (Calvert et al., 2008). The ratio of propanal to acetone has been measured to be 0.17 (Uintah Basin, 2012, Edwards et al. (2013)), 0.19 (Uintah Basin, 2014), and 0.22 (Denver-Julesburg basin, Gilman et al. (2013))

550 in oil and gas producing regions, compared to 0.06 in an urban area (Los Angeles, Borbon et al. (2013)). However, almost all of the signal at H_3O^+ ToF-CIMS m/z 59.049 $C_3H_6OH^+$ can be attributed to acetone. The PTR-MS reaction of H_3O^+ with propanal is dissociative, and the sensitivity of the H_3O^+ ToF-CIMS to propanal at m/z 59.049 is only 3% that of acetone. Acetone therefore dominates the signal at this mass, and we assume similar behavior for higher-mass

555 carbonyls.

Abundances of larger masses in the 1-double bond equivalent, 1-oxygen homologous series decrease rapidly with carbon number; these are most likely also carbonyls derived from alkanes. m/z 43.018 $C_2H_2OH^+$ was identified as a fragment of acetic acid from laboratory tests.

Glycolaldehyde (m/z 61.028 $C_2H_4O_2H^+$) and hydroxyacetone (m/z 75.044 $C_3H_6O_2H^+$) could be

560 interferences to acetic acid and propionic acid, respectively. However, these compounds have been mainly reported in environments affected by biogenic emissions and biomass burning, and acetic acid has been shown to be the dominant contributor to m/z 61.028 $C_2H_4O_2H^+$ in several environments (Karl et al., 2007;Fu et al., 2008;Haase et al., 2012).

3.4.2 Aromatic oxidation products

565 Aromatic oxidation products are of particular interest, as they have been shown to be an important source of radicals that drive ozone formation in oil and gas producing regions (Edwards et al., 2014). From laboratory and chamber studies, expected aromatic oxidation products include various diketones, phenols and nitrophenols, benzaldehyde-type compounds (from toluene and larger aromatics), and furanones (Wagner et al., 2003; Bloss et al., 2005; Wyche et al., 2009; Yuan et al.,

570 2016b). PTR-MS can detect phenols, benzaldehydes, and furanones (Müller et al., 2012); detection of some dicarbonyls, such as glyoxal and methylglyoxal, may be difficult due to fragmentation,



strong humidity dependence, and interference from other species (Pang et al., 2014; Stöner et al., 2017).

By far the most abundant aromatic oxidation product detected was m/z 107.049 $C_7H_6OH^+$,
575 benzaldehyde (max. 360 pptv). Phenol (C_6H_6O), cresol (C_7H_8O), and methylfuranone ($C_5H_6O_2$)
were detected at only an estimated 30-40 pptv maximum enhancement. Cresols, diketones, and
furanones are expected to have a much higher yield at the observed NO_x concentrations (average
1.1 ppb) (Smith et al., 1998; Müller et al., 2012), but are also much more reactive (Bierbach et al.,
1994; Atkinson and Arey, 2003).

580 Müller et al. (2012) reported several unidentified masses resulting from chamber oxidation of
trimethylbenzenes: m/z 87.044 $C_4H_6O_2H^+$ and m/z 101.060 $C_5H_8O_2H^+$. These masses were also
detected during the April 23 SONGNEX flight, and they were quite significant relative to other
oxygenates- m/z 87.044 at approximately 10% of the signal intensity of acetic acid. We classify
these as aromatic oxidation products but do not have enough information to suggest a structure.

585 3.4.3 Cycloalkane oxidation products

The H_3O^+ ToF-CIMS detected several potential cycloalkane oxidation products as shown in Figure
11. The 2-double bond equivalent, 1-oxygen homologous series ($C_nH_{2n-2}OH^+$) does not decrease
monotonically with carbon number: the species with four carbons (m/z 71.049 $C_4H_6OH^+$) and 6
carbons (m/z 99.080 $C_6H_{10}OH^+$) are the most abundant (Fig. 11a). In addition, m/z 71.049
590 $C_4H_6OH^+$ has a somewhat different spatial distribution than other oxygenates, including much
higher intensity in the central part and sections of the northern part of the field, and may have a
primary source (Section 3.4.4). The enhancement of the C6 oxygenate points to cycloalkane
precursors. Similar ion masses were measured by the PTR-ToF-MS instrument deployed in the
Uintah Basin, UT, in 2013 (Fig. 11b).

595

The smallest cycloalkane that exists in significant amounts in fossil fuels is cyclopentane (C5), but
we might expect the C6 products to be more abundant because C6 precursors can be more abundant
than C5 in oilfields (Simpson et al., 2010; Gilman et al., 2013). Additionally, OH reacts faster with
substituted cycloalkanes (starting at C-6, methylcyclopentane) than unsubstituted cycloalkanes,
600 although this probably has a smaller effect on ambient composition than the emissions composition
(Calvert et al., 2008). Figure 11 also shows the cycloalkane precursors in the Uintah Basin,



measured by GC techniques (Edwards et al., 2013; Edwards et al., 2014). The suggested oxygenated products have a similar distribution. Data are not available from iWAS/GC-MS to show a similar precursor distribution for the Permian flight.

605

The C6 cycloalkane oxidation product ($C_6H_{10}OH^+$) could be a ketone (methylcyclopentanone or cyclohexanone), or cyclopentyl aldehyde. Aldehyde ions often fragment in PTR-MS by loss of H_2O (Buhr et al., 2002). The fragmentary product of $C_6H_{10}OH^+$ dehydration, m/z 81.070 $C_6H_8H^+$, was significantly enhanced and correlated with photochemical species. Similar ions (m/z 95.086
610 $C_7H_{10}H^+$, m/z 109.101 $C_8H_{12}H^+$, etc.) also correlated with photochemical species and may also be fragments of cycloalkyl aldehydes. These ion masses were also seen by the PTR-ToF-MS in Utah in 2013, and showed behavior consistent with photochemical species (Fig. SI 6).

3.4.4 PAN and reactive nitrogen tracers

It has been previously reported that PTR-MS detects peroxy acetyl nitrate (PAN) at the protonated
615 parent mass m/z 122.008 $C_2H_3NO_5H^+$, at m/z 77.023 $C_2H_4O_3H^+$, and at m/z 45.992 NO_2^+ . Similarly, peroxy propionyl nitrate (PPN) is detected at m/z 91.039 $C_3H_6O_3H^+$ (Hansel and Wisthaler, 2000; de Gouw and Warneke, 2007; Kaser et al., 2013). However, m/z 77.023 may include contributions from another species such as acetone water cluster or peroxyacetic acid, and m/z 45.992 is not expected to be universally specific to PAN (de Gouw and Warneke, 2007; Kaser
620 et al., 2013).

During the April 23 SONGNEX flight over the Permian, there was no peak detected at m/z 122.008 $C_2H_3NO_5H^+$. m/z 77.023 $C_2H_4O_3H^+$ and 91.039 $C_3H_6O_3H^+$ were detected at moderate and low intensities (4 ncps and 2 ncps average enhancement, respectively) and m/z 45.992 NO_2^+ was one
625 of the most abundant ions detected (28 ncps average enhancement, similar to the average enhancement of benzene). M/z 77.023, m/z 91.039, and m/z 45.992 are compared to several independent measurements of reactive nitrogen: PAN, PPN, NO_y , NO_y-NO_x , and ethyl and propyl alkyl nitrates in Figure SI 8. When looking at all the SONGNEX flights, the slope of m/z 45.992 NO_2^+ versus NO_y-NO_x (which represents all NO_x oxidation products, including PAN), and the
630 slope of m/z 77.023 versus PAN, vary significantly between different environments. Figure 12 shows the relationship between m/z 77.023 and PAN. Comparison of m/z 45.992 NO_2^+ with PAN



is included in the supplemental information (Fig. SI 9). The slope of m/z 45.992 vs NO_z depends on the composition of NO_z (Fig. SI 10a). The range of slopes probably depends on the atmospheric variability of NO_z , and not instrument conditions: the slope of m/z 77.023 vs PAN is not dependent
635 on instrument conditions such as drift tube humidity (Fig. SI 10b), which is consistent with behavior reported by Hansel and Wisthaler (2000).

These measurements suggest that these product ions cannot be attributed to PAN-type compounds only. Product ions at m/z 45.992 NO_2^+ almost certainly derive from a number of NO_z species, with
640 a range of response factors; and there are probably at least two species that contribute to m/z 77.023 $\text{C}_2\text{H}_4\text{O}_3\text{H}^+$.

3.5 Other compounds with heteroatoms

VOC emissions from oil and gas operations are distinctly different from other commonly studied sources (urban areas, forests) with respect to the presence of non-photochemical species containing
645 nitrogen, sulfur, and oxygen heteroatoms. In this section we discuss cyclic nitrogen-containing species, H_2S , and two oxygenated species: methanol and m/z 71.049 $\text{C}_4\text{H}_6\text{OH}^+$.

3.5.1 Cyclic organic nitrogen species

An especially interesting observation is the broad enhancement of several nitrogen-containing organic species during the Permian flights. The most clearly enhanced ion, m/z 70.065 $\text{C}_4\text{H}_7\text{NH}^+$,
650 has signal intensity approximately 15% that of m/z 93.070 $\text{C}_7\text{H}_8\text{H}^+$, toluene (Fig. 13).

$\text{C}_4\text{H}_7\text{N}$ has several possible isomers, each with two degrees of unsaturation: a cyclic alkene structure (pyrroline) and several non-cyclic structures (C_4 nitriles). Using mass spectral context, we suggest a cyclic structure for this compound. Time series of the homologous series ($\text{C}_n\text{H}_{2n-1}\text{NH}^+$) including $\text{C}_4\text{H}_7\text{NH}^+$ are shown in Figure 13. The species having one, two, and three carbon
655 atoms are not enhanced above background, whereas species containing four or more carbon atoms are enhanced and correlated with aromatic compounds (Fig. 13, Fig. SI 11). A cyclic structure is the most likely explanation for this pattern. A similar argument can be made for the homologous series ($\text{C}_n\text{H}_{2n-3}\text{NH}^+$) having three degrees of unsaturation (pyrroles). The series with two
660 ($\text{C}_n\text{H}_{2n+1}\text{NH}^+$, pyrrolidines) and four ($\text{C}_n\text{H}_{2n-5}\text{NH}^+$, pyridines) degrees of unsaturation were not



significantly enhanced. Notably, m/z 84.081 $C_5H_9NH^+$ had a different distribution than other cyclic nitrogen compounds during the April 23 flight, with high mixing ratios in the eastern and western parts of the field, and overall correlated better with the photochemically produced species. We speculate that this mass may include a fragment of a photochemical product, analogous to the
665 interference with cycloalkanes discussed in section 3.2.2. Pyrroline (m/z 70.065) was enhanced by up to 200 pptv, and its concentration was typically comparable to C8 and C9 aromatics. Pyrrole (m/z 68.050) had a maximum enhancement of 40 pptv.

Organic nitrogen species detected over the Permian may have originated from the crude oil.
670 American shale oils can contain upwards of 2% nitrogen by weight, and a number of aromatic organic nitrogen species have been quantified in crude oil (Morandi and Jensen, 1966; Holmes and Thompson, 1983; Mushrush et al., 1999; Williams and Chishti, 2001). Porphyrins, nitrogen-containing geochemical fossils derived from chlorophyll, are known to exist in crude oil and contain both pyrrole and pyrroline as subunits (Fig. 14); aromatic nitrogen species may also form
675 through chemical processes in the oil reservoir. There are large differences in nitrogen content and speciation between reservoirs, which could explain why these species were detected in some regions and not in others (Baxby et al., 1994; Li et al., 1995; Oldenburg et al., 2007).

The reactivity of pyrroline is not known. We analyzed the relative decrease in C_4H_7N in the
680 western half of the field as a function of OH exposure (see section 3.1), and estimated the rate constant with OH to be in the range of $(1.3\text{--}2.0)\times 10^{-11}$ cm^3 molecule s^{-1} with a best estimate of 1.7×10^{-11} cm^3 molecule s^{-1} (details in SI). Saturated nitriles have much slower reactivity with OH – reported values are around 2×10^{-14} cm^3 molecule s^{-1} (Harris et al., 1981; Atkinson et al., 2006) – so this high OH reactivity is further evidence that C_4H_7N is not a nitrile. A similar analysis of
685 C_4H_5N gave a rate constant of 2.5×10^{-11} cm^3 molecule s^{-1} , which is much slower than the reported value for pyrrole (1×10^{-10} (Wallington, 1986)) but faster than butenenitrile (1.4×10^{-11} (Grosjean and Williams, 1992)). The pyrrole analysis is much less certain than that of pyrroline, due to the overall lower signal and higher noise. The rate constants for pyrroline and especially pyrrole could be significantly underestimated. In previous work, it has been shown that for fast-reacting species,
690 the derived rate constant was approximately constant and similar to the rate of the fastest-reacting aromatic used to calculate OH exposure (de Gouw et al., 2005). In our analysis, the fastest-reacting



aromatic considered was C10 aromatics, with a rate constant of $2.4 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This may explain why the derived rate constant for pyrrole was lower than expected.

695 Heterocycles are highly reactive with nitrate radicals, especially pyrrole ($k=4.6 \times 10^{-11} \text{ cm}^3$
molecule⁻¹ s⁻¹) (Atkinson et al., 1985; Cabañas et al., 2004). If pyrroline is similarly reactive, then
nitrogen heterocycles could potentially dominate VOC nitrate reactivity in oil and gas fields,
because most other VOCs measured during SONGNEX (aromatics and aliphatics) have low
reaction rates with nitrate radicals ($k \sim 10^{-18}$ - $10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$). A comparison of nitrate
700 reactivity for species measured during SONGNEX, and a comparison to nitrate loss rates reported
in the literature, are given in the supplementary information (Fig. SI 12, Section SI 4). We looked
for species with two nitrogen atoms, but at our instrument resolution, they are extremely difficult
to separate from isobaric hydrocarbon species unless they have very high signal intensity.

3.5.2 Hydrogen sulfide

705 Measurement of H₂S with PTR-MS has been described by Li et al. (2014). The H₃O⁺ ToF-CIMS
improves on the instrument described by Li et al. (2014) as the high mass resolution avoids the
isobaric background interference from isotopes of methanol and HO₂⁺. H₂S is detected at m/z
34.995 H₂SH⁺ and was calibrated directly using a standard cylinder.

710 A comparison with the Picarro H₂S instrument is shown in Figure 15. There is good quantitative
agreement between the two measurements. Compared to the Picarro instrument, the H₃O⁺ ToF-
CIMS H₂S measurement is more precise, and the 1s data are simultaneous with other H₃O⁺ ToF-
CIMS measurements, allowing easy comparison. The H₃O⁺ ToF-CIMS H₂S measurement had a
3σ detection limit of 2.3 ppbv for a 1-s measurement (or 0.8 ppbv over the 6s measurement period
715 of the Picarro instrument).

H₂S had a maximum concentration of 12.6 ppb during the April 23 Permian flight, comparable to
butanes. No other sulfur-containing species measurable with the H₃O⁺ ToF-CIMS were enhanced
above an estimated 1s 3σ detection limit of 30 pptv. (A few sulfur-containing species, such as m/z
720 121.032 C₄H₈SO₂H⁺, have non-zero intensity but are instrument contaminants).



3.5.3 Methanol

Methanol was the most abundant VOC detected by H_3O^+ ToF-CIMS during the April 23 Permian flight. In this section we discuss some possible sources.

725 Methanol is used by the oil and gas industry. Significant primary emissions, especially from
produced water storage infrastructure and storage containers on wellpads, have been measured in
the Uintah Basin (Warneke et al., 2014; Mansfield et al., 2016). Industry uses of methanol include
addition at well heads or further downstream in pipelines to prevent methane hydrate formation
(Anderson and Prausnitz, 1986), to inhibit corrosion and scaling, as a lubricant, and as a solvent
730 in other applications (Mansfield et al., 2016). A more detailed investigation of methanol sources
is outside the scope of this paper.

Globally, the dominant net source of methanol is direct biogenic emission, although there is a
substantial and poorly constrained source from secondary production and oceans (Jacob et al.,
735 2005; Millet et al., 2008). Primary biogenic emissions can explain the high mixing ratios over the
Haynesville region, but not over the Permian, given the absence of other biogenic VOCs.

Lewis et al. (2005) calculated rates of photochemical production of methanol from a set of VOC
precursors, the most important of which were methane, iso-butane, iso-pentane, and acetaldehyde.
740 The magnitude of photochemical methanol production in the Permian was estimated by scaling
the Lewis *et al.* precursor concentrations to the highest observed Permian VOC concentrations
during the April 23 flight. Only 1-2 ppb of methanol would have formed after 2-3 days of aging –
a small amount compared to the measured average 6 ppb and maximum 19 ppb. It should be noted
that the Lewis et al. (2005) calculations were in the remote marine boundary layer and the
745 photochemical production rate in the Permian could be different. Methanol does have a much
stronger correlation with photochemical species than with primary aromatic species (Fig. 7), but
it also has a relatively long atmospheric lifetime.



3.5.4 M/z 71.049 C₄H₆OH⁺

M/z 71.049 C₄H₆OH⁺ is typically interpreted as the sum of methyl vinyl ketone (MVK) and methacrolein, two carbonyl products of isoprene oxidation (de Gouw and Warneke, 2007), but isoprene was too low to produce the measured amount of C₄H₆OH⁺ in the Permian Basin. C₄H₆OH⁺ had a spatial distribution that differed from photochemically produced species, including a much larger enhancement in the central part of the flight and several north-south oriented plumes running the length of the surveyed area (Fig. 2). Additionally, C₄H₆OH⁺ was enhanced above what might be expected from the distribution of precursor species and the enhancements of similar oxygenates (Fig. 11). The enhancement in the more aged area of the field suggests that a large part of this ion signal is generated by a photochemical species, but there may be other contributions. Using the H₃O⁺ ToF-CIMS average sensitivity factor for methyl vinyl ketone (MVK) and methacrolein, the maximum boundary-layer enhancement of this species was 540 pptv.

760

The other carbonyl isomer, crotonaldehyde, has been reported in biomass burning emissions (Karl et al., 2007), which were not evident during this flight. It is possible that MVK, methacrolein, or crotonaldehyde could be directly emitted by anthropogenic sources, or photochemically derived from a non-biogenic species. The cyclic isomer, dihydrofuran, is the oxygenated analogue of the nitrogen heterocycle (pyrroline) discussed in section 3.4.1. Dihydrofurans are known to be products of OH oxidation of alkanes (Lim and Ziemann, 2005).

765

Preliminary iWAS/GC-MS measurements show strong correlation between MVK, methacrolein, and m/z 71.049 C₄H₆OH⁺. The appropriate mass for dihydrofuran was not included in the selected-ion-scan window of the iWAS GC quadrupole MS. Other evidence is needed to identify the predominant isomer(s), and determine if there is a significant interference to PTR-MS measurements of biogenic MVK and methacrolein in oil and gas producing regions.

770

4. Conclusions

We have analyzed PTR-ToF-MS mass spectra from aircraft measurements over several US oil and gas producing regions. Our analysis is supported by comparison to independent co-deployed instrumentation. We present a comparison between nine oil and gas basins of mixing ratios of

775



aromatics, major secondary species, methanol, and hydrogen sulfide. In every basin, measurements from H_3O^+ ToF-CIMS were dominated by small oxygenated compounds, especially C2-C4 photochemical products and methanol. Significant classes of hydrocarbon compounds detected included aromatics, cycloalkanes, and alkanes. The H_3O^+ ToF-CIMS measurements of aromatics, methanol, and H_2S agreed with independent measurements while methylcyclohexane and reactive nitrogen differed from independent measurement.

Between basins, there was large variation in the observed mixing ratios of aromatics, H_2S , and methanol. In every basin, narrow, highly concentrated plumes with high mixing ratios of aromatics were measured, and average mixing ratios in many basins were comparable to concentrations observed downwind of large metropolitan areas. However, the profile of aromatics is different from that in urban air. We demonstrated the ability of H_3O^+ ToF-CIMS to detect hydrogen sulfide, and measured significantly enhanced H_2S in the Permian and Haynesville regions. Methanol was the single most abundant VOC observed by H_3O^+ ToF-CIMS and may have industrial sources.

Compared to the variability in aromatics, methanol, and H_2S , photochemical compounds had similar mixing ratios in each basin. Additionally, the abundances of most oxygenates relative to acetone were similar between basins. This profile was also quite similar to that measured during a flight over the Los Angeles urban area during CALNEX 2010. The most highly variable compound was acetic acid, which can include primary emission from agriculture, especially in the Denver-Julesburg Basin.

The Permian Basin had the highest overall mixing ratios of all species reported here. This region is the largest oil field in the United States but has not been studied extensively from an air quality perspective. We conducted a detailed investigation of mass spectra recorded over Permian Basin. There are likely many chemically significant species, measureable by PTR-MS, in the atmosphere that are not currently routinely reported. This includes both primary species such as pyrroline and early-generation secondary species, such as the oxidation products of cycloalkanes. Pyrroline (m/z 70.065, $\text{C}_4\text{H}_7\text{NH}^+$) is especially interesting because it has not been previously reported as a VOC associated with oil and gas emissions, and may account for a substantial fraction of nitrate reactivity. The $\text{C}_{10}\text{H}_{16}$ measurement, which is most likely adamantane or an unusual monoterpene, indicates the presence of larger ($\text{C}_{10}+$) hydrocarbons emitted from oil and gas operations, which



are currently underexplored in the literature. The most important aromatic oxidation product detected was benzaldehyde; other products, including phenol and two unidentified oxygenates, were present at much smaller concentrations. Several ion masses that could be cycloalkane oxidation products were detected. Finally, we report several new interpretations of PTR-MS ion masses previously described in the literature.

Acknowledgments

A. Koss acknowledges funding from the NSF Graduate Fellowship Program. We thank the NOAA Aircraft Operations Center for their support with instrument installation on the NOAA WP-3D, research flights, and meteorological and aircraft data. We thank Ralf Staebler (Environment and Climate Change Canada) for the use of the Picarro H₂S instrument. We thank Andy Neuman for his scientific advice and thoughtful comments on the manuscript. T. Hanisco, G. Wolfe, J. St. Clair, M. Thayer, and F. Keutsch acknowledge NASA-GeoCAPE award number NNX15AH83G for funding.



References

- Adgate, J. L., Goldstein, B. D., and McKenzie, L. M.: Potential Public Health Hazards, Exposures and Health Effects from Unconventional Natural Gas Development, *Environ. Sci. Technol.*, 48, 8307-8320, 10.1021/es404621d, 2014.
- Anderson, F. E., and Prausnitz, J. M.: Inhibition of gas hydrates by methanol, *AIChE Journal*, 32, 1321-1333, 10.1002/aic.690320810, 1986.
- Araujo, P. L. B. d., Mansoori, G. A., and Araujo, E. S. d.: Diamondoids: occurrence in fossil fuels, applications in petroleum exploration and fouling in petroleum production. A review paper, *Int. J. Oil, Gas, and Coal Tech.*, 5, 316-367, 10.1504/ijogct.2012.048981, 2012.
- Arnold, S. T., Viggiano, A. A., and Morris, R. A.: Rate Constants and Product Branching Fractions for the Reactions of H₃O⁺ and NO⁺ with C₂-C₁₂ Alkanes, *J. Phys. Chem. A*, 102, 8881-8887, 10.1021/jp9815457, 1998.
- Atkinson, R., Aschmann, S. M., Winer, A. M., and Carter, W. P. L.: Rate constants for the gas-phase reactions of nitrate radicals with furan, thiophene, and pyrrole at 295 ± 1 K and atmospheric pressure, *Environ. Sci. Technol.*, 19, 87-90, 10.1021/es00131a010, 1985.
- Atkinson, R., and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605-4638, 2003.
- Atkinson, R., Baulch, D., Cox, R., Crowley, J., and Hampson Jr, R.: Summary of evaluated kinetic and photochemical data for atmospheric chemistry, IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry, 2006.
- Baxby, M., Patience, R. L., and Bartle, K. D.: The Origin and Diagenesis of Sedimentary Organic Nitrogen, *J. Petrol. Geol.*, 17, 211-230, 1994.
- Bierbach, A., Barnes, I., Becker, K. H., and Wiesen, E.: Atmospheric chemistry of unsaturated carbonyls: butenadiol, 4-oxo-2-pentenal, 3-hexene-2, 5-dione, maleic anhydride, 3H-furan-2-one, and 5-methyl-3H-furan-2-one, *Environ. Sci. Technol.*, 28, 715-729, 1994.
- Blake, R. S., Monks, P. S., and Ellis, A. M.: Proton-Transfer Reaction Mass Spectrometry, *Chem. Rev.*, 109, 861-896, 10.1021/cr800364q, 2009.
- Bloss, C., Wagner, V., Bonzanini, A., Jenkin, M. E., Wirtz, K., Martin-Reviejo, M., and Pilling, M. J.: Evaluation of detailed aromatic mechanisms (MCMv3 and MCMv3.1) against environmental chamber data, *Atmos. Chem. Phys.*, 5, 623-639, 10.5194/acp-5-623-2005, 2005.
- Borbon, A., Gilman, J. B., Kuster, W. C., Grand, N., Chevallier, S., Colomb, A., Dolgorouky, C., Gros, V., Lopez, M., Sarda-Esteve, R., Holloway, J., Stutz, J., Petetin, H., McKeen, S., Beekmann, M., Warneke, C., Parrish, D. D., and de Gouw, J. A.: Emission ratios of anthropogenic volatile organic compounds in northern mid-latitude megacities: Observations versus emission inventories in Los Angeles and Paris, *J. Geophys. Res.*, 118, 2041-2057, 10.1002/jgrd.50059, 2013.



Brandt, A. R., Heath, G. A., Kort, E. A., O'Sullivan, F., Pétron, G., Jordaan, S. M., Tans, P., Wilcox, J., Gopstein, A. M., Arent, D., Wofsy, S., Brown, N. J., Bradley, R., Stucky, G. D., Eardley, D., and Harriss, R.: Methane Leaks from North American Natural Gas Systems, *Science*, 343, 733-735, 2014.

860 Brillì, F., Gioli, B., Ciccioli, P., Zona, D., Loreto, F., Janssens, I. A., and Ceulemans, R.: Proton Transfer Reaction Time-of-Flight Mass Spectrometric (PTR-TOF-MS) determination of volatile organic compounds (VOCs) emitted from a biomass fire developed under stable nocturnal conditions, *Atmos. Environ.*, 97, 54-67, <http://dx.doi.org/10.1016/j.atmosenv.2014.08.007>, 2014.

865 Budzik, P., and Perrin, J.: Six formations are responsible for surge in Permian Basin crude oil production, in: *Today in Energy*, U.S. Energy Information Administration Independent Statistics and Analysis, U.S. Energy Information Administration, 2014.

Buhr, K., van Ruth, S., and Delahunty, C.: Analysis of volatile flavour compounds by Proton Transfer Reaction-Mass Spectrometry: fragmentation patterns and discrimination between isobaric and isomeric compounds, *Int. J. Mass Spectrom.*, 221, 1-7, [http://dx.doi.org/10.1016/S1387-3806\(02\)00896-5](http://dx.doi.org/10.1016/S1387-3806(02)00896-5), 2002.

870 Cabañas, B., Baeza, M. T., Salgado, S., Martín, P., Taccone, R., and Martínez, E.: Oxidation of Heterocycles in the Atmosphere: Kinetic Study of Their Reactions with NO₃ Radical, *J. Phys. Chem. A*, 108, 10818-10823, 10.1021/jp046524t, 2004.

Calvert, J. G., Derwent, R. G., Orlando, J. J., Tyndall, G. S., and Wallington, T. J.: *Mechanisms of Atmospheric Oxidation of the Alkanes*, Oxford University Press, 2008.

875 Cappellin, L., Biasioli, F., Granitto, P. M., Schuhfried, E., Soukoulis, C., Costa, F., Märk, T. D., and Gasperi, F.: On data analysis in PTR-TOF-MS: From raw spectra to data mining, *Sensor. Actuat. B-Chem.*, 155, 183-190, <http://dx.doi.org/10.1016/j.snb.2010.11.044>, 2011.

880 Cazorla, M., Wolfe, G. M., Bailey, S. A., Swanson, A. K., Arkinson, H. L., and Hanisco, T. F.: A new airborne laser-induced fluorescence instrument for in situ detection of formaldehyde throughout the troposphere and lower stratosphere, *Atmos. Meas. Tech.*, 8, 541-552, 10.5194/amt-8-541-2015, 2015.

885 Coggon, M. M., Veres, P. R., Yuan, B., Koss, A., Warneke, C., Gilman, J. B., Lerner, B. M., Peischl, J., Aikin, K. C., Stockwell, C. E., Hatch, L. E., Ryerson, T. B., Roberts, J. M., Yokelson, R. J., and de Gouw, J. A.: Emissions of nitrogen-containing organic compounds from the burning of herbaceous and arboraceous biomass: Fuel composition dependence and the variability of commonly used nitrile tracers, *Geophys. Res. Lett.*, 10.1002/2016GL070562, 2016.

Colorado Oil and Gas Conservation Commission: Colorado Oil and Gas Information System: <http://cogcc.state.co.us/data.html>, access: 1/31/2017, 2017.

890 Dahl, J. E., Moldowan, J. M., Peters, K. E., Claypool, G. E., Rooney, M. A., Michael, G. E., Mello, M. R., and Kohnen, M. L.: Diamondoid hydrocarbons as indicators of natural oil cracking, *Nature Geoscience*, 399, 54-57, 1999.

de Gouw, J., and Warneke, C.: Measurements of volatile organic compounds in the earth's atmosphere using proton-transfer-reaction mass spectrometry, *Mass. Spectrom. Rev.*, 26, 223-257, 10.1002/mas.20119, 2007.



895 de Gouw, J. A., Goldan, P. D., Warneke, C., Kuster, W. C., Roberts, J. M., Marchewka, M., Bertman, S. B.,
Pszenny, A. A. P., and Keene, W. C.: Validation of proton transfer reaction-mass spectrometry (PTR-MS)
measurements of gas-phase organic compounds in the atmosphere during the New England Air Quality
Study (NEAQS) in 2002, *J. Geophys. Res. Atmos.*, 108, n/a-n/a, 10.1029/2003JD003863, 2003.

900 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M., Fehsenfeld,
F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C., Marchewka, M., Bertman, S. B.,
and Bates, T. S.: Budget of organic carbon in a polluted atmosphere: Results from the New England Air
Quality Study in 2002, *J. Geophys. Res.: Atmos.*, 110, D16305, 10.1029/2004JD005623, 2005.

905 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Ahmadov, R., Atlas, E. L., Bahreini, R., Blake, D. R., Brock,
C. A., Brioude, J., Fahey, D. W., Fehsenfeld, F. C., Holloway, J. S., Le Henaff, M., Lueb, R. A., McKeen, S. A.,
Meagher, J. F., Murphy, D. M., Paris, C., Parrish, D. D., Perring, A. E., Pollack, I. B., Ravishankara, A. R.,
Robinson, A. L., Ryerson, T. B., Schwarz, J. P., Spackman, J. R., Srinivasan, A., and Watts, L. A.: Organic
Aerosol Formation Downwind from the Deepwater Horizon Oil Spill, *Science*, 331, 1295-1299,
10.1126/science.1200320, 2011.

910 Drozd, G. T., Worton, D. R., Aeppli, C., Reddy, C. M., Zhang, H., Variano, E., and Goldstein, A. H.: Modeling
comprehensive chemical composition of weathered oil following a marine spill to predict ozone and
potential secondary aerosol formation and constrain transport pathways, *J. Geophys. Res.*, 120, 7300-
7315, 10.1002/2015JC011093, 2015.

915 Edwards, P. M., Young, C. J., Aikin, K., deGouw, J., Dubé, W. P., Geiger, F., Gilman, J., Helmig, D., Holloway,
J. S., Kercher, J., Lerner, B., Martin, R., McLaren, R., Parrish, D. D., Peischl, J., Roberts, J. M., Ryerson, T. B.,
Thornton, J., Warneke, C., Williams, E. J., and Brown, S. S.: Ozone photochemistry in an oil and natural gas
extraction region during winter: simulations of a snow-free season in the Uintah Basin, Utah, *Atmos.*
Chem. Phys., 13, 8955-8971, 10.5194/acp-13-8955-2013, 2013.

920 Edwards, P. M., Brown, S. S., Roberts, J. M., Ahmadov, R., Banta, R. M., deGouw, J. A., Dube, W. P., Field,
R. A., Flynn, J. H., Gilman, J. B., Graus, M., Helmig, D., Koss, A., Langford, A. O., Lefer, B. L., Lerner, B. M.,
Li, R., Li, S.-M., McKeen, S. A., Murphy, S. M., Parrish, D. D., Senff, C. J., Soltis, J., Stutz, J., Sweeney, C.,
Thompson, C. R., Trainer, M. K., Tsai, C., Veres, P. R., Washenfelder, R. A., Warneke, C., Wild, R. J., Young,
C. J., Yuan, B., and Zamora, R.: High winter ozone pollution from carbonyl photolysis in an oil and gas basin,
Nature, 514, 351-354, 10.1038/nature13767, 2014.

925 Eilerman, S. J., Peischl, J., Neuman, J. A., Ryerson, T. B., Aikin, K. C., Holloway, M. W., Zondlo, M. A.,
Golston, L. M., Pan, D., Floerchinger, C., and Herndon, S.: Characterization of Ammonia, Methane, and
Nitrous Oxide Emissions from Concentrated Animal Feeding Operations in Northeastern Colorado,
Environ. Sci. Technol., 50, 10885-10893, 10.1021/acs.est.6b02851, 2016.

930 Eisele, A. P., Mukerjee, S., Smith, L. A., Thoma, E. D., Whitaker, D. A., Oliver, K. D., Wu, T., Colon, M., Alston,
L., Cousett, T. A., Miller, M. C., Smith, D. M., and Stallings, C.: Volatile organic compounds at two oil and
natural gas production well pads in Colorado and Texas using passive samplers, *JAPCA J. Air Waste Ma.*,
66, 412-419, 10.1080/10962247.2016.1141808, 2016.

Fall, R., Karl, T., Jordan, A., and Lindinger, W.: Biogenic C5 VOCs: release from leaves after freeze-thaw
wounding and occurrence in air at a high mountain observatory, *Atmos. Environ.*, 35, 3905-3916,
[http://dx.doi.org/10.1016/S1352-2310\(01\)00141-8](http://dx.doi.org/10.1016/S1352-2310(01)00141-8), 2001.



- 935 Feilberg, A., Bildsoe, P., and Nyord, T.: Application of PTR-MS for Measuring Odorant Emissions from Soil Application of Manure Slurry, *Sensors*, 15, 1148-1167, 2015.
- Field, R. A., Soltis, J., McCarthy, M. C., Murphy, S., and Montague, D. C.: Influence of oil and gas field operations on spatial and temporal distributions of atmospheric non-methane hydrocarbons and their effect on ozone formation in winter, *Atmos. Chem. Phys.*, 15, 3527-3542, 10.5194/acp-15-3527-2015, 2015.
- 940 Fu, T.-M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J. Geophys. Res.*, 113, 10.1029/2007JD009505, 2008.
- Gentner, D. R., Worton, D. R., Isaacman, G., Davis, L. C., Dallmann, T. R., Wood, E. C., Herndon, S. C., Goldstein, A. H., and Harley, R. A.: Chemical Composition of Gas-Phase Organic Carbon Emissions from Motor Vehicles and Implications for Ozone Production, *Environ. Sci. Technol.*, 47, 11837-11848, 10.1021/es401470e, 2013.
- 945 Gilman, J. B., Lerner, B. M., Kuster, W. C., and de Gouw, J. A.: Source signature of volatile organic compounds from oil and natural gas operations in northeastern Colorado, *Environ. Sci. Technol.*, 47, 1297-1305, 10.1021/es304119a, 2013.
- 950 Graus, M., Müller, M., and Hansel, A.: High Resolution PTR-TOF: Quantification and Formula Confirmation of VOC in Real Time, *J. Am. Soc. Mass Spectr.*, 21, 1037-1044, <http://dx.doi.org/10.1016/j.jasms.2010.02.006>, 2010.
- Grosjean, D., and Williams, E. L.: Environmental persistence of organic compounds estimated from structure-reactivity and linear free-energy relationships. Unsaturated aliphatics, *Atmos. Environ.*, 26, 1395-1405, [http://dx.doi.org/10.1016/0960-1686\(92\)90124-4](http://dx.doi.org/10.1016/0960-1686(92)90124-4), 1992.
- 955 Gueneron, M., Erickson, M. H., VanderSchelden, G. S., and Jobson, B. T.: PTR-MS fragmentation patterns of gasoline hydrocarbons, *Int. J. Mass Spectrom.*, 379, 97-109, <http://dx.doi.org/10.1016/j.ijms.2015.01.001>, 2015.
- 960 Haase, K. B., Keene, W. C., Pszenny, A. A. P., Mayne, H. R., Talbot, R. W., and Sive, B. C.: Calibration and intercomparison of acetic acid measurements using proton-transfer-reaction mass spectrometry (PTR-MS), *Atmos. Meas. Tech.*, 5, 2739-2750, 10.5194/amt-5-2739-2012, 2012.
- Hansel, A., and Wisthaler, A.: A method for real-time detection of PAN, PPN and MPAN in ambient air, *Geophys. Res. Lett.*, 27, 2000.
- 965 Harris, G. W., Kleindienst, T. E., and Pitts Jr, J. N.: Rate constants for the reaction of OH radicals with CH₃CN, C₂H₅CN AND CH₂CHCN in the temperature range 298–424 K, *Chem. Phys. Lett.*, 80, 479-483, [http://dx.doi.org/10.1016/0009-2614\(81\)85061-0](http://dx.doi.org/10.1016/0009-2614(81)85061-0), 1981.
- Heikes, B. G., Chang, W., Pilson, M. E. Q., Swift, E., Singh, H. B., Guenther, A., Jacob, D. J., Field, B. D., Fall, R., Riemer, D., and Brand, L.: Atmospheric methanol budget and ocean implication, *Global Biogeochem. Cy.*, 16, 80-81-80-13, 10.1029/2002GB001895, 2002.



- 970 Hendler, A., Nunn, J., and Lundeen, J.: VOC emissions from oil and condensate storage tanks, Houston Advanced Research Center, 2009.
- Henze, D. K., Seinfeld, J. H., Ng, N. L., Kroll, J. H., Fu, T. M., Jacob, D. J., and Heald, C. L.: Global modeling of secondary organic aerosol formation from aromatic hydrocarbons: high- vs. low-yield pathways, *Atmos. Chem. Phys.*, 8, 2405-2420, 10.5194/acp-8-2405-2008, 2008.
- 975 Holmes, S. A., and Thompson, L. F.: Nitrogen compound distributions in hydrotreated shale oil products from commercial-scale refining, *Fuel*, 62, 709-717, [http://dx.doi.org/10.1016/0016-2361\(83\)90312-5](http://dx.doi.org/10.1016/0016-2361(83)90312-5), 1983.
- Jacob, D. J., Field, B. D., Li, Q., Blake, D. R., de Gouw, J., Warneke, C., Hansel, A., Wisthaler, A., Singh, H. B., and Guenther, A.: Global budget of methanol: Constraints from atmospheric observations, *J. Geophys. Res.*: Atmos., 110, 10.1029/2004JD005172, 2005.
- 980 Jobson, B. T., Alexander, M. L., Maupin, G. D., and Muntean, G. G.: On-line analysis of organic compounds in diesel exhaust using a proton transfer reaction mass spectrometer (PTR-MS), *Int. J. Mass Spectrom.*, 245, 78-89, <http://dx.doi.org/10.1016/j.ijms.2005.05.009>, 2005.
- Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Märk, L., Seehauser, H., Schottkowsky, R., Sulzer, P., and Märk, T. D.: A high resolution and high sensitivity proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF-MS), *Int. J. Mass Spectrom.*, 286, 122-128, <http://dx.doi.org/10.1016/j.ijms.2009.07.005>, 2009.
- 985 Karl, T., Jobson, T., Kuster, W. C., Williams, E., Stutz, J., Shetter, R., Hall, S. R., Goldan, P., Fehsenfeld, F., and Lindinger, W.: Use of proton-transfer-reaction mass spectrometry to characterize volatile organic compound sources at the La Porte super site during the Texas Air Quality Study 2000, *J. Geophys. Res.*, 108, 10.1029/2002JD003333, 2003.
- 990 Karl, T. G., Christian, T. J., Yokelson, R. J., Artaxo, P., Hao, W. M., and Guenther, A.: The Tropical Forest and Fire Emissions Experiment: method evaluation of volatile organic compound emissions measured by PTR-MS, FTIR, and GC from tropical biomass burning, *Atmos. Chem. Phys.*, 7, 5883-5897, 10.5194/acp-7-5883-2007, 2007.
- 995 Kaser, L., Karl, T., Schnitzhofer, R., Graus, M., Herdlinger-Blatt, I. S., DiGangi, J. P., Sive, B., Turnipseed, A., Hornbrook, R. S., Zheng, W., Flocke, F. M., Guenther, A., Keutsch, F. N., Apel, E., and Hansel, A.: Comparison of different real time VOC measurement techniques in a ponderosa pine forest, *Atmos. Chem. Phys.*, 13, 2893-2906, 10.5194/acp-13-2893-2013, 2013.
- 1000 Kemball-Cook, S., Bar-Ilan, A., Grant, J., Parker, L., Jung, J., Santamaria, W., Mathews, J., and Yarwood, G.: Ozone Impacts of Natural Gas Development in the Haynesville Shale, *Environ. Sci. Technol.*, 44, 9357-9363, 10.1021/es1021137, 2010.
- 1005 Kim, S., Karl, T., Helmig, D., Daly, R., Rasmussen, R., and Guenther, A.: Measurement of atmospheric sesquiterpenes by proton transfer reaction-mass spectrometry (PTR-MS), *Atmos. Meas. Tech.*, 2, 99-112, 10.5194/amt-2-99-2009, 2009.
- Kim, S., Karl, T., Guenther, A., Tyndall, G., Orlando, J., Harley, P., Rasmussen, R., and Apel, E.: Emissions and ambient distributions of Biogenic Volatile Organic Compounds (BVOC) in a ponderosa pine ecosystem:



- interpretation of PTR-MS mass spectra, *Atmos. Chem. Phys.*, **10**, 1759-1771, 10.5194/acp-10-1759-2010, 2010.
- 1010 Knighton, W. B., Herndon, S. C., Franklin, J. F., Wood, E. C., Wormhoudt, J., Brooks, W., Fortner, E. C., and Allen, D. T.: Direct measurement of volatile organic compound emissions from industrial flares using real-time online techniques: Proton Transfer Reaction Mass Spectrometry and Tunable Infrared Laser Differential Absorption Spectroscopy, *Ind. Eng. Chem. Res.*, **51**, 12674-12684, 10.1021/ie202695v, 2012.
- 1015 Koss, A. R., de Gouw, J., Warneke, C., Gilman, J. B., Lerner, B. M., Graus, M., Yuan, B., Edwards, P., Brown, S. S., Wild, R., Roberts, J. M., Bates, T. S., and Quinn, P. K.: Photochemical aging of volatile organic compounds associated with oil and natural gas extraction in the Uintah Basin, UT, during a wintertime ozone formation event, *Atmos. Chem. Phys.*, **15**, 5727-5741, 10.5194/acp-15-5727-2015, 2015.
- 1020 Kuster, W. C., Jobson, B. T., Karl, T., Riemer, D., Apel, E., Goldan, P. D., and Fehsenfeld, F. C.: Intercomparison of Volatile Organic Carbon Measurement Techniques and Data at La Porte during the TexAQ52000 Air Quality Study, *Environ. Sci. Technol.*, **38**, 221-228, 10.1021/es034710r, 2004.
- 1025 Lerner, B. M., Gilman, J. B., Aikin, K. C., Atlas, E. L., Goldan, P. D., Graus, M., Hendershot, R., Isaacman-VanWertz, G. A., Koss, A., Kuster, W. C., Lueb, R. A., McLaughlin, R. J., Peischl, J., Sueper, D., Ryerson, T. B., Tokarek, T. W., Warneke, C., Yuan, B., and de Gouw, J. A.: An improved, automated whole air sampler and gas chromatography mass spectrometry analysis system for volatile organic compounds in the atmosphere, *Atmos. Meas. Tech.*, **10**, 291-313, 10.5194/amt-10-291-2017, 2017.
- Lewis, A. C., Hopkins, J. R., Carpenter, L. J., Stanton, J., Read, K. A., and Pilling, M. J.: Sources and sinks of acetone, methanol, and acetaldehyde in North Atlantic marine air, *Atmos. Chem. Phys.*, **5**, 1963-1974, 10.5194/acp-5-1963-2005, 2005.
- 1030 Li, M., Larter, S. R., Stoddart, D., and Bjørøy, M.: Fractionation of pyrrolic nitrogen compounds in petroleum during migration: derivation of migration-related geochemical parameters, Geological Society, London, Special Publications, **86**, 103-123, 10.1144/gsl.sp.1995.086.01.09, 1995.
- 1035 Li, R., Warneke, C., Graus, M., Field, R., Geiger, F., Veres, P. R., Soltis, J., Li, S. M., Murphy, S. M., Sweeney, C., Pétron, G., Roberts, J. M., and de Gouw, J.: Measurements of hydrogen sulfide (H₂S) using PTR-MS: calibration, humidity dependence, inter-comparison and results from field studies in an oil and gas production region, *Atmos. Meas. Tech.*, **7**, 3597-3610, 10.5194/amt-7-3597-2014, 2014.
- Lim, Y. B., and Ziemann, P. J.: Products and Mechanism of Secondary Organic Aerosol Formation from Reactions of n-Alkanes with OH Radicals in the Presence of NO_x, *Environ. Sci. Technol.*, **39**, 9229-9236, 10.1021/es051447g, 2005.
- 1040 Mansfield, M., Lyman, S., and Tran, H.: RPSEA Final Report: Measurement of hydrocarbons and greenhouse gases from uncharacterized area sources, Bingham Research Center, Utah State University, 2016.
- 1045 May, A. A., Nguyen, N. T., Presto, A. A., Gordon, T. D., Lipsky, E. M., Karve, M., Gutierrez, A., Robertson, W. H., Zhang, M., Brandow, C., Chang, O., Chen, S., Cicero-Fernandez, P., Dinkins, L., Fuentes, M., Huang, S.-M., Ling, R., Long, J., Maddox, C., Massetti, J., McCauley, E., Miguel, A., Na, K., Ong, R., Pang, Y., Rieger, P., Sax, T., Truong, T., Vo, T., Chattopadhyay, S., Maldonado, H., Maricq, M. M., and Robinson, A. L.: Gas-



- and particle-phase primary emissions from in-use, on-road gasoline and diesel vehicles, *Atmos. Environ.*, **48**, 247-260, <http://dx.doi.org/10.1016/j.atmosenv.2014.01.046>, 2014.
- McDuffie, E. E., Edwards, P. M., Gilman, J. B., Lerner, B. M., Dubé, W. P., Trainer, M., Wolfe, D. E., Angevine, W. M., deGouw, J., Williams, E. J., Tevlin, A. G., Murphy, J. G., Fischer, E. V., McKeen, S., Ryerson, T. B., Peischl, J., Holloway, J. S., Aikin, K., Langford, A. O., Senff, C. J., Alvarez, R. J., Hall, S. R., Ullmann, K., Lantz, K. O., and Brown, S. S.: Influence of oil and gas emissions on summertime ozone in the Colorado Northern Front Range, *J. Geophys. Res.: Atmos.*, **121**, 8712-8729, 10.1002/2016JD025265, 2016.
- McKenzie, L. M., Witter, R. Z., Newman, L. S., and Adgate, J. L.: Human health risk assessment of air emissions from development of unconventional natural gas resources, *Sci. Total Environ.*, **424**, 79-87, <http://dx.doi.org/10.1016/j.scitotenv.2012.02.018>, 2012.
- Midey, A. J., Williams, S., Miller, T. M., and Viggiano, A. A.: Reactions of O₂⁺, NO⁺ and H₃O⁺ with methylcyclohexane (C₇H₁₄) and cyclooctane (C₈H₁₆) from 298 to 700 K, *Int. J. Mass Spectrom.*, **222**, 413-430, [http://dx.doi.org/10.1016/S1387-3806\(02\)00996-X](http://dx.doi.org/10.1016/S1387-3806(02)00996-X), 2003.
- Miller, S. M., Wofsy, S. C., Michalak, A. M., Kort, E. A., Andrews, A. E., Biraud, S. C., Dlugokencky, E. J., Eluskiewicz, J., Fischer, M. L., Janssens-Maenhout, G., Miller, B. R., Miller, J. B., Montzka, S. A., Nehr Korn, T., and Sweeney, C.: Anthropogenic emissions of methane in the United States, *P. Natl. Acad. Sci. USA*, **110**, 20018-20022, 10.1073/pnas.1314392110, 2013.
- Millet, D. B., Jacob, D. J., Custer, T. G., de Gouw, J. A., Goldstein, A. H., Karl, T., Singh, H. B., Sive, B. C., Talbot, R. W., Warneke, C., and Williams, J.: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmos. Chem. Phys.*, **8**, 6887-6905, 10.5194/acp-8-6887-2008, 2008.
- Morandi, J. R., and Jensen, H. B.: Comparison of Porphyrins from Shale Oil, Oil Shale, and Petroleum by Absorption and Mass Spectroscopy, *J. Chem. Eng. Data*, **11**, 81-88, 10.1021/je60028a023, 1966.
- Müller, M., Graus, M., Wisthaler, A., Hansel, A., Metzger, A., Dommen, J., and Baltensperger, U.: Analysis of high mass resolution PTR-TOF mass spectra from 1,3,5-trimethylbenzene (TMB) environmental chamber experiments, *Atmos. Chem. Phys.*, **12**, 829-843, 10.5194/acp-12-829-2012, 2012.
- Mushrush, G. W., Beal, E. J., Hardy, D. R., and Hughes, J. M.: Nitrogen compound distribution in middle distillate fuels derived from petroleum, oil shale, and tar sand sources, *Fuel Process. Technol.*, **61**, 197-210, [http://dx.doi.org/10.1016/S0378-3820\(99\)00056-9](http://dx.doi.org/10.1016/S0378-3820(99)00056-9), 1999.
- National Research Council: Chemical Composition of Petroleum Hydrocarbon Sources, in: *Oil in the Sea: Inputs, Fates, and Effects*, National Academy Press, Washington, DC, 1985.
- Neuman, J. A., Trainer, M., Brown, S. S., Min, K. E., Nowak, J. B., Parrish, D. D., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., and Veres, P. R.: HONO emission and production determined from airborne measurements over the Southeast U.S, *J. Geophys. Res.: Atmos.*, **121**, 9237-9250, 10.1002/2016JD025197, 2016.
- Ngwabie, N. M., Schade, G. W., Custer, T. G., Linke, S., and Hinz, T.: Abundances and Flux Estimates of Volatile Organic Compounds from a Dairy Cowshed in Germany, *J. Environ. Qual.*, **37**, 565-573, 10.2134/jeq2006.0417, 2008.



- 1085 Oldenburg, T. B. P., Larter, S. R., and Huang, H.: Nitrogen isotope systematics of petroleum fractions of differing polarity – Neutral versus basic compounds, *Org. Geochem.*, 38, 1789-1794, <http://dx.doi.org/10.1016/j.orggeochem.2007.05.016>, 2007.
- Pang, X., Lewis, A. C., Rickard, A. R., Baeza-Romero, M. T., Adams, T. J., Ball, S. M., Daniels, M. J. S., Goodall, I. C. A., Monks, P. S., Peppe, S., Ródenas García, M., Sánchez, P., and Muñoz, A.: A smog chamber comparison of a microfluidic derivatisation measurement of gas-phase glyoxal and methylglyoxal with other analytical techniques, *Atmos. Meas. Tech.*, 7, 373-389, 10.5194/amt-7-373-2014, 2014.
- 1090 Park, J. H., Goldstein, A. H., Timkovsky, J., Fares, S., Weber, R., Karlik, J., and Holzinger, R.: Eddy covariance emission and deposition flux measurements using proton transfer reaction – time of flight – mass spectrometry (PTR-TOF-MS): comparison with PTR-MS measured vertical gradients and fluxes, *Atmos. Chem. Phys.*, 13, 1439-1456, 10.5194/acp-13-1439-2013, 2013.
- 1095 Peischl, J., Ryerson, T. B., Holloway, J. S., Trainer, M., Andrews, A. E., Atlas, E. L., Blake, D. R., Daube, B. C., Dlugokencky, E. J., Fischer, M. L., Goldstein, A. H., Guha, A., Karl, T., Kofler, J., Kosciuch, E., Misztal, P. K., Perring, A. E., Pollack, I. B., Santoni, G. W., Schwarz, J. P., Spackman, J. R., Wofsy, S. C., and Parrish, D. D.: Airborne observations of methane emissions from rice cultivation in the Sacramento Valley of California, *J. Geophys. Res.*, 117, n/a-n/a, 10.1029/2012JD017994, 2012.
- 1100 Peischl, J., Ryerson, T. B., Aikin, K. C., de Gouw, J. A., Gilman, J. B., Holloway, J. S., Lerner, B. M., Nadkarni, R., Neuman, J. A., Nowak, J. B., Trainer, M., Warneke, C., and Parrish, D. D.: Quantifying atmospheric methane emissions from the Haynesville, Fayetteville, and northeastern Marcellus shale gas production regions, *J. Geophys. Res.: Atmos.*, 120, 2119-2139, 10.1002/2014JD022697, 2015.
- 1105 Pétron, G., Karion, A., Sweeney, C., Miller, B. R., Montzka, S. A., Frost, G. J., Trainer, M., Tans, P., Andrews, A., Kofler, J., Helmig, D., Guenther, D., Dlugokencky, E., Lang, P., Newberger, T., Wolter, S., Hall, B., Novelli, P., Brewer, A., Conley, S., Hardesty, M., Banta, R., White, A., Noone, D., Wolfe, D., and Schnell, R.: A new look at methane and nonmethane hydrocarbon emissions from oil and natural gas operations in the Colorado Denver-Julesburg Basin, *J. Geophys. Res.*, 119, 6836-6852, 10.1002/2013JD021272, 2014.
- 1110 Pollack, I. B., Lerner, B. M., and Ryerson, T. B.: Evaluation of ultraviolet light-emitting diodes for detection of atmospheric NO₂ by photolysis - chemiluminescence, *J. Atmos. Chem.*, 65, 111-125, 10.1007/s10874-011-9184-3, 2010.
- Pring, M.: Condensate Tank Oil and Gas Activities, Eastern Research Group Inc., 2011.
- Railroad Commission of Texas: Oil and Gas Research and Statistics: <http://www.rrc.state.tx.us/oil-gas/research-and-statistics/>, access: 1/31/2017, 2017.
- 1115 Rich, A., Grover, J. P., and Sattler, M. L.: An exploratory study of air emissions associated with shale gas development and production in the Barnett Shale, *JAPCA J. Waste Ma.*, 64, 61-72, 10.1080/10962247.2013.832713, 2014.
- Rinne, J., Ruuskanen, T. M., Reissell, A., Taipale, R., Hakola, H., and Kulmala, M.: On-line PTR-MS measurements of atmospheric concentrations of volatile organic compounds in a European boreal forest ecosystem, *Boreal environment research*, 10, 425-436, 2005.



- 1120 Ruuskanen, T., Müller, M., Schnitzhofer, R., Karl, T., Graus, M., Bamberger, I., Hörtnagl, L., Brill, F., Wohlfahrt, G., and Hansel, A.: Eddy covariance VOC emission and deposition fluxes above grassland using PTR-TOF, *Atmos. Chem. Phys.*, **11**, 611-625, 2011.
- Ryerson, T. B., Williams, E. J., and Fehsenfeld, F. C.: An efficient photolysis system for fast-response NO₂ measurements, *J. Geophys. Res.: Atmos.*, **105**, 26447-26461, 10.1029/2000JD900389, 2000.
- 1125 Schade, G. W., and Roest, G.: Analysis of non-methane hydrocarbon data from a monitoring station affected by oil and gas development in the Eagle Ford shale, Texas, *Elementa*, **4**, 2016.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of Emissions from Air Pollution Sources. 5. C₁-C₃₂ Organic Compounds from Gasoline-Powered Motor Vehicles, *Environ. Sci. Technol.*, **36**, 1169-1180, 10.1021/es0108077, 2002.
- 1130 Sekimoto, K., Li, S.-M., Yuan, B., Koss, A., Coggon, M. M., Warnke, C., and de Gouw, J.: Calculation of the sensitivity of proton-transfer-reaction mass spectrometry (PTR-MS) for organic trace gases, *Int. J. Mass Spectrom.*, in review, 2017.
- Simpson, I. J., Blake, N. J., Barletta, B., Diskin, G. S., Fuelberg, H. E., Gorham, K., Huey, L. G., Meinardi, S., Rowland, F. S., Vay, S. A., Weinheimer, A. J., Yang, M., and Blake, D. R.: Characterization of trace gases measured over Alberta oil sands mining operations: 76 speciated C₂-C₁₀ volatile organic compounds (VOCs), CO₂, CH₄, CO, NO, NO₂, NO_y, O₃ and SO₂, *Atmos. Chem. Phys.*, **10**, 11931-11954, 10.5194/acp-10-11931-2010, 2010.
- 1135 Singh, H. B., Kanakidou, M., Crutzen, P. J., and Jacob, D. J.: High concentrations and photochemical fate of oxygenated hydrocarbons in the global troposphere, *Nature*, **378**, 50-54, 1995.
- 1140 Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal dissociation-chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of peroxyacetyl nitrates and dinitrogen pentoxide, *J. Geophys. Res.: Atmos.*, **109**, D19315, 10.1029/2004JD004670, 2004.
- Smith, D. F., McIver, C. D., and Kleindienst, T. E.: Primary Product Distribution from the Reaction of Hydroxyl Radicals with Toluene at ppb NO_x Mixing Ratios, *J. Atmos. Chem.*, **30**, 209-228, 10.1023/a:1005980301720, 1998.
- 1145 Smith, H. M.: *Qualitative and Quantitative Aspects of Crude Oil Composition*, Washington, DC, 1968.
- State of New Mexico Oil Conservation Division: Oil Conservation Division Data and Statistics: <http://www.emnrd.state.nm.us/OCD/statistics.html>, access: 2/1/2017, 2017.
- State of Utah Division of Oil Gas and Mining: Online Oil and Gas Information System: https://oilgas.ogm.utah.gov/Data_Center, access: 11/31/2017, 2017.
- 1150 Stockwell, C. E., Veres, P. R., Williams, J., and Yokelson, R. J.: Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, **15**, 845-865, 10.5194/acp-15-845-2015, 2015.



- 1155 Stönner, C., Derstroff, B., Klüpfel, T., Crowley, J. N., and Williams, J.: Glyoxal measurement with a proton transfer reaction time of flight mass spectrometer (PTR-TOF-MS): characterization and calibration, *J. Mass Spectrom.*, 52, 30-35, 10.1002/jms.3893, 2017.
- 1160 Stout, S. A., and Douglas, G. S.: Diamondoid Hydrocarbons—Application in the Chemical Fingerprinting of Natural Gas Condensate and Gasoline, *Environ. Forensics*, 5, 225-235, 10.1080/15275920490886734, 2004.
- Swarthout, R. F., Russo, R. S., Zhou, Y., Hart, A. H., and Sive, B. C.: Volatile organic compound distributions during the NACHTT campaign at the Boulder Atmospheric Observatory: Influence of urban and natural gas sources, *J. Geophys. Res.: Atmos.*, 118, 614-610,637, 10.1002/jgrd.50722, 2013.
- 1165 Tarver, G. A., and Dasgupta, P. K.: Oil Field Hydrogen Sulfide in Texas: Emission Estimates and Fate, *Environ. Sci. Technol.*, 31, 3669-3676, 10.1021/es970406k, 1997.
- Tissot, B. P., and Welte, D. H.: *Geochemical Fossils and Their Significance in Petroleum Formation, in: Petroleum Formation and Occurrence*, Springer Berlin Heidelberg, Berlin, Heidelberg, 93-130, 1984.
- US Energy Information Administration: Drilling Productivity Report: <http://www.eia.gov/petroleum/drilling/>, access: 1/31/2017, 2017.
- 1170 Utah Division of Air Quality: Final Report: Uinta Basin Winter Ozone Study, 2014.
- Veres, P. R., Roberts, J. M., Cochran, A. K., Gilman, J. B., Kuster, W. C., Holloway, J. S., Graus, M., Flynn, J., Lefer, B., Warneke, C., and de Gouw, J.: Evidence of rapid production of organic acids in an urban air mass, 38, n/a-n/a, 10.1029/2011GL048420, 2011.
- 1175 Verma, D. K., and Tombe, K. d.: Benzene in Gasoline and Crude Oil: Occupational and Environmental Implications, *AIHA Journal*, 63, 225-230, 10.1080/15428110208984708, 2002.
- Wagner, N. L., Dubé, W. P., Washenfelder, R. A., Young, C. J., Pollack, I. B., Ryerson, T. B., and Brown, S. S.: Diode laser-based cavity ring-down instrument for NO₃, N₂O₅, NO, NO₂ and O₃ from aircraft, *Atmos. Meas. Tech.*, 4, 1227-1240, 10.5194/amt-4-1227-2011, 2011.
- 1180 Wagner, V., Jenkin, M. E., Saunders, S. M., Stanton, J., Wirtz, K., and Pilling, M. J.: Modelling of the photooxidation of toluene: conceptual ideas for validating detailed mechanisms, *Atmos. Chem. Phys.*, 3, 89-106, 10.5194/acp-3-89-2003, 2003.
- Wallington, T. J.: Kinetics of the gas phase reaction of OH radicals with pyrrole and thiophene, *Int. J. Chem. Kinet.*, 18, 487-496, 10.1002/kin.550180407, 1986.
- 1185 Wang, Z., Yang, C., Hollebone, B., and Fingas, M.: Forensic Fingerprinting of Diamondoids for Correlation and Differentiation of Spilled Oil and Petroleum Products, *Environ. Sci. Technol.*, 40, 5636-5646, 10.1021/es060675n, 2006.
- Warneke, C., de Gouw, J. A., Kuster, W. C., Goldan, P. D., and Fall, R.: Validation of Atmospheric VOC Measurements by Proton-Transfer- Reaction Mass Spectrometry Using a Gas-Chromatographic Preseparation Method, *Environ. Sci. Technol.*, 37, 2494-2501, 10.1021/es026266i, 2003.



- 1190 Warneke, C., de Gouw, J. A., Edwards, P. M., Holloway, J. S., Gilman, J. B., Kuster, W. C., Graus, M., Atlas, E., Blake, D., Gentner, D. R., Goldstein, A. H., Harley, R. A., Alvarez, S., Rappenglueck, B., Trainer, M., and Parrish, D. D.: Photochemical aging of volatile organic compounds in the Los Angeles basin: Weekday-weekend effect, *J. Geophys. Res.*, 118, 5018-5028, 10.1002/jgrd.50423, 2013.
- 1195 Warneke, C., Geiger, F., Edwards, P. M., Dube, W., Pétron, G., Kofler, J., Zahn, A., Brown, S. S., Graus, M., Gilman, J. B., Lerner, B. M., Peischl, J., Ryerson, T. B., de Gouw, J. A., and Roberts, J. M.: Volatile organic compound emissions from the oil and natural gas industry in the Uintah Basin, Utah: oil and gas well pad emissions compared to ambient air composition, *Atmos. Chem. Phys.*, 14, 10977-10988, 10.5194/acp-14-10977-2014, 2014.
- 1200 Warneke, C., Veres, P., Murphy, S. M., Soltis, J., Field, R. A., Graus, M. G., Koss, A., Li, S. M., Li, R., Yuan, B., Roberts, J. M., and de Gouw, J. A.: PTR-QMS versus PTR-TOF comparison in a region with oil and natural gas extraction industry in the Uintah Basin in 2013, *Atmos. Meas. Tech.*, 8, 411-420, 10.5194/amt-8-411-2015, 2015.
- Williams, P. T., and Chishti, H. M.: Reaction of nitrogen and sulphur compounds during catalytic hydrotreatment of shale oil, *Fuel*, 80, 957-963, 2001.
- 1205 Wyche, K. P., Monks, P. S., Ellis, A. M., Cordell, R. L., Parker, A. E., Whyte, C., Metzger, A., Dommen, J., Duplissy, J., Prevot, A. S. H., Baltensperger, U., Rickard, A. R., and Wulfert, F.: Gas phase precursors to anthropogenic secondary organic aerosol: detailed observations of 1,3,5-trimethylbenzene photooxidation, *Atmos. Chem. Phys.*, 9, 635-665, 10.5194/acp-9-635-2009, 2009.
- 1210 Wyoming Oil and Gas Conservation Commission: Production Data: <http://wogcc.state.wy.us/>, access: 1/31/2017, 2017.
- Yacovitch, T. I., Herndon, S. C., Roscioli, J. R., Floerchinger, C., McGovern, R. M., Agnese, M., Pétron, G., Kofler, J., Sweeney, C., Karion, A., Conley, S. A., Kort, E. A., Nähle, L., Fischer, M., Hildebrandt, L., Koeth, J., McManus, J. B., Nelson, D. D., Zahniser, M. S., and Kolb, C. E.: Demonstration of an Ethane Spectrometer for Methane Source Identification, *Environ. Sci. Technol.*, 48, 8028-8034, 10.1021/es501475q, 2014.
- 1215 Yuan, B., Warneke, C., Shao, M., and de Gouw, J. A.: Interpretation of volatile organic compound measurements by proton-transfer-reaction mass spectrometry over the deepwater horizon oil spill, *Int. J. Mass Spectrom.*, 358, 43-48, <http://dx.doi.org/10.1016/j.ijms.2013.11.006>, 2014.
- 1220 Yuan, B., Veres, P. R., Warneke, C., Roberts, J. M., Gilman, J. B., Koss, A., Edwards, P. M., Graus, M., Kuster, W. C., Li, S. M., Wild, R. J., Brown, S. S., Dubé, W. P., Lerner, B. M., Williams, E. J., Johnson, J. E., Quinn, P. K., Bates, T. S., Lefer, B., Hayes, P. L., Jimenez, J. L., Weber, R. J., Zamora, R., Ervens, B., Millet, D. B., Rappenglück, B., and de Gouw, J. A.: Investigation of secondary formation of formic acid: urban environment vs. oil and gas producing region, *Atmos. Chem. Phys.*, 15, 1975-1993, 10.5194/acp-15-1975-2015, 2015.
- 1225 Yuan, B., Koss, A., Warneke, C., Gilman, J. B., Lerner, B. M., Stark, H., and de Gouw, J. A.: A high-resolution time-of-flight chemical ionization mass spectrometer utilizing hydronium ions (H₃O⁺ ToF-CIMS) for measurements of volatile organic compounds in the atmosphere, *Atmos. Meas. Tech.*, 9, 2735-2752, 10.5194/amt-9-2735-2016, 2016a.



1230 Yuan, B., Liggio, J., Wentzell, J., Li, S. M., Stark, H., Roberts, J. M., Gilman, J., Lerner, B., Warneke, C., Li, R.,
Leithead, A., Osthoff, H. D., Wild, R., Brown, S. S., and de Gouw, J. A.: Secondary formation of nitrated
phenols: insights from observations during the Uintah Basin Winter Ozone Study (UBWOS) 2014, Atmos.
Chem. Phys., 16, 2139-2153, 10.5194/acp-16-2139-2016, 2016b.

1235 Yuan, B., Coggon, M. M., Koss, A. R., Warneke, C., Eilerman, S., Peischl, J., Aikiin, K. C., Ryerson, T. B., and
de Gouw, J. A.: Emissions of volatile organic compounds (VOCs) from concentrated animal feeding
operations (CAFOs): chemical compositions and separation of sources, Atmos. Chem. Phys. Discuss., 2017,
1-25, 10.5194/acp-2016-1148, 2017.

**Table 1.** SONGNEX chemical instrumentation

Name of instrument	Species measured	Measurement technique	Citation or details
iWAS/GC-MS (improved Whole Air Sampler)	C2-C6 alkanes Select cycloalkanes C6-C8 aromatics Small alkyl nitrates Isoprene Monoterpenes Ethene Ethyne Methanol	Whole-air canister sampling followed by offline GC-MS analysis	(Lerner et al., 2017)
Aerodyne C ₂ H ₆ instrument	Ethane	Tunable infrared laser direct absorption spectroscopy	(Yacovitch et al., 2014)
Picarro CO ₂ /CH ₄ instrument	Methane CO ₂	Cavity ring-down spectroscopy	(Peischl et al., 2012)
PAN CIMS	Peroxyacyl nitrates: PAN PPN MPAN APAN	Thermal dissociation/ I ⁻ chemical ionization mass spectrometry	(Slusher et al., 2004)
In-Situ Airborne Formaldehyde (ISAF)	Formaldehyde	Laser-induced fluorescence	(Cazorla et al., 2015)
Picarro G1204 H ₂ S instrument	H ₂ S	Cavity ring-down spectroscopy	Manufacturer specifications: 10 ppb 5-s LoD (1σ) ~5 second measurement interval May experience interference from organics
NO/NO ₂ /NO _y Chemi- luminescence	NO NO ₂ NO _y	NO by NO/O ₃ chemiluminescence NO ₂ by photolysis and NO/ O ₃ chemiluminescence NO _y by Au converter and NO/O ₃ chemiluminescence	(Ryerson et al., 2000;Pollack et al., 2010)
HNO ₃ /HCOOH CIMS	Nitric acid Formic acid	I ⁻ chemical ionization mass spectrometry	(Neuman et al., 2016)
NO _x CaRDS	NO NO ₂ NO _y	Cavity ring-down spectroscopy	(Wagner et al., 2011)



Table 2. Significant ion masses and interpretation. Mixing ratio (ppbv) enhancements are listed only for non-fragmentary ions of relatively certain identification. Ion masses with VOC interpretation of particular interest are highlighted using bold text.

Ion Exact Mass (Th)	Ion formula	Previously reported interpretations	Interpretation in oil and gas producing regions	Max boundary layer enhancement (ppbv) during April 23 flight
33.0335	CH ₄ OH ⁺	Methanol (de Gouw and Warneke, 2007;Blake et al., 2009)	Methanol	19.09
34.9950	H ₂ SH ⁺	Hydrogen sulfide (Li et al., 2014)	Hydrogen sulfide	12.6
41.0386	C ₃ H ₅ ⁺	General alkane/VOC fragment (Gueneron et al., 2015) MBO fragment (Kim et al., 2010); propyne (Stockwell et al., 2015)	General alkane/VOC fragment	
43.0178	C ₂ H ₂ OH ⁺	Biogenic aldehyde fragment (Kim et al., 2010;Ruuskanen et al., 2011) Acetic acid fragment (de Gouw et al., 2003;Müller et al., 2012)	Fragment of acetic acid	
43.0542	C ₃ H ₇ ⁺	General alkane/VOC fragment (Gueneron et al., 2015) Propene (Kuster et al., 2004;Knighton et al., 2012;Stockwell et al., 2015)	General alkane/VOC fragment	
45.0335	C ₂ H ₄ OH ⁺	Acetaldehyde (de Gouw and Warneke, 2007;Blake et al., 2009)	Acetaldehyde	3.60
45.9924	NO₂⁺	PAN (de Gouw et al., 2003;Müller et al., 2012;Kaser et al., 2013)	Unresolvable NO₂ species	
57.0699	C ₄ H ₉ ⁺	General alkane/VOC fragment (Gueneron et al., 2015) Butenes (Karl et al., 2003)	General alkane/VOC fragment	
59.0491	C ₃ H ₆ OH ⁺	Acetone (de Gouw and Warneke, 2007;Blake et al., 2009)	Acetone (negligible contribution from propanal)	6.17
61.0284	C ₂ H ₄ O ₂ H ⁺	Acetic acid (de Gouw et al., 2003)	Acetic acid	1.57
68.0495	C₄H₅NH⁺	Pyrrrole (Brilli et al., 2014;Stockwell et al., 2015)	Pyrrrole	0.04
69.0699	C₅H₉⁺	Isoprene (Blake et al., 2009) Cycloalkane fragment (Gueneron et al., 2015) MBO fragment (Kim et al., 2010)	Cycloalkane fragment + secondary species fragment	
70.0651	C₄H₇NH⁺	Butane nitrile (Brilli et al., 2014)	Pyrroline (dihydropyrrrole)	0.21
71.0491	C₄H₆OH⁺	Biogenic MVK/methacrolein (Blake et al., 2009)	MVK/methacrolein /dihydrofuran	0.54
71.0855	C ₅ H ₁₁ ⁺	General alkane/VOC fragment (Yuan et al., 2014;Gueneron et al., 2015)	General alkane/VOC fragment	
73.0648	C ₄ H ₈ OH ⁺	2-butanone (MEK) (de Gouw and Warneke, 2007;Blake et al., 2009)	2-butanone (MEK) (negligible contribution from butanals)	1.96
75.0441	C ₃ H ₆ O ₂ H ⁺	Propionic acid (Ngwabie et al., 2008;Feilberg et al., 2015) Hydroxyacetone (Kim et al., 2010)	Propionic acid	0.48
77.0233	C₂H₄O₃H⁺	PAN (Hansel and Wisthaler, 2000)	Unknown species + PAN	
79.0542	C ₆ H ₆ H ⁺	Benzene (de Gouw and Warneke, 2007;Blake et al., 2009)	Benzene	4.54
81.0699	C₆H₈H⁺	Monoterpene fragment (Kim et al., 2010) PAH fragment (Gueneron et al., 2015)	Cyclopentyl aldehyde fragment (tentative)	



83.0855	C₆H₁₁⁺	Methylcyclopentane (Yuan et al., 2014;Gueneron et al., 2015)	Methylcyclopentane + secondary species fragment	
85.0648	C₅H₈OH⁺		Cyclopentanone (tentative)	0.11
87.0441	C ₄ H ₆ O ₂ H ⁺	Aromatic oxidation product (Müller et al., 2012) 2-3,butadione (Stockwell et al., 2015)	Aromatic oxidation product	
87.0804	C ₅ H ₁₀ OH ⁺	C5 carbonyls (Fall et al., 2001) MBO (Kim et al., 2010;Fall et al., 2001)	C5 carbonyls	0.34
93.0699	C ₇ H ₈ H ⁺	Toluene (de Gouw and Warneke, 2007;Blake et al., 2009)	Toluene	1.69
95.0855	C ₇ H ₁₀ H ⁺	Terpene fragment (Kim et al., 2009;Kim et al., 2010)	C7 cycloaldehyde fragment (tentative)	
97.1012	C₇H₁₃⁺	C7 cycloalkanes (Yuan et al., 2014;Warneke et al., 2015;Gueneron et al., 2015)	Methylcyclohexane + secondary species fragment	
99.0804	C₆H₁₀OH⁺	Hexenal (Fall et al., 2001;Ruuskanen et al., 2011;Park et al., 2013)	C6 cycloalkane oxidation products (tentative)	0.12
101.0597	C ₅ H ₈ O ₂ H ⁺	Aromatic oxidation product (Müller et al., 2012)	Aromatic oxidation product	
101.0961	C ₆ H ₁₂ OH ⁺	Hexanal (Rinne et al., 2005;Brilli et al., 2014)	C6 carbonyls	0.08
105.0699	C ₈ H ₈ H ⁺	Styrene (Kuster et al., 2004)	Styrene	0.03
107.0491	C ₇ H ₆ OH ⁺	Benzaldehyde (de Gouw et al., 2003)	Benzaldehyde	0.37
107.0855	C ₈ H ₁₀ H ⁺	C8 aromatics (de Gouw and Warneke, 2007;Blake et al., 2009)	C8 aromatics	0.65
111.1168	C₈H₁₅⁺	C8 cycloalkanes (Yuan et al., 2014;Warneke et al., 2015;Gueneron et al., 2015)	C8 cycloalkanes + secondary species fragment	
113.0961	C ₇ H ₁₂ OH ⁺	Heptenal (Brilli et al., 2014)	Cycloalkane oxidation product (tentative)	0.06
115.1117	C ₇ H ₁₄ OH ⁺		C7 carbonyls	0.03
121.1012	C ₉ H ₁₂ H ⁺	C9 aromatics (de Gouw and Warneke, 2007;Blake et al., 2009)	C9 aromatics	0.25
125.1325	C₉H₁₇⁺	C9 cycloalkanes (Yuan et al., 2014;Warneke et al., 2015;Gueneron et al., 2015)	C9 cycloalkanes + secondary species fragment	
135.1168	C ₁₀ H ₁₄ H ⁺	C10 aromatics (de Gouw and Warneke, 2007;Blake et al., 2009)	C10 aromatics	0.08
137.1325	C₁₀H₁₆H⁺	Monoterpenes (de Gouw and Warneke, 2007;Blake et al., 2009)	Adamantane or mystery monoterpene	0.09
151.1481	C₁₁H₁₈H⁺		Methyl adamantane (tentative)	




Figure 1. (a) SONGNEX study regions and P3 flight tracks. (b) Oil and natural gas production over time in various regions. The vertical dashed gray line marks the time period of the SONGNEX measurements. Production data from US Energy Information Administration (2017), Colorado Oil and Gas Conservation Commission (2017), State of New Mexico Oil Conservation Division (2017), Railroad Commission of Texas (2017), Wyoming Oil and Gas Conservation Commission (2017), and State of Utah Division of Oil Gas and Mining (2017).




Figure 2. (a) April 23 flight track; wind direction; locations of oil and gas wells, NEI 2011 point sources, and the Midland/Odessa urban area. NEI point sources are sized by total reported VOC emission, ranging in the area shown from 0-400 ton/year. (b) Flight track colored and sized by toluene (0-1.7 ppbv). (c) Flight track colored and sized by acetaldehyde (0-3.6 ppbv). (d) Flight track colored and sized by H₂S (0-8 ppbv). (e) Flight track colored and sized by m/z 71.049 C₄H₆OH⁺ (0-86 normalized counts per second).




Figure 3. Mixing ratios observed within the boundary layer during SONGNEX flights. (a) Observed mixing ratios of aromatics (note log scale). (b) Mixing ratios of acetone. Tropospheric



background is marked by dashed line (Singh et al., 1995). (c) Mixing ratios of H₂S. Note split axes to show highest mixing ratios. Estimated limits of detection for 1 second measurement (relevant to maximum mixing ratio) and 5-hour average (relevant to basin average) are marked as dashed lines. (d) Mixing ratios of methanol. Note split axes. Estimates of tropospheric background are within the shaded gray box (Heikes et al., 2002). (e) Average mixing ratios of methane. The shaded box marks typical concentrations measured upwind of oil and gas basins, from Peischl et al. (2015); Pétron et al. (2014). (f) Average mixing ratios of $\alpha+\beta$ pinene from iWAS/GC-MS. The SENEX and CalNex data are sum of monoterpenes from PTR-qMS. (g) Average wind speeds in the boundary layer. (h) Average mixing ratios of NO_x.




Figure 4. Average mixing ratios of oxygenates relative to acetone. Formaldehyde is from LIF instrument and formic acid from HNO₃/HCOOH CIMS. (For CalNex and SENEX instrumentation information, see <https://www.esrl.noaa.gov/csd/field.html>).




Figure 5. Overview of H_3O^+ ToF-CIMS mass spectra measured during the April 23 Permian flight. The categories “HC”, “HCO₁”, “HCO₂”, etc. mean hydrocarbon species (without S or N) with no oxygen, with one oxygen atom, with two oxygen atoms, etc. Top: average mass spectrum, colored by elemental composition. Large pie chart: overall VOC composition. Color code is the same as mass spectrum. Small pie charts: composition of oxygenated (top), HC-only (middle), and N- and S- containing species (bottom). For all pie charts, the composition in terms of both relative instrument signal (ncps, left column) and mixing ratio (ppb, right column) is shown. See text for



a description of which species were included in the mixing ratio charts, and how concentrations were calculated.




Figure 6. Comparison of H_3O^+ ToF-CIMS measurement of VOCs over the Permian Basin to VOCs evaporated from West Texas Crude oil. The values reported are the average boundary-layer enhancement in signal (ncps) during the SONGNEX April 23 flight, and the signal (ncps) relative to the most abundant mass (m/z 57.070 C_4H_9^+) in the West Texas Crude oil headspace.




Figure 7. H_3O^+ ToF-CIMS VOC ion mass correlation with toluene, representative of less aged emissions, and with acetaldehyde, representative of more aged emissions. Toluene and acetaldehyde were selected for this analysis because they have high signal-to-noise ratio and unambiguous VOC ion interpretation.




Figure 8. Comparison of C6-C10 aromatics distribution. Y-axis (molar ratio to benzene) is the same in each panel. (a) Measurements by H_3O^+ ToF-CIMS: SONGNEX flights and West Texas crude oil headspace. (b) Aromatic profiles from published studies of oil and gas fields, condensate tanks, oil tanks, and water storage tanks. (c) Aromatics profiles from published studies of urban areas and vehicle exhaust. (d) Average profiles of all oil and gas sources (panels a and b) and urban sources (panel c).




Figure 9. Comparison between PTR-ToF-MS and GC measurements of methylcyclohexane. A. SONGNEX 2015. B. UBWOS 2013.




Figure 10. Time series of hydrocarbon masses. (a) m/z 69 $C_5H_9^+$, converted to ppb using isoprene sensitivity. (b) m/z 137 $C_{10}H_{16}H^+$, converted to ppb using monoterpenes sensitivity. (c) m/z 151 $C_{11}H_{18}H^+$, in ncps. The structures of adamantane and a methyl adamantane isomer (both isomers exist in oil) are included in panels (b) and (c), respectively.




Figure 11. Ion masses related to cycloalkane oxidation products. The x-axis is the number of carbon in the molecule. (a) $C_xH_{2x-2}O$ oxygenates measured in the Permian basin during SONGNEX. (b) Cycloalkane precursors (measured by GC) and $C_xH_{2x-2}O$ oxygenates measured in the Uintah basin during UBWOS. Panels a and b show the same ion masses. (c) Possible dehydration masses ($C_xH_{2x-4}O$) of the oxygenates shown in (a), measured in the Permian basin. (d) Possible dehydration masses of the oxygenates shown in (b), measured in the Uintah basin.




Figure 12. Relationship between m/z 77.023 $C_2H_4O_3H^+$ measured by H_3O^+ ToF-CIMS and PAN.




Figure 13. Time series of the homologous series of nitrogen-containing masses $C_nH_{2n-1}NH^+$ from $n=2$ to $n=6$ during the Permian basin flights on April 23, April 6, and April 9. Toluene and pyrrole are also shown. The y-axis is in units of normalized counts per second (instrument signal) for all ion masses (including toluene, $C_7H_8H^+$). Ten-second box averages of all species are shown for clarity. Data collected during vertical profiles are included in these time series, to show low



concentrations outside of the boundary layer. Right: some possible isomers for each mass. An example of a cyclic structure is given in the left column, and an example of a non-cyclic structure in the right column.




Figure 14. Structure of porphyrin with pyrrolic and pyrroline subunits highlighted.




Figure 15. Scatterplot and time-series H_3O^+ comparison between H_3O^+ ToF-CIMS and Picarro CaRDS instruments. (a) Scatterplot of H_3O^+ ToF-CIMS vs Picarro H_2S for April 23 flight (Permian). Data points when the aircraft was ascending (red) and descending (blue) are highlighted. (b) Scatterplot of H_3O^+ ToF-CIMS vs Picarro H_2S for April 4 flight (Haynesville). (c) Time series during the April 23 flight. The H_3O^+ ToF-CIMS measurement has been averaged to a 6 second time basis. The Picarro H_2S is offset by 2 ppb (the intercept in panel a). The 1-s and 6-s detection limits for the H_3O^+ ToF-CIMS are shown by dashed red lines.