

Response to Referee #1

Lars Norin, Abhay Devasthale, and Tristan L'Ecuyer

July 11, 2017

We thank the referee for the time and effort devoted to review this manuscript as well as for the very constructive comments and suggestions. Below, please find a point-by-point reply to the comments (reproduced in *italics*).

General comments to the authors:

1. This seems like a precipitation science paper, but has been submitted to a journal that generally deals with advances to methods of atmospheric measurement. If you can make a few additions that steer the paper slightly in that direction — for example by describing how your methodology could (or could not) be applied in analysis of other snowy regions — that would make it a better fit for the journal.

We agree that the manuscript is more scientific in nature. But one of the main reasons for submitting in AMTD was to demonstrate the usefulness of long-term radar observations in Sweden and how the synergy with the latest space-based observations (specifically CloudSat) can be exploited to study snowfall. We believe such study can also be done for the other regions that have long-term radar observations.

2. In a few parts of the paper, the effects of the frequency/intensity of snowfall in a given weather state on one hand, and the frequency of that weather state on the other, are mixed. Maybe you could add a table that shows the relative frequencies of the different weather states; that would help decouple those effects.

We thanks for the suggestion. A table listing the number of days containing the different weather states (as well as how many other weather states occurred for those days) has been added to the revised manuscript.

3. Related to point 2, the statistical analysis is somewhat muddled by the fact that the occurrences of the different weather states are likely not independent. The prevailing winds, the NAO index and the occurrence of high/low pressure systems are probably correlated with each other at least to some degree. This should also be quantified, and if there are strong correlations, the impact of those on the results should be discussed.

As mentioned above, a table listing dates with simultaneous weather states has been added. We agree that there are correlations between some of the different weather states. However, within the different groups of weather states (wind directions, high and low MSLP, and different NAO indices) there are no overlapping dates. Nevertheless, the stronger correlations are pointed out in the revised manuscript.

4. *The analysis relies on the “prevailing” wind and pressure conditions, but Sweden is a big country and there can be quite some variability in those conditions between the southern and northern extremes. For example, Figs. 1b and 1d show that there can be about 90 deg difference in the 850 hPa wind direction between the north and the south. In fact, in “northeasterly” wind conditions in Fig. 1b, the prevailing winds in the far north appear to be from the northwest! And if one follows the streamlines in that figure, it appears that the flow to most of Sweden appears to originate from the Norwegian Sea in the (north)west rather than from the Barents Sea in the northeast. I wonder how much effect this has on the results, especially as the authors discuss differences between northern and southern Sweden in several places in the paper.*

The weather states were calculated in a paper by Thomas and Devasthale (Atmos. Chem. Phys., 14, 11545–11555, 2014). In their paper they evaluate the weather states using measurements from an area positioned over southern Sweden. This is the reason why certain wind directions can appear differently in the northernmost part of the country. The description of their methodology has been expanded in the manuscript. The area in which the weather states were calculated has been added to figures 1 and 2.

5. *Since there are already studies that investigated the connections of rainfall patterns to weather states, it would be reasonable for the authors to devote a paragraph or two to discussing the similarities and differences that can be found between the responses of rain and snow events to large-scale weather states.*

Following the referee suggestion, a paragraph discussing difference in circulation regimes during summer and winter seasons is added in the revised manuscript. Perhaps the most comprehensive evaluation of monthly circulation types over Sweden is done by Chen (2000) and Busuioc et al. (2001). Chen (2000) has shown that while the large-scale circulation patterns over North East Atlantic favouring westerly and northwesterly winds dominate 35–50% of the time during the summer months (June through August), the wintertime circulation patterns are characterized by nearly equal dominance of cyclonic and anticyclonic conditions. Similarly, Busuioc et al. (2001) showed that during winter months (especially beginning of year), NAO explained 41% of MSLP variability followed by the dipole structure with centres located over eastern Scandinavia and East Atlantic explained about 24% variability in January and the third mode represented cyclonic/anticyclonic conditions. However, during the summer rainfall months, the cyclonic/anticyclonic conditions dominated circulation types representing about 37% variability.

Specific comments:

Page 2:

Line 1: “Selected” previous studies? Should this say “several” instead?

Done.

Page 5:

Lines 7-8: Did you investigate if there were systematic differences in surface snowfall between 2CSNOW and the ground-based radar?

Yes, a comparison of the surface snowfall from 2CSNOW and Nordrad was investigated and described in the paper by Norin et al. (Atmos. Meas. Tech., 8, 5009–5021, 2015). A short paragraph has been added to further describe the results of that study.

Line 14: I think OCO-2 is currently the A-Train lead satellite.

True. This is now corrected in the manuscript.

*Lines 27-28: This should probably say “more than one standard deviation *above/below the mean*”*

Done.

Page 6:

Line 22: I think that the average is only for cases where it is snowing, i.e. zero snowfall is not included in the average? Or am I mistaken? In any case, this should be clarified.

Correct. This is now clarified.

Page 7:

Lines 23-25: “However, snowfall intensities are...” This is a confusing sentence, I am not sure what you are trying to say.

The sentence has been rewritten.

Line 25: Temperature anomalies relative to what?

The temperatures anomalies are relative to the average temperature for the investigated years. This is now clarified in the manuscript.

Page 8:

Line 3: See my previous comment: what is the reference for calculating specific humidity anomalies?

The specific humidity anomalies are relative to the average specific humidity for the investigated years. This is now clarified in the manuscript.

Page 9:

Line 16: Comparing the mean lines Figs. 9a and 9b, it is rather hard to see any difference between them near the surface.

Agreed. The sentence has been rewritten.

Page 10:

Line 10: “Relative strengths of snowfall” — again, relative to what?

Here relative snowfall means that the sum of all four snowfall rates are equal to one. A sentence has been added to clarify this.

Figs. 1 and 2: Since the wind speed is already encoded in the arrows, the using the color to also show that is somewhat redundant. On the other hand, the pressure patterns might be interesting. How about using color to show the pressure instead?

We thank the referee for the suggestion. Although it would likely be an improvement it is unfortunately not possible for us within this project.

Response to Referee #2

Lars Norin, Abhay Devasthale, and Tristan L'Ecuyer

June 26, 2017

We thank the referee for the time and effort spent to review the manuscript as well as for the very constructive comments and suggestions. Below, please find a point-by-point reply to the comments (reproduced in italics).

General comments

1. Previous research has identified the 8 states the paper analyzes. While investigating snowfall relative to these states is appropriate, I think the authors should note that these states are not independent from one another. I would like to know how many events comprise each category, and if individual dates can be included in multiple categories.

A table listing the number of days containing the different weather states (as well as how many other other weather states occurred for those days) has been added to the manuscript.

2. Are there no snowfall observations over Sweden that can be used to further validate the remotely sensed data? At times the authors refer to “snowfall accumulation”. As I understand the analysis performed, no statements can be made about snowfall accumulation, as the duration of the snowfall events does not appear to be considered.

We agree that no quantitative statements can be made about snowfall accumulation. To find a qualitative estimation of the relative contribution of the different weather states to the total snowfall amount, the average snowfall intensities were multiplied with the number of days for each weather state. The text in Section 4.3 has been adjusted in the revised manuscript.

3. Figures 1 and 2 — are these composites (means) of all events? Or individual events that are representative of the composites?

These are individual events, representative of the composites. This has now been clarified in the manuscript.

4. On the use of CloudSat — As you discuss, data represents a “snapshot” taken once or twice per day. Is snowfall occurring during all of these snapshots? How many snapshots are there per event? I am concerned about how representative these snapshots are for the events analyzed using the other more frequent data sources, as comparisons are made between them.

Yes, only passes during which CloudSat detected snowfall over Sweden were used in the analysis. The number of data points with identified snowfall per pass varied from 1 up to 1007. It is true that measurements from the ground-based radars are more frequent as well as made over spatially smaller areas. However, only CloudSat can provide a detailed vertical structure of the snowfall. The impact of CloudSat's lower temporal and spatial resolution is now discussed in the manuscript.

5. I would like to know more about how the wind directions were defined. The NAO is straightforward and I can see how identifying the presence of a pressure center over land is tractable. But wind direction can be highly variable over the domains shown in the figures.

The method for identifying the weather states was presented in a paper by Thomas and Devasthale (Atmos. Chem. Phys., 14, 11545–11555, 2014). The wind directions were determined by analysing the average wind direction over an area near the centre of Sweden. For this reason wind direction can be different when far from this area (e.g. in the northernmost part of the country). The description of this method has now been expanded in the manuscript. The area in which the weather states were calculated has been added to figures 1 and 2.

6. I think it would be useful to the reader to see figures that show the radar coverages referred to in the text (48–82 km from a Swedish radar station), so we can have an idea of how representative the data is with respect to the entire country.

Thanks for the suggestion. The radar coverages used in the analysis have now been added to figure 3 and 10.

7. Related to the above comment, were the intensities for each event defined by averages over each radar volume? How many times were used in each event?

Yes, for each radar composite the average snowfall intensity was determined using all radar cells detecting snowfall. All snowfall measurements detected during each day (24 h) with an identified weather state were used.

8. I see that Norin et al. (2015) describes how the snowfall product is generated. Given the fact that your conclusions are heavily dependent upon the quality of the data that results from the methods used to define the product, I think it would be worthwhile to further discuss the limitations of the radar-based data.

A paragraph discussing the limitations of the radar-based snowfall product has been added to the revised manuscript.

9. Section 4.2, line 15. I see what you are referring to in Figure 9a and b, but I am not sure how this resolves the apparent paradox discussed in the previous paragraph. Perhaps you could be more explicit.

Fig. 2a shows that during High MSLP conditions, anticyclonic winds coming from Northern Europe further pick up moisture from the North Sea before making precipitation over western coast of Norway and southern parts of Sweden. This leads to increase in specific humidity over these regions (Figs. 7a and 8a). These snowfall

events, whenever they do occur, also seem to be well vertically developed as evident in Fig. 9a. However, during low MSLP conditions, the location of low pressure (Fig. 2b) is such that more northerly, colder winds result in lower specific humidity anomalies (Fig. 8b). Notice however that over the southernmost tip of Sweden, humidity anomalies are slightly positive during low MSLP conditions due to warmer southwesterly winds. Since the centre of low pressure is slightly northward (compared to the centre of high pressure), the occurrence of snowfall events with differing intensities, over northern and southern regions of Sweden leads to more broader vertical distribution of snowfall as observed by CloudSat (Fig. 9b).

Minor comments

1. In line 5, it is stated that the NAO correlates strongly with precipitation, but explains only 32 to 54% of precipitation variability locally (Busuioc et al. 2001). In line 17, Linderson (2001) further observes a robust correspondence between local precipitation and large-scale circulation during winter months over southern Sweden. As written, it seems these results are contradictory. If this is the case, it should be highlighted as motivation for this study. If this isn't the case, perhaps the text could more clearly explain the differences in the studies.

We agree that the current formulation of these sentences is confusing. They are therefore clarified in the revised manuscript. During winter months, NAO can explain a significant part of precipitation variability. But in a year as a whole, the other circulation types, such as cyclonic systems, persistent westerly winds and even anticyclonic systems, can be more important, especially over the southern parts of Sweden.

2. Line 22 — Such a characterization also provides guidance concerning the performance of NWP models by quantifying their strengths and limitations in wintertime regimes — could you state more clearly how your work relates to NWP verification?

This rather abstract sentence is also rephrased in the revised manuscript. Our intention was to point out the fact that NWP models need to show similar response of snow to weather states as we have presented in this study, in order to capture snowfall events realistically. Therefore, the observationally based results from our study can be used for process oriented evaluation of NWP models to test the fidelity of NWP models.

3. Line 30 — snowfall distribution and frequency is very inhomogeneous meridionally across Sweden. How does it vary?

The snowfall distribution across Sweden is shown in Fig. 1. The southern parts of Sweden receives the smallest amounts of snow. The snow depth increases steadily with increasing latitudes but the largest snow depths are found in the mountains to the west, along the border to Norway. This description has been added to the manuscript.




Figure 1: Average largest snow depth during the period 1961–1990.

Response to Referee #3

Lars Norin, Abhay Devasthale, and Tristan L'Ecuyer

June 24, 2017

We thank the referee for the time and effort dedicated to review the manuscript as well as for the very constructive comments and suggestions. Below, please find a point-by-point reply to the comments (reproduced in italics).

1. Obtaining reliable quantitative estimates of snow rates and accumulation from radar is challenging. Have the snow rates and accumulation been evaluated?

We agree that obtaining reliable quantitative estimates of snow rates and accumulation from radar is challenging. As a first step toward improving the ground-based radar snowfall measurements a comparison of snow rates from ground-based radar and CloudSat was examined in a paper by Norin et al. (Atmos. Meas. Tech., 8, 5009–5021, 2015). A paragraph discussing the results of that paper in more detail has been added to the manuscript.

2. It would be valuable to systematically break down snow accumulation into frequency of occurrence and snow rate, and frequency of weather state.

A table listing the number of days containing the different weather states (as well as how many other weather states occurred for those days) has been added to the manuscript.

3. Partitioning rain and snow based on 2m air temperatures is subject to uncertainties (Harpold et al. 2017), specifically in complex terrain. Other variables such as atmospheric humidity could be used to estimate the phase of precipitation. Can the authors elaborate on this aspect?

We agree that it is difficult to determine the phase of precipitation, especially when 2-m air temperatures are close to 0° C. However, for this paper we have chosen to classify precipitation as snow for temperatures below freezing and as rain otherwise. A short paragraph has been added to the revised manuscript mentioning the difficulties in obtaining accurate snowfall products based on radar data and surface air temperatures.

The sensitivity of snowfall to weather states over Sweden

Lars Norin¹, Abhay Devasthale¹, and Tristan S. L’Ecuyer²

¹Atmospheric Remote Sensing Unit, Research Department, Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

²Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

Correspondence to: Lars Norin (lars.norin@smhi.se)

Abstract. For a high latitude country like Sweden snowfall is an important contributor to the regional water cycle. Furthermore, snowfall impacts surface properties, affects atmospheric thermodynamics, has implications for traffic and logistics management, disaster preparedness, and also impacts climate through changes in surface albedo and turbulent heat fluxes. For Sweden it has been shown that large-scale atmospheric circulation patterns, or weather states, are important for precipitation variability.

5 Although the link between atmospheric circulation patterns and precipitation has been investigated for rainfall there are no studies focused on the sensitivity of snowfall to weather states over Sweden.

In this work we investigate the response of snowfall to eight selected weather states. These weather states consist of four dominant wind directions together with cyclonic and anti-cyclonic circulation patterns and enhanced positive and negative phases of the North Atlantic oscillation. The presented analysis is based on multiple data sources, such as ground-based
10 radar measurements, satellite observations, spatially-interpolated in situ observations, and reanalysis data. The data from these sources converge to underline the sensitivity of falling snow over Sweden to the different weather states.

In this paper we examine both average snowfall intensities and snowfall accumulations associated with the different weather states. It is shown that even though the heaviest snowfall intensities occur during conditions with winds from the southwest, the largest contribution to snowfall accumulation arrives from winds from the southeast. Large differences in snowfall due
15 to variations in the North Atlantic oscillation are shown as well as a strong effect of cyclonic and anti-cyclonic circulation patterns. Satellite observations are used to reveal the vertical structures of snowfall during the different weather states.

1 Introduction

At high latitudes, solid precipitation is an important contributor to the regional water cycle (Levizzani et al., 2011; Waliser et al., 2011), it impacts surface properties, affects atmospheric thermodynamics, has implications for traffic and logistics management, disaster preparedness, and also impacts climate through changes in surface albedo and turbulent heat fluxes. In a country like Sweden with strong north-south gradients in meteorological, topographical, and surface conditions, understanding the response of snowfall to large-scale weather states is even more important. Such a characterisation also provides guidance concerning the performance of for numerical weather prediction models by quantifying their strengths and limitations in wintertime regimes, which should show a similar response of snow to large-scale weather states, in order to capture snowfall events realistically.
25

Precipitation variability over Sweden has been investigated in ~~seleected~~several previous studies (Busuioc et al., 2001; Johansson and Chen, 2003; Hellström, 2005; Gustafsson et al., 2010; Devasthale and Norin, 2014). Since the local circulation patterns (here termed “weather states”) often decide the precipitation response, it is important to study dominant weather states and their link to precipitation. The dominant weather states responsible for rainfall and snowfall can be quite different over different
5 regions of Sweden. Furthermore, the dominant weather state may not always explain a significant part of the precipitation variability. Perhaps the most comprehensive evaluation of monthly circulation types over Sweden is done by Chen (2000), Linderson (2001), and Busuioc et al. (2001). Among 25 different circulation patterns studied by Chen (2000) over Sweden, four weather states are shown to generally dominate. These are characterised by either cyclonic or anti-cyclonic conditions or persistent large-scale westerly or southwesterly wind regimes. While the large-scale circulation patterns over North East
10 Atlantic favouring westerly and northwesterly winds dominate 35–50 % of the time during the summer months (June through August), the wintertime circulation patterns are often characterized by nearly equal dominance of cyclonic and anticyclonic conditions.

Using data from 33 ground-based stations between the years ~~1980–1990~~1980—1990, Busuioc et al. (2001) found that the link between precipitation variability and the dominant circulation patterns can be very complex over Sweden. For example,
15 while the North Atlantic oscillation (NAO) is the principle mode of sea-level pressure variability during cold months and correlates strongly (0.86) with precipitation, it explained only 32 % to 54 % of precipitation variability locally. ~~Hellström (2005) pointed out the importance of cyclonic and anti-cyclonic circulations in regulating the summertime extreme precipitation events while Gustafsson et al. (2010) further argued for the crucial role of continental Europe and the Baltic Sea in controlling the moisture transport during extreme precipitation events. But the majority of these studies focus on liquid precipitation (rain) and the~~
20 ~~summer months. Far fewer studies directly address the factors that govern snowfall and its covariability with the other regional impacts of the different wintertime weather states that occur across Sweden and those that do lack sufficient detail to inform forecasts.~~

~~Snowfall over Sweden can occur under the influence of a range of circulations patterns (here termed “weather states”). Using Lamb classification (Lamb, 1950), Chen (2000) investigated the monthly circulation climatology over Sweden. He showed~~
25 ~~that among 25 different circulation patterns observed over Sweden, four states dominate. These can be characterised by either cyclonic or anti-cyclonic conditions or persistent large-scale westerly or southwesterly wind regimes. During winter months, westerly and southwesterly winds occur 21 % and 17 % respectively, followed by anti-cyclonic conditions that occur about 11 % of the time. Using a similar classification methodology, After NAO, the dipole structure with centres located over eastern Scandinavia and the East Atlantic explained about 24 % variability in January and the third mode represented~~
30 cyclonic/anticyclonic conditions. However, during the summer rainfall months, the cyclonic/anticyclonic conditions dominated circulation types representing about 37 % variability. Linderson (2001) further observed a robust correspondence between local precipitation and large-scale circulation during winter months over southern parts of Sweden. She showed that over southern Sweden anti-cyclonic conditions dominated (16 %) all circulation types.

The link between weather states and precipitation (especially rainfall) during extreme events can be different than that
35 observed during average conditions. For example, Hellström (2005) pointed out the importance of cyclonic and anti-cyclonic

circulations in regulating the summertime extreme precipitation events while Gustafsson et al. (2010) further argued for the crucial role of continental Europe and the Baltic Sea in controlling the moisture transport during extreme precipitation events. More recently, ~~Thomas and Devasthale (2014)~~ Thomas and Devasthale (2014) used this weather state classification to investigate temperature and humidity response and pollutant variability over Scandinavia. They showed that enhanced positive phase of the NAO and southeasterly winds warm the entire troposphere and the surface. These studies form the basis for the investigation and selection of weather states used in the present study. However, it is to be noted that the majority of the studies linking weather states to precipitation over Sweden focus mainly on rainfall and the summer months. Far fewer studies directly address the factors that govern snowfall and its covariability with the other regional impacts of the different wintertime weather states that occur across Sweden and those that do lack sufficient detail to inform forecasts.

The main aim of this work is to exploit the synergy between two remote sensing systems, ground-based weather radar and space-based cloud radar to further document the character of falling snow during each weather state and to quantify the influence of the key meteorological factors that modulate snowfall differences between the different weather states. We seek to answer two specific questions:

1. How does falling snow respond to different atmospheric weather states? The aim here is to investigate to what extent snowfall is sensitive to weather states (different wind directions, mean sea level pressures (MSLP), and phases of the NAO) and to characterise differences in this dependence between southern and northern parts of the country. The latter is especially important since the snowfall distribution and frequency is very inhomogeneous meridionally across Sweden, with average snow depths increasing with increasing latitude (Busuioc et al., 2001; Devasthale and Norin, 2014).
2. What are the typical meteorological conditions associated with each weather state and how do they contribute to the observed differences in snowfall between weather states? Here we quantify the extent to which similar co-variability between meteorological parameters and snowfall exists across all weather states.

2 Datasets and weather states

In this work we have investigated the response of snowfall to eight different weather states. These weather states consist of four dominant wind direction (northeast, northwest, southeast, and southwest) as well as conditions of high and low MSLP and enhanced positive or negative phases of the NAO. To study the response of snowfall to the different weather states we made use of a combination of spatially-interpolated in situ observations, reanalyses, ground-based radar measurements, and satellite observations. All dates satisfying any of these conditions were identified in reanalyses. Mean radar-derived surface snowfall intensities and meteorological analyses of surface air temperatures were then composited for these dates. Finally, vertical profiles of reflectivity from satellite observations were extracted to document the vertical structure and implied microphysical properties of snowfall in each weather state.

Below follows a description of all datasets used in this study together with a definition of the selected weather states.

2.1 ERA-Interim reanalysis

ERA-Interim provides global atmospheric reanalysis, updated continuously since 1979 (Dee et al., 2011). Gridded data products include a large variety of 3-hourly surface parameters, describing weather as well as ocean-wave and land-surface conditions, and 6-hourly upper-air parameters covering the troposphere and stratosphere. Vertical columns of monthly averages for many atmospheric parameters and other derived fields are also available. Berrisford et al. (2009) provide a detailed description of the ERA-Interim archive.

The data assimilation system used to produce ERA-Interim is based on a 2006 release of the European Centre for Medium-range Weather Forecasting (ECMWF) Integrated Forecast Model. The system includes a 4-dimensional variational analysis with a 12-hour analysis window. The spatial resolution of the data set is approximately 80 km on 60 vertical levels from the surface up to 0.1 hPa.

In this study u and v components of winds at 850 hPa were used to identify the dominant weather states. The wind fields have a spatial resolution of about 80 km and are available every 3 hours.

2.2 Mesan

To characterise the phase of precipitation 2 m air temperatures were obtained from Mesan (Häggmark et al., 1997, 2000), a mesoscale analysis system that has been operational at the Swedish Meteorological and Hydrological Institute (SMHI) since 1996. Mesan creates spatially continuous fields of selected meteorological parameters (such as precipitation, temperature, humidity, visibility, wind, and clouds) on a model grid using observations from synoptic stations, automatic stations, satellites, and weather radars. In addition to these observations, a first guess field from SMHI's operational numerical weather prediction model, HIRLAM (Källén, 1996), is used. The field values of the meteorological parameters are calculated in Mesan using the optimal interpolation technique (Daley, 1993). The Mesan 2 m temperatures are further adjusted for both the fraction of land in the first guess field's grid square and for the altitude of the temperature observations.

Mesan covers Scandinavia and the Baltic Sea and has a spatial resolution of $0.1^\circ \times 0.1^\circ$ (11 km \times 11 km). The temporal resolution is one hour. Archived data exist from 1998 to present and is continuously updated.

2.3 Nordrad

The meteorological offices in the Nordic countries (Danish Meteorological Institute, Estonian Environment Agency, Finnish Meteorological Institute, Latvian Environment, Geology and Meteorology Agency, Norwegian Meteorological Institute, and SMHI) collaborate by exchanging radar data in real time. The collaboration is called Nordrad (Carlsson, 1995) and consists today of 35 operational weather radars.

The SMHI uses data from all available weather radars within Nordrad to generate composite radar images covering the Nordic countries. These data are corrected for beam blockage (Bech et al., 2003). Non-meteorological echoes, originating for example from ground clutter or clear-air targets, are removed by a satellite-based filter (Michelson, 2006). Furthermore, in order to correct the radar measurements for distance-dependence, the radar data is adjusted to precipitation measurements from

rain gauges by fitting a second degree polynomial to the logarithmic gauge-to-radar ratio (Michelson and Koistinen, 2000). The conversion from reflectivity to precipitation rate R (mm h^{-1}) is made using the empirical relationship $Z = aR^b$ (e.g. Battan, 1973), where $a = 200$ and $b = 1.5$. For a more detailed description of the radar processing see, e.g., Michelson (2006); Norin et al. (2015).

5 The quality-adjusted radar data are merged into a precipitation composite, Nordrad, which is generated every 15 min and has a spatial resolution of $2 \text{ km} \times 2 \text{ km}$. Archived Nordrad composite images exist at the SMHI from 2005 to present. However, in late 2007 a hardware upgrade was installed at the Swedish weather radars to enable Doppler processing for all scans. In September 2014 a modernisation process began to upgrade all Swedish weather radars to dual polarisation. As a result, the present analysis is restricted to the homogeneous processing period from January 2008 through December 2012.

10 2.4 Snowfall product from the CloudSat Cloud Profiling Radar

CloudSat snowfall estimates were obtained from the Release 04 2C-SNOW-PROFILE (hereafter, 2CSNOW) data product (Wood, 2011; Wood et al., 2013). The product provides vertically-resolved snowfall estimates consistent with profiles of W-band (94 GHz) reflectivities measured by CloudSat’s Cloud Profiling Radar (CPR). CloudSat observes falling snow between 82° N and 82° S latitude along a ground track that repeats every 16 days providing moderately dense spatial sampling at high
15 latitudes relative to more equatorial locations (Kulie et al., 2016; McIlhattan and L’Ecuyer, 2016). Snowfall rates are produced at the full 1.7 km along-track by 1.4 km cross-track spatial resolution of the CPR. The radar has an intrinsic vertical resolution of 485 m, but measurements are oversampled to yield profiles at an effective vertical resolution of 239 m.

Rather than assuming a fixed relationship between reflectivity and snowfall rate (a so-called $Z-S$ relationship), the 2CSNOW retrieval algorithm estimates vertical profiles of the probability density functions of snow particle size distribution parameters.
20 Information collected during recent field campaigns are used to supply a priori estimates of the environmental distributions of these parameters as well as snow particle microphysical and scattering properties in the retrieval (Wood et al., 2015). Attenuation and multiple-scattering by snow particles are also modelled. A snowfall rate is generated whenever the CloudSat 2C-PRECIP-COLUMN product (Haynes et al., 2009) indicates the presence of frozen precipitation at the surface defined as a scene with a melted mass fraction of less than 10 %. The melted mass fraction is estimated based on the height of the freezing
25 level and a melting model that accounts for the environmental lapse rate below the freezing level derived from ECMWF analyses.

Ground clutter affects the CPR measurements in the radar range bins nearest the surface so these near-surface bins are not explicitly included in the reflectivity profiles when retrievals are performed, creating what is sometimes called a blind zone. Consequently 2CSNOW estimates the surface snowfall rate as the rate retrieved in the radar bin immediately above the blind
30 zone. Over land, this zone extends about 1 km above the surface (Smalley et al., 2014). This surface snowfall rate is assigned a confidence value from “None” to “High” depending on the expected performance of the forward model and the reliability of the temperature-based estimate of the precipitation phase, among other factors.

2.5 Humidity data from the Atmospheric Infrared Sounder

Humidity profiles from the Atmospheric Infrared Sounder (AIRS) instrument flying onboard NASA's Aqua satellite since 2002 are further used to investigate differences in the moisture advection during weather states (Chahine et al., 2006; Susskind et al., 2014; Devasthale et al., 2016). Aqua ~~leads-is part of~~ the A-Train constellation of satellites ~~wherein CloudSat follows closely~~
5 ~~behind the Aqua satellite-, and travels in front of CloudSat~~ in the same orbit. The retrievals provided in the standard daily gridded Version 6 product (AIRX3STD) at 850 hPa are analysed (Susskind et al., 2014). The estimated accuracy of humidity profiles is 15 % per 2 km. In this study, AIRS provides an independent information on the moisture variability during the different weather states.

2.6 Weather states

- 10 In the present study we have, following Thomas and Devasthale (2014), selected eight different weather states which frequently occur over the Nordic countries. The eight weather states are divided into two groups: wind regimes, consisting of conditions dominated by four wind directions (northwest, northeast, southeast and southwest), and synoptic states, during which either cyclonic or anti-cyclonic conditions or enhanced positive or negative phases of the NAO prevailed.

- To find days corresponding to the first four weather states the average daily wind speed and wind direction at 850 hPa from
15 the ERA-Interim reanalysis were extracted for an area centred over southern Sweden (55–60° N, 12–20° E). Based on these daily averages, days during which a particular wind direction prevailed and persisted for at least 3 days were selected in order to capture large-scale precipitation events. Cyclonic and anti-cyclonic conditions were found in a similar way by analysing the MSLP ~~over-central-in the above mentioned area over southern~~ Sweden. Anti-cyclonic conditions were identified when the MSLP was ~~higher-than-more than than~~ one standard deviation above the mean for 3 consecutive days. If the MSLP was
20 ~~lower-more~~ than one standard deviation below the mean, cyclonic conditions prevailed. The NAO indices were obtained from NOAA's Climate Prediction Center website (NOAA, 2016). Enhanced positive and negative phases of NAO correspond to absolute values of the NAO index exceeding one standard deviation during those phases. Enhanced positive or negative NAO conditions were defined as 3 consecutive days with an enhanced positive or negative NAO index.

- ~~Examples-of-mean-~~ Table 1 shows the number of days identified for the different weather states during the period January
25 2008 to December 2012. From the table it is clear that not all weather states are mutually exclusive. For example, it can be seen that more than half of the days characterised by southeasterly winds are also identified as days with with high MSLP whereas days with low MSLP to a large degree coincide with days with enhanced negative NAO.

- Representative examples of circulation patterns for the weather states characterised by dominant wind directions are shown in Fig. 1 whereas representative examples of circulation patterns for high and low MSLP together with enhanced positive
30 and negative phases of the NAO are displayed in Fig. 2. In both figures the area over which the wind speed and the MSLP were calculated is depicted by a white rectangle. Because the wind directions are evaluated over southern Sweden the wind directions can be different in the northern part of the country.

3 Data processing

The data sets from ERA-Interim and Mesan extend from 1979 and 1998 to present, respectively. Nordrad data are nominally available since 2005 but due to modifications to the Swedish weather radar network, a homogeneous Nordrad dataset can only be compiled from 2008 to 2014. Data from CloudSat are available from 2006 to present. Thus in this study we have restricted the study period to 1 January 2008 to 31 December 2012 when all required datasets are available.

Snowfall products can be generated by combining ground-based radar measurements with near-surface air temperature measurements. While obtaining an accurate and reliable snowfall product is difficult (Harpold et al., 2017), a simple but robust snowfall product is sufficient for the purpose of this study.

As described in Norin et al. (2015) a simple but robust radar-based snowfall product can be generated by combining Nordrad's precipitation composite with 2 m temperatures from Mesan. For every Nordrad composite image the 2 m temperature field from Mesan nearest in time and space was used to identify frozen precipitation. If the 2 m temperature was below freezing (0°C) the corresponding precipitation from Nordrad was classified as snow, otherwise as rain. In this work we have used snowfall estimates from the Nordrad composites but in order to provide a uniformly calibrated data set as well as to have full knowledge of the measurements we have only selected radar data originating from areas covered by Swedish radars.

~~Measurements from ground-based radars generally have lower quality far away from the radar station.~~ Comparing ground-based radar measurements of snowfall intensities to CloudSat snowfall measurements, Norin et al. (2015) found that Nordrad snowfall estimates are optimal at distances between 46–82 km from a Swedish radar station. Outside this range the disagreements were larger due to a combination of different reasons. The sensitivity of the ground-based radars decreases for increasing distances making it more difficult to detect light snowfall at large distances. At large distances Nordrad also systematically overshoots low-level precipitating systems. This leads to an underestimation of Nordrad's measured snowfall rates and occasionally to Nordrad missing precipitation altogether. Close to the radar stations Nordrad detects non-precipitating, low-intensity echoes that can be misclassified as light snowfall.

Snowfall intensities from Nordrad were therefore selected within ~~this~~ the optimal distance range from the 12 Swedish weather radar stations together with the corresponding 2 m temperatures from Mesan, for each of the 8 identified weather states. The average number of radar cells within the optimal measurement zone for any radar is approximately 3620 while Mesan has approximately 120 data gridboxes covering the same area.

Average ~~snowfall intensity and the~~ intensity of falling snow and the corresponding average 2 m temperature were calculated for measurements containing precipitation for all days assigned to each weather state. Corresponding temperature anomalies were defined relative to the daily average 2 m temperatures from Mesan for all regions within the optimum distance over the period from 2008–2012.

~~Each day CloudSat passes over Sweden twice and~~ CloudSat samples data every 1.4 km and passes over Sweden twice each day. The CloudSat satellite has been operational since 2006 but in April 2011, the satellite experienced a battery anomaly that caused a ten month gap in observations. Measurements resumed in February 2012 but the satellite has only operated during the daytime portion of its orbit since that time. In this study, all available ascending and descending passes of CloudSat

from 1 January 2006 to 31 December 2012 in the latitude band between 54° N and 70° N were used (excluding the summer months, June to September, in which no snowfall was observed). To ensure high quality, only CloudSat snowfall retrievals with a confidence flag corresponding to “Moderate” or “High” were used in the analysis yielding between 0 and 1007 observed snowfall profiles per day. CloudSat vertical snowfall profiles together with the estimated surface height over Sweden were
5 composited for each weather state to characterise the vertical structure of snowfall in each regime. Due to the lower temporal resolution of CloudSat, average vertical profiles were calculated using data from the entire country to obtain sufficient amounts of data for the different weather states. Snowfall intensities below 1250 m above the surface were excluded to avoid CloudSat’s blind zone.

The humidity profiles at 850 hPa available at the $1^\circ \times 1^\circ$ resolution from the AIRX3STD were analysed in the same way
10 as the surface temperature data from MESAN, i.e. by taking into account the only those cases where surface temperatures (also derived from AIRS for the sake of consistency) were colder than 0° C. The data from the ascending passes of AIRS were analysed.

4 Results and discussion

The response of snowfall to the eight selected weather states were examined using the snowfall products from Nordrad and
15 CloudSat, together with temperature observations from Mesan and measurements of specific humidity from AIRS. Below, the results for the two different groups of weather states are presented and the implications for snowfall accumulations are discussed.

4.1 Snowfall during dominant wind regimes

Average snowfall intensity for the weather states characterised by the four dominant wind directions (northwest, northeast, southwest, and southeast, as illustrated in Fig. 1) from all of Sweden’s 12 weather radars are shown in Fig. 3a. In general
20 the average snowfall intensities are seen to decrease with increasing latitude. The smallest average snowfall intensities, below 0.2 mm h^{-1} , are seen to occur for winds from the southeast at northern radar stations (above 60° N). It is also found that the largest average snowfall intensities occur when winds originate from the southwest. This is valid for all radar stations, except for one radar located in the northwestern region which is slightly more influenced by northwesterly frontal systems arriving
25 from the Norwegian Sea. For the southernmost radar stations (below 59° N), where this effect is most evident, average snowfall intensities are near 0.5 mm h^{-1} .

The corresponding 2 m temperatures, shown in Fig. 3c, reveal that the most intense snowfall occurs at warmer temperatures, ranging from approximately -1° C to -4° C , whereas the lightest snowfall intensities correspond to colder temperatures, between -6° C and -11° C . For northerly winds the snowfall intensities are similar, around $0.15\text{--}0.25 \text{ mm h}^{-1}$. However,
30 snowfall intensities are slightly larger for winds from the northwest for ~~most radar stations when~~ the southernmost radar stations (below 59° N). The corresponding 2 m temperatures are on average $1\text{--}3^\circ \text{ C}$ warmer compared to the northeasterly wind regime. Figure 3e shows that temperature anomalies (relative the daily average temperature, see Sect. 3) during snowfall

are, in general, largest for the southernmost stations (below 59°N), for all wind directions. However, the largest temperature anomalies for all station occur for winds from the northeast, ranging from -2°C to -7°C while the smallest temperature anomalies (0°C to -3°C) occur for winds from the southeast.

Average values of specific humidity, observed by AIRS, are shown in Fig. 4. In agreement with the snowfall observations it is seen that winds from the southwest advect sufficient moisture over all of Sweden which in combination with warm temperatures lead to large snowfall intensities. During winds from the southeast, moist air originating from continental Europe passes over the southern and central parts of Sweden. Together with warm temperatures this leads to increased snowfall intensities. On the other hand, this moist air does not reach the northern parts of the country and this lack of moisture in combination with cold temperatures results in small snowfall intensities. For northerly winds moist air from the Atlantic is blocked by the mountains on the border to Norway and the result is moderate snowfall over Sweden. Anomalies of specific humidity ([relative the daily average specific humidity, see Sect. 3](#)) are presented in Fig. 5. The anomalies show that unusually moist air covers Sweden during southwesterly winds while the opposite occurs during winds from the north. During winds from the southeast, southern Sweden receives moist air while the air in the northern part of the country is drier than usual.

Figure 6 shows the vertical structure of snowfall, as observed by CloudSat, for the four weather states characterised by dominant wind directions. Mean snowfall rates as a function of altitude are shown together with 2D histograms of snowfall intensities and altitudes. In addition, the altitudes at which 50 % and 95 % of the snowfall rates were observed are displayed. It is seen that the largest mean near-surface snowfall rates, above 0.2 mm h^{-1} , occur during westerly wind regimes. These snow events are characterised by deeper snow columns and stronger vertical intensity gradients that may be indicative of enhanced aggregation processes relative to easterly snow events. The smallest mean near-surface snowfall intensity, below 0.1 mm h^{-1} , occurs during winds from the southeast, consistent with the observations from the ground-based radars. It is also interesting to note that the mean snowfall intensity (approximately 0.1 mm h^{-1}) from the ground up to 5 km above the surface in northeasterly regimes indicating an absence of aggregation in the prevailing colder, drier conditions. Near the surface, the distribution of snowfall intensities peaks around 0.5 mm h^{-1} when winds are from the southwest. For winds from the east, the distribution of snowfall rates peaks at a much lower intensity, below 0.01 mm h^{-1} . [Since the histograms of the vertical snowfall structure were obtained using data from the entire country \(see Sect. 3\) the spatial information in Fig. 6 has been lost. However, when we compared vertical profiles from the northern and the southern parts of the country, no noticeable difference in the vertical structure could be found.](#)

The different response of the snowfall intensities to these weather states can be understood by examining the circulation patterns in Fig. 1. Winds from the southwest originate from the Atlantic and the North Sea bringing moist, warm air over Sweden resulting in heavy snowfall once the temperature falls and sustains below 0°C . Winds from the northwest also originate from the Atlantic but pass over the mountainous border to Norway before arriving in Sweden, thereby bringing less snow at slightly colder temperatures compared to southwesterly winds. These northwesterly winds have the second largest influence on the central Swedish regions (after southwesterly winds). Winds from the northeast, on the other hand, arrive from the Norwegian/Barent's Sea and bring relatively colder air and less moisture, have the second largest impact (after southwesterly winds) on the northernmost Swedish regions. Finally, southeasterly winds originating from continental Europe pick up moisture

from the southern Baltic Sea resulting in moderate snowfall intensities over southern Sweden. Their influence on the northern parts of the country is minimal. This is mainly due to the fact that southeasterly frontal systems dry up before reaching the northernmost parts of Sweden. Furthermore, the Bay of Bothnia is frozen and covered by sea ice during core winter months closing the moisture source in the Baltic Sea. These results underline the varied meridional nature of the response of snowfall to dominant wind directions.

4.2 Snowfall during different synoptic states

The principles behind the results for the dominant wind regimes are also manifested in the observed sensitivity of snowfall to MSLP and enhanced positive or negative NAO index.

Figure 3b shows that for the weather state characterised by anti-cyclonic conditions (high MSLP), corresponding to an atmospheric blocking-like circulation pattern that stagnates over the entire southern Scandinavia, average snowfall intensities are similar for all radar stations, around $0.2\text{--}0.25\text{ mm h}^{-1}$. Temperatures range between -6°C to -2°C with the colder temperatures occurring at the northern stations (Fig. 3d, f). When cyclonic conditions prevail (low MSLP) the snowfall intensities are higher, as expected, ranging from 0.25 to 0.40 mm h^{-1} for the southern radar stations. On the other hand, at the northern stations snowfall rates are $0.2\text{--}0.3\text{ mm h}^{-1}$ for both cyclonic and anti-cyclonic regimes. The average values for the specific humidity, as observed by AIRS, for high and low MSLP are presented in Fig. 7a and b. The corresponding anomalies are shown in Fig. 8. It is seen that during high MSLP the air is more humid than for conditions with low MSLP. This is in contrast to the observations of snowfall intensities. To understand this apparent paradox we must examine the vertical structure of snowfall.

Figure 9a and b show the vertical structure of snowfall during high and low MSLP, respectively, as observed by CloudSat. It can be seen that the mean snowfall intensity near the surface is around 0.2 mm h^{-1} during both weather states, agreeing with ground-based observations (cf. Fig. 3b). CloudSat observations suggest, however, that more vertically-developed snowfall events occur predominantly during anti-cyclonic conditions. When MSLP is high, snowfall rates are confined around $0.1\text{--}0.2\text{ mm h}^{-1}$ at all heights from the surface up to 3 km whereas the vertical distributions of snowfall during low MSLP events are more widespread, ranging from 0.01 mm h^{-1} to 0.5 mm h^{-1} . ~~This explains-~~

It can now be explained why the specific humidity is higher during anti-cyclonic conditions while at the same time the average surface snowfall rates are ~~slightly lower~~, similar. Figure 2a shows that during high MSLP conditions, anti-cyclonic winds coming from central Europe further pick up moisture from the North Sea before releasing precipitation over western coast of Norway and southern parts of Sweden. This leads to increase in specific humidity over these regions (cf. Fig. 7a and Fig. 8a). These snowfall events, whenever they do occur, also seem to be well vertically developed as evident in Fig. 9a. However, during low MSLP conditions, the location of low pressure (Fig. 2b) is such that more northerly, colder winds result in lower specific humidity anomalies (Fig. 8b). Notice however that over the southernmost tip of Sweden, humidity anomalies are slightly positive during low MSLP conditions due to warmer southwesterly winds. Since the centre of the low pressure is slightly northward (compared to the centre of the high pressure), the occurrence of snowfall events with differing intensities, over northern and southern regions of Sweden leads to more broader vertical distribution of snowfall as observed by CloudSat (Fig. 9b).

The apparent insensitivity of snowfall to change in MSLP over the northern parts of Sweden can be understood by noting that during anti-cyclonic conditions these areas receive similar amount of average snowfall from winds carrying moisture from the northern part of the Norwegian Sea. Very low differences in average temperatures over the northern regions during anti-cyclonic and cyclonic conditions are further observed (Fig. 3d). At 850 hPa the air is more humid for high MSLP leading to the observed differences in the vertical structure of snowfall. However, even though the distribution of snowfall intensities are very different during these weather states the average snowfall rates are similar at the surface.

Ground-based observations of snowfall intensities during weather states with enhanced positive NAO indices, shown in Fig. 3b, range between $0.20\text{--}0.35\text{ mm h}^{-1}$ and are quite similar to those observed during enhanced negative NAO phase. Except for the southeastern coast, snowfall intensities during positive NAO indices are slightly larger. The corresponding average temperatures are shown in Fig. 3d. It is seen that temperatures decrease with increasing latitude but also that temperatures are $1\text{--}3^\circ\text{C}$ colder for enhanced negative phase of the NAO. The average values for the specific humidity are depicted in Fig. 7c and d. During enhanced positive NAO conditions the air is more humid compared to conditions of enhanced negative NAO.

By examining CloudSat's observations of the vertical structure of snowfall during these weather states, shown in Fig. 9c and d, it can be seen that the mean snowfall rate near the surface is slightly higher for the enhanced positive NAO phase, near 0.25 mm h^{-1} , compared to the enhanced negative NAO phase, 0.2 mm h^{-1} . The median heights of the snowfall intensities are similar for both conditions. A more obvious difference can be seen in the distribution of the snowfall intensities. For enhanced positive NAO phase the snowfall distribution peaks around 0.4 mm h^{-1} for heights from the surface up to 2 km whereas the distribution of snowfall intensities during enhanced negative NAO phase is more widespread, dominated by snowfall intensities from 0.005 mm h^{-1} to 0.1 mm h^{-1} .

The similarity in snowfall intensities during enhanced negative and enhanced positive NAO phases can be explained by temperature anomalies (Fig. 3f) and circulation patterns (Fig. 2). During the enhanced positive NAO phase, only the persistent southwesterly winds can bring sufficient moisture to colder northern parts of Sweden to create snowfall events with average intensities similar to those during the negative phase of NAO. While in the case of enhanced negative NAO phase over the southern parts of country, the circulation pattern favours conditions wherein southerly moist winds are confronted by colder and drier northerly winds (Fig. 8d) resulting in snowfall events with average intensities similar to those during the enhanced positive NAO phase. These results underline the complex interplay between circulation patterns and snowfall by affecting moisture transport and meridional temperature gradients.

4.3 Implications for snowfall accumulations

The relative strengths of the snowfall intensities for the different weather states are shown in Fig. 10a and b. In [order to more easily compare the snowfall intensities they have been normalised in Fig. 10a and b so that the sum of the snowfall intensities is equal to one for each of the different groups of weather states \(wind directions, MSPL, and NAO\)](#). In Fig. 10a it can be seen that the highest snowfall intensities occur for winds from the southwest for almost all stations in Sweden, which is a result of the warm temperatures and moist air that characterises this weather state. The other dominant wind directions all lead to similar snowfall intensities, albeit due to different reasons which were discussed in the preceding sections.

From the presented analysis no quantitative statement can be made for snowfall accumulation. However, the relative contribution of the different weather states to the total snowfall amount can be estimated by multiplying the average snowfall intensities with the number of days for each weather state. Even though snowfall intensities are highest for winds from the southwest this weather state does not contribute most to the total snowfall amount. The relative contribution to snowfall accumulation for

5 weather states characterised by dominant wind directions are shown in Fig. 10c. Here it is seen that for southern radar stations easterly winds lead to the largest amount of snow whereas for northern stations winds from the northwest lead to the largest snowfall accumulation. These results reflect the frequency with which snowfall occur during the different wind regimes. For example, even though southwesterly winds are the most common wind direction in Sweden they are usually accompanied by warm temperatures. Hence precipitation normally falls as rain during this weather state but if temperatures happen to be below
10 0° C, the snowfall intensities become large. On the other hand, during easterly winds temperatures are more often freezing and these weather states therefore contribute most to the total accumulation of snow.

For the other weather states (high and low MSLP and enhanced positive or negative NAO indices) the relative strength of the snowfall intensities are shown in Fig. 10b. The absolute differences in snowfall intensities are smaller for these weather states compared to those characterised by the dominant wind directions. However, it is seen that higher snowfall intensities occur for
15 low MSLP for all Sweden (except at the very north) while snowfall intensities are slightly higher for enhanced positive NAO for almost the entire country, except for the southernmost part.

For high and low MSLP it is clear that the contribution to total snowfall accumulation is larger for weather states with low MSLP for all stations. This is partly a reflection of the higher snowfall intensities associated with low MSLP but also due to the higher frequency with which it snows during these conditions. For enhanced positive and negative NAO indices it is clear
20 that in southern Sweden the contribution to total snow accumulation is dominated by conditions with enhanced negative NAO. This is because snowfall is relatively uncommon during weather states characterised by enhanced positive NAO, due to the warm temperatures. For northern Sweden both weather states contribute with approximately the same amount.

5 Conclusions

In this work we have examined the sensitivity of snowfall to eight different atmospheric circulation patterns, or weather states,
25 commonly occurring over Sweden. The circulation patterns include four dominant wind directions, high and low MSLP, and enhanced positive and negative NAO indices. To this end we have used a combination of different data sources: spatially-interpolated in situ observations, ground-based radar data, satellite measurements, and reanalyses products.

It was found that falling snow over Sweden is indeed sensitive to different atmospheric weather states. A clear variation in average snowfall intensity was observed, depending on circulation pattern but also depending on latitude. Snowfall over
30 southern Sweden responds to different weather states in a different way compared to over the northern parts of the country. The largest snowfall intensities were observed for southwesterly winds, which bring warm, moist air from the Atlantic over the entire country. The smallest average snowfall rates for northern Sweden were observed during southeasterly winds while the southern parts of the country received the smallest snowfall rates northeasterly winds.

Using satellite observations the vertical snowfall structure of snowfall were examined. Clear differences between the various circulation patterns could be observed. During easterly winds snowfall was seen to be much more shallow compared to snowfall during westerly winds. During high MSLP and positive phases of the NAO snowfall tended to be concentrated to larger intensities compared to conditions with low MSLP or negative phases of the NAO.

- 5 Snowfall accumulations depend on the observed snowfall rates but also on the frequency with which the weather states occur. It was found that easterly winds contribute most to snowfall accumulation over southern Sweden whereas westerly winds bring the largest snow amounts to the northern parts of the country. Further, low MSLP and negative phases of the NAO contribute more to snowfall accumulation than high MSLP and positive phases of the NAO.

- 10 These results indicate a complex interplay between circulation patterns, heat and moisture transport, and snowfall and under-
lines the varied meridional nature of this interplay. The study also demonstrates the advantage of the synergetic use of different observing systems to investigate snowfall variability. [This study was performed over Sweden but a similar study could easily be performed for other regions that have long-term radar observations available.](#)

6 Data availability

- 15 ERA-Interim data can be accessed from the website of the ECMWF, `apps.ecmwf.int/datasets/`. Nordrad data for research are available on request from SMHI. Composites based only on Swedish radars, very similar to Nordrad, are available from `opendata-catalog.smhi.se/explore/` where also Mesan data can be downloaded. CloudSat data are available from `cloudsat.atmos.colostate.edu/data`.

Author contributions. The authors designed the study and prepared the manuscript together. L. Norin and A. Devasthale performed the data analyses.

- 20 *Competing interests.* The authors declare that they have no conflict of interest.

References

- Battan, L. J.: Radar observation of the atmosphere, University of Chicago Press, 1973.
- Bech, J., Codina, B., Lorente, J., and Bebbington, D.: The sensitivity of single polarization weather radar beam blockage correction to variability in the vertical refractivity gradient, *J. Atmos. Oceanic Technol.*, 20, 845–855, doi:10.1175/1520-0426(2003)020<0845:TSOSPW>2.0.CO;2, 2003.
- 5 Berrisford, P., Dee, D., Poli, P., Brugge, R., Fielding, K., Fuentes, M., Kållberg, P., Kobayashi, S., Uppala, S., and Simmons, A.: The ERA-Interim archive, Era report series, European Centre for Medium Range Weather Forecasts, 2009.
- Busuioc, A., Chen, D., and Hellström, C.: Temporal and spatial variability of precipitation in Sweden and its link with the large-scale atmospheric circulation, *Tellus A*, 53, 348–367, doi:10.1034/j.1600-0870.2001.01152.x, 2001.
- 10 Carlsson, I.: NORDRAD — weather radar network, in: COST 75 Weather Radar Systems, edited by Collier, C. G., pp. 45–52, European Commission, 1995.
- Chahine, M., Pagano, T., Aumann, H., Atlas, R., Barnet, C., Blaisdell, J., Chen, L., Divakarla, M., Fetzer, E., Goldberg, M., Gautier, C., Granger, S., Hannon, S., Irion, F., Kakar, R., Kalnay, E., Lambrigtsen, B., Lee, S., Marshall, J. L., McMillan, W., McMillin, L., Olsen, E., Revercomb, H., Rosenkranz, P., Smith, W., Staelin, D., Strow, L., Susskind, J., Tobin, D., Wolf, W., and Zhou, L.: AIRS: Improving
- 15 Weather Forecasting and Providing New Data on Greenhouse Gases, *Bull. Amer. Meteor. Soc.*, 87, 911–926, doi:10.1175/BAMS-87-7-911, 2006.
- Chen, D.: A monthly circulation climatology for Sweden and its application to a winter temperature case study, *Int. J. Climatol.*, 20, 1067–1076, doi:10.1002/1097-0088(200008)20:10<1067::AID-JOC528>3.0.CO;2-Q, 2000.
- Daley, R.: Atmospheric Data Analysis, Cambridge Atmospheric and Space Science Series, Cambridge University Press, 2nd edn., 1993.
- 20 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q.J.R. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- 25 Devasthale, A. and Norin, L.: The large-scale spatio-temporal variability of precipitation over Sweden observed from the weather radar network, *Atmos. Meas. Tech.*, 7, 1605–1617, doi:10.5194/amt-7-1605-2014, 2014.
- Devasthale, A., Sedlar, J., Kahn, B., Tjernström, M., Fetzer, E., Tian, B., Teixeira, J., and Pagano, T.: A Decade of Spaceborne Observations of the Arctic Atmosphere: Novel Insights from NASA’s AIRS Instrument, *Bull. Amer. Meteor. Soc.*, 97, 2163–2176, doi:10.1175/BAMS-D-14-00202.1, 2016.
- 30 Gustafsson, M., Rayner, D., and Chen, D.: Extreme rainfall events in southern Sweden: where does the moisture come from?, *Tellus A*, 62, 605–616, doi:10.1111/j.1600-0870.2010.00456.x, 2010.
- Häggmark, L., Ivarsson, K.-I., and Olofsson, P.-O.: MESAN — mesoskalig analys, Reports Meteorology and Climatology RMK 75, SMHI, in Swedish, 1997.
- Häggmark, L., Ivarsson, K.-I., Gollvik, S., and Olofsson, P.-O.: Mesan, an operational mesoscale analysis system, *Tellus A*, 52, 2–20, doi:10.1034/j.1600-0870.2000.520102.x, 2000.
- 35 Harpold, A. A., Kaplan, M. L., Klos, P. Z., Link, T., McNamara, J. P., Rajagopal, S., Schumer, R., and Steele, C. M.: Rain or snow: hydrologic processes, observations, prediction, and research needs, *Hydrol. Earth Syst. Sci.*, 21, 1–22, doi:10.5194/hess-21-1-2017, 2017.

- Haynes, J. M., L'Ecuyer, T. S., Stephens, G. L., Miller, S. D., Mitrescu, C., Wood, N. B., and Tanelli, S.: Rainfall retrieval over the ocean with spaceborne W-band radar, *J. Geophys. Res.*, 114, doi:10.1029/2008JD009974, 2009.
- Hellström, C.: Atmospheric conditions during extreme and non-extreme precipitation events in Sweden, *Int. J. Climatol.*, 25, 631–648, doi:10.1002/joc.1119, 2005.
- 5 Johansson, B. and Chen, D.: The influence of wind and topography on precipitation distribution in Sweden: statistical analysis and modelling, *Int. J. Climatol.*, 23, 1523–1535, doi:10.1002/joc.951, 2003.
- Källén, E.: HIRLAM Documentation Manual: System 2.5, Swedish Meteorological and Hydrological Institute, 1996.
- Kulie, M. S., Milani, L., Wood, N. B., Tushaus, S. A., and L'Ecuyer, T. S.: A Shallow Snowfall Census Using Spaceborne Radar, *J. Hydrometeor.*, 17, 1261–1279, doi:10.1175/JHM-D-15-0123.1, 2016.
- 10 Lamb, H. H.: Types and spells of weather around the year in the British Isles : Annual trends, seasonal structure of the year, singularities, *Q. J. Roy. Meteor. Soc.*, 76, 393–429, doi:10.1002/qj.49707633005, 1950.
- Levizzani, V., Laviola, S., and Cattani, E.: Detection and measurement of snowfall from space, *Remote Sensing*, 3, 145–166, doi:10.3390/rs3010145, 2011.
- Linderson, M.-L.: Objective classification of atmospheric circulation over southern Scandinavia, *Int. J. Climatol.*, 21, 155–169, doi:10.1002/joc.604, 2001.
- 15 McIlhattan and L'Ecuyer, T. S.: Observational Evidence Linking GCM Arctic Supercooled Liquid Cloud Biases to Snowfall Processes, Submitted to *J. Climate*, 2016.
- Michelson, D.: The Swedish weather radar production chain, in: *Proceedings of the Fourth European Conference on Radar in Meteorology and Hydrology*, Barcelona, Spain, pp. 382–385, ERAD, 2006.
- 20 Michelson, D. B. and Koistinen, J.: Gauge-Radar network adjustment for the Baltic sea experiment, *Phys. Chem. Earth Pt. B*, 25, 915–920, doi:http://dx.doi.org/10.1016/S1464-1909(00)00125-8, 2000.
- NOAA: <http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml>, accessed 16 January 2017, 2016.
- Norin, L., Devasthale, A., L'Ecuyer, T. S., Wood, N. B., and Smalley, M.: Intercomparison of snowfall estimates derived from the CloudSat Cloud Profiling Radar and the ground-based weather radar network over Sweden, *Atmos. Meas. Tech.*, 8, 5009–5021, doi:10.5194/amt-8-5009-2015, 2015.
- 25 Smalley, M., L'Ecuyer, T., Lebsock, M., and Haynes, J.: A comparison of precipitation occurrence from the NCEP Stage IV QPE product and the CloudSat Cloud Profiling Radar, *J. Hydrometeor.*, 15, 444–458, doi:10.1175/JHM-D-13-048.1, 2014.
- Susskind, J., Blaisdell, J. M., and Iredell, L.: Improved methodology for surface and atmospheric soundings, error estimates and quality control procedures: the atmospheric infrared sounder science team version-6 retrieval algorithm, *J. Appl. Remote Sens.*, 8, 084994, doi:10.1117/1.JRS.8.084994, 2014.
- 30 Thomas, M. A. and Devasthale, A.: Sensitivity of free tropospheric carbon monoxide to atmospheric weather states and their persistency: an observational assessment over the Nordic countries, *Atmos. Chem. Phys.*, 14, 11 545–11 555, doi:10.5194/acp-14-11545-2014, 2014.
- Waliser, D. E., Li, J.-L. F., L'Ecuyer, T. S., and Chen, W.-T.: The impact of precipitating ice and snow on the radiation balance in global climate models, *Geophys. Res. Letters*, 38, doi:10.1029/2010GL046478, 2011.
- 35 Wood, N. B.: Estimation of snow microphysical properties with application to millimeter-wavelength radar retrievals for snowfall rate, Ph.D. thesis, Colorado State University, Fort Collins, Colorado, <http://hdl.handle.net/10217/48170>, 2011.

Wood, N. B., L'Ecuyer, T., Vane, D. G., Stephens, G. L., and Partain, P.: Level 2C snow profile process description and interface control document, Tech. rep., Colorado State University, http://www.cloudsat.cira.colostate.edu/ICD/2C-SNOW-PROFILE/2C-SNOW-PROFILE_PDICD_P_R04.pdf, 2013.

5 Wood, N. B., L'Ecuyer, T. S., Heymsfield, A. J., and Stephens, G. L.: Microphysical constraints on millimeter-wavelength scattering properties of snow particles, *J. Appl. Meteorol. Clim.*, 54, 909–931, doi:10.1175/JAMC-D-14-0137.1, 2015.

Table 1. Number of days during which any of the eight different weather states was found for the period (2008–2012). Note that not all weather states are mutually exclusive.

	<u>NE</u>	<u>NW</u>	<u>SE</u>	<u>SW</u>	<u>HI</u>	<u>LO</u>	<u>EN</u>	<u>EP</u>
<u>NE</u>	<u>75</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>38</u>	<u>17</u>	<u>36</u>	<u>20</u>
<u>NW</u>	<u>0</u>	<u>191</u>	<u>0</u>	<u>0</u>	<u>58</u>	<u>55</u>	<u>68</u>	<u>85</u>
<u>SE</u>	<u>0</u>	<u>0</u>	<u>77</u>	<u>0</u>	<u>46</u>	<u>7</u>	<u>44</u>	<u>14</u>
<u>SW</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>283</u>	<u>53</u>	<u>127</u>	<u>132</u>	<u>67</u>
<u>HI</u>	<u>38</u>	<u>58</u>	<u>46</u>	<u>53</u>	<u>588</u>	<u>0</u>	<u>211</u>	<u>185</u>
<u>LO</u>	<u>17</u>	<u>55</u>	<u>7</u>	<u>127</u>	<u>0</u>	<u>624</u>	<u>330</u>	<u>157</u>
<u>EN</u>	<u>36</u>	<u>68</u>	<u>44</u>	<u>132</u>	<u>211</u>	<u>330</u>	<u>797</u>	<u>0</u>
<u>EP</u>	<u>20</u>	<u>85</u>	<u>14</u>	<u>67</u>	<u>185</u>	<u>157</u>	<u>0</u>	<u>531</u>




Figure 1. Atmospheric circulation patterns at 850 hPa leading to four persistent wind directions over Sweden. The wind directions were evaluated over the area enclosed by the white rectangle. Panels (a) to (d) show representative examples of circulation patterns resulting in winds from northwest, northeast, southwest, and southeast, respectively. The colourbar indicates wind strength (in m s^{-1}).




Figure 2. Atmospheric circulation patterns at 850 hPa for four different weather states. Panels (a) and (b) show representative examples of circulation patterns during high and low MSLP conditions (evaluated over the area enclosed by the white rectangle) whereas panels (c) and (d) depict patterns for enhanced positive and negative phases of the NAO. The colourbar shows wind strength (in m s^{-1}).




Figure 3. Average snowfall intensities (in mm h^{-1}) observed by the Swedish weather radar network for winds from northwest, northeast, southwest, and southeast (a) and during high and low MSLP and enhanced positive and negative NAO conditions (b). Average 2 m temperatures (in $^{\circ}\text{C}$), obtained from the mesoscale analysis system Mesan, for winds from northwest, northeast, southwest, and southeast (c) and during high and low MSLP and enhanced positive and negative NAO conditions (d). Average temperature anomalies (in $^{\circ}\text{C}$), obtained from the mesoscale analysis system Mesan, for winds from northwest, northeast, southwest, and southeast (e) and during high and low MSLP and enhanced positive and negative NAO conditions (f). [The coloured rings in panels \(a\), \(c\), and \(e\) correspond to the radar coverage area used in the analysis.](#)




Figure 4. Average values of specific humidity, observed by AIRS at 850 hPa. Panel (a) shows the specific humidity for atmospheric conditions dominated by winds from the northwest. In panels (b) to (d) the winds are from northeast, southwest and southeast, respectively.




Figure 5. Anomalies of specific humidity, observed by AIRS at 850 hPa. Panel (a) shows anomalies in the specific humidity for atmospheric conditions dominated by winds from the northwest. In panels (b) to (d) the winds are from northeast, southwest and southeast, respectively.




Figure 6. Histograms showing relative frequency of snowfall rates as a function of altitude above the surface, using data from CloudSat. The different panels show the vertical structure of snowfall for weather states dominated by different wind directions (northwest, northeast, southwest, and southeast). The red solid lines show the mean snowfall rates as a function of altitude. The two red dashed lines show the altitudes where 50 % and 95 % of the snowfall have occurred, as a function of snowfall intensity.




Figure 7. Average values of specific humidity, observed by AIRS at 850 hPa. Panels (a) and (b) show average values of the specific humidity for high and low MSLP while panels (c) and (d) show the corresponding variable for enhanced positive and negative phases of the NAO.




Figure 8. Anomalies of specific humidity, observed by AIRS at 850 hPa. Panels (a) and (b) show anomalies in the specific humidity for high and low MSLP while panels (c) and (d) show the corresponding variable for enhanced positive and negative phases of the NAO.




Figure 9. Histograms showing relative frequency of snowfall rates as a function of altitude above the surface, using data from CloudSat. The different panels show the vertical structure of snowfall for weather states with high and low MSLP (**a** and **b**), and enhanced positive and negative NAO indices (**c** and **d**). The red solid line shows the mean snowfall rate as a function of altitude. The two red dashed lines show the altitudes where 50 % and 95 % of the snowfall have occurred, as a function of snowfall intensity.




Figure 10. Relative strength of snowfall intensities for winds from northwest, northeast, southwest, and southeast **(a)**. For every radar station the sum of the relative strength from the four wind directions is equal to one. Relative strength of snowfall intensities during high and low MSLP and enhanced positive and negative NAO conditions **(b)**. Here the sum of the relative strength during high and low MSLP for every station is equal to one as well as the sum of the relative strength during enhanced positive and negative NAO conditions is equal to one. Relative contribution of total snowfall for winds from northwest, northeast, southwest, and southeast **(c)**. The sum for every radar station is equal to one. Relative contribution of total snowfall during high and low MSLP and enhanced positive and negative NAO conditions **(d)**. The sum of the relative contribution during high and low MSLP for every station is equal to one and the sum of the relative contribution during enhanced positive and negative NAO conditions is equal to one. The coloured rings in panels (a) and (c) correspond to the radar coverage area used in the analysis.