

1 **Examination on total ozone column retrievals by Brewer spectrophotometry**
2 **using different processing software.**

3
4 Anna Maria Siani^{1*}, **Francesca Frasca**², Francesco Scarlatti³, Arianna Religi⁴, Henri Diémoz⁵,
5 Giuseppe.R. Casale¹, Massimiliano Pedone⁶, Volodya Savastiouk⁷

6
7 ¹Department of Physics, Sapienza Università di Roma, Rome, Italy

8 ²Department of Earth Sciences, Sapienza Università di Roma, Rome, Italy

9 ³Independent scientist

10 ⁴Institute of Services Science, University of Geneva, Geneva, Switzerland

11 ⁵Aosta Valley Regional Environmental Protection Agency (ARPA), Saint-Christophe, Italy

12 ⁶Infosapienza Settore per i Sistemi Centrali e per l'Office Automation, Sapienza Università di Roma, Rome, Italy

13 ⁷International Ozone Services Inc., Toronto, Ontario, Canada

14
15 *Correspondence to: Anna Maria Siani (annamaria.siani@uniroma1.it)

16
17
18 **Abstract.** The availability of long-term records of the total ozone content (TOC) represents a
19 valuable source of information in studies on the assessment of short and long-term atmospheric
20 changes and their impact on the terrestrial ecosystem. In *particular*, ground-based observations
21 represent a valuable tool to validate satellite-derived products. To our knowledge, details about
22 ~~processing~~ software packages to process Brewer spectrophotometer measurements and to retrieve
23 the TOC are seldom specified in studies using such datasets. *The sources of the differences among*
24 *retrieved TOCs from the Brewer instruments located at the Italian stations Rome and Aosta, using*
25 *three freely available codes* (Brewer Processing Software, O3Brewer software and EUBREWNET
26 *Level 1.5 products*) are here investigated. Ground-based TOCs are also compared with the Ozone
27 Monitoring Instrument (OMI) TOC retrievals used as an independent dataset since no other
28 instruments near the Brewer sites, are available.

29 *The overall agreement of the BPS and O3Brewer TOC data with EUBREWNET data is within the*
30 *estimated total uncertainty in the retrieval of total ozone from Brewer spectrophotometer (1%).*
31 *However, differences can be found depending on the software in use. Such differences become*
32 *larger when the instrumental sensitivity exhibits a fast and dramatic drift which can affect the*
33 *ozone retrievals significantly. Moreover, if daily mean values are directly generated by the*

34 *software, differences can be observed due to the configuration set by the users to process single*
35 *ozone measurement and the rejection conditions applied to data to calculate the daily value.*

36 This work aims to provide useful information both for scientists engaged in ozone measurements
37 with Brewer spectrophotometry and for stakeholders of the Brewer data products available at
38 web-based platforms.

39
40 **Key words:** ozone, Brewer spectrophotometry, standard lamp correction, processing software,
41 calibration

42

43

44 1.INTRODUCTION

45

46 Although ozone (O₃) is present in small amounts in the terrestrial atmosphere, it plays a
47 crucial role in the attenuation of solar ultraviolet (UV) radiation (200 - 400 nm) reaching the
48 surface and in radiative processes controlling the energy balance on the Earth (Ramanathan and
49 Dickinson, 1979; Dessler, 2000; Bordi et al., 2012; WMO, 2015).

50 The cumulative amount of stratospheric and tropospheric ozone represents the total ozone
51 column (TOC). The most common ground-based instruments to measure TOC are
52 spectrophotometers which are designed to measure *ground level spectral intensities of solar*
53 *ultraviolet radiation attenuated by ozone absorption. From these spectra, it is possible to* retrieve
54 the TOCs. The first TOC observations were recorded using the Dobson spectrophotometer
55 (Dobson and Harrison, 1926) in the late 1920s. Since then, a growing number of sites were
56 equipped with the Dobson spectrophotometer and later in the 1980s with the automated Brewer
57 spectrophotometer (Brewer, 1973). Nowadays, both the Dobson and the Brewer
58 spectrophotometers are used all over the world and *the accuracy of measurements taken with a*
59 *well-maintained Brewer spectrophotometer is 1% in the direct sun (DS) mode (Vanicek, 2006).*

60 It should be pointed out that high-quality TOC retrievals from ground-based stations are
61 necessary not only in support of the validation of satellite-derived products (Tzortziou et al.,
62 2012) but also for the assessment of the long-term ozone trend and the verification of the
63 effectiveness of the Montreal Protocol on substances that deplete the ozone layer. Moreover,
64 ground-based TOC data are also necessary to calibrate the parameters in the global climate
65 models used to predict the expected behaviour of the ozone layer in the future (Stübi et al., 2017).

66 The above issues show the importance to measure the ozone amount from ground-based
67 stations with a very good performance.

68 *Even though all available processing software packages use the same TOC retrieval*
69 *algorithm, which is based on the Bouguer-Lambert-Beer law, slightly different implementations*
70 *potentially trigger some differences in the processed TOC data.*

71 The largest part of the total ozone column data analysed in the current/available scientific
72 literature is extracted from the WOUDC data archive (World Ozone and Ultraviolet Radiation
73 Data Centre). To our knowledge, the processing software of Brewer TOC data varies from site to
74 site, the processing algorithm *and the data rejection rules are seldom specified. WOUDC ozone*
75 *files (2017) do not include information on the software used to process ozone data, the version of*

76 *such software as well as the adopted data rejection rules. The same information is usually not*
77 *reported in the studies related to ozone monitoring, trend detection and satellite validation.* This
78 can be due to the fact that a standard processing software of Brewer raw data has currently not
79 been adopted. *For this reason,* the COST Action ES1207 “A European Brewer Network”
80 (EUBREWNET) was established aiming at defining, among the others, a standard procedure for
81 processing the raw Brewer data, thus ensuring the quality of the data and harmonizing the
82 products from the European *Brewers* (EUBREWNET, 2017).

83 *The purpose of the present study is to investigate the differences among the TOCs retrieved*
84 *by three different processing software packages: the Brewer Processing Software, hereafter*
85 *called BPS, developed by Dr Fioletov V. and Ogyu A. (Environment Canada), O3Brewer software*
86 *developed by Ing Stanek M. (Solar and Ozone Observatory of CHMI/International Ozone*
87 *Service) and the EUBREWNET level 1.5 ozone product. To the purpose of an intercomparison*
88 *exercise, we tested the mentioned software on the datasets collected by the Brewer instruments*
89 *installed at Rome and Aosta, Italy. Then, Brewer ozone recalculations were also compared with*
90 *the Ozone Monitoring Instrument (OMI) TOC retrievals. The OMI data were used since no other*
91 *independent collocated TOC measurements were available.*

92 *This paper is structured as follows: Section 2.1 briefly describes the theory on the ozone*
93 *estimates from Brewer direct sun (DS) measurements. In Section 2.3, the procedure used by three*
94 *software packages to process ozone data is presented. Section 2.4 describes the Brewer stations*
95 *under study. Section 3 is dedicated to the comparison among the three TOC data retrievals and to*
96 *understand the causes responsible for the differences among processed ozone values. Additional*
97 *comparison between ground-based data and OMI products is also carried out. Moreover ozone*
98 *trends are estimated to investigate if using specific software could affect the results. Finally,*
99 *conclusions are drawn in the last section.*

100

101 **2. DATA AND METHOD**

102 **2.1 Theory of direct sun measurements with Brewer spectrophotometer**

103 The Brewer spectrophotometer is an instrument designed to retrieve the total ozone
104 column *by measuring irradiances of both direct sunlight (Kerr et al., 1981) and polarized*

105 *radiation scattered from the zenith sky (Brewer and Kerr, 1973, Muthama et al., 1995). Total*
106 *ozone can be also derived from focused sun measurements, commonly employed at high latitudes*
107 *(Josefsson, 1992). It is also possible to measure total ozone by using the moon (Kerr et al., 1990),*
108 *or the global irradiance method in the UV region (Kerr and Davis, 2007), as a light source.*

109 The most accurate method to determine the total column amount of an atmospheric gas is based
110 on the direct sun (DS) measurements. *It was shown (Vanicek, 2006) that the accuracy of*
111 *measurements taken with a well-maintained Brewer spectrophotometer is 1% in the DS mode and*
112 *3-4% in the ZS mode. The random errors of individual measurements were found to be within ±*
113 *1% for all measurements (Fioletov et al., 2005).*

114 The algorithm to retrieve the total ozone column from the Brewer in DS mode is based on
115 a differential measurement method involving 4 selected wavelengths in the ozone absorption
116 spectra, nominally 310.1, 313.5, 316.8 and 320.1 nm.

117 *The wavelengths are selected by a rapidly rotating slit-mask and raw photon counts for*
118 *each slit-mask wavelength position (from 3 to 6) are registered by a photomultiplier. During each*
119 *measurement run cycle the slit-mask is rotated 20 times.* The raw photon counts are then
120 converted into count rates *and are corrected for the characteristics of the photomultiplier (dark*
121 *count and dead time) and for the internal Brewer temperature (Kerr, 2010).* In addition, a
122 correction for the spectral transmittance of the attenuation filters can be added depending on the
123 filter used, if the respective characterisation is available.

124 *A linear combination (F) of the base-ten logarithms of the count rates (F_i) measured*
125 *during the direct sun spectral irradiance observations for the i-th slit is computed by weighting*
126 *the F_i with coefficients (w_i=-1, +0.5, +2.2, -1.7). The weighting coefficients are chosen in order*
127 *to minimize the effect of the aerosol extinction, to eliminate the effect of the sulphur dioxide*
128 *absorption (Kerr et al., 1981; Kerr, 2010) and all factors independent of the wavelength (flat*
129 *factors):*

130

131
$$F = \sum_{i=1}^4 w_i \log F_i \quad (1)$$

132

133 F_i is also compensated for the effect of the Rayleigh scattering by subtracting:

134

$$135 \quad \frac{p}{p_o} \mu_R \sum_{i=1}^4 w_i \beta_i \quad (2)$$

136

137 where p is the climatological pressure at the measurement site and p_o is the pressure at the sea
138 level; μ_R is the Rayleigh air mass factor (i.e. the slant path of direct radiation through air),
139 calculated for a thin layer at 5 km altitude, β_i is the Rayleigh scattering coefficient at the
140 wavelength, λ_i .

141 According to the Bouguer-Lambert-Beer law, it is possible to retrieve the total ozone
142 column (TOC) as:

143

$$144 \quad TOC = \frac{F_o - F}{\Delta\alpha\mu} \quad (3)$$

145

146 where $\Delta\alpha$ is the *differential ozone absorption coefficient*, i.e. the linear combination of the ozone
147 cross sections using the same weighting coefficients employed for F . $\Delta\alpha$ is calculated after
148 performing a specific test using spectral lamps providing the precise operational wavelengths and
149 applying the convolution with the slit function characterised for each individual
150 spectrophotometer. Then $\Delta\alpha$ is obtained for these wavelengths using Bass-Paur ozone absorption
151 spectrum (Bass and Paur, 1985) at the fixed temperature of -45°C (Kerr, 2010).

152 The standard Brewer algorithm assumes that the ozone is concentrated in a thin layer at
153 the altitude of 22 km, thus the air mass factor (μ) is expressed by:

$$154 \quad \mu = \sec \left[\arcsin \left(\frac{R_E}{R_E + 22} \sin Z \right) \right] \quad (4)$$

155 where R_E is the Earth's radius and Z is the solar zenith angle. F_o is also expressed as the linear
156 combination of the extraterrestrial irradiance at the operational Brewer wavelengths with the same
157 weighting coefficients used for F . F_o corresponds to F at the top of the atmosphere and it is usually
158 named "extraterrestrial constant" (ETC), *a specific factor different for each Brewer, and*
159 *determined through a calibration procedure.*

160 There are two methods to determine the ETC. The first is based on the use of the Langley
161 plot technique i.e. plotting F versus μ , and then the ETC value is extrapolated at zero air mass.
162 This method is used for the calibration of primary standards and requires to be carried out under
163 stable atmospheric conditions and low pollution concentrations. The second method is based on
164 transferring the calibration from a reference Brewer instrument with a known ETC to a candidate
165 instrument during field campaigns. This latter technique is the most common way for regularly
166 calibrating the instruments which belong to the Brewer network. In between the calibration audits
167 with a travelling standard, the TOC data are processed adjusting the ETC according to the
168 changes of the radiometric sensitivity of the instrument, if needed. The correction uses *time series*
169 *of the internal standard lamp tests, described in the Section 2.2.*

170 Direct-sun measurements are carried out at specific solar zenith angles throughout the day
171 depending on the user schedule (a sequence of commands written by the operator), allowing the
172 Brewer to make observations continuously and automatically. During a DS measurement, *a group*
173 *of five consecutive sub-measurements* are taken in less than five minutes. Then the mean and the
174 standard deviation of the five ozone values are computed and associated to that DS measurement.

175 *The standard deviation is used to determine the acceptability of each TOC measurement.*
176 *An individual TOC value is normally considered acceptable if the standard deviation of the five*
177 *measurements is lower than 2.5 DU or 3 DU.*

178

179 **2.2 Standard lamp correction**

180 Several tests are performed on a daily and weekly basis *to verify that* the Brewer operates
181 correctly and *to track* the changes in instrumental properties. The main standard tests included in
182 the diurnal operational schedule are: shutter motor run/stop (RS), photomultiplier dead time (DT),
183 mercury lamp (Hg) and standard lamp (SL).

184 The RS test verifies that the slit-mask motor is operating properly. It calculates the ratio of
185 irradiances at the operational wavelength using an internal 20 W quartz-halogen *lamp as the light*
186 *source* in a dynamic mode and in a static mode. This ratio should be as close as possible to unity.

187 The DT test measures the dead-time of the photomultiplier and the photon-counting
188 circuitry and the result of the test value should be within 5 ns with respect to the instrument
189 constant. Also during the DT test, the halogen lamp is turned on.

190 *For the Hg test a mercury lamp is used. This test ensures the correct wavelength*
191 *alignment of the Brewer due to the internal temperature changes.* This test is usually carried out
192 several times every day.

193 The SL test is used to monitor the stability of the instrument response after the calibration
194 with the reference spectrophotometer. The test is performed *using the internal quartz-halogen*
195 *lamp* as the light source. The photon counts are recorded at the same operational wavelengths
196 employed in the DS measurement and the result of the SL test, the so-called R6 ratio which
197 corresponds to a fictitious value of ozone column density, is determined using Eq.(1). In this way
198 changes *with respect to the reference R6 value ($R6_{ref}$), determined during the calibration with the*
199 *reference instrument, are constantly tracked.*

200 If a change in R6 is experienced, this results in a corresponding change in the ETC
201 assuming that the relative lamp intensities at the four wavelengths do not change. *Consequently, a*
202 *correction in the reference ETC should be applied to determine the ozone values in between each*
203 *calibration, as follows:*

204

$$205 \quad TOC = \frac{ETC - F + \Delta SL}{\Delta \alpha \mu} \quad (5)$$

206

207 where ΔSL is the correction factor measuring the difference between $R6_{ref}$ *which is determined at*
208 *every calibration* and R6 for a specific day.

209 Depending on the processing software used by the station operator, ΔSL is computed in
210 different ways, *not always clearly explained by the software documentation:*

211 *•In the BPS, the reference value $R6_{ref}$ is determined with a triangular smoothing filter of*
212 *SL-test values over the 15- days period immediately following the calibration date. There*
213 *should be at least one good SL-test value per day. If the corresponding B-files are not*
214 *available, the program is not able to establish the reference SL level and the ETC will be*

215 *not adjusted. Notice that for other processing software $R6_{ref}$ is based on the SL-test values*
216 *during the calibration campaign. If the $abs(R6_{ref} - R6) \leq 250$ units, then the median of*
217 *daily averages from all R6 data before 15 days and after 15 days for a particular day is*
218 *used for the correction. The median is used because it is less influenced by single invalid*
219 *R6s. If the $abs(R6_{ref} - R6)$ is above 250 units then ETC is adjusted taking into account the*
220 *difference between the $R6_{ref}$ and the present daily mean values of R6. That correction is*
221 *reported in the file named “o3data” produced by the BPS. The threshold and the time window*
222 *are however not adjustable by the users (Fioletov personal communication, 2018).*

223 • O3Brewer adjusts the ETC using a Gaussian smoothing filter on R6 values (Stanek M.,
224 2016). *There should be SL measurements 10 days before and 10 days after the selected*
225 *date period.* The software creates the smoothed R6 time series (hereafter named $R6_{smooth}$)
226 which is used for ETC adjustment. *It means that there should be at least one SL test per*
227 *day. The application of the ETC correction is done when the difference between the*
228 *reference $R6_{ref}$ and R6 from SL test results, does not exceed is a certain value (the default*
229 *value is 500 units) . This threshold is now configurable in the latest version 6.0 (Stanek*
230 *personal communication, 2018). The time window is however not adjustable by the users. If this*
231 *difference exceeds the threshold, then the software can remember the last day with good*
232 *SL test and will apply that correction (Stanek personal communication, 2018). This option can*
233 *be turned off and then the daily mean values for SL are used for the correction of the ETC.*

234 • *Level 1.5 total ozone column data from EUBREWNET are recalculated with the ΔSL*
235 *correction determined applying a triangular moving average over the daily median values*
236 *of R6 within a seven days window (default time window). The correction is applied if the*
237 *difference between $R6_{ref}$ and the calculated value exceeds 5 units. Level 2.0 are 1.5*
238 *observations validated with a posterior calibration. If the reference constants of a*
239 *posteriori calibration do not differ significantly from the values in use then level 1.5*
240 *products are not reprocessed and represent the most reliable products*
241 *(<http://rbcce.aemet.es/dokuwiki/doku.php>).*

242 At the present time, tools for Level 2.0 are developed but not yet implemented. A
243 complete description of the processing can be found on the EUBREWNET website (2017).

244

245

246

247 2.3 Measuring instruments and sites

248 Brewers MKIV serial numbers 067 and 066 have been operating at the Solar Radiometry
249 Observatory of Sapienza University of Rome (hereafter Rome) and at the headquarter of Aosta
250 Valley Regional Environmental Protection Agency (ARPA) at Aosta-Saint Christophe (hereafter
251 Aosta), respectively. The former has been recording TOCs since 1992 (Siani et al., 2002) whereas
252 the latter since 2007 (Siani et al., 2013).

253 In this study the above-mentioned sites were selected because both Brewers belong to
254 Sapienza University of Rome and have been calibrated with the same reference
255 spectrophotometer since their installation, submitting regularly data to the WOUDC and taking
256 part to the COST Action ES1207 EUBREWNET. The station characteristics are reported in Table
257 1.

258 **Table 1.** Characteristics of the two Italian Brewer sites
259

Station name (GAW ID)	Brewer Serial number	Coordinates (latitude, longitude, elevation (m above sea level))	Observation period	Environmental context
Aosta (AST)	066	45.7°N, 7.4°E, 569 m a.s.l.	29/01/2007 - 31/12/2015	semi-rural
Rome University (ROM)	067	41.9°N, 12.5°E, 75 m a.s.l.	01/01/1992 - 31/12/2015	urban

260

261 Since their installation, both Italian Brewers have been calibrated every one or two years
262 by intercomparison with the traveling reference Brewer 017 from International Ozone Services
263 Inc. (IOS), (2017). This Brewer is in turn calibrated against the World Brewer Reference Triad in
264 Toronto (Fioletov et al., 2005). In this way the ozone calibration of Italian spectrophotometers is
265 also traceable to the Brewer Reference Triad.

266 The calibration history of the two instruments used in this study is reported in Table 2.
267 Although zenith sky and global irradiance measurements were available, only DS measurements
268 were selected in this study because they have a lower uncertainty compared to the other types of
269 measurements (Fioletov et al., 2005).

270 *In this study we analysed individual DS values and daily averages of Rome and Aosta stations,*
271 *generated by BPS version 2.1.1 updated to 2017/02/14 (Fioletov and Ogyu, 2007), by O3Brewer*
272 *software packages version 6.0 updated to 2018/03/14, and EUBREWNET level 1.5 ozone*

273 *products. Level 1.5 individual TOC values are discarded when the standard deviation is above 2.5*
 274 *DU and the maximum ozone air mass is above 3.5. In addition ozone values less than 100 DU and*
 275 *greater than 500 DU are also rejected. The stray light* correction was not applied because it
 276 requires a calibration against a double monochromator Brewer and an instrumental
 277 characterization (Karppinen et al., 2015, Redondas et al., 2016) which *was* not available. Level
 278 1.5 TOC values were downloaded from EUBREWNET platform over the period 2005-2015 at
 279 Rome and 2007-2015 at Aosta.

280
 281 **Table 2.** Calibration history of Brewer 066 and 067. In brackets it is reported the month of the calibration for Brewer
 282 067 (*The recalculation of the constants was performed by IOS after the calibration on July 2009). In one case the
 283 calibration of Italian Brewers was performed in Arosa (Switzerland) at the Lichtklimatisches Observatorium during
 284 the Seventh Intercomparison campaign of the Regional Brewer Calibration Center Europe (WMO-GAW, 2015). In
 285 2013 the calibration of both Brewers was carried out at Aosta.

Year	Period	Location (Brewer 066)	Location (Brewer 067)
1992	January		Rome
1993	September		Rome
1995	May		Rome
1996	April		Rome
1997	May		Rome
1998	July		Rome
1999	September		Rome
2000	September		Rome
2002	March		Rome
2003	September		Rome
2006	September		Rome
2007	April	Aosta	Rome
2009	July	Aosta	Rome
2010*	January	Aosta	Rome
2011	August (July)	Aosta	Rome
2012	August (July)	Arosa	Arosa
2013	May (June)	Aosta	Aosta
2014	July		Rome
2015	July	Aosta	Rome

288
 289 *We set in the configuration file of BPS and O3Brewer software, where it is suitable, the*
 290 *same rejection criteria used in EUBREWNET, i.e. maximum standard deviation of 2.5 DU and*
 291 *maximum ozone air mass of 3.5. TOC.*

292 *The rejection criteria of ozone values are hardcoded in the BPS software and consist on*
 293 *three sequential checks: 1) if raw counts are less than 2500, the value is rejected; 2) if calculated*

294 *ozone for DS/ZS is less than 50 DU, the value is rejected 3) if observation is in the DS mode and*
295 *the calculated ozone is between 50 and 100 DU, the value is rejected (Ogyu, personal*
296 *communication 2018).*

297 *The maximum calculated ozone is indeed configurable in the BPS setup and was set to 500 DU.*

298 *The limits on the calculated ozone are not configurable in the O3Brewer setup. In the*
299 *latest version used in this study, the standard lamp maximum for applying of ETC correction from*
300 *SL test results is now configurable. Here we used the default limit of 500 units for the difference*
301 *between R6 and the reference R6_{ref}.*

302

303 **2.4 Satellite TOC data**

304 The Ozone Monitoring Instrument (OMI) products were used as an ancillary dataset with
305 the purpose of helping understand the difference among the investigated Brewer retrievals and the
306 comparison should not be regarded as exhaustive validation exercises of satellite total ozone data.
307 Daily averages of the Brewer TOC were compared with satellite ozone values obtained during the
308 overpass. The use of daily means instead of Brewer TOC observations taken close to the OMI
309 overpass is reasonable since it allows to compare a large number of pair measurements (Antón et
310 al., 2009; Vaz Peres et al., 2017) because there are only one or two daily satellite values.

311 Satellite overpass data at Rome and Aosta were derived from OMI, on board NASA EOS-
312 Aura spacecraft launched in July 2004. The OMI instrument is a nadir-viewing spectrometer
313 measuring solar reflected and backscattered light from the Earth atmosphere and surface in the
314 wavelength range from 270 nm to 500 nm, providing global daily coverage with a spatial
315 resolution of $13 \times 24 \text{ km}^2$ in nadir. The Aura satellite ~~describes~~ *travels in* a sun-synchronous polar
316 orbit, crossing the equator at 13:45 local time. *Two algorithms, OMI-TOMS (Total Ozone Mapping*
317 *Spectrometer) and OMI-DOAS (Differential Optical Absorption Spectroscopy), are used to produce*
318 *OMI daily total ozone datasets.*

319 In our study OMI-TOMS ozone overpasses based on TOMS V8.5 algorithm (Bhartia and
320 Wellemeyer, 2002) at the stations under study over the period 01/10/2004-31/12/2015 were
321 downloaded from the NASA –AURA validation data center platform. *Here we used OMI-TOMS since it*

322 *has a better agreement with the ground-based Brewer and Dobson instruments (Balis et al.,*
323 *2007).*

324

325 **2.5 Statistical parameters**

326 The following statistical parameters are used with the aim to quantify the differences
327 among the TOC series: nonparametric Spearman coefficient (RHO), Mean Bias (MB), Mean
328 Percentage Error (MPE), Root Mean Square Error (RMSE). RHO was used to measure the
329 correlation between two variables without making any assumption about their distribution. MB
330 represents the systematic differences between two selected datasets; MPE provides the average of
331 percentage errors with respect to TOC values taken as the reference. RMSE is an estimate of the
332 standard deviation of the difference (residuals) between two dataset.

$$333 \quad MB = \frac{1}{N} \sum_1^N (y_i - y'_i) \quad (6)$$

$$334 \quad MPE = 100 * \frac{1}{N} \sum_1^N \frac{(y_i - y'_i)}{y'_i} \quad (7)$$

$$335 \quad RMSE = \sqrt{\sum_1^N \frac{(y_i - y'_i)^2}{N}} \quad (8)$$

336 The previous equations show the formulas of the mentioned statistical parameters, where
337 y_i is the i-th TOC value (O3Brewer, or OMI) value, y'_i is the i-th TOC value of the BPS (or
338 EUBREWNET) series, N the number of all the possible data pairs analysed. *The uncertainty of*
339 *MB and MPE is characterized by the standard deviation.*

340 *In the comparison between Brewer and OMI data the scaled correlation (RHO) was calculated*
341 *(Diémoz et al.,2016) to exclude the possibility that the source of the correlation is a common*
342 *cycle (e.g. the annual cycle). That calculation is performed by splitting the series of the ozone*
343 *daily values in short intervals (here K=30 days) and for each interval RHO coefficient is*
344 *determined. Then RHOs are given by:*

345

$$346 \quad RHO_s = \frac{1}{K} \sum_{i=1}^K RHO_i \quad (9)$$

347 *In this way the high frequency component (<30 days) common to Brewer and OMI series are*
348 *revealed.*

349


350 **2.6 Trend analysis**

351 *To assess whether a specific software could affect the trend, we estimated the trend from the*
352 *annual mean anomalies. We applied the same methodology proposed by Fountoulakis et al.,*
353 *(2016). Climatological ozone values for each day were calculated over the period under study.*
354 *The daily anomaly with respect to the daily climatological value was calculated. Afterward the*
355 *monthly anomalies were determined by averaging the daily anomalies for each month provided*
356 *that at least 15 days of data were available. Finally the monthly anomalies were averaged to*
357 *determine the annual mean anomalies. The trend among the three codes was expressed as*
358 *percentage change per decade and used in their comparison. The statistical significance of the*
359 *trends is derived from the Mann–Kendall test with statistical significance set at $p \leq 5\%$.*

360

361 **3. RESULTS AND DISCUSSION**


362 The time series of TOC daily means *generated by* BPS, O3Brewer and calculated from
363 EUBREWNET individual ozone values, are presented in Figs. 1 (upper panel Rome, lower panel
364 Aosta). *Individual measurements are distinctly plotted for each site in Fig.2 and Fig.3.*



365


366 **Figure 1.** Time series of TOC daily means from BPS (black), O3Brewer (red) and EUBREWNET (blue) at
367 Rome (upper panel) and at Aosta (lower panel). *At Aosta the EUBREWNET L1.5 ozone values were not generated*
368 *between May 24 and September 8, 2008, because the standard lamp got burned out since May 2008 and was replaced*
369 *in September 2008.*

370



371
372
373
374

Figure 2. Individual TOC values generated by BPS (black), O3Brewer (red) and EUBREWNET (blue) at Rome.



375
376
377
378
379

Figure 3. Individual TOC values generated by BPS (black), O3Brewer (red) and EUBREWNET (blue) at Aosta.

380 It is worth noticing that ozone seasonal cycles show an overall similarity between the two
381 sites with maximum value in late spring and minimum in late autumn, both on daily means and on
382 individual ozone series. *The seasonal behaviour of O3Brewer is not easily distinguishable since*
383 *the y-axis range has flattened it due to negative recalculated ozone values. However it is clearly*
384 *visible that there are some periods in which TOC daily means as well as individual measurements*


385 *obtained by the three processing software, are different* (mainly between 2006 and 2007 at Rome
386 and at the end of 2011 at Aosta).

387 In order to understand *where the differences come from, we analysed both individual TOC*
388 *observations and the resulting daily values processed by BPS and O3Brewer*. Afterwards we
389 compared both TOC retrievals with EUBREWNET data. Finally, the processed Brewer data were
390 compared with OMI products.

391

392 **3.1 Comparison between BPS and O3Brewer TOC retrievals**

393 *Fig. 4 shows the temporal behaviour of the ozone differences between BPS and O3Brewer*
394 *taking into account both daily means whereas Fig. 5 shows individual values. It can be noticed*
395 *that in several cases large differences can be attributed to wrong negative ozone recalculations*
396 *by O3Brewer as also shown in Fig. 2 and 3. The minimum and maximum differences in the daily*
397 *means are -278.1 DU and 567.9 DU at Rome, -332.3 DU and 532.0 DU at Aosta, respectively.*
398 *The differences between BPS and O3Brewer individual ozone values range from a minimum of -*
399 *304.4 DU to a maximum of 90.6 DU at Rome, from -435.6 DU to -157.7 DU at Aosta.*




400

401

402

Figure 4. Time plot of the differences between BPS and O3Brewer daily means at Rome (upper panel) and at Aosta (bottom panel). Vertical lines represent the date of the calibration campaigns.




403

404 *Figure 5. Time plot of the differences between BPS and O3Brewer individual ozone data at Rome (upper panel) and*
 405 *at Aosta (bottom panel).*

406

407 *We took into consideration the spectral sensitivity of both Brewer instruments through the*
 408 *R6 ratio time behaviour (Fig. 6). In the same figure it is also plotted how each software ($R6_{BPS}$*
 409 *and $R6_{smooth}$) tracks changes in the spectral sensitivity of the instrument. $R6_{BPS}$ is obtained as the*
 410 *sum of BPS correction and $R6_{ref}$. $R6_{ref}$ values established during the calibration campaigns are*
 411 *also plotted. It is worth noticing that the number of standard lamp test per day is on average from*
 412 *4 to 6 at Rome, and from 2 to 4 in winter and from 8 to 10 in summer at Aosta and that only the*
 413 *daily means of BPS correction and $R6_{smooth}$ are stored. The latter is calculated if at least one*
 414 *standard lamp test is performed.*



415

416 **Figure 6.** Daily series of the ratios $R6$, $R6_{BPS}$ and $R6_{smooth}$ at Rome (upper panel) and at Aosta (bottom panel).
 417 Vertical lines represent $R6_{ref}$ established during each calibration campaign.


418

419 *Looking at R6 behaviour (Fig. 6 upper panel), it can be noticed that the sensitivity of the*
420 *instrument at Rome has changed mainly in two periods (between 1994 and 1995, and between*
421 *2006 and 2007). R6_{smooth} becomes a constant offset when the sensitivity of the instrument starts to*
422 *change. The cut off is not exactly equal to the threshold set in the configuration (in this case 500*
423 *units), but lower, because the filter looks 10 days before and 10 days after the date when SL R6 is*
424 *calculated. If the cut off remains constant, it means that the last calculated correction which*
425 *passes through rejection criteria, is taken into account, the same situation is experienced when*
426 *there is no valid SL test (Stanek personal communication, 2018). Consequently, the temporal*
427 *behaviour of R6_{smooth} during these time intervals appears as a plateau. In this case SL correction*
428 *is not applied since it is too high. Once a new calibration is performed (i.e. new references of R6*
429 *and the ETC are defined) R6 and R6_{smooth} show a similar behaviour again.*

430 Brewer 066 (Aosta) exhibited a better stability except for some R6 spikes (Fig. 6, bottom
431 panel) whereas R6_{smooth} time series shows a stable behaviour with respect to R6. R6_{BPS} shows a
432 similar behaviour to R6 at both stations due to the calculation method of the standard lamp
433 correction by the BPS.

434 A better visualization of the effect of the correction factor on TOCs is provided by plotting
435 the difference between the TOC daily means (BPS – O3Brewer) as a function of the difference
436 between R6_{BPS} and R6_{smooth} (Fig. 7). Large deviations between the two reprocessed TOC daily
437 means appear when there is a large difference between R6_{BPS} and R6_{smooth}. However large
438 differences occur even if R6_{BPS} does not differ too much from R6_{smooth} .

439



440

441

Figure 7. Differences between BPS and O3Brewer TOC daily means vs $R6_{BPS}-R6_{smooth}$ at Rome (upper panel) and at Aosta (bottom panel).

442

443

444

Three circumstances are here analysed when differences between BPS and O3Brewer ozone data exceed the value of the declared DS accuracy: 1. $R6_{BPS}$ lower than $R6_{smooth}$, 2. $R6_{BPS}$ higher than $R6_{smooth}$, 3. $R6_{BPS}$ similar to $R6_{smooth}$.

446

447I. $R6_{BPS}$ lower than $R6_{smooth}$.

448

Slight ozone difference took place when $R6_{BPS}$ was lower than $R6_{smooth}$ (at least 100 units), then the difference in ozone daily means was between -3% and 21% and in case of individual values from -3% up to 27 %, at Rome. At Aosta there was only one episode (2011/6/18) in which the O3Brewer daily mean differed about 30% from BPS. In that case, O3Brewer average was derived by three individual ozone values that showed the same difference with respect to the BPS ones. In this case, a large negative correction was applied to ozone values, thus generating a falsely high ozone case. The spike in the R6 value was originated by the two wrong SL test carried in that day caused perhaps by the micrometer in a wrong position, noisy communication, incorrect zenith drive position, or lamp aging. Consequently, the negative BPS correction generated high ozone values with a large standard deviation, whereas $R6_{smooth}$ was not applied to individual TOC data that resulted consistent with ozone values before and after that date.

459

At Rome the conditions in which $R6_{BPS}$ was lower than $R6_{smooth}$ occurred during the calibrations in 1995, 2006, 2007 and 2014. The discrepancy between the two codes could have been caused by the offset introduced by the way BPS determines the R6 reference value as for the other code the $R6_{ref}$ is obtained during the calibration campaign and set manually in the

460

461

462

463 configuration. The BPS $R6_{ref}$ is computed with a triangular smoothing filter of SL-test over the 15
464 day period after the calibration and it is calculated "on the fly" from daily mean SL values and it
465 is not stored (Fioletov, personal communication 2018).

466 To look into the possible effect of the BPS offset we estimated $R6_{ref_BPS}$, for each day over
467 the 15 days after the calibration by subtracting the correction (reported in the file o3data.txt) to
468 the corresponding $R6$ value. Then the average over the 15 $R6_{ref_BPS}$ values was compared with
469 $R6_{ref}$ (given by hand after the calibration). The estimated offset introduced by BPS with respect to
470 $R6_{ref}$ is very small, ranging between -19 to 6 units at Rome and between -10 to 2 units at Aosta.
471 Consequently, the BPS offset appears not to be responsible for the ozone difference that can be
472 attributed to the calculation method of the standard lamp correction.

473


474.2. $R6_{BPS}$ higher than $R6_{smooth}$

475 Large negative ozone differences occur when $R6_{BPS}$ is higher than $R6_{smooth}$ (at least >100
476 units). This causes a variation between the daily means generated by the codes from -5% till -50%
477 at Rome and from -51% till -91% at Aosta: Considering the individual values a mean percentage
478 difference between -3.1% and -57% was found at Rome, and of the same magnitude as that of
479 daily means at Aosta.

480 Two long periods were found at Rome belonging to this condition (29st October 1994 - 5th
481 May 1995; 26th June 2006 - 16th April 2007). The large drift in $R6$ turned out to be the
482 deterioration of the filter (NiSO₄/UG11) which was replaced during the calibration visits both in
483 1995 and 2007. In both cases it can be observed the cut off in $R6_{smooth}$ and hence the O3Brewer
484 recalculation provides uncommon TOC values. Then, we processed Rome ozone data using
485 O3Brewer by setting the SL maximal limit to higher value to assess whether the smooth correction
486 can properly process ozone data when large changes occurred in the instrumental response. The
487 SL maximal correction limit was to 3000 units keeping identical conditions for the air mass and
488 the standard deviation of the previous processing. In addition, a further ozone processing was
489 carried out by switching off the smoothing filter. Fig. 8 shows the time series of the ratios $R6$,
490 $R6_{BPS}$ and $R6_{smooth}$ (setting the limit to 3000 units) at Rome. It can be noticed that the $R6_{smooth}$ has
491 now similar behaviour as $R6_{BPS}$, nevertheless in some circumstances its behaviour is noisier than
492 both $R6_{smooth}$ (setting the limit to 500 units) and $R6_{BPS}$.

493 Fig.9 shows individual TOC data processed by O3Brewer 1) without applying the
494 smoothing filter, 2) with the SL limit correction set to 500 and 3) with the SL limit correction set
495 to 3000 units over the period of the $R6$ drift (2006-2007) at Rome. In the same figure, individual


496 *BPS recalculations without modifying the set up are also plotted. A better agreement with BPS*
 497 *ozone data is visible when ozone data were processed without the smoothing filter and with*
 498 *higher cut off in R6, however there are still anomalous ozone values due SL correction, whereas*
 499 *ozone values calculated without the correction seem not be not affected.*



500

501 *Figure 8. Daily series of the ratios $R6$, $R6_{BPS}$ and $R6_{smooth}$ at Rome. Vertical lines represent $R6_{ref}$ established during*
 502 *each calibration campaign.*

503



504

505 *Figure 9. Individual ozone values calculated by the BPS (black), O3Brewer without the filter smoothing correction*
 506 *(blue), O3Brewer with the cut off at 500 units (red), O3Brewer with the cut off at 3000 units (green) over the period*
 507 *of the drift in 2006 - 2007 at Rome.*


508

509 *The* occasional anomalous R6 ratios occurred at Aosta, most of them in 2011 and at the
510 beginning of 2012. Wrong wavelength selection by the micrometer, communication problems or
511 incorrect zenith drive position in relation to the lamp could have caused the R6 spikes. In this case
512 the algorithm of O3Brewer (with the cut off at 500 units) did not follow the abrupt change. The
513 correction was not applied resulting in large over- or under-estimation of TOC or with uncertain
514 data quality.

515

5163. *R6_{BPS} similar to R6_{smooth}*

517 *A different number of observations taken into account in the determination of the daily*
518 *means by the two codes can generate significant differences in some cases. The total number of*
519 *individual calculated total ozone values by O3Brewer is 104666 at Rome and 50088 at Aosta, the*
520 *number of those calculated by BPS is 100352 at Rome and 46617 at Aosta. Fig. 10 shows the*
521 *number of individual ozone values calculated by O3Brewer which is, in some days s, higher than*
522 *that of BPS.*




523

524 **Figure 10.** Time plot of the number of individual ozone values per day.

525

526 *Such difference can be due to the fact that there are no filter conditions on the minimum*
527 *and the maximum ozone values calculated by O3Brewer. Consequently, the daily means*
528 *generated by this software are determined including anomalous values. The case of R6_{BPS} similar*
529 *to R6_{smooth} responsible for significant ozone differences in the daily means (>5%) falls in this*
530 *conditions.*

531 As a specific example of the above case, we showed individual ozone values generated by
 532 both codes on 23/06/2001 at Rome with a daily average of 335 DU for BPS and 375.4 DU for
 533 O3Brewer (Fig.11, upper panel). The high individual ozone value generated by O3Brewer (618.7
 534 DU) affecting the daily average is clearly visible. Another example is provided for Aosta (Fig. 11,
 535 lower panel). On 5/1/2010 the daily average is 323.5 DU for BPS whereas it is 208.4 DU for
 536 O3Brewer. It is found that very low ozone values generated by O3Brewer, not discarded in the
 537 determination of the daily means, affect the quality of its value.



538

539

540 **Figure 11.** Individual TOC values generated by BPS and O3Brewer on 23/06/2001 at Rome (upper panel) and on
 541 5/1/2010 at Aosta (bottom panel) taken as examples where large difference between occurred although the SL
 542 correction is similar. Horizontal lines (dashed for BPS; solid for O3Brewer) represent the daily average (avg).

543

544


545 Table 3 shows the statistical comparison between O3Brewer (with cut off at 500 units) and
 546 BPS individual data and daily means, after data belonging to three groups described in the
 547 previous section, have been discarded. TOC data without R6 values (no SL test was performed in
 548 that day) were also discarded. The temporal behaviour of the differences between O3Brewer
 549 and BPS individual calculated ozone values, are plotted in Figure 12 showing a variability
 550 in general within ± 25 DU at Rome and ± 10 DU at Aosta.

551 A good overall agreement is found both on individual values and daily means when
 552 data belonging to the above conditions were removed, the correlation is close the unity at
 553 both stations; MPE is not significant on both individual values and daily means at Rome as
 554 well as at Aosta.

555 **Table 3.** Summary of the statistics O3Brewer vs BPS at both sites (*N*= number of data; *RHO*= Spearman
 556 correlation; *MB* =Mean Bias, *MPE*=Mean Percentage Error, *RMSE* =Root Mean Square Error, the
 557 uncertainty of *MB* and *MPE* is characterized by the standard deviation).
 558
 559

<i>O3Brewer_vs_BPS</i>	<i>N</i>	<i>RHO</i>	<i>MB (DU)</i>	<i>MPE (%)</i>	<i>RMSE (DU)</i>
Rome					
<i>Individual values</i>	89273	0.997	-0.6±2.1	-0.2±0.7	2.18
<i>Daily averages</i>	6304	0.997	-0.8±2.4	-0.2±0.7	2.47
Aosta					
<i>Individual values</i>	44117	0.999	0.1±0.8	0.03±0.30	0.83
<i>Daily averages</i>	2381	0.999	0.004±1.700	0.001±0.600	1.70

560



561

562 **Figure 12.** Difference between individual TOC values generated by BPS and O3Brewer at Rome (upper panel) and at
 563 Aosta (bottom panel) when anomalous values were discarded.
 564

565 3.2 Comparison of BPS and O3Brewer TOC retrievals with EUBREWNET data

566 The TOC individual values and daily means retrieved by O3Brewer and BPS data were
 567 compared with those derived from EUBREWNET retrievals. *The analysis was performed by*
 568 *removing data belonging to the three periods mentioned in Section 3.1 from all series.*


569 Table 4 shows the statistical results of the two processed TOC data sets against the
 570 EUBREWNET data set. It was found that the difference among the TOC retrievals is less than
 571 1%.

572 **Table 4.** Summary of the statistics O3Brewer vs BPS at both sites (N= number of data; RHO= Spearman
 573 correlation; MB =Mean Bias, MPE=Mean Percentage Error, RMSE =Root Mean Square Error, the
 574 uncertainty of MB and MPE is characterized by the standard deviation).
 575
 576


	N	RHO	MB (DU)	MPE (%)	RMSE (DU)
O3Brewer vs EUBREWNET					
Rome					
Individual values	38227	0.996	-0.2±3.8	- 0.05±1.00	3.80
Daily averages	2972	0.996	-0.1±4.6	- 0.02±1.20	4.60
Aosta					
Individual values	35746	0.997	0.3±5.3	0.2±2.4	5.33
Daily averages	2186	0.994	0.5±7.6	0.2±3.2	7.76
BPS vs EUBREWNET					
Rome					
Individual values	38227	0.995	1.0±4.1	0.3±1.1	4.27
Daily averages	2972	0.995	1.2±5.0	0.4±1.3	5.11
Aosta					
Individual values	35746	0.997	0.2±5.3	0.1±2.4	5.34
Daily averages	2186	0.994	0.5±7.6	0.2±3.2	7.59

577
 578 *However looking at Figs. 13-14 the differences between the individual ozone values*
 579 *calculated by BPS and EUBREWNET (Fig.13) and, by O3Brewer and EUBREWNET (Fig.14) are*
 580 *in some cases relevant. It seems that problems of the standard lamp values not properly filtered*
 581 *by the currently applied 7-days window smoothing, generate results less reliable (see the*
 582 *temporal behaviour of R6_{EUBREWNET} in Fig.15). This problem could be solved in the level 2 data, in*
 583 *which the setting a filter in the R6 values and smoothing the R6 time series is planned to be taken*


584 into account in the EUBREWNET algorithm (Fountoulakis, personal communication 2018).
585 However, although these options exist in the configuration form they are still inactive.
586



587 **Figure 13.** Difference between individual TOC values generated by BPS and EUBREWNET (Rome upper panel and
588 Aosta lower panel).
589
590



591
592 **Figure 14.** Difference between individual TOC values generated by O3Brewer and EUBREWNET (Rome upper panel
593 and Aosta lower panels).




594
 595 **Figure 15.** Daily series of the ratios $R6$, $R6_{EUBREWNET}$ at Rome (upper panel) and at Aosta (lower panel). Periods with
 596 the $R6$ drift or spikes were removed. Vertical lines represent $R6_{ref}$ established during each calibration campaign.
 597

598
 599 **3.3 Comparison of BPS, O3Brewer and EUBREWNET TOC retrievals with OMI data**

600 OMI overpasses were also compared with the processed Brewer TOC retrievals. The
 601 comparison was performed taking into account the same design criteria described in the previous
 602 section. The scatterplots of OMI vs Brewer data are shown in Fig. 16. However depending on the
 603 Brewer processing software a different behaviour is visible, even when only “good” data are
 604 considered. *It can be observed that EUBREWNET data show larger deviations from the bisectrix*
 605 *with respect to the other retrievals.*

606 The results of the statistical analysis are summarized in Table 5. The results of the
 607 statistical analysis are summarized in Table 5. *In general, the scaled correlation is, for both sites,*
 608 *on average $RHOs = 0.8$ which represents how the series are well connected in the short term.*

609 OMI products show a systematic underestimation with respect to ground-based data. *At*
 610 *Rome satellite data are less than 1 % for both O3Brewer and EUBREWNET whereas at Aosta*
 611 *about 2.5%; 1.2% (Rome) and 2.5% (Aosta) in the case of BPS data.* These results are in
 612 agreement with previous studies on validation of the OMI total ozone column by Brewer
 613 spectrophotometry conducted at the same latitudes (Ialongo et al., 2008; Anton et al., 2009).



614
615
616
617
618
619
620
621
622

Figure 16. Scatterplots OMI versus Brewer total ozone column at Rome (upper panel) and Aosta (lower panel). The solid line represents the bisectrix, The comparison is carried out with O3Brewer (green), EUBREWNET (blue) and BPS (red) data.

Table 5. Summary of the statistics of the comparison between OMI versus BPS, O3Brewer and EUBREWNET (N= number of data; **RHOs**= Spearman scaled correlation; MB =Mean Bias, MPE=Mean Percentage Error, RMSE =Root Mean Square Error, the uncertainty of MB and MPE is characterized by the standard deviation).

Rome	N	RHOs	MB (DU)	MPE (%)	RMSE (DU)	Aosta	N	RHOs	MB (DU))	MPE (%)	RMSE (DU)
OMI vs BPS											
	2622	0.841	-4.0±7.8	-1.2±2.3	8.63		2022	0.9	-8.6±10.4	-2.5±4.4	13.45
OMI vs O3Brewer											
	2622	0.843	-2.8±8.4	-0.8±2.5	8.85		2022	0.882	-8.6±10.7	-2.5±4.8	13.74
OMI vs EUBREWNET											
	2522	0.814	-2.8±9.6	-0.8±-2.7	9.99		1849	0.835	-8.2±10.5	-2.4±3.5	13.30

623
624
625
626
627
628
629

When comparing RMSE values it can be noticed that RMSE at Rome is lower than that found at Aosta, which supports the observed scatter plot shown in Fig. 16.

Besides, systematic differences between ozone estimated from OMI and from Brewer at Aosta could be related to the ground pixel size which can affect ozone amounts probed by the satellite, due to the complex orography of the valley.

630
631

3.4 Comparison among the trends estimated by the three processing software ozone retrievals

632 *The detected trends in ozone series calculated by using the three processing software are*
633 *reported in Table 6. The trends were quantified over the period 2005-2015 for Rome to be*
634 *consistent with the EUBREWNET ozone data coverage, and 2007 -2015 for Aosta. Ozone data*
635 *showing large differences among the codes, were not included in the trend analysis.*

636 *The QBO and solar cycle effects were not filtered in the ozone series. The former was*
637 *found small at mid-latitude stations (Fountoulakis et al., 2016), whereas the latter was not taken*
638 *into account due the short length of the analysed ozone series (< 11 years). All trends were found*
639 *to be not statistically significant (p-value is 0.05).*

640 *It is clear from Table 6 that there are not significant differences in the trends expressed in*
641 *terms of percentage variation per decade among the three codes, when data affected by rapidly*
642 *changes in R6 or the spectral response of the instrument shows a persistent drift, were removed.*

643

644 **Table 6.** *The total ozone linear trends derived by the processed ozone values using three different processing codes*

645

	<i>period</i>	<i>BPS</i> <i>(% per decade)</i>	<i>O3Brewer</i> <i>(% per decade)</i>	<i>EUBREWNET</i> <i>(% per decade)</i>
<i>Rome</i>	<i>2005-2015</i>	<i>-0.23 ± 0.18</i>	<i>-0.32 ± 0.20</i>	<i>-0.34 ± 0.21</i>
<i>Aosta</i>	<i>2007-2015</i>	<i>0.07 ± 0.35</i>	<i>0.04 ± 0.34</i>	<i>0.00 ± 0.38</i>

646

647 **4. Conclusions**

648

649 This study analyzed the total ozone column recalculations at Rome and Aosta using three
650 different software packages. We found that large differences in total ozone column retrievals can
651 be experienced when the instrumental sensitivity exhibits *a fast and dramatic drift between two*
652 *consecutive calibrations* or spikes. These conditions can affect TOCs retrievals due to the
653 algorithm of the standard lamp correction applied.

654 *When R6 exceed the default value of the cut off (500 units) set in the configuration of the*
655 *O3Brewer software during an occasional spike, the correction is not applied, whereas the BPS*
656 *correction does. This could generate false high/low ozone values. In latest version of O3Brewer it*
657 *is possible to set the cut off to higher value that is useful when there a large R6 drift is*
658 *experienced. However, anomalous ozone values can be still observed, since in O3Brewer there*
659 *are no filter conditions on the minimum and the maximum ozone values. Similarly, the current*

660 Level 1.5 in the EUBREWNET can produce erroneous ozone recalculations when anomalous R6
661 values are experienced. The issue is expected to be solved in Level 2.0 products, when they will
662 be released. The BPS ozone recalculations seem to be less affected in the case of R6 drift.

663 However when serious changes in the spectral sensitivity of instrument is experienced, a
664 solution consists in dividing the periods of R6 drifts into shorter time intervals and for that period
665 a new set of constants ($R6_{ref}$ and ETC) could be established by the user as the averages of R6
666 ratios in that time interval. This process (“synthetic calibration”) allows the user to introduce
667 standard lamp corrections larger than the software hardcoded thresholds. In any case the synthetic
668 constants in use must be confirmed at the next calibration with the reference instrument.

669 *Here we decided to discard the periods with drifts or occasional abrupt changes in R6,*
670 *and a good overall agreement is found between BPS, O3Brewer and EUBREWNET (MPE about*
671 *<1%).* However a spread among the **EUBREWNET individual ozone values** and those retrieved
672 by the other twos codes is still found, probably due to the standard lamp values not filtered
673 properly by the currently applied 7-day window smoothing, generating results less reliable.

674 The analysis of the differences between recalculated TOCs and OMI overpasses showed
675 that the latter dataset underestimate less than 2% ground –based total ozone columns at Rome and
676 less than 3% at Aosta (using “good” cases). Yet, the estimate of the trends using the retrievals
677 from the three different codes resulted not be affected when ozone data with anomalous R6 values
678 are removed.

679 The operators should constantly monitor the sensitivity of the instrument and know
680 carefully the processing software used to recalculate the total ozone. This means that the quality-
681 controlled data cannot be assured only by automatic data rejection rules of the adopted software,
682 but a rigorous manual data inspection is always necessary to prevent inconsistent data produced
683 by the processing software package in use.

684 *As a final remark, it is important to underline that for sake of consistency and*
685 *comparability between the results from different stations which send ozone products to*
686 *international data centres such as WOUDC or others, it is important to know the processing*
687 *software used to generate individual ozone values, the time behaviour of the instrumental*

688 *stability, the method applied for the standard lamp correction as well as the adopted rejection*
689 *criteria to determine the daily means.*

690

691 **Data availability.** The data used for the present study can be asked to the authors of the present
692 paper.

693 **Competing interests.** The authors declare that they have no conflict of interest.

694

695 **Acknowledgments:** We thank the European Brewer Network (<http://rbcce.aemet.es/eubrewnet/>)
696 for providing access to the data used in this investigation, and the COST Action ES1207 “A
697 European Brewer Network (EUBREWNET)”, supported by COST (European Cooperation in
698 Science and Technology). We also thank NASA Goddard Space Flight Center for OMI data
699 available (<https://avdc.gsfc.nasa.gov/>).

700 The authors are grateful to M. Stanek, V. Fioletov, A. Ogyu and I. Fountoulakis for their helpful
701 clarifications on the processing software.

702 The authors thank the anonymous reviewers for their valuable suggestions and comments.

703 This paper is dedicated in memory of Ken Lamb, founder of International Ozone Services Inc.
704 (IOS), who zealously delivered accurate ozone and UV calibrations to the worldwide Brewer
705 community.

706

707 **Author Contributions:** All authors have helped to develop the paper. A.M. S played the major
708 role supervising and coordinating the whole work; G.R. C. and H. D. have equally provided
709 helpful comments on the draft. F.F, F. S. and A. R. have contributed in the elaboration of the
710 Brewer and satellite data. A.M. S and G.R. C. are responsible for establishing and maintaining
711 Brewer 067; H. D. has contributed with data of Brewer 066 and in establishing and maintaining
712 the site; F.F and M. P. have given Matlab support; V. S. has given support with the Brewer
713 processing software.

714

715 **References**

716 Antón, M., López, M., Vilaplana, J. M., Kroon, M., McPeters, R., Bañón, M., and Serrano, A.:
717 Validation of OMI-TOMS and OMI-DOAS total ozone column using five Brewer

718 spectroradiometers at the Iberian Peninsula, J. Geophys. Res.-Atmos., 114, D14307,
719 doi:10.1029/2009JD012003, 2009.

720

721 Balis, D. Kroon, M., Koukouli, M. E., Brinksma, E. J., Labow, G., Veefkind, J. P., and McPeters,
722 R. D.: Validation of Ozone Monitoring Instrument total ozone column measurements using
723 Brewer and Dobson spectrophotometer ground-based observations, J. Geophys. Res., 112,
724 D24S46, doi:10.1029/2007JD008796, 2007.

725

726 Bass, A.M., and Paur, R.J.: The ultraviolet cross-sections of ozone, I, The measurements, in
727 atmospheric ozone. In: Zerefos CS, Ghazi A, Reidel D (eds) Proceedings of the Quadrennial
728 Ozone Symposium, Halkidiki, Greece, 1984. Dordrecht, Holland,606- 610,1985.

729

730 Bhartia, K. and Wellemeyer, C.:Toms-v8 total O3 algorithm, OMI Algorithm Theoretical Basis
731 Document, 2, 15–31, 2002.

732

733 Bordi, I., Fraedrich K., Sutera A., Zhu X.: On the climate response to zero ozone. Theor. Appl.
734 Climatol. 109, 253 – 259, 2012.

735

736 Brewer, A.W.: A replacement for the Dobson spectrophotometer? Pure App.Geophys.106-108,
737 919 -927, 1973.

738

739 *Brewer, A.W. and Kerr, J. B.: Total ozone measurements in cloudy weather, Pure appl. Geophys.*
740 *106-108, 928-937, 1973.*

741

742 EUBREWNET (A European Brewer Network) COST Action ES 1207
743 (<http://rbce.aemet.es/eubrewnet>, <http://www.eubrewnet.org/cost1207>, last accessed on June
744 2017; <http://www.eubrewnet.org/cost1207/> last accessed on November 2017).

745

746 Dessler, A.: Chemistry and Physics of Stratospheric Ozone, International Geophysics Series 74.
747 Academic press, London U.K., 2000.

748

749 *Diémoz, H., Eleftheratos, K., Kazadzis, S., Amiridis, V., Zerefos, C. S.: Retrieval of aerosol*
750 *optical depth in the visible range with a Brewer spectrophotometer in Athens, Atmos. Meas.*
751 *Tech., 9, 1871–1888, 2016*

752

753 Dobson G.M.B. and Harrison D.N.: Measurements of the amount of ozone in the Earth's
754 atmosphere and its relation to other geophysical conditions. Proc. R. Soc. London Ser. A,110,
755 660-692, 1926

756

757 Fioletov, V. E., Kerr, J. B., McElroy, C. T., Wardle, D. I., Savastiouk, V., and Grajnar, T. S.: The
758 Brewer reference triad, Geophys. Res. Lett., 32, doi:10.1029/2005GL024244, 2005.

759

760 Fioletov V.E. and Ogyu A.: Brewer Processing Software, [http://exp-](http://exp-studies.tor.ec.gc.ca/pub/Brewer%20Processing%20Software/brewer_processing_software.pdf)
761 [studies.tor.ec.gc.ca/pub/Brewer Processing Software/brewer processing software.pdf](http://exp-studies.tor.ec.gc.ca/pub/Brewer Processing Software/brewer processing software.pdf), last
762 accessed on 2007.

763

764 *Fountoulakis, I., Bais, A. F., Fragkos, K., Meleti, C., Tourpali, K., Zempila, M. M.: Short- and*
765 *long-term variability of spectral solar UV irradiance at Thessaloniki, Greece: effects of changes*
766 *in aerosols, total ozone and clouds, Atmos. Chem. Phys., 16, 2493–2505, 2016*

767

768 Ialongo, I., Casale G.R., and Siani A.M.: Comparison of total ozone and erythemal UV data from
769 OMI with ground-based measurements at Rome station, *Atmos. Chem. Phys.*, 8, 3283–3289, 2008.

770

771 International Ozone Service (IOS) <http://www.io3.ca/>, last accessed on September 2017

772

773 *Josefsson, W. A. P.: Focused sun observations using a Brewer ozone spectrophotometer, J.*
774 *Geophys. Res.,97(D14), 15813–15817, doi:10.1029/92JD01030,1992.*

775

776 Karppinen, T., Lakkala, K., Karhu, J.M., Heikkinen, P., Kivi, R., Kyrö, E.: Brewer spectrometer
777 total ozone column measurements in Sodankylä. *Geosci. Instrum. Method. Data Syst.*, 5, 229–239,
778 2016.

779

780 Karppinen T., Redondas, A., García, R.D., Lakkala, K., McElroy, C. T., Esko K.: Compensating
781 for the Effects of Stray Light in Single-Monochromator Brewer Spectrophotometer Ozone
782 Retrieval, *Atmosphere-Ocean* 53 (1), 66 – 73, 2015.

783

784 Kerr, J. B., McElroy, C. T., and Olafson, R. A.: Measurements of total ozone with the Brewer
785 spectrophotometer, in *Procs. of the Quadrennial Ozone Symposium*, edited by J. London, 74–79,
786 *Natl. Cent. for Atmos. Res., Boulder, Colo.*11., 1981.

787

788 *Kerr, J.B., McElroy C.T., Wardle D.I. and Dorokhov V.: Measurements of arctic total ozone*
789 *during the polar winter, Atmosphere-Ocean, 28:4, 383-392, 1990.*

790

791 Kerr, J. B. and Davis, J. M.: New methodology applied to deriving total ozone and other
792 atmospheric variables from global irradiance spectra, *J. Geophys. Res.*, 112, D21301,
793 doi:10.1029/2007JD008708, 2007.

794

795 Kerr, J. B.: The Brewer Spectrophotometer, in *UV Radiation in Global Climate Change*, edited by
796 W. Gao, D. Schmoltdt, and J. Slusser, Springer, Berlin, 160–191, 2010.

797

798 *Muthama N.J., Scimia U., Siani A.M., Palmieri S.: Toward optimizing Brewer zenith sky total*
799 *ozone measurements at the Italian stations of Rome and Ispra, J. Geophys. Res.,100, 3017-3022,*
800 *1995.*

801

802 Ramanathan, V., Dickinson R.E.: The role of stratospheric ozone in the zonal and seasonal
803 radiative energy balance of the earth-troposphere system, *J. Atmos. Sci.* 36:1084 – 1104, 1979.

804

805 Redondas A., Franco, J.R., Lopez-Solano, J., Carreno, V., León-Luis, S.F., Hernández-Cruz,
806 B.:The Regional Brewer Calibration Center - Europe: an overview of the X Brewer
807 Intercomparison Campaign, WMO Commission for Instruments and Methods of Observation,
808 TECO 2016.
809

810 Siani A.M, Casale, G.R., Galliani A.: Investigation on a low ozone episode at the end of
811 November 2000 and its effect on ultraviolet radiation, *Opt. Eng.* 41(12), 3082-3089, 2002.
812

813 Siani, A. M., Modesti, S., Casale, G.R., Diemoz, H., and A. Colosimo: Biologically effective
814 surface UV climatology at Rome and Aosta. *AIP Conf. Proc.* 1531, 903-906, 2013.
815

816 Stanek, M.: O3Brewer, <http://www.o3soft.eu/o3brewer.html>, last accessed on January, 2016.
817

818 Stübi, R., Schill, H., Klausen, J., Vuilleumier, L., and Ruffieux D.: Reproducibility of total ozone
819 column monitoring by the Arosa Brewer spectrophotometer triad, *Geophys. Res. Atmos.*, 122,
820 4735–4745, 2017.
821

822 Tzortziou, M., Herman, J. R., Cede, A., and Abuhassan, N.: High precision, absolute total column
823 ozone measurements from the Pandora spectrometer system: Comparisons with data from a
824 Brewer double monochromator and Aura OMI, *J. Geophys.,Res.*, 117, D16303,
825 doi:10.1029/2012JD017814, 2012.
826

827 Vanicek, K.: Differences between ground Dobson, Brewer and satellite TOMS-8, GOME-
828 WFDOAS total ozone observations at Hradec Kralove, Czech. *Atmos. Chem. Phys.*, 6, 5163–
829 5171, 2006.
830

831 Vaz Peres, L., Bencherif, H., Mbatha, N., Passaglia Schuch, A., Tohir, A. M., Bègue, N.,
832 Portafaix, T., Anabor, V., Pinheiro, D. K., Paes Leme, N. M., Bageston, J. V., Schuch, N. J.:
833 Measurements of the total ozone column using a Brewer spectrophotometer and TOMS and OMI
834 satellite instruments over the Southern Space Observatory in Brazil, *Ann. Geophys.*, 35, 25–37,
835 2017.
836

837 WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2014,
838 Global Ozone Research and Monitoring Project, Report No. 55, Geneva, Switzerland, 2015.
839

840 WMO-GAW Seventh Intercomparison Campaign of the Regional Brewer Calibration Center
841 Europe (RBCC-E) Lichtklimatisches Observatorium, Arosa, Switzerland, 16-27 July 2012, Report
842 n.216, 2015.
843

844 WOUDC (World Ozone and Ultraviolet Data Centre), <http://woudc.org>, last accessed on June
845 2017.
846