

Evaluation of OAFlux datasets based on in situ air-sea flux tower observations over the Yongxing Islands in 2016

Fenghua Zhou^{1,2}, Rongwang Zhang¹, Rui Shi¹, Ju Chen¹, Yunkai He¹, Dongxiao Wang^{1*}, Qiang Xie^{3,4*}

¹State Key Laboratory of Tropical Oceanography, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510300, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³Institute of Deep-sea Science and Engineering, Chinese Academy of Sciences, Sanya 572000, China

⁴Laboratory for Regional Oceanography and Numerical Modeling, Qingdao National Laboratory for Marine Science and Technology, Qiangdao, 266237, China

10 *Correspondence to: Dongxiao Wang (dxwang@scsio.ac.cn); Qiang Xie (gordonxie@idsse.ac.cn)

Abstract. The Yongxing air-sea flux tower (YXASFT), which was specially designed for air-sea boundary layer observations, was constructed on Yongxing Island in the South China Sea (SCS). Surface bulk variable measurements were collected during a one-year period from 2016/02/01 to 2017/01/31. The sensible heat flux (*SHF*) and latent heat flux (*LHF*) were further derived via the Coupled Ocean-Atmosphere Response Experiment version 3.0 (COARE3.0). This study employed the YXASFT in situ observations to evaluate the Woods Hole Oceanographic Institute (WHOI) OAFlux reanalysis data products.

First, the reliability of COARE3.0 data in the SCS was validated using direct turbulent heat flux measurements via an eddy covariance flux (ECF) system. The *LHF* data derived from COARE3.0 are highly consistent with the ECF with a coefficient of determination (R^2) of 0.78. Second, the overall reliabilities of the bulk OAFlux variables were diminished in order from T_a (air temperature), U (wind speed), Q_a (air humidity) to T_s (sea surface temperature) based on a combination of R^2 values and biases. OAFlux overestimates (underestimates) U (Q_a) throughout the year and provides better estimates of both variables in the winter and spring than in the summer_autumn period, which seems to be highly correlated with the monsoon climate in the SCS. The lowest R^2 between the OAFlux-estimated and YXASFT-observed T_s , indicating that T_s is the least reliable dataset and should thus be used with considerable caution. In terms of the heat fluxes, OAFlux considerably overestimates *LHF* with an ocean heat loss bias of 52 w/m² in the spring, and the seasonal OAFlux *LHF* performance is consistent with U and Q_a . The OAFlux-estimated *SHF* appears to be poorly representative with enormous overestimations in the spring and winter, while its performance is much better during the summer_autumn period. Third, analysis reveals that the biases in Q_a are the most dominant factor on the *LHF* biases in the spring and winter and that the biases in both Q_a and U are responsible for controlling the biases in *LHF* during the summer_autumn period. The biases in T_s are responsible for controlling the *SHF* biases, and the effects of biases in T_s on the biases in *SHF* during the spring and winter are much greater than that in the summer_autumn period.

1 Introduction

Exchanges of momentum, heat and water vapor fluxes at the air-sea interface constitute a significant component of air-sea interactions, which affect weather processes and climate change at all scales (Zhu et al., 2002; Persson et al., 2002; Frenger et al., 2013). As the surface that lies beneath the atmosphere, the ocean influences the stability of the atmospheric layer and the evolution of the atmospheric boundary layer through turbulent exchange (Chelton and Xie, 2010). In addition, sensible heat flux (*SHF*) and latent heat flux (*LHF*) at the air-sea interface are both important factors that affect changes in the mixing layer and thermocline (Hogg et al., 2009).

Accurate calculations of air-sea fluxes play a crucial role in driving marine and atmospheric circulation models, understanding atmosphere-ocean interactions, and evaluating and assessing numerical weather forecast models (Sun et al., 2003). Currently available air-sea flux datasets (including satellite remote sensing inversion data and reanalysis data) are quite uncertain, as they are mainly derived from inaccurate flux modeling algorithms and uncertainties in the turbulent exchange coefficient were also involved in the fluxes calculations (Zeng et al., 1998; Josey, 2001; Smith et al., 2001). In turn, these intrinsic uncertainties limit the ability to assess numerical models based on reanalysis flux datasets (Yu et al., 2006).

The South China Sea (SCS) is mainly controlled by various monsoon systems, it is connected with the western Pacific Ocean and the Indian Ocean through marine and atmospheric processes, and thus, the SCS exhibits potential influences on global climate change as well as regional climate regimes (Wang et al., 2006; Shi et al., 2015). Air-sea interactions in the SCS induce many marine meteorological hazards and greatly affect the transfer of heat and water vapor in regions throughout South China and Southeast Asia (Yang et al., 2015). Acquiring long-term observations of air-sea fluxes in the SCS can therefore help us to better understand the characteristics and evolutionary behavior of air-sea interactions in the SCS, optimize the parameterization schemes in atmospheric models, and improve long-term weather forecasts and extreme hazardous weather alerts.

To achieve the abovementioned scientific goals, a mesoscale observation network in the Xisha sea area in the northern SCS was initiated in 2008 (Yang et al., 2015) with the primary ambition of researching air-sea interactions. At present, the observation network includes a surface mooring buoy array, a system of shore-based wave-tide gauges, an automatic weather station, a shore-based boundary layer air-sea flux tower, and a submerged mooring buoy array. A large dataset comprising in situ observational data was obtained to serve as “ground truth” reference data to quantify the uncertainties within regional model flux products for the SCS.

Many in situ observations and model analysis comparisons have been studied in different oceans around the world, including the Arabian Sea (Weller et al., 1998; Swain et al., 2009), the tropical Pacific Ocean (Weller and Anderson, 1996; Wang and McPhaden, 2001), the northeast Atlantic Ocean (Sun et al., 2003; Yu et al., 2004), the Indian Ocean (Goswami, 2003) and the SCS (Zeng et al., 2009; Wang et al., 2013). Unfortunately, due to limited field observations of flux-related variables, detailed evaluation studies in the SCS are scarce.

In this study, turbulent *SHF* and *LHF* variations as well as numerous bulk variables, including the air temperature (T_a), sea surface temperature (T_s), air humidity (Q_a) and wind speed (U), from the Woods Hole Oceanographic Institution (WHOI) Objectively Analyzed Air-Sea Fluxes (OAFlux) project are compared with YXASFT measurements in the northern SCS. This investigation spans a full year from 2016/02/01 to 2017/01/31. Seasonal comparisons of the bulk variables and heat fluxes are described in **Sect. 3**. An overview of the instrumentation on the Yongxing air-sea flux tower (YXASFT) in addition to the data and methodology employed in this paper are introduced in **Sect. 2**. Finally, the summary and conclusions are provided in **Sect. 4**.

2 Instrumentation, data and methods

2.1 Yongxing air-sea flux tower (YXASFT)

The 20-m-tall YXASFT (**Fig. 1a**), which was specially designed for the observation of air-sea boundary layer fluxes, is located approximately 100 m off the northeastern coastline of Yongxing Island ($16.84^\circ N$, $112.33^\circ E$; **Fig.1d and 1e**). A gradient meteorological system (GMS) and an eddy covariance flux (ECF) system were mounted on the tower (**Fig. 1b**). A CR3000 data logger manufactured by Campbell Scientific Company, USA, is used for data sampling, preprocessing, storage, and transmission. The real-time observation data from the YXASFT are open for access at the website <http://mabl.scso.ac.cn:8040> (login: CSL-CER and password: ruhuna). A data sharing agreement must be signed by the user before being authorized to download the data.

The sensor wiring and data acquisition diagram for the YXASFT is shown in **Fig. 2**. The observational variables within the GMS include U , the wind direction (W_d), T_a , Q_a , the air pressure (P_a), the net radiation (R_n) and T_s . Each parameter is sampled once every second, and 1-, 10- and 30-min averages are recorded and transmitted to the data center in real-time. The ECF system can collect high-frequency turbulent data with a 10-Hz sampling frequency. Successive 30-min fully corrected fluxes of the momentum (T_{au}), *SHF*, *LHF*, and CO_2 (F_c) can be calculated using the online program Easy_flux. The sensors in the YXASFT and their respective measurement specifications are listed in **Table 1**. All of the sensors (**Fig. 1c**) have been checked via pre- and post-installment calibrations by the National Center of Ocean Standards and Metrology.

2.2 Data

The data employed in this study originate from two sources. The in situ observations obtained by YXASFT, and the reanalysis datasets are derived from the OAFlux project. Table 2 shows various information, including the variable height, time period, interval and location, regarding the data adopted in this study.

2.2.1 In situ data

High-frequency turbulent data (u, v, w, t, ρ_v) were collected by the ECF system installed at a height of 12 m from 2016/02/01 to 2016/03/29. Direct measurements of turbulent data were further used to calculate the fluxes using the eddy covariance (EC) method in a specified time period (30 min or 60 min). Meanwhile, direct measurements of turbulent fluxes using the ECF system were used only to verify the applicability of version 3.0 of the Coupled Ocean-Atmosphere Response Experiment (COARE3.0) over the SCS.

The selected 30-min averages of the bulk variables (U measured at a height of 10 m and T_a, Q_a and T_s measured at a height of 5 m) used for the bulk flux calculations range from 02/01/2016 to 01/31/2017. Note that T_s was measured using an SI-112 infrared radiation thermometer manufactured by Campbell Scientific Company, USA, installed at a height of 5 m, and therefore, we consider T_s as representative of the sea surface temperature at a depth of 0.05 m. The value of Q_a was derived using **Eq. 1** as described in COARE3.0 using T_a , the relative humidity (R_h) and the air pressure (P_a). Furthermore, this paper also adopts SHF and LHF averages within 30-min intervals derived via COARE3.0 using the input observed bulk variables. The heights of T_a and Q_a in the OAFflux dataset are both 2 m, while the measurement heights for these two parameters on the YXASFT are both 5 m. Thus, prior to conducting a comparison, we corrected the corresponding heights of the in situ data to correspond to the heights in the OAFflux dataset using COARE3.0. In addition, downward longwave radiation (DLR) data measured using an NR01 net radiometer manufactured by Hukseflux were used in this paper as an indirect variable to infer the cloud cover in the sky.

$$e_s = 6.112e^{\frac{17.502T_a}{T_a+240.97}(1.0007+3.46P_a*10^{-6})}$$
$$Q_a = \frac{621.97e_s}{(P_a - 0.378)R_h / 100} \quad (1)$$

2.2.2 Reanalysis data

In this paper, the OAFflux reanalysis data were selected for two reasons. First, a previous study showed that the OAFflux dataset is the most preferable among five different products (i.e., ERA-1, NCEPS, JRA55, TropFlux and OAFflux) with regard to LHF data over the SCS (Wang et al., 2017; Zhang et al., 2018). Second, OAFflux represents the most recently updated data product (as of July 2017) accessible for the study period. OAFflux is an ongoing global flux product compiled by WHOI with a spatial resolution of $1^\circ \times 1^\circ$. OAFflux utilizes an integrated analysis method to combine satellite data with modeling and reanalysis data, and it employs COARE3.0 to calculate heat fluxes (Yu et al., 2008). In this study, the daily mean OAFflux datasets include U, Q_a, T_s, T_a, LHF and SHF , and YXASFT observations during the same time period were used for a comparison.

2.3 Methods

2.3.1 Bulk algorithm

The bulk algorithm utilized in this study is based on the Monin-Obukhov similarity theory, which is widely considered to be an advanced bulk algorithm (Fairall et al., 1996). To keep consistent with the bulk method of calculating flux in OAFflux, the COARE3.0 was used to calculate the heat fluxes using the in-situ observation bulk parameters in this paper. Compared to COARE2.5, the updated COARE3.0 has some noted improvements as follows. First, the range of wind speed validity is now extended to 0–20 m/s after modifying roughness representation. Second, the COARE 3.0 is shown to be accurate within 5% for wind speeds of 0–10 m/s and 10% for wind speeds between 10 and 20 m/s (Fairall et al., 2003). In this method, the calculation equations for the SHF and LHF can be written as follows:

$$SHF = \rho_a C_p C_h U (T_s - T_a) \quad (2)$$

$$LHF = \rho_a L_e C_e U (Q_s - Q_a) \quad (3)$$

where ρ_a represents the air density, L_e represents the latent heat of evaporation, C_p represents the constant-pressure specific heat, U represents the sea surface wind speed (measured at a height of 10 m in this study), C_e and C_h correspond to the turbulence exchange coefficients for the latent heat and sensible heat, respectively, Q_s and Q_a correspond to the air saturation specific humidity at the sea surface and the air specific humidity near the sea surface, respectively, and T_s and T_a correspond to the sea surface skin temperature and the air temperature near the sea surface, respectively. In **Eqs. 2** and **3**, only U , T_s , T_a and Q_a are independent measurement variables, while the remainder of the variables must be calculated based on the four independent variables.

2.3.2 Eddy covariance method

The EC method is one of the most direct ways to measure and calculate turbulent fluxes (Crawford et al., 1993). Reynolds decomposition is utilized to break raw data down into their means and deviations. Furthermore, the values of SHF and LHF can be calculated as the covariance between w and scalar values (t , ρ_v) using the following formulas, respectively:

$$SHF = \rho C_p \overline{w't'} \quad (4)$$

$$LHF = \lambda \overline{w'\rho'_v} \quad (5)$$

where ρ is the dry air density, C_p is the specific heat of dry air at a constant pressure (where $1004.67 \text{ Jkg}^{-1}\text{K}^{-1}$ is used in the calculation), and λ is the latent heat ratio of water vapor evaporation. The overbar represents the Reynolds–ensemble average, and the prime symbol denotes the instantaneous deviation from the ensemble average.

2.3.3 Data processing

To match the timescale of the OAFflux daily data, we derived the daily means of the YXASFT-observed bulk variables and heat fluxes by averaging all of the 30-min datasets from each day. In addition, we used bilinearly interpolated OAFflux values (inversely weighted by the distance) from the surrounding four grid points ($111.5^\circ E, 16.5^\circ N$; $112.5^\circ E, 16.5^\circ N$; $112.5^\circ E, 15.5^\circ N$; $111.5^\circ E, 15.5^\circ N$) to represent the corresponding OAFflux value at the YXASFT observation site.

The comparison between the YXASFT and OAFflux datasets (described in **Sects. 3** and **4**) was quantitatively analyzed by using the mean bias (*Bias*, defined in **Eq. 6**), root mean squared error (*RMSE*, defined in **Eq. 7**), coefficient of determination (R^2) and linear regressions, respectively.

$$Bias = \frac{1}{N} \sum_{i=1}^N (x_i - y_i) \quad (6)$$

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - y_i)^2} \quad (7)$$

where x and y denote the OAFflux values and YXASFT observations, respectively.

3 Results and discussion

3.1 Validation of COARE3.0 using direct ECF measurements

The heat fluxes from both YXASFT and OAFflux used for the comparison herein were derived from COARE3.0. However, the COARE algorithm was originally developed for the Tropical Ocean Global Atmosphere-COARE (TOGA-COARE) experiment in tropical oceans (Fairall et al., 1996), while the reliability of COARE3.0 was verified by (Brunke et al., 2003) using 12 ship cruises over tropical and mid-latitude oceans (between $5^\circ S$ and $60^\circ N$). The adaptability of OAFflux in the SCS must be verified due to its unique geographical location (i.e., it is the largest marginal sea in the northwestern Pacific Ocean) and its monsoon climate system. In this study, the EC fluxes directly measured by the IRGASON ECF system were used to validate the performance of COARE3.0 in the SCS. The EC method is mathematically complex, significant care is required to set up different processing steps for different sites, measurements and study purposes. In this paper, the EC program running on CR3000 was based on the processing steps shown in **Fig. 3**. The daily *LHF* time series in COARE3.0 are basically consistent with those in ECF (**Fig. 4a**) with an R^2 value of 0.78 (**Fig. 4c**). COARE3.0 underestimates the *LHF* with a mean bias of 18.55 w/m^2 . A larger difference in the *LHF* measurement occurs when relatively larger *LHF* values are observed (e.g., 2016/02/07 and 2016/02/25), which can be readily observed in **Fig. 4a**. The precipitation on these days is the most likely explanation for the overestimation in the *LHF* by the ECF system (Mauder et al., 2006; Zhang et al., 2006). Although the YXASFT possesses a lack of field precipitation observations, we can speculate that precipitation may have occurred on 2016/02/07 based on a $1.8^\circ C$ drop in the air temperature and an increase of 13% in the relative humidity within the daily mean. In addition, we spot similar trends on 2016/02/25. In contrast, the *SHF* data pair is far from agreement with

an R^2 value of 0.03 (**Fig. 4d**). The large variation in the SHF observed using the ECF is not detected within the COARE3.0-derived time series (**Fig. 4b**). Direct heat flux measurements with a 60-day interval obtained using the ECF system show that SHF (with a mean of 23.5 w/m^2) is significantly smaller than LHF (with a mean of 93.3 w/m^2). A small SHF magnitude may amplify variations in the time series and reduce the R^2 values in scatter plots under the same deviation values. In this comparison, we were more concerned about the magnitude of correlation in the LHF data. Thus, COARE3.0 was considered to be receptive and was used as an appropriate bulk flux algorithm over the SCS.

3.2 Evaluation of the OAFflux datasets

OAFflux is a flux product based on a composite algorithm that improves the calculation accuracies of flux-related variables by using a weighting method for target analysis. However, this method could lead to a time-scale mismatch if the data variables have different data sources (Fairall et al., 2010). It is therefore necessary to evaluate the OAFflux dataset to assess its applicability in the SCS before further application.

3.2.1 Time series of the YXASFT observations and OAFflux reanalysis data

Time series of the bulk variables and heat fluxes are given in **Figs. 5** and **6**, respectively. As shown in **Fig. 6**, there is an obvious overestimation in both SHF and LHF in OAFflux compared with the YXASFT observations, and this overestimation demonstrates an evident seasonal variation. The time series of LHF from the YXASFT observations and OAFflux data show essentially consistent variation trends and agree with one another better during the spring (February to March) and winter (December to January) than during the summer and autumn (April to November) (**Fig. 6b**). The SHF variation trend appears to be opposite to that of LHF , since the deviations during the winter and spring are clearly larger than those during the summer and autumn (**Fig. 6a**). For the bulk variables in **Fig. 5**, the OAFflux data maintained a higher consistency with the YXASFT observations with regard to the overall variation trend. Furthermore, U and Q_a seemed to match better during the winter and spring periods, while an overestimation (underestimation) in U (Q_a) is more evident during the summer and autumn periods (**Fig. 5a** and **5b**). Some abrupt drops (i.e., variations of 3 to 5 days) in the YXASFT T_s observations were obviously not captured by OAFflux (**Fig. 5d**). In the next section, we divide the annual study period into three periods, namely, spring (2016/02/01-2016/03/31), summer_autumn (2016/04/01-2016/11/31) and winter (2016/12/01-2017/01/31), to conduct a detailed comparison of their seasonal variations.

3.2.2 Comparison of the bulk variables

The heat fluxes from both OAFflux and YXASFT were derived using COARE3.0. Thus, we can further analyze the origin of the seasonal deviations in the heat fluxes by conducting seasonal comparisons of the bulk variables. The scatter plots of U , Q_a , T_s , and T_a constructed using the YXASFT and OAFflux data for the three separate periods are shown in **Fig. 7**, and a quantitative statistical summary for each variable is listed in **Table 3**.

U : The spring, summer_autumn, and winter periods in the Yongxing Islands represent the monsoon transition, southwest monsoon and northeast monsoon periods, respectively. Previous studies indicated that the northeast monsoon in the northern

SCS is much stronger than the southwest monsoon (Yan et al., 2005). In this study, the observed mean wind speeds during the three periods were 6.40, 4.97 and 9.40 m/s. It can be seen from **Fig. 7 (first row)** that the R^2 values of U between the OAFflux and YXASFT data during the three periods are 0.90, 0.79 and 0.92. OAFflux overestimates the values of U in the spring, summer_autumn, and winter periods with mean biases of 0.96 (15% of the YXASFT-observed mean value), 1.19
5 (24%) and 0.67 m/s (7%), respectively.

Q_a : The southwest monsoon is often accompanied by a high water vapor and cloudy skies (Chen et al., 2012). Therefore, the Q_a value during the summer_autumn period was the highest throughout the year with an observed mean of 21.08 g/kg. The R^2 values of Q_a between the OAFflux and YXASFT data during the three periods are 0.81, 0.68 and 0.80 (**Fig. 7 (second row)**). In contrast to U , OAFflux exhibits an overall underestimation of Q_a in the spring, summer_autumn, and winter periods
10 with dry biases of 0.33 (2%), 0.75 (4%) and 0.11 g/kg (1%), respectively.

T_a : The OAFflux T_a values are highly consistent with the YXASFT observations with R^2 values of 0.92, 0.84 and 0.89 in the spring, summer_autumn, and winter periods, respectively (**Fig. 7 (fourth row)**). As shown in **Fig. 5c**, both the seasonal trends and day-to-day variations are effectively captured in the OAFflux data. The OAFflux reanalyzed T_a data have a warmer bias of 0.52 °C (2%) in the spring and colder biases of 0.10 (0.3%) and 0.57 °C (2%) in the summer_autumn and winter
15 periods, respectively. Consequently, the OAFflux-estimated T_a can be considered as the most reliable variable in this study.

T_s : The OAFflux-estimated T_s captures only the seasonal trend, and the estimates exclude some special synoptic signals, such as abrupt drops during cold air temperatures and typhoons or gradual temperature increases induced by the passage of a warm eddy. The R^2 values of T_s between the OAFflux and YXASFT data are relatively small when compared with those of U , Q_a and T_a , suggesting that the reliability of the OAFflux-analyzed T_s is generally low. In contrast to U and Q_a , the OAFflux T_s
20 performance better in the summer_autumn period ($R^2=0.70$) than in the spring ($R^2=0.47$) and winter ($R^2=0.54$) periods, as shown in **Fig. 7 (third row)**.

In summary, the seasonal performances of the OAFflux-estimated U and Q_a seem to be highly correlated with the monsoon system in the SCS. This manifests a better performance of the OAFflux-estimated U (Q_a) during the spring and winter periods characterized by a stronger (drier) northeast monsoon than during the summer_autumn period characterized by a relatively
25 weaker (wetter) southwest monsoon. The significant difference between the T_s estimates may stem largely from the fact that the OAFflux T_s estimates are retrieved using the Advanced Very High Resolution Radiometer (AVHRR), which is easily affected by the presence of clouds. Therefore, the available OAFflux T_s estimates were dramatically reduced during the abovementioned special synoptic processes. With the onset of the southwest monsoon, the average total cloud cover, low cloud cover and precipitation all increase throughout the SCS (Yan et al., 2003), and the T_s retrieved via the AVHRR should
30 correspondingly exhibit a lower quality. However, this trend is not observed in the results of this paper. We further utilized in situ observations of the *DLR* to infer the sky cloud cover. There is an evidently greater fluctuation in the *DLR* during the winter and spring periods than in the summer_autumn period, indicating that the winter and spring seasons possess greater probabilities of cloudy days (**Fig. 8**). This interesting phenomenon may be caused by the fact that the intensity of the summer monsoon in 2016 was weaker than those in preceding years; this hypothesis will be further explored hereafter.

3.2.3 Comparison of heat fluxes

The scatter plots of the *LHF* and *SHF* estimates obtained from the YXASFT and from OAFlux during the three periods are shown in **Fig. 9**, and a quantitative statistical summary of each variable is also listed in **Table 3**. Note that an upward (downward) heat flux is positive (negative) in this paper, and a positive (negative) value represents the loss (gain) of ocean heat to (from) the atmosphere.

LHF: Compared with the YXASFT observations, the OAFlux-estimated *LHF* is overestimated by a mean bias of 50.95 (70%) in the spring, 42.43 (76%) in the summer_autumn and 63.29 w/m^2 (74%) in the winter. The R^2 values are 0.80 in the spring, 0.66 in the winter and 0.40 in the summer_autumn (**Fig. 9 (first row)**). This is also consistent with the R^2 values for U and Q_a , which are the two key input factors in the *LHF* calculations.

SHF: Large *SHF* variations during the spring and winter are not evident in the YXASFT-derived *SHF* time series (**Fig. 5e**). Compared to *LHF*, the OAFlux-estimated *SHF* has the smallest R^2 values for all three individual periods, as shown in **Table 3** for the spring (0.01), summer_autumn (0.31) and winter (0.14). In comparison, the OAFlux-estimated *SHF* is more reliable during the summer_autumn with a mean bias of 1.07 w/m^2 than in the spring (16.83 w/m^2) or winter (23.56 w/m^2).

Overall, we can infer that the OAFlux-estimated *LHF* product is more reliable during the spring and winter periods than during the summer_autumn period, which is consistent with the key input variables U and Q_a , and that the product is further affected by the monsoon system in the SCS. Meanwhile, the *SHF* estimates exhibit opposite characteristics relative to those of *LHF*, as the OAFlux *SHF* product is more credible during the summer_autumn than during the spring and winter periods, which is consistent with the seasonal OAFlux T_s performance and is highly correlated with the cloud cover.

3.3 Possible effects of bulk variables on the biases in the *SHF* and *LHF*

The values of *SHF* and *LHF* were calculated using **Eqs. 2** and **3**. Thus, possible biases in the *LHF* and *SHF* results are mainly associated with the input bulk variables and the parameterization of the turbulent exchange coefficients in the equations. In this paper, the parameterization scheme is not discussed due to limited space. The relationships among the OAFlux *LHF* bias with U , Q_a and T_s were studied extensively by a previous study through years of moored buoy data, automatic weather station (AWS) data and cruise data over different regions in the SCS; it was found that the biases in Q_a dominated the *LHF* biases, followed by the biases in U (Wang et al., 2017). To determine whether similar conclusions exist in this study and to quantify the relationships among the heat flux biases and the bulk variable biases, we constructed scatter plots of the biases in *LHF* (**Fig. 10**) and *SHF* (**Fig. 11**) against the biases in U , Q_a , T_s and T_a . All of the biased data were normalized first to understand their relative importance.

ALHF: The biases in Q_a are the most dominant factor in determining the biases in *LHF* during the spring with relatively high R^2 values of 0.38 compared with the other biased bulk variables (**Fig. 10 (first column)**). Both of the Q_a and U biases are responsible for controlling the biases in *LHF* during the summer_autumn period with R^2 values of 0.36 and 0.32, respectively (**Fig. 10 (second column)**). Both of the Q_a and T_a biases are the dominate factors in determining the biased in *LHF* during the winter period with R^2 values of 0.43 and 0.16, respectively (**Fig. 9 (third column)**). The biases in T_s is negligible control

factors on the biases in *LHF*, since their R^2 values are all relatively small during the three periods compared with those of Q_a (**Fig. 9 (third and fourth rows)**). In general, the result revealed that the Q_a is the most dominated factor controlling the biases in *LHF* throughout the year is similar to those reported in previous studies (Wang et al., 2013, 2017). Additional, these dominate factors that cause the seasonal biases in *LHF* are new findings in this article. *ASHF*: During the observational period, the biases in T_s were the key factor dominating the biases in *SHF*. The effects of T_s biases on the biased *SHF* during the spring ($R^2=0.79$) and winter ($R^2=0.72$) periods were much larger than that during the summer_autumn period ($R^2=0.38$), which is also consistent with the fact that OAFflux better estimates T_s in the summer_autumn than in the spring and winter (**Fig. 7 third row**). From **Eq. 2**, *SHF* is largely determined by T_s-T_a , as shown in **Fig. 6**. OAFflux is unable (able) to capture the variations in T_s (T_a) during the spring and winter, thereby causing large fluctuations in T_s-T_a and further leading to large variabilities in the OAFflux *SHF* time series.

4 Summary and conclusions

Successive air-sea heat flux-related observational data were acquired over the course of a year (2016/02/01-2017/01/31) at the YXASFT in the Yongxing Islands. In this paper, we first used direct heat flux measurements from a high-frequency (10 Hz) ECF system to validate the reliability of the COARE3.0 bulk algorithm in the SCS. Then, seasonal comparisons were conducted for the daily mean surface bulk variables and heat fluxes between the WHOI OAFflux products and YXASFT observations. Finally, the effects of biased bulk variables on the biases in the heat fluxes were presented to determine the possible sources of the biases in *LHF* and *SHF*. The conclusions are summarized as follows.

The magnitude of the mean of the directly measured *SHF* is small compared with that of *LHF* and can even be ignored in air-sea heat flux interactions during the ECF measurement period. Therefore, we were more concerned with the *LHF* estimation differences between COARE3.0 and the ECF system in this validation. The daily mean *LHF* from COARE3.0 was basically consistent with the ECF measurements with a high R^2 and an acceptable bias. Furthermore, if possible precipitation periods were excluded, the consistency between the COARE3.0 and ECF *LHF* data were better. Thus, the COARE3.0 bulk algorithm was considered to be reliable in this study.

Comparisons of the bulk variables revealed that the reliabilities of the OAFflux datasets diminished in order from T_a , U , Q_a to T_s based on a combination of R^2 values and biases. The performances of the OAFflux-estimated U and Q_a seem to be highly correlated with the monsoon system in the SCS; OAFflux provides a better estimation of U (Q_a) in the spring and winter characterized by a stronger (drier) northeast monsoon than in the summer_autumn characterized by a relatively weaker (wetter) southwest monsoon. Similar to a previous study, this study also indicated that T_s is the least reliable OAFflux product (Sun et al., 2003). The T_s signals during special synoptic process were poorly captured by OAFflux due to the presence of clouds, which affect the recorded AVHRR data. The performance of the OAFflux-estimated T_s is better during the summer_autumn than in the winter or spring due to a reduced cloud cover during the summer monsoon period, which could be attributable to the fact that the summer monsoon in 2016 was weaker than those in preceding years. With respect to a

comparison of the heat fluxes, OAFlux considerably overestimates LHF with ocean heat loss biases of 50.95 w/m^2 (70%) in the spring, 42.43 w/m^2 (76%) in the summer_autumn and 63.29 w/m^2 (74%) in the winter. Consistent with the key input variables U and Q_a , the OAFlux LHF performance is better during the spring and winter than in the summer_autumn, which is further associated with the monsoon climate in the SCS. The seasonal SHF reliability is coincident with that of T_s , as the most poorly reliable T_s estimates lead to the most unreliable SHF estimates with enormous overestimations throughout the year. An analysis of the possible sources of biases in the heat fluxes show that biases in Q_a are the most dominant factor in determining the biases in LHF during the spring and winter. Meanwhile, both of the biases in Q_a and U are responsible for controlling the biases in LHF during the summer_autumn period. Biases in T_s are responsible for controlling the biases in SHF , and the effects of biases in T_s on the biases in SHF during the spring and winter are much greater than that in the summer_autumn period.

In summary, both T_s and SHF in OAFlux should be utilized with considerable caution in further research. Additionally, U , Q_a and LHF should be used with proper consideration due to their seasonal reliability variations. Researchers should feel more at ease using these data during the northeast monsoon than in the southwest monsoon. The performance of the OAFlux-estimated T_a seems to change little with the seasons and is highly consistent with the YXASFT observations throughout the year. Improving the observation capability of the AVHRR sensor under cloudy conditions is necessary for improving the accuracy of T_s estimates and the reliability of calculating SHF . Larger quantities of in situ bulk variable observations and direct turbulent heat flux measurements as well as improvements in the parameterization of variables in different regions of the SCS are also essential for improving the reliability of OAFlux datasets in the SCS.

Author contribution

Fenghua Zhou designed the experiments and Rui Shi, Ju Chen and Yunkai He carried them out. Fenghua Zhou and Rongwang Zhang write the Matlab program and performed the data processing and analysis. Qiangxie and Dongxiao Wang give help on useful discussions and data collection. Fenghua Zhou and Qiang Xie prepared the manuscript with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest

Acknowledgements

This study was funded by the Key Research Program of Frontier Sciences, Chinese Academy of Sciences (CAS) (QYZDJ-SSW-DQC022), the National Natural Science Foundation of China (41706102), the Chinese Academy of Sciences (CAS)

Key Technology Talent Program of 2016, the Station Network Construction Project-Xisha Marine Observatory of the CAS (KZCX2-YW-Y202). Rongwang Zhang was also supported by the National Key R&D Program of China (Grant No. 2017YFA0603200). The YXASFT data were provided by the Xisha Deep Sea Marine Environment Observation Station, South China Sea Institute of Oceanology, CAS. All of the in situ data adopted in this study can be obtained by contacting the first author, Fenghua Zhou (zhoufh@scsio.ac.cn). The authors would like to express their gratitude for the reanalysis data products comprising global ocean heat flux and evaporation data provided by the WHOI OaFlux project (<http://oaflux.whoi.edu>) funded by the NOAA Climate Observations and Monitoring (COM) program. The source code for the COARE 3.0 algorithm is freely available at http://coaps.fsu.edu/COARE/flux_algor/. The first author would like to thank the engineers from Campbell Scientific Company, USA, for their help with the observation system integration and data acquisition on the YXASFT. Finally, the authors thank the anonymous reviewers for their valuable comments and suggestions that improved the quality of this paper.

References

- Brunke, M. A., Fairall, C. W., Zeng, X., Eymard, L., and Curry, J. A.: Which bulk aerodynamic algorithms are least problematic in computing ocean surface turbulent fluxes?, *J. Climate*, 16, 619-635, doi:10.1175/1520-0442(2003)016<0619:WBAAAL>2.0.CO;2, 2003.
- Campbell Scientific, Inc. 2018: CR3000 Micrologger User's Manual. Accessed 21 March 2018, <https://s.campbellsci.com/documents/us/manuals/cr3000.pdf>.
- Chelton, D. and Xie, S.-P.: Coupled Ocean-atmosphere interaction at oceanic mesoscales, *Oceanography*, 23, 52–69, doi:10.5670/oceanog.2010.05, 2010.
- Chen, J., Zuo, T., and Wang, H.: Variation of latent heat flux over the Bengal Bay-South China Sea area and its relationship with South China Sea summer monsoon onset, in: IEEE International Geoscience and Remote Sensing Symposium, Munich, Germany, 22–27 July 2012, 856–859, 2012.
- Crawford, T. L., McMillen, R. T., Meyers, T. P., and Hicks, B. B.: Spatial and temporal variability of heat, water vapor, carbon dioxide, and momentum air-sea exchange in a coastal environment, *J. Geophys. Res.*, 98, 12869–12880, doi:10.1029/93jd00628, 1993.
- de Cosmo, J., Katsaros, K. B., Smith, S. D., Anderson, R. J., Oost, W. A., Bumke, K., and Chadwick, H.: Air-sea exchange of water vapor and sensible heat: the Humidity Exchange Over the Sea (HEXOS) results, *J. Geophys. Res.*, 101, 12001–12016, doi:10.1029/95jc03796, 1996.
- Fairall, C. W., Bradley, E. F., Rogers, D. P., Edson, J. B., and Young, G. S.: Bulk parameterization of air-sea fluxes for tropical ocean-global atmosphere coupled-ocean atmosphere response experiment, *J. Geophys. Res.*, 101, 3747–3764, doi:10.1029/95jc03205, 1996.

- Fairall, C. W., Barnier, B., Berry, D. I., Bourassa, M. A., Bradley, E. F., Clayson, C. A., de Leeuw, G., Drennan, W. M., Gille, S. T., Gulev, S. K., Kent, E. C., McGillis, W. R., Quartly, G. D., Ryabinin, V., Smith, S. R., Weller, R. A., Yelland, M. J., and Zhang, H.-M.: Observations to quantify air-sea fluxes and their role in climate variability and predictability, in: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*, edited by: Hall, J., Harrison, D. E., and Stammer, D., European Space Agency, Noordwijk, The Netherlands, 299-313, 2010.
- 5 Frenger, I., Gruber, N., Knutti, R., and Münnich, M.: Imprint of Southern Ocean eddies on winds, clouds and rainfall, *Nat. Geosci.*, 6, 608–612, doi:10.1038/ngeo1863, 2013.
- Goswami, B. N.: A note on the deficiency of NCEP/NCAR reanalysis surface winds over the equatorial Indian Ocean, *J. Geophys. Res.*, 108, 3124, doi:10.1029/2002jc001497, 2003.
- 10 Hogg, A. M. C., Dewar, W. K., Berloff, P., Kravtsov, S., and Hutchinson, D. K.: The effects of mesoscale ocean–atmosphere coupling on the large-scale ocean circulation, *J. Climate*, 22, 4066–4082, doi:10.1175/2009JCLI2629.1, 2009.
- Josey, S. A.: A Comparison of ECMWF, NCEP–NCAR, and SOC Surface heat fluxes with moored buoy measurements in the subduction region of the Northeast Atlantic, *J. Climate*, 14, 1780–1789, doi:10.1175/1520-0442(2001)014<1780:acoenn>2.0.co;2, 2001.
- 15 Mauder, M., Liebenthal, C., Göckede, M., Leps, J.-P., Beyrich, F., and Foken, T.: Processing and quality control of flux data during LITFASS-2003, *Boundary-Layer Meteorology*, 121, 67–88, doi:10.1007/s10546-006-9094-0, 2006.
- Persson, P. O. G., Fairall, C. W., Andreas, E. L., Guest, P. S., and Perovich, D. K.: Measurements near the Atmospheric Surface Flux Group tower at SHEBA: near-surface conditions and surface energy budget, *J. Geophys. Res.*, 107, 8045, doi:10.1029/2000JC000705, 2002.
- 20 Shi, R., Zeng, L., Chen, J., Yang, L., Dong, L., He, Y., Li, D., and Yao, J.: Observation and numerical simulation of the marine meteorology elements and air-sea fluxes at Yongxing Island in September 2013, *Aquat. Ecosyst. Health*, 18, 394–402, doi:10.1080/14634988.2015.1108822, 2015.
- Smith, S. R., Legler, D. M., and Verzone, K. V.: Quantifying uncertainties in NCEP reanalyses using high-quality research vessel observations, *J. Climate*, 14, 4062–4072, doi:10.1175/1520-0442(2001)014<4062:quinru>2.0.co;2, 2001.
- 25 Sun, B., Yu, L., and Weller, R. A.: Comparisons of surface meteorology and turbulent heat fluxes over the Atlantic: NWP model analyses versus moored buoy observations, *J. Climate*, 16, 679–695, doi:10.1175/1520-0442(2003)016<0679:COSMAT>2.0.CO;2, 2003.
- Swain, D., Rahman, S. H., and Ravichandran, M.: Comparison of NCEP turbulent heat fluxes with in situ observations over the south-eastern Arabian Sea, *Meteorol. Atmos. Phys.*, 104, 163–175, doi:10.1007/s00703-009-0023-x, 2009.
- 30 Wang, D., Liu, Q., Huang, R. X., Du, Y., and Qu, T.: Interannual variability of the South China Sea throughflow inferred from wind data and an ocean data assimilation product, *Geophys. Res. Lett.*, 33, L14605, doi:10.1029/2006gl026316, 2006.

- Wang, D., Zeng, L., Li, X., and Shi, P.: Validation of satellite-derived daily latent heat flux over the South China Sea, compared with observations and five products, *J. Atmos. Ocean. Tech.*, 30, 1820–1832, doi:10.1175/jtech-d-12-00153.1, 2013.
- Wang, W. and McPhaden, M. J.: What is the mean seasonal cycle of surface heat flux in the equatorial Pacific? *J. Geophys. Res.*, 106, 837–857, doi:10.1029/1999jc000076, 2001.
- Wang, X., Zhang, R., Huang, J., Zeng, L., and Huang, F.: Biases of five latent heat flux products and their impacts on mixed-layer temperature estimates in the South China Sea, *J. Geophys. Res.*, 122, 5088–5104, doi:10.1002/2016jc012332, 2017.
- Weller, R. A. and Anderson, S. P.: Surface meteorology and air-sea fluxes in the Western equatorial Pacific warm pool during the TOGA Coupled Ocean-Atmosphere Response Experiment, *J. Climate*, 9, 1959–1990, doi:10.1175/1520-0442(1996)009<1959:smaasf>2.0.co;2, 1996.
- Weller, R. A., Baumgartner, M. F., Josey, S. A., Fischer, A. S., and Kindle, J. C.: Atmospheric forcing in the Arabian Sea during 1994–1995: observations and comparisons with climatology and models, *Deep Sea Research, Part II: Topical Studies in Oceanography*, 45B, 1961–1999, doi:10.1016/s0967-0645(98)00060-5, 1998.
- Yan, J.-Y., Tang, Z.-Y., Yao, H.-D., Li, J.-L., Xiao, Y.-G., and Chen, Y.-D.: Air-sea flux exchange over the Xisha Sea area before and after the onset of southwest monsoon in 2002, *Chinese J. Geophys.*, 48, 1078–1090, doi:10.1002/cjg2.751, 2005.
- Yan, J. Y., Yao, H. D., Li, J. L., Tang, Z. Y., Jiang, G. R., Sha, W. Y., Li, X. Q., and Xiao, Y. G.: Air-sea heat flux exchange over the South China Sea under different weather conditions before and after southwest monsoon onset in 2000, *Acta Oceanol. Sin.*, 22, 369–383, 2003.
- Yang, L., Wang, D., Huang, J., Wang, X., Zeng, L., Shi, R., He, Y., Xie, Q., Wang, S., Chen, R., Yuan, J., Wang, Q., Chen, J., Zu, T., Li, J., Sui, D., and Peng, S.: Toward a mesoscale hydrological and marine meteorological observation network in the South China Sea, *B. Am. Meteorol. Soc.*, 96, 1117–1135, doi:10.1175/bams-d-14-00159.1, 2015.
- Yu, L., Weller, R. A., and Sun, B.: Mean and variability of the WHOI daily latent and sensible heat fluxes at in situ flux measurement sites in the Atlantic Ocean, *J. Climate*, 17, 2096–2118, doi:10.1175/1520-0442(2004)017<2096:mavotw>2.0.co;2, 2004.
- Yu, L., Jin, X., and Weller, R. A.: Multidecade global flux datasets from the objectively analyzed air-sea fluxes (OAFlux) project: latent and sensible heat fluxes, ocean evaporation, and related surface meteorological variables. http://oafux.whoi.edu/pdfs/OAFlux_TechReport_3rd_release.pdf, last access: 19 September 2017, 2008.
- Yu, L., Jin, X., and Weller, R. A.: Role of net surface heat flux in seasonal variations of sea surface temperature in the tropical Atlantic Ocean, *J. Climate*, 19, 6153–6169, doi:10.1175/jcli3970.1, 2006.
- Zeng, L., Shi, P., Liu, W. T., and Wang, D.: Evaluation of a satellite-derived latent heat flux product in the South China Sea: a comparison with moored buoy data and various products, *Atmos. Res.*, 94, 91–105, doi:10.1016/j.atmosres.2008.12.007, 2009.

- Zeng, X., Zhao, M., and Dickinson, R. E.: Intercomparison of bulk aerodynamic algorithms for the computation of sea surface fluxes using TOGA COARE and TAO data, *J. Climate*, 11, 2628–2644, doi:10.1175/1520-0442(1998)011<2628:iobaaf>2.0.co;2, 1998.
- 5 Zhang, R., J. Huang, X. Wang, J. A. Zhang, and F. Huang, 2016: Effects of Precipitation on Sonic Anemometer Measurements of Turbulent Fluxes in the Atmospheric Surface Layer. *J. Ocean Univ. China*, 15(3), 389–398. doi:10.1007/s11802-016-2804-4.
- Zhang, R., X. Wang, and C. Wang, 2018: On the Simulations of Global Oceanic Latent Heat Flux in the CMIP5 Multimodel Ensemble. *J. Climate*, 31, 7111–7128. doi: 10.1175/jcli-d-17-0713.1.
- 10 Zhu, J., Kamachi, M., and Wang, D.: Estimation of air-sea heat flux from ocean measurements: an ill-posed problem, *J. Geophys. Res.*, 107, 3159, doi:10.1029/2001jc000995, 2002.

Figure captions

Figure 1. (a) Yongxing Island air-sea flux tower (YXASFT).

(b) Instrumentation and data acquisition system mounted on the YXASFT.

(c) Pictures of some sensors on the YXASFT.

5 (d) Google satellite image of Yongxing Island. The red triangle indicates the location of the YXASFT.

(e) Map of the northern SCS. The black star indicates the location of Yongxing Island.

Figure 2. Diagram of the real-time data acquisition system and the sensor wiring scheme on the YXASFT (SEx: single-ended channel; VXx: voltage excitation channel; Px: pulse-input channel; IXx: current excitation channel; SDM: SDM channel; (Campbell Scientific, Inc. 2018) GPRS: General Packet Radio Service; CDMA: code-division multiple access).

10 Figure 3. EC turbulence data processing and quality control flow chart

Figure 4. Daily means of the *LHF* and *SHF* time series (top panels) and scatter plots (bottom panels) of COARE3.0 versus ECF (from 2016/02/01 to 2016/03/29). The R^2 values, linear regressions and numbers of matched pairs (N) are given in the bottom panels. The solid red line refers to the linear regression of the matched pairs. The solid green line $y=x$ indicates a 1:1 correspondence.

15 Figure 5. Daily mean time series plots of the YXASFT-observed (red solid lines) and OAFlux-analyzed (blue solid lines) U , Q_a , T_a , and T_s values over the study period (2016/02/01-2017/01/31).

Figure 6. Daily mean time series plots of the YXASFT-observed (red solid lines) and OAFlux-analyzed (blue solid lines) *SHF* and *LHF* over the study period (2016/02/01-2017/01/31).

20 Figure 7. Scatter plots of the YXASFT and OAFlux wind speeds at 10 m (U), air specific humidity at 2 m (Q_a), and sea surface temperatures (T_s) and air temperatures at 2 m (T_a) during the spring (left column), summer _autumn (middle column), and winter (right column) periods. The units for U , Q_a , T_s and T_a are m/s , g/kg , $^{\circ}C$ and $^{\circ}C$, respectively. The linear regression equation, coefficient of determination (R^2), and number of matched pairs (N) are given in each panel. The solid red line refers to the linear regression of the matched pairs.

25 Figure 8. Daily mean time series plots of the YXASFT observed downward long radiation (*DLR*) over the study period (2016/02/01-2017/01/31).

Figure 9. Same as Fig. 5 but for *LHF* (first row) and *SHF* (second row).

Figure 10. Scatter plots for the biases of U (ΔU), Q_a (ΔQ_a), T_s (ΔT_s), and T_a (ΔT_a) with respect to the biases of *LHF* (ΔLHF). All of the data are normalized to the range of -10 to 10 in this paper. The linear regression equation and coefficient of determination (R^2) are given in each panel. The solid red line refers to the linear regression of the matched pairs.

30 Figure11. Same as Fig.9 but for the biases in *SHF*.

Tables

Table 1. List of sensors installed on the YXASFT and their specifications

Parameters	Sensor	Scan interval (Hz)	Averaging interval (min)	Installation height (m)
Wind speed and direction	Young 05106	1	1, 10, 30	5, 10, 15, 20
Air temperature and humidity	Vaisala HMP155A	1	1, 10, 30	5, 10, 15, 20
Four-component radiation	Hukseflux NR01	1	1, 10, 30	8
Sea surface temperature	Campbell SI-112	1	1, 10, 30	5
Eddy turbulent fluxes ($u, v, w, t, \rho_v, T_{au}, SHF, LHF, F_c$)	Campbell IRGASON	10	30	12

Table 2. Information regarding the adopted in situ and reanalysis data*

Data	Variables	Location	Height (m)	Interval	Period (day)
In situ bulk variables	U	112.33° E, 16.84° N	10	30 min	366
	Q_a		5	30 min	
	T_s		0.05	30 min	
	T_a		5	30 min	
	DLR		8	30 min	
In situ bulk heat fluxes	SHF	112.33° E, 16.84° N	10	30 min	
	LHF		10	30 min	
In situ ECF turbulent data	u	111.5° E, 16.5° N	12	0.1 sec	57
	v		12	0.1 sec	
	w		12	0.1 sec	
	t		12	0.1 sec	
	ρ_v		12	0.1 sec	
	SHF		12	30 min	
	LHF		12	30 min	
OAFlux bulk variables And heat fluxes	U	111.5° E, 16.5° N	10	1 day	366
	Q_a		2		
	T_s		0.05		
	T_a		2		
	SHF		10		
LHF	10				

u : wind speed along the sonic x-axis, v : wind speed along the sonic y-axis, w : wind speed along the sonic z-axis, t : sonic temperature, ρ_v : water vapor density. The height of the bulk fluxes derived via COARE3.0 for both in situ data and OAFlux are considered at 10 m.

Table 3. Quantitative statistical summary based on comparisons between daily YXASFT measurements and daily OAFflux products in the spring, summer_ autumn, and winter Periods

Season	Variable	OAFflux mean	YXASFT mean	<i>RMSE</i>	<i>Bias</i>	<i>R</i> ²	Regression	
							<i>C</i> ₁	<i>C</i> ₂
Spring	<i>U</i> (m/s)	7.36	6.40	1.36	0.96	0.90	0.89	1.66
	<i>Q</i> _a (g/kg)	15.29	15.63	1.27	-0.33	0.81	0.57	6.42
	<i>T</i> _a (°C)	24.10	24.62	0.68	0.52	0.92	0.90	2.06
	<i>T</i> _s (°C)	25.12	24.65	1.29	0.46	0.47	0.32	17.27
	<i>SHF</i> (w/m ²)	15.46	-1.37	25.64	16.83	0.01	-0.45	14.84
	<i>LHF</i> (w/m ²)	123.87	72.92	63.23	50.95	0.80	1.42	20.39
Summer_Autumn	<i>U</i> (m/s)	6.16	4.97	1.67	1.19	0.79	0.85	1.93
	<i>Q</i> _a (g/kg)	20.33	21.08	1.09	-0.75	0.68	0.66	6.47
	<i>T</i> _a (°C)	28.86	28.95	0.43	-0.10	0.84	1.00	-0.09
	<i>T</i> _s (°C)	29.04	29.11	0.61	-0.07	0.70	0.70	8.62
	<i>SHF</i> (w/m ²)	1.65	0.51	6.33	1.07	0.31	1.10	1.02
	<i>LHF</i> (w/m ²)	97.97	55.98	50.49	42.43	0.40	0.94	46.04
Winter	<i>U</i> (m/s)	10.07	9.40	0.93	0.67	0.92	0.95	1.14
	<i>Q</i> _a (g/kg)	16.35	16.47	0.67	-0.11	0.80	0.71	4.60
	<i>T</i> _a (°C)	24.91	25.48	0.67	-0.57	0.89	0.90	1.95
	<i>T</i> _s (°C)	25.72	25.67	0.68	0.05	0.54	0.50	12.90
	<i>SHF</i> (w/m ²)	13.83	9.73	28.85	23.56	0.14	-1.59	-1.62
	<i>LHF</i> (w/m ²)	148.32	85.03	72.35	63.29	0.66	1.30	37.45

*OAFflux = *C*₁ × YXASFT + *C*₂

Figures




Figure 1

5




Figure 2

u, v, w, t, ρ_v

Raw data split

Split the raw TOB3 data file into a series of 30-min TOA5 data files by
LoggerNet 4.3 with naming format of
TOA5_6703.Time_Series_??_yyyy_mm_dd_hhmm.dat

Preprocessing

Filter data with diagnostic flags and signal strengths and unacceptable ranges
Despiking

Flux calculation and necessary correction

Axis rotation for tilt correction
High and low frequency spectral corrections
Compensation of density fluctuations

Fully corrected *SHF, LHF*

Quality control and evaluation

Steady state test
Developed turbulent conditions test
Flux data quality grading

5

Figure 3

10

15

20




Figure 4




Figure 5

5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11