# Reply to RC1

General Comments: This paper deals with rage resolved observations of aerosol properties, over Tsukuba, Japan, and a comprehensive study on how they affect the atmospheric boundary layer evolution. For the latter, the authors used the remote sensing observations as initial inputs in a 1-D atmospheric model. The paper rightly acknowledges previously related studies. The manuscript is well written, but in order to be improved, I would suggest to the authors to take into consideration the following comments.

Thank you very much for your time, evaluation and useful comments. We reflected your suggestions in the revised manuscript as well as the comments from the other reviewer, which improved the manuscript significantly. Point-by-point responses to your comments are written hereinafter. Please note that we added two figures (Figs. 1 and 10), one table (Table 4) and Sect. 3.2.3 in the revised paper.

The changes to meet your comments are marked with blue in the revised manuscript.

#### Minor Comments:

1. The Abstract section is well written. However, I would like to draw the attention to the authors to consider stating with numbers the main outcomes of their study.

We revised the structure of the abstract.

2. Page 1, line 21: "compared to".

We corrected it. Page 1, line 27 in the revised paper.

3. Page 1, lines 24-26: This sentence is too long. The authors are kindly requested to rephrase it, and make their statement clearer.

We revised it.

Page 1, lines 29-32 in the revised paper.

4. Page 3, line 8: Apart from the coordinates, please provide also the elevation of Tsukuba station. It would be useful also for the reader, if you could provide the link of the used station which operates under SKYNET network. With a quick search I was not able to find this station here <u>http://www.skynetisdc.org/quicklooks.php</u>.

We added the elevation to the text.

Unfortunately, our data of the sky radiometer at Tsukuba is not transfer to the International SKYNET Data Center due to the policy of our institute. Therefore, the data at Tsukuba is not available online. Page 4, lines 10-12 in the revised paper.

5. Page 3, line 12: The authors are mentioning that among other aerosol properties the lidar data obtained by the AD-Net, contains also the depolarization ratio for particle and molecular scattering. Maybe the authors are referring to the physical quantity of volume depolarization ratio, which includes the contribution of molecular and particle depolarization. Please clarify this.

"depolarization ratio for particle and molecular scattering" is "volume depolarization ratio" including the contribution of molecular and particle depolarization. We changed "depolarization ratio" to "volume depolarization ratio".

Page 4, lines 15-16 in the revised paper.

It is not so clear in the text, the contribution of the auxiliary data mentioned by the authors (page 3 lines 13-18). I suppose that this dataset was used in the radiative transfer module of the 1-D atmospheric model, but it would be nice if this is mentioned here.

These data were used for the calculation of Rayleigh scattering and gas absorption in the SKYLIDAR retrieval. These are necessary for the forward calculation of the sky radiometer and lidar data in the retrieval. We described this in the revised paper.

Page 4, lines 21-22 in the revised paper.

7. Moreover, I would suggest to the authors to consider producing a flowchart diagram as the very first figure of their manuscript, in order to clearly demonstrate there the inputs and outputs of their approach/methodology.

We added a flowchart (Fig. 1) to the revised manuscript.

Page 3, line 1, and page 24 in the revised paper.

8. Page 4 line 21: Define here the % of successful retrievals of ABL estimation from your dataset.

The successful retrievals of ABL estimation was about 95 % for 2,305 profiles under the clear sky conditions. In the 5 %, the base height of the transported aerosols was used as the ABL height. We added this to the manuscript.

Page 6, lines 23-24 in the revised paper.

9. Page 4 line 23: What is the time window used for smoothing the time series of ABL height?

One hour.

Page 6, line 15 in the revised paper.

9. Page 7 line 4: Consider providing the appropriate reference for Ångström exponent namely: "Ångström, A., On the atmospheric transmission of Sun radiation and on dust in the air, Geogr. Ann., 11, 156–166, 1929". Moreover, it is not clear, if this quantity refers to the entire atmospheric column, or is the range resolved backscatter or extinction related exponent calculated from the 532-1064 nm pair of wavelengths from the lidar measurements. Please try to clarify this in the manuscript.

We are afraid that we were not able to get the paper of Ångström 1929 and cannot add the reference to the manuscript. We calculated the Ångström exponent from the retrieved extinction coefficients at 532 and 1064 nm. We added this explanation to the text. Page 9, lines 11-13 in the revised paper.

Page 9, lines 11-13 in the revised paper.

10. Figure 3: Title "Ångström Exponent". Please go through the entire manuscript (tables, figures and text) to correct "Ångströme" to "Ångström".

We checked all the text and corrected.

11. Page 7 lines 14-15: The authors are kindly requested to provide some indicative studies related to the single scattering albedo of dust and black carbon.

We added the references to the revised paper. Page 9, line 23.

12. Page 7 lines 29-31: The authors are commenting in lidar ratio values. But it is confusing since in 2.1.1 they mention that they operated a two wavelength Mie-scattering lidar from AD-Net. Since this system is not a Raman or HSRL to obtain independently the lidar ratio, please comment from where these values came from.

The lidar ratio was calculated from the retrieved single-scattering albedo and phase function. We added this explanation to the revised manuscript. Page 10, lines 14-15 in the revised paper.

13. Figure 6 (c): The authors are demonstrating two-days of air mass back-trajectories, in order to identify the aerosol source. Even though they provide some stars to indicate the starting altitude, the final height of the air mass arriving over the station as well as the height through the entire travel path is not shown. The authors are kindly requested to update this figure or at least mention these heights in the text.

We added the altitude vs time plot as Fig. 7d to the revised manuscript. Page 30.

14. I noticed that during the entire paragraph 3.1.2 the authors are describing qualitatively (e.g. "values were small; large; was large; was small" etc.) the aerosol optical and physical properties. They are kindly requested to provide also some numbers in the text.

We described our results with numbers and compared them with other studies in the revised manuscript. Pages 10-12 in the revised paper. 15. Page 9 lines 5-6: Please specify also the warming range of the FA due to the direct aerosol heating.

The warming range is from 0.0 to 0.4 K at noon (Fig. 8d). We added this to the text. Page 12, line 24 in the revised paper.

16. Page 9 line 21: "... the decrease in the ABL height was smaller...". Please specify how much.

We described the results with numbers in the revised manuscript. Page 13, lines 9-11 in the revised paper.

17. Figures 7 & 8: Please consider to make this graphs clearer. I think that it will help if: (1) units exist next to the parameters EXP0 and EXP1- EXP0 (2) provide altitude (y-axis in each graph) in km for being consistent with the previous figures. Finally I think that in the primary x-axis of (d) and (e) the x-title should be Potential Temperature (K) and Specific Humidity (g kg-1) respectively, and not EXP0.

We revised the figures in accordance with your suggestions, (1) and (2). However, we were not able to delete "EXP0" from the x-title in the primary x-axis of (d) and (e) because we need to indicate that the primary x-axis is valid for only the result in EXP0.

Pages 31-32 in the revised paper.

18. Please update the link related to SKYNET, as provided in Page 11 line 3.

We updated the link. Page 16, line 3.

# Major Comments:

1. A general comment to the authors. Select some indicative numbers from the tables, representing the major findings of this study, and provide them into the manuscript (sections: Abstract, Results, Conclusions).

Thank you for your suggestion. We checked all the text and revised the descriptions.

2. The authors apart from the mean statistical values are also using values observed from 5 case studies. The three of them where characterized as dust, one as smoke and one as mixture of dust and smoke. From their findings, the case studies differ only at the Ångström exponent and the aerosol load. I wonder if during their study, they can also draw any conclusion related to the influence of each individual aerosol type on the ABL evolution.

We added a new section 3.2.3 and discussed the dependencies of the 2 m temperature and ABL height on the optical thickness and Ångström exponent. When the Ångström exponent was small (dust cases), the decrease rates of the 2 m temperature and ABL height were large (Fig. 10 in the revised paper). The dust particles have the small values of the Ångström exponent and the large optical thickness in the near infrared wavelength region. Therefore, the impacts of aerosols on the 2 m temperature and ABL height were large in the dust cases. However, the decrease of the ABL height in 9 May 2013 (smoke and dust case) was larger than that in 2 April 2012 (dust case). In both cases, the aerosol optical thickness in FA had similar values, about 0.33. The geometric thickness of the aerosol layer in the FA was smaller in 9 May 2013 than in 2 April 2012, and the extinction coefficient in the FA was larger in 9 May 2013 than in 2 April 2012 (Fig. 7a in the revised paper). Therefore, the temperature increase in the FA was large in 9 May 2013 (Fig. 8d in the revised paper), the capping inversion became strong, and the ABL height was less developed. We found the dependencies of the 2 m temperature and ABL height on the Ångström exponent, but the most important factor was the vertical profile of the extinction coefficient above the ABL.

Pages 13-14 in the revised paper.

3. Page 4 lines 26-27: How much uncertainty this assumption may introduce? This maybe implies an overestimation of the aerosol load inside the ABL and an underestimation in the free troposphere. In any case the authors are kindly requested to comment on this.

As you mentioned, our method may overestimate the aerosol load inside the ABL around 3 UTC in Fig. 2b in the revised paper. Our assumption can affect the results of aerosol optical properties in ABL and FA. However, it is very difficult to determine the ABL height of the case that the aerosols in the ABL and FA are mixed. Since we do not know the true ABL height, we cannot evaluate the uncertainties of our method. However, the number of the case, in which the base height was used as the ABL height, was 5 %. We believe the influence of our assumption would be small. I added this

explanation in the revised paper.

Page 6, lines 20-25 in the revised paper.

4. Page 5 lines 21-22: Please provide an estimation of the error that is introduced especially in the all wavelength net radiation that is discussed later, by ignoring the influence of aerosols on the infrared region (λ>1064nm), since dust particles are mostly in the coarse mode with diameters > 2 um. Moreover, the linear interpolation of the aerosol optical properties between 532 and 1064 nm can also influence the final retrievals. Please elaborate more on this.

We consider the influences of aerosols in the wavelength region from 300 nm to 3.0 um. We calculated the real and imaginary parts of the refractive index between 532 and 1064 nm by the linear interpolation in a log-log plane. For wavelengths less than 532 nm and greater than 1064nm, the refractive index at 532 and 1064 nm were used. The extinction coefficient, single-scattering albedo, and phase function from 300 nm to 3.0 um were calculated from these refractive indices, the volume size distributions, and the volume ratio of the non-spherical particles in the coarse mode. Then, the radiative fluxes and heating rates were calculated by the radiative transfer model. We revised the text. Page 7, lines 16-25 in the revised paper.

5. Page 7 lines 2-3: The authors are giving an explanation of their findings regarding the seasonal variation of the mean ABL values, presented in Figure 2. However, I wonder if they compared these mean seasonal ABL values with corresponding measurements from radiosondes. In the same figure, it seems that there are a few cases that aerosol layers are detected up to 9 km height. Are these smoke particles advected from Russia?

We determined the ABL height from the lidar data in order to distinguish the locally emitted aerosols and the transported aerosols. For this purpose, we should evaluate the ABL height from the lidar data, which directly relates to the aerosol vertical profile. The ABL height estimated from the radiosonde may not be consistent with the aerosol vertical distribution. Therefore, we did not compare the ABL height with that by radiosonde. Moreover, the radiosonde launched only twice in a day (09 and 21 JST. JST is Japan Standard Time.) in Japan. The ABL height around the noon cannot be obtained from the radiosonde.

The smoke from Russia in 8 May 2013 was not detected near 9 km height (Fig. 7a in the revised paper). The aerosols around 9 km is sometimes observed in Tsukuba, Japan. However, we do not know the source of aerosols at 9 km altitude because our backward trajectory analysis was conducted for only

the five cases of high aerosol load in the FA. In the future study, we will calculate a lot of backward trajectories for investigating the statistical relations between the aerosol optical properties and their sources.

6. Page 7 lines 23-25: Since particles located in FA and ABL may have totally different optical and chemical properties how comes and the refractive index is the same for the aforementioned atmospheric regions? Is this a bias coming from the retrieval method of SKYLIDAR? In any case the authors are kindly requested to comment more on this.

In the work of Kudo et al. 2016, we conducted the sensitivity tests using the simulated data of lidar and sky radiometer for the transported dust and pollution cases to investigate the capability of the SKYLIDAR. The pollution aerosol was defined as small-sized and light-absorbing particle. The columnar values of the refractive index were estimated well in both cases. In the transported dust case, the vertical profile of the imaginary part of the refractive index was estimated successfully, but the that of the real part was not. In the transported pollution case, the vertical profile of the real part was estimated, but that of the imaginary part was not. In the failed cases, the vertical profile of the refractive index was constant with their vertical means. Therefore, the similar values of the refractive index in the ABL and FA may be caused by the errors of the SKYLIDAR. I added these explanation to the text. Page 5, lines 8-25, and page 10, lines 2-8 in the revised paper.

7. Page 8 line 15: Even though the aerosol type detected in 08 May2013, was characterized as smoke, no clear differences compared to the dust and dust smoke days can be found, regarding the values of single-scattering albedo and asymmetry factor (Table 2). Is this coming from the retrieval method of the SKYLIDAR?

In the numerical sensitivity tests of the SKYLIDAR (Kudo et al. 2016), the vertical profiles of singlescattering albedo and asymmetry factor for the transported dust were estimated well. Our retrievals in the transported dust cases were from 0.95 to 0.98 for the single-scattering albedo and from 0.65 to 0.71 for the asymmetry factor. These were consistent with the reports of the AERONET retrievals in the desert regions in the world, and the sky radiometer retrievals for the Asian dust.

The SKYLIDAR fails to retrieve the vertical profile of the single-scattering albedo of the transported smoke, but the estimated vertical profile is constant with the vertical mean (Kudo et al. 2016). Therefore, our estimated single-scattering albedo can be compared with that of the AERONET retrievals. The comparisons with other studies showed that the our estimated single-scattering albedo

and asymmetry factor lied among those reported in the other studies. We added the explanation of the retrieval errors of the SKYLIDAR and the comparisons of our retrievals with other studies to the revised paper.

Pages 10-12 in the revised paper.

### Reply to RC2

Thank you very much for your time, evaluation and useful comments. We reflected your suggestions in the revised manuscript as well as the comments from the other reviewer, which improved the manuscript significantly. Point-by-point responses to your comments are written hereinafter. Please note that we added two figures (Figs. 1 and 10), one table (Table 4) and Sect. 3.2.3 in the revised paper.

The changes to meet your comments are marked with blue in the revised manuscript.

#### General Comments

In this paper, the authors apply an inversion method that combines radiometric and lidar measurements and obtain vertical profiles of multiple aerosol optical and microphysical properties. The climatological behavior of these properties is then examined. Moreover, some of these products are used as input to an 1D atmospheric model in order to investigate the effect of the aerosol presence as well as their vertical distribution to the boundary layer height. In general the paper, is rather interesting and well organized. My main concern, however, is about the uncertainties of the inversion method applied, especially for cases where the aerosol load is small. Typically such inversion methods require a certain level of confidence to the input parameters. This is usually achieved for the columnar properties when a sufficient amount of aerosol is present in the atmosphere. The lidar profiles, on the other hand, always include regions where the extinction coefficient is low, e.g. the last part of the profile. How accurate is the inversion for those cases? For example, are the retrievals for an extinction coefficient below 0.02 km<sup>-1</sup> (Figures 2,3 and 4) trustworthy? This issues should be discussed in the manuscript. If the error of the retrieval has not been quantified, then the authors should consider applying thresholds to the AOD and extinction coefficients in order to exclude profiles, or parts of the profiles with potentially high uncertainties from their analysis.

We conducted the intensive sensitivity tests using the simulated data of lidar and sky radiometer with random noises in the work of Kudo et al. 2016. The retrieval errors increased with a decrease of AOD or extinction coefficient. In the case that the AOD was about 0.05 and the extinction coefficient was about 0.01 km<sup>-1</sup>, the retrieval errors were  $\pm 0.003$  km<sup>-1</sup> for the extinction coefficient, and  $\pm 0.05$  for single-scattering albedo and asymmetry factor. We added the explanation of the retrieval errors of the SKYLIDAR in Sect. 2.1.1 of the revised paper. And, we discussed our results of the optical properties and the retrieval errors in Sect. 3.1 of the revised paper.

Page 5, lines 8-25 in the revised paper.

# Specific Comments

Section 2.1.2 (page 4, lines 11-12)

Why do you use the wavelet covariance transform (WCT) at 532nm instead of 1064nm for the cloud and aerosol base retrievals when both are available during the day? The aerosol layers tend to appear more clearly in the 1064nm channel.

In the beginning of this study, We used the 1064 nm channel, and many tests were conducted. However, the base height was sometimes too low. The signal noises in the 1064 nm channel were very large in the layers where the aerosol load was small. The influences of the signal noises remained in the vertical distribution of the WCT at 1064 nm and affected the estimation of the ABL height. On the other hand, the signal noises in the 532 nm channel were small. After the many tests, we decided to use the 532 nm channel at first. When the base height is not detected by the 532 nm channel, we try to detect the base height by the 1064 nm channel.

### Section 2.2.1 (page 5, lines 18-22)

It is mentioned here that "we determined the optical properties between 532 and 1064 nm by linear interpolation and used the optical properties at 532 and 1064 nm for wavelengths of less than 532 nm and greater than 1064 nm". Is this true for all the aforementioned optical properties? The aerosol extinction coefficient, in particular, usually has a strong wavelength dependence, even within the visible spectrum. Since the spectral range is much larger here, the authors should consider applying a correction, e.g. with the use of a constant angstrom exponent, for the extinction profiles in wavelengths below 532nm and above 1064nm.

We calculated the real and imaginary parts of the refractive index between 532 and 1064 nm by the linear interpolation in a log-log plane. For wavelengths less than 532 nm and greater than 1064nm, the refractive index at 532 and 1064 nm were used. The extinction coefficient, single-scattering albedo, and phase function were calculated from these refractive indices, the volume size distributions, and the volume ratio of the non-spherical particles in the coarse mode. Then, the radiative fluxes and heating rates were calculated by the radiative transfer model. We revised the text. Page 7, lines 16-25 in the revised paper.

Section 3.1.1 (page 6, lines 22-23) The authors mention that "In all the seasons, the extinction coefficient was large in three layers: from the surface to 1.5 km; from 1.5 to 3.5 km; and from 3.5 to 6 km altitude." The upper layer, however, only appears during winter and spring (Figure 2). Please rephrase. Since only the boundary layer is provided in the figures the discussion should be focused mainly on the differences between the free atmosphere and the boundary layer.

I revised the text.

Page 8, lines 28-31 in the revised paper.

Section 3.1.1 (page 6, lines 31-32)

The number of daily mean profiles is considerably lower during summer (only 7 days - Figure 2). Is this related to cloudy or rainy conditions? The constant rainy conditions could explain the lower extinction values.

Yes. Summer in Japan is hot and humid. The cumulus clouds develop almost every day. The small number of daily mean profiles is due to the cloudy or rainy conditions. We added this explanation in the text.

Page 8, lines 26-28 in the revised paper.

Section 3.1.2 (page 8) Please provide specific values from the analysis (Table 2). A short comparison with estimates from other scientific studies would be interesting here as well.

We provided the specific values and compared them with other studies. Pages 10-12 in the revised paper.

Figure 4 The winter profiles exhibit high extinction and high SSA values while the summer profiles show low extinction and low SSA values. During spring both kinds of profile are observed. Do you observe any specific seasonal pattern within the spring months. For example, do the winter-like profiles tend occur in early spring while the summer-like profiles in late spring?

We did not make the monthly profiles because the data number was not enough to make them. However, from the daily figures, the spring-like profile was observed sometimes in late winter. The observation period of two years was too short to investigate the monthly pattern. Technical Comments - Suggestions Table 1 Consider defining abbreviations for the table properties in the caption and include the variability with

+- in the same line with the mean value. Currently, each table property occupies 2 lines and sometimes this is difficult to follow.

I revised the table. Page 20 in the revised paper.

Figure 6

A small height versus time plot per trajectory could be included along with the map. Regions where the trajectories pass closer to the ground are more probable to affect the layer aerosol content.

I added the height versus time plot as Fig. 7d. Page 30 in the revised paper.

Section 3.1.1 (page 7, lines 14-15) Add citation

I added a citation. Page 9, line 23 in the revised paper.

Section 3.1.1 (page 7, lines 23-24) typo - replace "almost same values" with "almost the same"

We changed the contents in the paragraph to the comparisons of our results with other papers. Page 10, lines 1-16.

# Characteristics of aerosol vertical profiles in Tsukuba, Japan, and their impacts on the evolution of the atmospheric boundary layer

Rei Kudo<sup>1</sup>, Toshinori Aoyagi<sup>2</sup>, and Tomoaki Nishizawa<sup>3</sup>

<sup>1</sup>Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, 305-0052, Japan 5

<sup>2</sup>Japan Meteorological Agency, Tokyo, 100-8122, Japan

<sup>3</sup>National Institute for Environmental Studies, Tsukuba, 305-0053, Japan

Correspondence to: Rei Kudo (reikudo@mri-jma.go.jp)

- Abstract. Vertical profiles of the aerosol physical and optical properties, with a focus on seasonal means and on transport events, were investigated in Tsukuba, Japan, by a synergistic remote sensing method that uses lidar and sky radiometer data. 10 The retrieved aerosol vertical profiles of the springtime mean and five transport events were input into our developed onedimensional atmospheric model, and the impacts of the aerosol vertical profiles on the evolution of the atmospheric boundary layer (ABL) were studied by numerical sensitivity experiments. The characteristics of the aerosol vertical profiles in Tsukuba are as follows: (1) The retrieval results in the spring showed that aerosol optical thickness at 532 nm in the free
- 15 atmosphere (FA) was 0.13, greater than 0.08 in the ABL owing to the frequent occurrence of transported aerosols in the FA. In other seasons, optical thickness in the FA was almost the same as that in the ABL. (2) The aerosol optical and physical properties in the ABL showed a dependency on the extinction coefficient. With an increase of the extinction coefficient from 0.00 to 0.24 km<sup>-1</sup>, Ångström exponent increased from 0.0 to 2.0, the single-scattering albedo increased from 0.87 to 0.99, and the asymmetry factor decreased from 0.75 to 0.50. (3) The large variability in the physical and optical properties of
- aerosols in the FA were attributed to transport events, during which the transported aerosols consisted of varying amounts of 20 dust and smoke particles depending on where they originated (China, Mongolia, or Russia). The results of the numerical sensitivity experiments using the aerosol vertical profiles of the springtime mean and five transport events in the FA are as follows: (1) Numerical sensitivity experiments based on simulations conducted with and without aerosols showed that aerosols caused the net downward radiation and the sensible and latent heat fluxes at the surface to decrease. The decease of
- 25 the temperature in the ABL (-0.2 to -0.6 K) and the direct heating of aerosols in the FA (0.0 to 0.4 K) strengthened the capping inversion around the top of the ABL. Consequently, the ABL height was decreased by 133 to 208 m in simulations with aerosols compared to simulations without aerosols. (2) We also conducted simulations in which all aerosols were compressed into the ABL but in which the columnar properties were the same and compared with the simulation results for uncompressed aerosol profiles. The results showed that the reductions in net downward radiation and in sensible and latent
- 30 heat fluxes were the same in both types of simulations. However, the capping inversion in the simulations with compression was weakened owing to aerosol direct heating in the ABL and the lack of direct heating in the FA. This resulted in an increase of the ABL height, compared with that in the simulations without compression. (3) The dependencies of the 2 m



temperature and ABL height on the optical thickness and Ångström exponent in the FA was investigated using the results of the numerical sensitivity tests. The 2 m temperature and ABL height was decrease with an increase of the optical thickness, and their decrease rates increase with a decrease of Ångström exponent because the optical thickness in the near infrared wavelength region was large when the Ångström exponent is small. However, there was a case that the Ångström exponent

5 was large but the decrease of the ABL height was largest in all the simulation results. In this case, the strong capping inversion due to the large extinction coefficient around the top of the ABL was an import factor. These results suggest that the vertical profiles of the aerosol physical and optical properties, and the resulting direct heating has important effects on the ABL evolution.

10

#### **1** Introduction

Solar radiation heats the Earth's surface, thereby causing thermal instability and evaporation. The thermal energy and water vapor are transported into the atmosphere through turbulent mixing in the atmospheric boundary layer (ABL). These processes in the ABL have important implications for global energy and water circulation. Aerosols have significant impacts

- 5 on the radiation budget of the Earth because they scatter and absorb solar radiation (aerosol-radiation interaction) and modify cloud physical properties (aerosol-cloud interaction) (IPCC, 2013). In this study, we focus on aerosol-radiation interaction and did not consider clouds and precipitation processes. Direct scattering and absorption of solar radiation by aerosols decrease the amount of solar radiation that reaches the Earth's surface and sensible and latent heat fluxes, heat the atmosphere, and modify atmospheric stability. These effects have significant impacts on the evolution of the ABL, but the impacts differ depending on the aerosol optical properties (Yu et al., 2002; Pandithurai et al., 2008).
  - Yu et al. (2002) and Pandithurai, et al. (2008) investigated the influences of aerosol optical properties on the ABL structure by sensitivity experiments with a high resolution ABL model coupled with an accurate radiative transfer model. They showed that the light absorption characteristics of aerosols are important determinates of their impact on ABL evolution. However, these studies focused on aerosols only in the ABL. Tsunematsu et al. (2006) examined sounding data
- 15 obtained by frequently launched sondes and showed that direct heating of transported dust in the free atmosphere (FA) strengthened the capping inversion at the top of the ABL. Therefore, it is also necessary to study the influences of aerosols in the FA on ABL evolution, especially because aerosols in the FA can be transported both regionally and globally (Uno et al., 2009).
- Ground-based remote sensing has the advantage that it can be used for continuous monitoring of aerosol vertical profiles. We developed a synergistic method, SKYLIDAR, that retrieves vertical profiles of aerosol optical properties from lidar and sun/sky photometer data (Kudo et al., 2016). SKYLIDAR provides vertical profiles of the extinction coefficient, single-scattering albedo, and phase function, and with these products the solar heating rate can be evaluated (Kudo et al., 2016). Then, by inputting the retrieved aerosol optical properties into an ABL model, it is possible to investigate the influences of aerosols in the ABL and FA on the evolution of the ABL.
- 25 This study comprises two parts (Fig. 1). We first evaluated the vertical profiles of aerosol physical and optical properties in a two-year lidar and sun/sky photometer data set collected at Tsukuba, Japan, in a rural area located near the mega-city of Tokyo. The ABL height was determined from the lidar data to distinguish the locally emitted aerosols in the ABL and the transported aerosols in the FA, and the characteristics of the physical and optical properties in ABL and FA were investigated. The columnar properties of aerosols at Tsukuba have been investigated by many researchers (e.g., Nishizawa et al., 2004; Kudo et al., 2010a, 2010b, 2011), but these previous studies did not investigate their vertical profiles statistically. Second, we investigated the impact of aerosol vertical profiles on the evolution of the ABL by conducting
- numerical sensitivity experiments with our developed one-dimensional (1-D) atmospheric model which consists of the ABL and radiative transfer schemes. Details of the data set and methodologies are described in Sect. 2. The characteristics of the

aerosol vertical profiles and results of the sensitivity experiments conducted with the 1-D atmospheric model are presented in Sect. 3. Our findings are summarized in Sect. 4.

#### 2 Data and methodology

#### 2.1 Remote sensing of aerosol vertical profiles

#### 5 2.1.1 Data retrieval

The vertical profiles of aerosol optical and physical properties were estimated from sky radiometer and lidar data obtained by the SKYLIDAR remote sensing method (Kudo et al., 2016). The sky radiometer (Prede Co., Ltd, Tokyo, Japan), deployed in the SKYNET (Takamura and Nakajima, 2004), is a scanning photometer that measures direct solar radiation and the angular distribution of diffuse radiation. In this study, we used observation data at the Meteorological Research Institute (MRI) of

- 10 Japan Meteorological Agency (JMA) in Tsukuba (36.05°N, 140.12°E, about 25 m above sea level). Note that our data of the sky radiometer at Tsukuba is not transferred to the International SKYNET Data Center (<u>http://www.skynet-isdc.org/index.php</u>). The wavelengths of the sky radiometer data used in this study are 340, 380, 400, 500, 675, 870, and 1020 nm. We also used data from a two-wavelength Mie scattering lidar deployed by AD-Net (Sugimoto et al., 2015) at the National Institute for Environmental Studies (NIES) near MRI. The lidar data consisted of the attenuated backscatter
- 15 coefficients for particle and molecular scattering at 532 and 1064 nm, and the volume depolarization ratio including the contributions of particle and molecular depolarization at 532 nm. The sky radiometer and lidar observation data were collected during 2012 and 2013. As auxiliary data, we used vertical profiles of pressure and temperature from the U.S. National Centers for Environmental Prediction (NCEP) 6-hourly reanalysis data set (Kalnay et al., 1996), total ozone from observations made at the JMA Aerological Observatory (AO) near MRI, and surface albedo from the 5-year climatology of
- 20 the Filled Land Surface Albedo Product, which was generated from the official Terra/MODIS-derived Land Surface Albedo Product (Moody et al., 2005, 2007; Moody 2008). These auxiliary data were used for the calculation of Rayleigh scattering and gas absorption in the SKYLIDAR retrieval. The MRI, NIES and AO instruments are all located within a circle with a radius of 1 km.
- SKYLIDAR estimates aerosol vertical profiles by the following two steps, based on a maximum a posteriori scheme (Kudo et al., 2016). In the first step, the columnar values of the aerosol physical and optical properties (optical thickness, single-scattering albedo, etc.) are estimated by optimizing real and imaginary parts of the refractive index, volume size distribution, and volume ratio of non-spherical particles in the coarse mode to all of the sky radiometer data and the vertical mean of the depolarization ratio of the lidar data. Volume size distribution is assumed to follow a bi-modal lognormal distribution, and the volumes, mode radii, and widths of the fine and coarse modes are estimated. The optical
- 30 properties of non-spherical particles are calculated from a data table of randomly oriented spheroids (Dubovik et al., 2006). In the second step, the vertical profiles of the volume concentrations of fine and coarse modes, the volume ratio of non-

spherical particles in the coarse mode, and the real and imaginary parts of the refractive index are optimized to all of the lidar data and to the optical thickness and single-scattering albedo obtained in the first step. The final outputs are vertical profiles of the extinction coefficient, single-scattering albedo, phase function, the real and imaginary parts of the refractive index, the bi-modal volume size distribution, and the volume ratio of non-spherical particles in the coarse mode. The output

5

wavelengths of the optical properties are 532 and 1064 nm. Note that the mode radii and width of the fine and coarse modes in the second step are fixed by the columnar values obtained in the first step. These outputs enable us to use the radiative transfer model to calculate the vertical profile of the solar heating rate (Kudo et al., 2016).

In the work of Kudo et al. 2016, we conducted the sensitivity tests of the SKYLIDAR using the simulated data of lidar and sky radiometer for the transported dust and pollution aerosol. The pollution aerosol was defined as small-sized and light-absorbing particles. The aerosol optical thickness at 500 nm in the simulation was from 0.05 to 1.2. The random errors were added to the simulated data. The errors were ±2 % for the direct solar radiation, ±3 % for the diffuse radiation, ± 10 % for the attenuated backscatter coefficient, and ±15 % for the volume depolarization ratio. The SKYLIDAR successfully retrieved the columnar values (integrated values or vertical means) of the optical thickness, single-scattering albedo, asymmetry factor, real and imaginary parts of the refractive index, and volume size distribution in all the tests.

- 15 However, the retrieval errors of the vertical profiles increased with a decrease of aerosol optical thickness. In the case that the aerosol optical thickness at 532 nm was 0.05, the retrieval errors were  $\pm 0.003$  km<sup>-1</sup> for the extinction coefficient at 532 nm, and  $\pm 0.05$  for the single-scattering albedo and asymmetry factor at 532 nm. The vertical profiles of the retrieved parameters other than the extinction coefficient had large oscillations due to the random errors of the lidar. In the sensitivity tests for the optical thickness more than 0.1, the vertical profiles of the size distribution, imaginary part of the refractive
- 20 index, extinction coefficient, single-scattering albedo, and asymmetry factor of the transported dust were successfully retrieved, but that of the real part of the refractive index was not. In the transported pollution aerosol case, the vertical profiles of the size distribution, real part of the refractive index, extinction coefficient, and asymmetry factor were estimated well, but those of the imaginary parts of the refractive index and single-scattering albedo were not. When the SKYLIDAR failed to retrieve the vertical profiles of above mentioned parameters, the estimated vertical profiles were uniform, and the
- 25 values were their vertical means.

30

#### 2.1.2 Determination of ABL height

Locally emitted aerosols in the ABL and the transported aerosols in the FA can have different optical properties, which can be evaluated separately after the ABL height has been determined. We estimated the ABL height from the lidar data by the method of Baars et al. (2008), which is based on the wavelet covariance transform (WCT) with the Haar function. The WCT method is less affected by signal noise than the gradient and variance methods. The local maximum of the WCT vertical profile corresponds to the ABL height, and the local minimum corresponds to the base height of clouds or of transported

5

aerosols. In this study, the ABL height was determined by the following procedure:

(1) The attenuated backscatter coefficients at 532 and 1064 nm were normalized by their maximum values below 1000 m, and the WCTs for data at 532 and 1064 nm were calculated.

(2) The local minima and maxima of the WCT vertical profiles at 532 and 1064 nm were detected.

(3) The base height of clouds (or transported aerosols) was searched by using a threshold of -0.1 for the local minimum of

#### 5 the WCT at 532 nm.

10

15

(4) If the base height was not detected in step (3), it was repeated using the WCT at 1064 nm. If the base height was still not detected, it was considered absent or unclear.

(5) The ABL height was searched by using a threshold of 0.05 for the local maximum of the WCT at 532 nm in the day-time, and of the WCT at 1064 nm at night. The top height of the search range was below the base height, if the base height of

clouds or transported aerosols was detected in steps (3) or (4). (6) If the ABL height was not detected, the threshold in step (5) was decreased by -0.01, and the search was repeated until the threshold reached to 0.01.

(7) If the ABL height was not detected in step (6), the search was repeated using the WCT at another wavelength.

(8) If the ABL height was not detected in step (7), the ABL height was considered undetermined.

(9) The time series of ABL height was smoothed by the running-mean with the time window of one hour.

Figure 2 shows the examples of the determined ABL height together with the extinction coefficient estimated by SKYLIDAR. The ABL height could be determined very well when transported aerosols were well above the ABL (Fig 2a). However, when transported aerosols become mixed with the aerosols in the ABL, the ABL height could not be detected (see from 0 to 7 UTC 2 April in Fig. 2b). In this case, we considered aerosols below base height to be in the ABL and those

- above to be in the FA. This assumption causes the uncertainties in evaluating the aerosol optical and physical properties in 20 the FA and ABL. However, it is very difficult to evaluate the uncertainties because the mixing of the transported aerosols with those in the ABL makes the ABL height ambiguous, and the ABL height cannot be detected by the lidar data and our eyes. The successful retrievals of the ABL height by the above procedures from (1) to (9) was about 95 % of the 2,305 profiles under the clear sky conditions, and the base height was used as the ABL height in the remaining profiles. Therefore,
- the influences by our assumption would be small. 25

# 2.2 Model simulation

#### 2.2.1 1-D atmospheric model

We developed a 1-D atmospheric model, consisting of ABL and radiative transfer (RT) schemes, and conducted sensitivity experiments to investigate the radiative impact of aerosols on the evolution of the ABL. The ABL scheme in the model is

based on the ABL model used as the JMA operational mesoscale model for weather forecasting in Japan. The RT scheme is 30 an RT model developed in our laboratory for the remote sensing of aerosols and clouds, and their impacts on the radiative balance in the solar and infrared wavelength regions (Asano and Shiobara, 1989; Nishizawa et al., 2004; Kudo et al., 2011).

The 1-D atmospheric model has a high resolution atmospheric vertical grid with 70 layers from the surface to 40 km. The thickness of the bottom layer is 5 m. Turbulent mixing is calculated by the Mellow-Yamada-Nakanishi-Niino Level 3 scheme (Nakanishi, 2001; Nakanishi and Niino, 2004, 2006), and calculations of surface fluxes are based on the Monin-Obukhov similarity using the universal function of Beljaars and Holtslag (1991). The vertical grid in the soil has 10 layers

5 from the surface to 2 m depth, and the soil temperature is calculated by solving the diffusion equation. The water content in the soil layers was fixed in this study.

In the 1-D atmospheric model, vertical diffusion terms for turbulent mixing and vertical advection by a prescribed vertical motion field are considered for the vertical mixing of potential temperature, specific humidity and the horizontal component of wind. Neither cloud formation nor precipitation is included. In addition, vertical diffusion of aerosols is not considered in the model; aerosol vertical profiles are fixed by the initially given ones.

In the RT scheme, the solar spectrum from 300 nm to 3.0 µm and the infrared spectrum from 4.0 to 50.0 µm are divided into 54 and 19 intervals, respectively. The downward and upward fluxes and the heating rate are calculated by the doubling and adding method (Lacis and Hansen, 1974). Gaseous absorption of water vapor, carbon dioxide, oxygen, and ozone are calculated by the correlated k-distribution method. Scattering at the ground surface is assumed to be Lambert reflection.

15

10

The aerosol parameters input to the RT scheme are the vertical profiles of the extinction coefficient, singlescattering albedo, and the phase function at wavelengths from 300 nm to 3.0 µm. However, the wavelengths of the SKYLIDAR retrievals are limited to 532 and 1064 nm. We determined the refractive index between 532 and 1064 nm by linear interpolation in a log-log plane and used the refractive index at 532 and 1064 nm for wavelengths of less than 532 nm

and greater than 1064 nm, respectively (Kudo et al., 2016). The extinction coefficient, single-scattering albedo, and phase 20 function from 300 nm to 3.0 µm were calculated from these refractive index, the volume size distribution, and the volume ratio of the non-spherical particles in the coarse mode. The influences of aerosols on the infrared wavelength region of more than 3.0 µm were ignored. The heating ratio estimated by this procedure was now validated now, but the surface solar radiation was compared with the measurements of the pyranometer. The difference was small, about 10 W m<sup>-2</sup> (Kudo et al. 2016). 25

#### 2.2.2 Sensitivity experiments

We conducted three types of simulation experiments to investigate the impact of aerosols on the evolution of the ABL. The first type was simulations without aerosols (EXP0), the second was simulations using the observed aerosol vertical profile (EXP1), and the third was the same as the second one but with the entire aerosol vertical profile was compressed into the bottom 1 km (EXP2). Note that the columnar optical properties in EXP1 and EXP2 simulations were the same; only the

30 vertical profile differed between them. Thus, the influences of aerosols can be evaluated from the difference between EXP0 and EXP1 simulations, and the influences of the aerosol vertical profile can be investigated by comparing the results of EXP2 and EXP1 simulations. We conducted experiments using the springtime mean of the aerosol vertical profile and the aerosol vertical profiles observed in the spring during five aerosol transport events in the FA.

To set up the model parameter, we referred to the sensitivity experiments conducted by Yu et al. (2002) and Pandithurai et al. (2008). For our sensitivity experiments, we used the following specified parameters in the 1-D atmospheric

- 5 model. The integration time of all simulations was 24 hours with a time step of 1 minute. The solar orientation was set to that on 5 April 2012 at 36.05°N. The surface albedo was set to the spring mean of the 5-year climatology of the Filled Land Surface Albedo Product (Moody et al., 2005, 2007; Moody 2008). The vertical motion was set to the spring mean of NCEP 6-hourly reanalysis data set. The initial vertical profiles of pressure, temperature, specific humidity, and horizontal wind were also set to the spring means of the NCEP 6-hourly reanalysis data set. The soil surface was assumed to be bare, and the
- 10 heat capacity and thermal conductivity in the soil layers were set to 1.3 × 10<sup>6</sup> J m<sup>-3</sup> K<sup>-1</sup> and 0.3 W m<sup>-1</sup> K<sup>-1</sup>, respectively, based on values for dry sandy clay (Kondo, 1994). The initial temperatures in the soil layers were based on the spring mean of the soil temperature observed at the weather observation field of Mito Meteorological Observatory (Ministry of Agriculture, Forestry, and Fisheries, and Japan Meteorological Agency, 1982), which is 60 km north of the MRI.
- Because the sensible and latent heat fluxes at the surface depend on the water content of the soil, we performed sensitivity experiments for both dry and wet soils. The daily means of the sensible and latent heat fluxes in EXP0 for the dry soil case (volumetric water content 0.1) were 88 and 78 W m<sup>-2</sup>, respectively. For the wet soil case (volumetric water content 0.2), the sensible heat flux was decreased by 22 W m<sup>-2</sup>, and the latent heat flux was increased by 32 W m<sup>-2</sup>. These differences affected the ABL structures (e.g., temperature and specific humidity) in the EXP1 and EXP2 experiments, but not the impacts of aerosols (i.e., the signs of differences, EXP1 – EXP0 or EXP2 – EXP0 were either both positive or both negative in the dry and wet soil cases, and their absolute values were not significantly different). Therefore, we do not show
- the results for the wet soil case in this paper. Thus, the volumetric water content in the soil layers was fixed at 0.1.

#### **3** Results

#### 3.1 Characteristics of aerosol vertical profiles

#### 3.1.1 Seasonal characteristics

- 25 Frequency distributions of the extinction coefficient at 532 nm, based on daily means, were obtained for spring (43 analyzed days), summer (7 days), autumn (35 days), and winter (59 days) (Fig. 3). The small number in summer is due to the lack of completely clear-sky conditions. Summer in Japan is hot and humid, and the cumulous clouds develops almost every day. The SKYLIDAR can be applied to only the clear sky condition. In all the seasons, the extinction coefficient was large in the layer from the surface to 1.5 km altitude. This layer is the ABL, and the aerosols in this layer originate primarily from local
- 30 emissions. In spring and winter, the two large peaks of the extinction coefficients were observed in the layers from 1.5 to 3.5 km and from 3.5 to 6 km altitude. These layers are in the FA, and most aerosols in these layers have been transported long

distances. Transported aerosols are frequently observed in the FA in spring, autumn, and winter, when low-pressure systems carrying aerosols emitted in the eastern region of the Eurasian continent frequently pass over Japan. In summer, Japan is dominated by a high-pressure system, so it receives fewer transported aerosols. In our data, the optical thickness in summer, autumn, and winter were almost the same in the ABL and FA, but in spring, optical thickness in the FA was 0.13, larger than

5 0.08 in the ABL (Table 1).

10

In general, the ABL height is high in summer and low in winter, but in our results, the ABL height was higher in winter and spring, and lower in summer and autumn (Fig. 3). The higher ABL heights in winter and spring can be attributed to the mixing of aerosols between the ABL and FA, which makes it difficult to determine the ABL height (see Sect. 2.1.2). The low ABL height in summer and autumn may be influenced by clouds, which form near the top of the convective mixed layer. Under these circumstances, the ABL height cannot be determined from only lidar data.

The Ångström exponent is a parameter related to particle size: a smaller value indicates a larger particle size. We calculated the vertical profile of the Ångström exponent from the retrievals of the extinction coefficients at 532 and 1064 nm (Fig. 4). The Ångström exponent in the ABL was from 0.0 to 2.0 in all the seasons and increased as the extinction coefficient increased. This result suggests that large extinction coefficients were mainly due to small particles, such as sulfate, nitrate,

- and organics. Conversely, background aerosols consist of large particles, such as locally emitted mineral dust, likely derived 15 from the large areas of bare soil exposed by agriculture and urban development in Tsukuba. The Ångström exponent in the FA ranged from 0.0 to 2.5. This large variability can be attributed to differences in the composition of aerosols, in particular, the proportions of dust and smoke particles, during transport events. The characteristics of five transport events are described in Sect. 3.1.2.
- 20 Single-scattering albedo is an important parameter related to light absorption. In the FA, single-scattering albedo was around 0.95 with small variability (Fig. 5 and Table 1), but in the ABL, it was from 0.87 to 0.99 and decreased as the extinction coefficient decreased. In general, the single-scattering albedo of dust particles is small, whereas that of small particles, other than black carbon, is large (Hess et al. 1998, Aoki et al. 2005). The dependency of the single-scattering albedo on the extinction coefficient in the ABL is therefore consistent with the particle size result shown in Fig. 4.
- 25 The asymmetry factor is an indicator of how much solar energy reaches the surface: a large asymmetry factor value indicates strong forward scattering, which means that more solar energy reaches to surface. In addition, the value of the asymmetry factor is inversely proportional to that of the Angström exponent. In our results, large variation of the asymmetry factor, from 0.4 to 0.8, was observed in the FA (Fig. 6). The asymmetry factor in the ABL was from 0.5 to 0.75, and a dependency of the asymmetry factor on the extinction coefficient was observed.
- 30
- In Sect. 2.1.1, we described that the vertical profiles of the single-scattering albedo and asymmetry factor in the case of the small aerosol optical thickness, less than 0.1, contain large retrieval errors due to the signal noises of the lidar data. We should note that the single-scattering albedo and asymmetry factor where the extinction coefficient was less than 0.02 km<sup>-1</sup> in Figs. 5 and 6 might be contaminated with the retrieval errors.

The values of other important optical and physical parameters are shown in Table 1. These values are particularly useful for comparisons of aerosols in different areas. The values of the real and imaginary parts of the refractive index were from 1.41 to 1.45 and from 0.002 to 0.008, and they were similar values in both the ABL and FA and in all the seasons. When the SKYLIDAR fails to retrieve the vertical profiles of the real part of the refractive index in the transported dust case

- 5 and the imaginary parts of the refractive index in the transported pollution case, the estimated vertical profiles are uniform and the values are their vertical means (Sect. 2.1). This may cause the similar values of the refractive index in the ABL and FA. The values of the mode radius were from 0.11 to 0.14 for the fine mode and from 2.83 to 5.89 for the coarse mode, respectively. They mostly did not differ among seasons, although the coarse mode radius was smaller in spring. The smaller coarse mode radius in spring reflects the relatively small coarse mode radius of the transported aerosols, which ranged from
- 10 1.93 to 3.61 µm (Table 2). In each season, the volume ratio of non-spherical particles in the coarse mode in the ABL was larger than that in the FA, owing to the presence of local dust in the ABL; the smallest value in the ABL was observed in summer, when the ground surface is generally covered with grasses and few dust particles are emitted from the surface. The lidar ratio (extinction to backscatter ratio) is an important parameter for estimating the extinction coefficient, particularly when only the lidar data are available for that purpose. We can calculate the lidar ratio from the single-scattering albedo and
- 15 phase function in the SKYLIDAR retrievals. In our results, no clear seasonal difference was observed in the lidar ratio, and their values were around 60.

#### 3.1.2 Aerosol transport events in the FA

The optical thickness in the FA was largest in spring among all seasons because of presence of transported aerosols. From our results obtained over two years, we selected for further examination five events, which occurred on 2 April 2012 and 16

- 20 April and 8, 9, and 14 May 2013, characterized by large optical thickness in the FA. The daily means of the optical and physical properties of transported aerosols in the FA on these five dates are summarized in Fig. 7 and Table 2. Large extinction coefficients were observed in the FA during these five transport events (Fig. 7a). In addition, we inferred that the aerosols during the events on 2 April 2012, 16 April and 14 May 2013 consisted primarily of transported dust, because on these dates the volume of coarse-mode particles was particularly large (Fig. 7b), Ångström exponent values were small, less
- 25 than 1.0, and the volume ratio of non-spherical particles was large, from 0.84 to 0.99 (Table 2). The two-day backward trajectories for those events (Figs. 7c and d) suggest that the transported dust originated in desert areas of China and Mongolia. On 8 May 2013, the volume of fine-mode particles was very large (Fig. 7b), the Ångström exponent was also large, 1.82, and the volume ratio of non-spherical particles was smallest in all the cases (Table 2); these results indicate that the aerosols consisted dominantly of small and spherical particles. The backward trajectory (Figs. 7c and d) indicated that the
- 30 source region was in Russia, to the southeast area of Lake Baikal, where a forest fire had been observed in early May 2013. Therefore, we interpreted this transported aerosol to consist of transported smoke particles were from that forest fire. The following day, 9 May 2013, the source had moved to northeastern China (Figs. 7c and d), and the volume of coarse-mode

particles was large (Fig. 7b); these results suggest that this aerosol may have consisted of transported smoke and dust particles.

The single-scattering albedo and asymmetry factor at 532 nm of the dust cases (2 April 2012, and 16 April and 14 May 2013), and the smoke and dust mixture case (9 May 2013) were from 0.95 to 0.98 and from 0.65 to 0.71, respectively

- 5 (Table 2). Dubovik et al. (2002) summarized the AERONET retrievals in the world and showed the single-scattering albedo and asymmetry factor at visible wavelengths of 0.44 and 0.69 μm in the desert regions were from 0.92 to 0.98 and from 0.66 to 0.73, respectively. Moreover, the single-scattering albedo estimated from the sky radiometer for the Asian dust was from 0.91 to 0.97 (Uchiyama et al., 2005). These retrievals were the columnar values, but the cases that the coarse mode was dominant were selected. Our results were consistent with these results. The refractive index in this study was from 1.43 to
- 10 1.53 for the real part, and from 0.001 to 0.004 for the imaginary part, respectively. Aoki et al. (2005) summarised the refractive index of the dust from the reports of the various works, and showed that the real and imaginary parts at 500 nm are from 1.45 to 1.55, and from 0.0005 to 0.008, respectively. The mode radius for the coarse particles in this study was from 1.93 to 3.61 µm, and the AERONET retrievals in the desert regions were from 1.9 to 2.7 µm (Dubovik et al., 2002). The lidar ratio at 532 nm in this study was from 47 to 56, and the results of High Spectral Resolution Lidar or Raman lidar were
- 15 from 20 to 70 (Burton et al., 2012; Groß et al., 2015). Consequently, our retrieved physical and optical properties of the transported dust were consistent with those reported in other studies.

For the transported smoke on 9 May 2013, the single-scattering albedo and asymmetry factor were 0.97 and 0.64, respectively. The SKYLIDAR fails to retrieve the vertical profile of the single-scattering albedo of the transported pollution aerosol (small-sized and light-absorbing particle). However, the estimated vertical profile is uniform, and the estimated value

- 20 is the vertical mean (Sect. 2.1.1). Therefore, our estimated single-scattering albedo can be compared with that of the AERONET retrievals. Furthermore, since the extinction coefficient in the ABL was much smaller than that in the FA (Fig. 7a), the vertical mean of the single-scattering albedo would represent the transported smoke in the FA. The AERONET retrievals at visible wavelengths for the biomass-burning aerosols in Amazonian forest (Brazil), South American cerrado (Brazil), African savanna (Zambia), and Boreal forest (United Saes and Canada) were from 0.84 to 0.94 for the single-
- 25 scattering albedo, and from 0.53 to 0.69 for the asymmetry factor (Dubovik et al., 2002). Our estimated asymmetry factor of 0.64 lied among these values, but the single scattering albedo of 0.97 was higher than the AERONET retrievals. In general, the smoke from the biomass-burning is composed of black carbon, organic carbon, and inorganic materials (Reid et al., 2005). The single-scattering albedo strongly depends on the fuel type and the burning conditions and ranges from 0.2 to 1.0 by depending on the ratio of black carbon (or elemental carbon) to organic carbon in the FLAME-4 experiment (Liu et al.,
- 30 2013; Pokhrel et al., 2016). The AERONET retrievals for the boreal biomass-burning aerosols in the Alaska showed the single-scattering albedo in 2004 and 2005 was about 0.96 (Eck et al., 2009). They suggested a significant amount of smoldering combustion of woody fuels and peat/soil layers that would result in relatively low black carbon mass fractions for smoke particles. The black carbon fraction of the our analysed transported smoke also might be low. The refractive index of the smoke in this study was 1.42 for the real part and 0.003 for the imaginary part, respectively. These values were smaller
  - 11

than those of the AERONET retrievals in above mentioned regions, from 1.47 to 1.52 for the real part and from 0.00093 to 0.021 for the imaginary part, respectively. Moreover, our results of the refractive index were similar to those of the water-soluble aerosols in the OPAC, which originates from gas to particle conversion and consists of various kind of sulfates, nitrates, and other, also organic, water-soluble substances (Hess et al., 1998). This supports the low black carbon fraction and

5 large single-scattering albedo for the smoke in this study. It is possible that the black carbon fraction was decreased in the long-range transport from Russia to Japan by the increase of the water-soluble aerosols. The lidar ration of 61 at 532 nm in this study lied among the range of the observations by High Spectral Resolution Lidar or Raman lidar, from 50 to 100 (Burton et al. 2012; Groß et al. 2015).

#### 3.2 Sensitivity experiment results

#### 10 **3.2.1 Impact of aerosols on the evolution of the ABL**

Figure 8 and Table 3 show the results of EXP0 and EXP1 – EXP0. The net downward surface radiation in the solar and infrared wavelength regions, as well as the sensible and latent heat fluxes, were decreased in the EXP1 simulations (with aerosols) compared with EXP0 simulation (without aerosols) (Figs. 7a–c). The change in the daily mean ranged from -14 to - 23 W m<sup>-2</sup> for the net downward radiation, from -7 to -11 W m<sup>-2</sup> for the sensible heat flux, and from -6 to -10 W m<sup>-2</sup> for the

- 15 latent heat flux (Table 3). Absorption by the ground was also decreased; the change in the daily ranged from -1.3 to -2.2 Wm<sup>-2</sup> (Table 3). In general, the downward surface solar radiation becomes small when optical thickness is large, single-scattering albedo is small, and the asymmetry factor is small, (Kudo et al., 2010b). The single-scattering albedo and asymmetry factor were not very different between the springtime mean and the five transport events (Tables 1 and 2), so the reduction of the net downward radiation (Fig. 8a) mainly reflects the optical thickness of the column (Table 3), and the
- 20 reductions of the sensible and latent heat fluxes were caused by the decrease of the net downward radiation. The potential temperature profile at noon local time decreased in the ABL owing to the decline in the sensible heat flux (Fig. 8d). Note that in the 1-D atmospheric model results, the latent heat flux could not warm the atmosphere in the ABL because condensation is not included in the model. The daily maximum 2 m temperature was decreased by 0.2 to 0.6 K (Table 3). In contrast, the potential temperature was increased by 0.0 to 0.4 K in the FA owing to the direct heating of transported aerosols (Fig. 8d).
- 25 The vertical profiles of direct heating depended on the profiles of the extinction coefficient (Fig. 7a). The warming of the FA and the cooling of the ABL stabilized the atmosphere and strengthened the capping inversion around the top of the ABL. The strengthened capping inversion and the decline of the sensible heat flux caused the ABL height to decease by -133 to 208 m (Fig. 8f and Table 3).

The latent heat flux, that is, the water vapor flux, was apparently decreased by aerosols, but the change in the 30 amount of surface evaporation was small, from -0.21 to -0.36 kg m<sup>-2</sup> day<sup>-1</sup> (Fig. 8c and Table 3). The change in the vertical profile of specific humidity was very small, but the specific humidity around the top of the ABL was decreased as a result of the decrease in the ABL height and the dry air in the FA (Fig. 8e).

### 3.2.2 Impact of the aerosol vertical profile on the evolution of the ABL

Figure 9 and Table 3 show the results of EXT0 and EXP2 – EXP0. Note that the entire aerosol vertical profile was compressed in the bottom 1 km in the EXP2 simulations, but the optical thickness of the column was the same as that in the EXP1 simulations. The influence of only the aerosol vertical profile can thus be investigated by comparing Figs. 8 and 9.

- 5 The amounts of decrease in the net downward radiation and the sensible and latent heat fluxes in the EXP2 simulations were almost the same as those in the EXP1 simulations (Figs. 9a-c). However, the decrease in the potential temperature in the ABL was about -0.1 K at noon and was smaller in EXP2 than in EXP1 (Fig. 9d), because aerosol direct heating in the ABL was stronger in EXP2 than in EXP1. The changes in surface evaporation (Table 3) and specific humidity (Fig. 9e) in EXP2 were similar to those in EXP1. The aerosol direct heating in the ABL, together with the lack of direct heating in the FA,
- 10 weakened the capping inversion around the top of the ABL. Therefore, the decrease in the ABL height was from -208 to -133 m in EXP1 but those in EXP2 was from -90 to -24 m (Table 3). Thus, the evolution of the ABL was changed by the aerosol vertical profile, even though the columnar characteristics of the aerosol optical properties were the same. The impacts of aerosols on the ABL structure, that is, reductions of the temperature in the ABL and of the ABL height, were larger when aerosols were present in the FA.

#### 15 3.2.3 Relations between the aerosol physical and optical properties in the FA and the evolution of the ABL

The relations between the aerosol physical and optical properties in the FA and the evolution of the ABL was investigated using the EXP1 results for the springtime mean and five transport events. Fig. 10 shows the dependencies of the decreases in the daily maximums of the 2 m temperature and ABL height on the optical thickness and Ångström exponent in the FA. We focused on the influences of Ångström exponent to the ABL evolution because the single-scattering albedo and asymmetry

- 20 factor in the springtime mean and 5 events had similar values (Table 2). The solid lines in Fig. 10 are the simulation results for different optical thickness and Ångström exponent in the FA with the simplified aerosol vertical profile, where the vertical profiles of the aerosol physical and optical properties are uniform in the ABL (from the surface to 2 km altitude) and FA (from 2 to 6 km altitude), respectively. The physical and optical properties used in the simulations are summarized in Table 4 and are the means calculated from the results of the springtime mean and five transport events. The optical thickness
- 25 and Ångström exponent in the FA were changed by using the different values of the total volume and volume ratio of the fine and coarse modes for the size distribution.

The 2 m temperature and the ABL height was decreased with an increase of the optical thickness (Fig. 10). This influence of aerosols was described in Sect. 3.2.1. We found that the decrease rates of the 2 m temperature and ABL height increased with a decrease of the Ångström exponent. The small value of Ångström exponent indicates the large optical thickness in the near infrared wavelengths region. Therefore, the large particles such as dust weakened the ABL evolution

efficiently due to the influences for both the visible and near infrared wavelength regions. However, the plots for the springtime mean and five transport events in Fig. 10 were not completely consistent with the solid lines for the simulation



results because the aerosol vertical profiles used in the simplified simulations were different from those for the springtime mean and five transport events. Particularly, the decrease in the ABL height in 9 May 2013 (smoke and dust case) was larger than that in 2 April 2012 (dust case). This result is opposite to above mentioned influence of Ångström exponent. In both cases, the optical thickness in the FA was similar value, about 0.33. However, the geometric thickness of the aerosol layer in

- 5 the FA was about 3 km, smaller 5 km in 9 May 2013 than in 2 April 2012, and the extinction coefficient from 1.0 to 3.5 km altitude was larger in 9 May 2013 than in 2 April 2012 (Fig. 7a). This resulted in the largest temperature increase in the FA and the strong capping inversion in 9 May 2013 (Fig. 8d). Consequently, the ABL height was low in 9 May 2013. The dependencies of the 2 m temperature and ABL height on the Ångström exponent were found in this study, but the most important factor was the vertical profile of the extinction coefficient, in particular the extinction coefficient around the top of
- 10 the ABL.

15

#### **4** Conclusion

We first investigated vertical profiles of aerosol physical and optical properties at Tsukuba, Japan, with focus on the seasonal means and on five aerosol transport events, by a synergistic remote sensing method (SKYLIDAR) using sky radiometer and lidar data in two years (2012 and 2013). Second, we investigated the impact of the aerosol vertical profile on the evolution of the ABL by conducting sensitivity experiments with our 1-D atmospheric model.

The vertical profiles of the seasonal mean extinction coefficients showed high loads of the locally emitted aerosols in the ABL, from the surface to 1.5 km altitude, and the transported aerosols in the FA, from 1.5 to 6 km altitude. In summer, autumn, and winter, the aerosol optical thickness was almost same in the ABL and FA. In spring, the optical thickness at 532 nm in the FA was 0.13 and was larger than 0.08 in the ABL.

- 20 The physical and optical properties of the aerosols in the ABL were dependent on the extinction coefficient: As the extinction coefficient increased from 0.02 to 0.24 km<sup>-1</sup>, the Ångström exponent was increased from 0.0 to 2.0, the single-scattering albedo increased from 0.87 to 0.99, and the asymmetry factor decreased from 0.75 to 0.5. These characteristics suggests that the background aerosols consisted of the locally emitted dust particles, and the large extinction coefficient was attributed to an increase of the small and non-absorbing particles.
- 25 The optical and physical properties in the FA varied greatly owing to the presence of transported aerosols. We investigated the vertical profiles and backward trajectories of five transport events. In three events, the aerosol consisted of dust particles transported from desert regions of China and Mongolia. In one event, the aerosol consisted of small smoke particles transported from a forest fire in Russia. The aerosols of a fifth event consisted of both small and large particles which we interpreted as smoke and dust particles, respectively. The single-scattering albedo and asymmetry factor of the
- 30 transported dust particles and mixture of dust and smoke particles were from 0.95 to 0.98 and from 0.65 to 0.71, respectively. These values were consistent with those reported in the other works for the Asian dust and the desert regions in the world. In the transported smoke case, the asymmetry factor was 0.64 and was consistent with the reports for the biomass-burning

aerosols in the world. However, the single-scattering albedo was 0.97 and was higher than the other reports. It is supposed that the black carbon fraction was low in the source of the smoke, or the black carbon fraction decreased in the long-range transport from Russia to Japan.

We conducted sensitivity experiments in which the aerosol vertical profiles of the springtime mean and the five
transport events were input into our 1-D atmospheric model. The sensitivity experiments with (EXP1) and without aerosols (EXP0) showed that the aerosols decreased net downward surface radiation (-14 to -23 W m<sup>-2</sup>) and sensible and latent heat fluxes (-7 to -11, and -6 to -10 W m<sup>-2</sup>, respectively). These resulted in the decrease of the maximum 2 m temperature (-0.2 to -0.6 K). The decrease of the temperature in the ABL and the direct heating of the transported aerosols in the FA strengthened the capping inversion near the top of the ABL. Consequently, the ABL height was less developed in EXP1 simulations than
in the EXP0 simulations, and the decrease of the ABL height due to aerosols was from -133 to -208 m.

To investigate the impact of only the aerosol vertical profile on the evolution of the ABL, we conducted simulations (EXP2) in which all aerosols were compressed into the ABL (0-1 km altitude), but in which the columnar optical thickness was the same as that in the EXP1 simulations. The net downward radiation and the sensible and latent heat fluxes were not changed, but the ABL height was increased, in EXP2 simulations compared with EXP1 simulations. This increase in the ABL height resulted from a weakened capping inversion caused by aerosol direct heating in the ABL and the lack of direct

15 ABL height resulted from a weakened capping inversion caused by aerosol direct heating in the ABI heating in the FA.

Using the results of EXP1 simulations for the springtime mean and five transport events, the dependencies of the decreases in the 2 m temperature and ABL height on the optical thickness and Ångström exponent in the FA were investigated. The 2 m temperature and ABL height were decreased with an increase of the optical thickness, and their decrease rates depended on Ångström exponent. The 2 m temperature and ABL height efficiently decreased in the case that

- 20 decrease rates depended on Ångström exponent. The 2 m temperature and ABL height efficiently decreased in the case that the Ångström exponent was small because of the large optical thickness in the near infrared wavelength region. However, in the case of the smoke and dust mixture event, the Ångström exponent was large, but the decrease of the ABL height was largest in the springtime mean and five events. The extinction coefficient around the top of the ABL was largest in the five transport events, and the strong capping inversion resulted in the lowest ABL height.
- 25

These sensitivity experiment results suggest that the vertical profiles of the aerosol physical and optical properties, and resulting direct heating are essential factors in the evolution of the ABL. Moreover, it is particularly important to characterize aerosol optical properties in the FA because aerosols in the FA can be transported widely and therefore affect the ABL both regionally and globally.

Our 1-D atmospheric model did not consider cloud formation or precipitation, although both of these can be 30 affected by aerosol-induced modification of atmospheric stability. In the future, we plan to develop a 1-D or 3-D model that includes these processes and investigate aerosol-cloud interactions by inputting the observed aerosol data into the models.

# 5 Data availability

The lidar data are available from the AD-Net (<u>http://www-lidar.nies.go.jp</u>). The sky radiometer data are available from the International SKYNET Data Center (<u>http://www.skynet-isdc.org/index.php</u>), but the sky radiometer data at Tsukuba, Japan are available on request by contacting the first author of the paper.

5

#### Acknowledgements.

This work was supported by the Japan Society for the Promotion of Science KAKENHI grant nos. 24510026, 15H01728, and 15H02808. NCEP reanalysis data were provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from its website at <a href="http://www.esrl.noaa.gov/psd/">http://www.esrl.noaa.gov/psd/</a>. The MODIS MCD12C1 product was retrieved from the online data pool, courtesy

10 of the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota (<u>https://lpdaac.usgs.gov/data\_access/data\_pool</u>).

# References

- Aoki, T., Tanaka, T. Y., Uchiyama, A., Chiba, M., Mikami, M., Yabuki, S., and Key, J. R.: Sensitivity experiments of direct forcing caused by mineral dust simulated with a chemical transport model, J. Meteorol. Soc. Jpn., 83A, 315-331, 2005.
- Asano, S. and Shiobara, M.: Aircraft measurements of the radiative effects of tropospheric aerosols: I. Observational Results

5 of the Radiation Budget, J. Meteorol. Soc. Japan, 847–861, 1989.

- Baars, H., Ansmann, A., Engelmann, R. and Althausen, D.: Continuous monitoring of the boundary-layer top with lidar, Atmos. Chem. Phys., 8, 7281–7296, doi:10.5194/acpd-8-10749-2008, 2008.
- Beljaars, A. C. M., Holtslag, A. A. M.: Flux Parameterization over Land Surfaces for Atmospheric Models, J. Appl. Meteorol., 30, 327–341, 1991.
- Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Bulter, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne high spectral resolution lidar measurements methodology and examples, Atmos. Meas. Tech., 5, 73-98, 2012.
  - Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.: Varaiability of absorption and optical properties of key aerosol types observed in worldwide locations, J. Atmos., Sci., 59, 590-608, 2002.
- 15 2002
  - Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W. J., Leon, J. F., Sorokin, M. and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, J. Geophys. Res. Atmos., 111(11), 1–34, doi:10.1029/2005JD006619, 2006.
- 20 Eck, T. E., Holben, B. N., Reid, J. S., Sinyuk, A., Hyer, E. J., O'Neill, N. T., Shaw, G. E., Vande Castle, J. R., Chapin, F. S., Dubovik, O., Smirnov, A., Vermote, E., Schafer, J. S., Giles, D., Slutsker, I., Sorokine, M., and Newcomb, W. W.: Optical properties of boreal region biomass burning aerosols in central Alaska and seasonal variation of aerosol optical depth at an Arctic coastal site, J. Geophys. Res., 114, D11201, doi:10.1029/2008JD010870, 2009.
- Groß, S., Freudenthaler, V., Schepanski, K., Toledano, C., Schäfler, A., Ansman, A., and Weinzierl, B.: Optical properties of
- 25 long-range transported saharan dust over Barbados as measured by dual-wavelength depolarization Raman Idiar measurements, Atmos. Chem. Phys., 15, 11067-11080, 2015.
  - Hess, M., Koepke, P., and Schult, I.: Optical properties of aerosols and clouds: the software package OPAC, B. Am. Meteorol. Soc., 79, 831-844, 1998.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report
   of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
  - Boschung, A. Nauels, Y. Xia, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.

- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project, Bull. Am. Meteorol. Soc., 437– 471, 1996.
- Kondo, J.: Mizukankyo-no-kisyougaku (in Japanese), 2nd editio., Asakura Publishing Co. Ltd., Tokyo., 1994.
   Kudo, R., Uchiyama, A., Yamazaki, A., Sakami, T. and Kobayashi, E.: From solar radiation measurements to optical properties: 1998-2008 trends in Japan, Geophys. Res. Lett., 37(4), 1–6, doi:10.1029/2009GL041794, 2010a.

Kudo, R., Uchiyama, A., Yamazaki, A. and Kobayashi, E.: Seasonal characteristics of aerosol radiative effect estimated from ground-based solar radiation measurements in Tsukuba, Japan, J. Geophys. Res. Atmos., 115(1), 1–10,

- 10 doi:10.1029/2009JD012487, 2010b.
- Kudo, R., Uchiyama, A., Yamazaki, A., Sakami, T. and Ijima, O.: Decadal changes in aerosol optical thickness and single scattering albedo estimated from ground-based broadband radiometers: A case study in Japan, J. Geophys. Res. Atmos., 116(3), 1–14, doi:10.1029/2010JD014911, 2011.

Kudo, R., Nishizawa, T. and Aoyagi, T.: Vertical profiles of aerosol optical properties and the solar heating rate estimated by

- combining sky radiometer and lidar measurements, Atmos. Meas. Tech., 9(7), 3223–3243, doi:10.5194/amt-9-3223-2016, 2016.
  - Lacis, A., A., Hansen, J. E.: A Parameterization for the Absorption of Solar Radiation in the Earth's Atmosphere, , 31, 118–133, 1974.

- 20 L., DeMott, P. J., and Kreidenweis, S. M.: Aerosol single scattering albedo dependence on biomass combustion efficiency: laboratory and field studies, Geophys. Res. Lett., doi:10.1002/2013GL058392, 2013.
  - LP DAAC: Land Processes Distributed Active Archive Center: The Land Cover Type Climate Modeling Grid (CMG) product (MCD12C1). Version 051. NASA EOSDIS Land Processes DAAC, USGS Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota, [online] Available from:
- 25 https://lpdaac.usgs.gov/dataset\_discovery/modis/modis\_products\_table/mcd12c1 (Accessed 20 May 2007), 2013.
  - Ministry of Agriculture, Forestry, and Fisheries, and J. M. A.: Chichu-ondo-nado-ni-kansuru-shiryou [Report about the soil temperature] (in Japanese), Tokyo., 1982.
    - Moody, E. G.: MODIS-derived spatially complete surface albedo products: Spatial and temporal pixel and zonal averages, J. Appl. Meteorol. Climatol., 47(11), 2879–2894, doi:10.1175/2008JAMC1795.1, 2008.
- 30 Moody, E. G., King, M. D., Platnick, S., Schaaf, C. B. and Gao, F.: Spatially complete global spectral surface albedos: Value-added datasets derived from terra MODIS land products, IEEE Trans. Geosci. Remote Sens., 43(1), 144–157, doi:10.1109/TGRS.2004.838359, 2005.

Liu S., Aiken, A. C., Arata, C., Dubey, M. K., Stockwell, C. E., Yokelson, R. J., Stone, E. A., Jayarathne, T., Robinson, A.

- Moody, E. G., King, M. D., Schaaf, C. B., Hall, D. K. and Platnick, S.: Northern Hemisphere five-year average (2000-2004) spectral albedos of surfaces in the presence of snow: Statistics computed from Terra MODIS land products, Remote Sens. Environ., 111(2), 337–345, doi:10.1016/j.rse.2007.03.026, 2007.
- Nakanishi, M.: Closure Model Based on Large-Eddy Simulation Data, , 349-378, 2001.
- 5 Nakanishi, M. and Niino, H.: An improved Mellor-Yamada Level-3 model with condensation physics: Its design and verification, Boundary-Layer Meteorol., 112(1), 1–31, doi:10.1023/B:BOUN.0000020164.04146.98, 2004.
  - Nakanishi, M. and Niino, H.: An improved Mellor-Yamada Level-3 model: Its numerical stability and application to a regional prediction of advection fog, Boundary-Layer Meteorol., 119(2), 397–407, doi:10.1007/s10546-005-9030-8, 2006.
- 10 NEO: NASA Earth Observatory: The imagery of MODIS active fire, [online] Available from: http://neo.sci.gsfc.nasa.gov/ (Accessed 16 July 2016), 2016.
  - Nishizawa, T., Asano, S., Uchiyama, A. and Yamazaki, A.: Seasonal variation of aerosol direct radiative forcing and optical properties estimated from ground based radiative measurements, J. Atmos. Sci., 61, 57–72, 2004.
- Pandithurai, G., Seethala, C., Murthy, B. S. and Devara, P. C. S.: Investigation of atmospheric boundary layer characteristics
  for different aerosol absorptions: Case studies using CAPS model, Atmos. Environ., 42(19), 4755–4768,
  doi:10.1016/j.atmosenv.2008.01.038, 2008.
  - Pokhrel, R. P., Wagner, N. L., Langridge, J. M., Lack, D. A., Jayarathne, T., Stone, E. A., Stockwell, C. E., Yokeison, R. J., and Murphy, S. M.: Pramererization of single-scattering albedo (SSA) and absorption Ångström exponent (AAE) with EC / OC for aerosol emissions from biomass buring, Atmos. Chem. Phys., 16, 9549-9561, doi:10.5194/acp-16-9549-
- 20 2016, 2016
  - Sugimoto, N., Nishizawa, T., Shimizu, A., Matsui, I., Jin, Y., Higurashi, A., Uno, I., Hara, Y., Yumimoto, K. and Kudo, R.: Continuous observations of atmospheric aerosols across East Asia, SPIE Newsroom, 2–5, doi:10.1117/2.1201510.006178, 2015.

Takamura, T. and Nakajima, T.: Overview of SKYNET and its activities, Opt. Pura Apl, 37(3), 3303-3308, 2004.

- 25 Tsunematsu, N., Sakai, T. and Nagai, T.: Strong capping inversion over the Tokyo metropolitan area associated with airborne Asian dust, Geophys. Res. Lett., 33(19), 1–5, doi:10.1029/2006GL026645, 2006.
  - Uchiyama, A., Yamazaki, A., Togawa, H., Asano, J., and Shi, G.: Single scattering albedo of aeolian dust as inferred from sky-dradiometer and in situ ground-based measurement, SOLA, 1, 209-212, doi:10.2151/sola2005-054, 2005.
- Uno, I., Eguchi, K., Yumimoto, K., Takemura, T., Shimizu, A., Uematsu, M., Liu, Z., Wang, Z., Hara, Y. and Sugimoto, N.:
  Asian dust transported one full circuit around the globe, Nat. Geosci., 2(8), 557–560, doi:10.1038/ngeo583, 2009.
- Yu, H., Liu, S. C. and Dickinson, R. E.: Radiative effects of aerosols on the evolution of the atmospheric boundary layer, J. Geophys. Res., 107, doi:10.1029/2001JD000754, 2002.

Table 1. Seasonal means and standard deviations of aerosol optical and physical properties.

| Physical and optical properties       |                | Spring            |                   | Summer            |                   | Autumn            |                   | Winter            |                   |
|---------------------------------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
|                                       |                | ABL               | FA                | ABL               | FA                | ABL               | FA                | ABL               | FA                |
| Optical thickness <sup>a</sup>        |                | $0.08 \pm 0.03$   | $0.13 \pm 0.08$   | $0.07 \pm 0.02$   | $0.07 \pm 0.05$   | $0.05 \pm 0.02$   | $0.05 \pm 0.02$   | $0.05 \pm 0.02$   | $0.06 \pm 0.04$   |
| Ångström exponent                     |                | $0.81 \pm 0.36$   | $0.97 \pm 0.52$   | $0.84 \pm 0.46$   | $1.53 \pm 0.17$   | $1.18 \pm 0.35$   | $1.05 \pm 0.26$   | $1.06 \pm 0.40$   | $1.11 \pm 0.38$   |
| Single scattering albedo <sup>a</sup> |                | $0.93 \pm 0.03$   | $0.96 \pm 0.01$   | $0.92 \pm 0.05$   | $0.92 \pm 0.06$   | $0.96 \pm 0.03$   | $0.95 \pm 0.03$   | $0.96 \pm 0.02$   | $0.96 \pm 0.03$   |
| Asymmetry facto                       | r <sup>a</sup> | $0.70 \pm 0.03$   | $0.68 \pm 0.03$   | $0.71 \pm 0.04$   | $0.66 \pm 0.04$   | $0.66 \pm 0.03$   | $0.67 \pm 0.03$   | $0.66 \pm 0.04$   | $0.66 \pm 0.04$   |
| Refractive                            | Real part      | $1.44 \pm 0.05$   | $1.46 \pm 0.04$   | $1.41 \pm 0.03$   | $1.41 \pm 0.02$   | $1.42 \pm 0.04$   | $1.41 \pm 0.03$   | $1.42 \pm 0.05$   | $1.42 \pm 0.03$   |
| index <sup>a</sup>                    | Imaginary part | $0.006 \pm 0.004$ | $0.003 \pm 0.002$ | $0.006 \pm 0.004$ | $0.008 \pm 0.006$ | $0.003 \pm 0.002$ | $0.003 \pm 0.002$ | $0.002 \pm 0.002$ | $0.002 \pm 0.002$ |
| Mode radius (µm)                      | ) fine         | 0.14=             | ±0.02             | 0.14              | ±0.04             | 0.12              | ±0.03             | 0.11              | ±0.02             |
|                                       | coarse         | 2.83              | ±1.45             | 4.59              | ±1.37             | 4.70              | ±2.00             | 5.89              | ±2.30             |
| Mode width fine coarse                |                | $0.46 \pm 0.13$   |                   | $0.59 \pm 0.09$   |                   | $0.53 \pm 0.13$   |                   | $0.60 \pm 0.13$   |                   |
|                                       |                | $0.92 \pm 0.10$   |                   | $0.98 \pm 0.02$   |                   | $0.98 \pm 0.02$   |                   | $0.97 \pm 0.06$   |                   |
| Volume ratio of non-spherical         |                | 0.96+0.06         | $0.85 \pm 0.22$   | $0.79 \pm 0.20$   | $0.68 \pm 0.21$   | 0.95 + 0.08       | $0.91 \pm 0.09$   | $0.97 \pm 0.07$   | $0.86 \pm 0.14$   |
| particles in the coarse mode          |                | 0.90±0.00         | 0.85 ± 0.22       | 0.79±0.20         | 0.08±0.21         | 0.95 ± 0.08       | 0.91 ± 0.09       | 0.97±0.07         | 0.80±0.14         |
| Lidar ratio <sup>a</sup>              |                | 69±10             | 58±7              | 68±23             | 65±13             | $57 \pm 10$       | 63±10             | 56±9              | $56 \pm 10$       |

| Physical and op  | otical  |              |               |            |            |             |
|--|---|--------------|---------------|------------|------------|-------------|
| properties   |   | 2 April 2012 | 16 April 2013 | 8 May 2013 | 9 May 2013 | 14 May 2013 |
| Optical thickness<br>at 532 nm                         |   | 0.33         | 0.24          | 0.27       | 0.33       | 0.25        |
| Ångström expo  | Ångström exponent                                 |              | 0.47          | 1.82       | 1.28       | 0.78        |
| Single-scatterin<br>at 532 nm                          | Single-scattering albedo<br>at 532 nm             |              | 0.97          | 0.97       | 0.96       | 0.95        |
| Asymmetry fac<br>at 532 nm                             | Asymmetry factor<br>at 532 nm                     |              | 0.71          | 0.64       | 0.65       | 0.68        |
| Real part<br>of the refractiv<br>at 532 nm             | Real part<br>of the refractive index<br>at 532 nm |              | 1.43          | 1.42       | 1.53       | 1.48        |
| Imaginary part<br>of the refractive index<br>at 532 nm |   | 0.001        | 0.001         | 0.003      | 0.004      | 0.004       |
| Mode radius  | fine  | 0.15         | 0.13          | 0.14       | 0.15       | 0.15        |
| (µm)   | coarse  | 2.43         | 2.28          | 2.15       | 3.61       | 1.93        |
| Mode width   | fine  | 0.31         | 0.46          | 0.43       | 0.44       | 0.48        |
|  | coarse  | 0.90         | 0.89          | 0.98       | 0.98       | 0.77        |
| Volume ratio o   | f   |              |               |            |            |             |
| non-spherical particles                                |   | 0.99         | 0.97          | 0.34       | 0.96       | 0.84        |
| in the coarse m  | ode   |              |               |            |            |             |
| Lidar ratio at 5.                                      | Lidar ratio at 532 nm                             |              | 56            | 61         | 55         | 56          |

Table 2. Daily means of optical and physical properties of transported aerosols in the FA

|             | Aerosol      | Daily mean           | Daily                | Daily                | Daily                |             | Daily      | Daily                                 |            |
|-------------|--------------|----------------------|----------------------|----------------------|----------------------|-------------|------------|---------------------------------------|------------|
|             | optical      | net                  | mean                 | mean                 | mean                 | Daily mean  | maximum    | integrated                            | Daily      |
|             | thickness in | downward             | sensible             | latent               | ground               | of 2 m      | of 2 m     | surface                               | maximum    |
|             | the column   | radiation            | heat flux            | heat flux            | absorption           | temperature | temperatur | evaporation                           | ABL        |
|             | (532 nm)     | (W m <sup>-2</sup> ) | (K)         | e (K)      | $(\text{kg m}^{-2} \text{ day}^{-1})$ | height (m) |
| EXP0        |              |                      |                      |                      |                      |             |            |                                       |            |
| Spring mean | 0.0          | 166                  | 88                   | 78                   | -0.6                 | 285         | 293        | 2.68                                  | 2352       |
| EXP1 – EXP0 |              |                      |                      |                      |                      |             |            |                                       |            |
| Spring mean | 0.21         | -14                  | -7                   | -6                   | -1.3                 | -0.3        | -0.2       | -0.21                                 | -133       |
| 2 Apr 2012  | 0.38         | -22                  | -10                  | -10                  | -2.2                 | -0.5        | -0.6       | -0.34                                 | -186       |
| 16 Apr 2013 | 0.35         | -19                  | -9                   | -8                   | -1.8                 | -0.4        | -0.4       | -0.29                                 | -162       |
| 8 May 2013  | 0.32         | -18                  | -8                   | -8                   | -1.7                 | -0.4        | -0.4       | -0.27                                 | -150       |
| 9 May 2013  | 0.42         | -23                  | -11                  | -10                  | -2.2                 | -0.5        | -0.5       | -0.36                                 | -208       |
| 14 May 2013 | 0.37         | -22                  | -11                  | -10                  | -2.1                 | -0.5        | -0.5       | -0.33                                 | -163       |
| EXP2 - EXP0 |              |                      |                      |                      |                      |             |            |                                       |            |
| Spring mean | 0.21         | -14                  | -7                   | -5                   | -1.2                 | -0.2        | -0.1       | -0.18                                 | -24        |
| 2 Apr 2012  | 0.38         | -22                  | -11                  | -9                   | -1.9                 | -0.4        | -0.2       | -0.30                                 | -72        |
| 16 Apr 2013 | 0.35         | -19                  | -9                   | -8                   | -1.7                 | -0.3        | -0.2       | -0.26                                 | -83        |
| 8 May 2013  | 0.32         | -18                  | -9                   | -7                   | -1.6                 | -0.3        | -0.2       | -0.25                                 | -77        |
| 9 May 2013  | 0.42         | -23                  | -12                  | -9                   | -2.0                 | -0.4        | -0.3       | -0.32                                 | -90        |
| 14 May 2013 | 0.37         | -22                  | -11                  | -9                   | -1.9                 | -0.3        | -0.2       | -0.30                                 | -70        |

Table 3: Results of EXP0, EXP1, and EXP2 sensitivity experiments

Table 4: Physical and optical properties used in the simulations for the simplified aerosol vertical profiles in Fig. 10

| Physical and optical p  | roperties                               | ABL                        | FA                   |  |
|-------------------------|---|----------------------------|----------------------|--|
|                         |   | (Surface to 2 km altitude) | (2 to 6 km altitude) |  |
| Optical thickness at 5  | 32 nm                                   | 0.083                      | 0.1 to 0.4           |  |
| Ångström exponent       |   | 0.86                       | 0.0 to 2.0           |  |
| Real part of the refrac | tive index at all the wavelengths       | 1.45                       | 1.48                 |  |
| Imaginary part of the   | refractive index at all the wavelengths | 0.005                      | 0.003                |  |
| Mode radius (µm)        | fine                                    | 0.15                       | 0.15                 |  |
|                         | coarse                                  | 2.5                        | 2.5                  |  |
| Mode width              | fine                                    | 0.40                       | 0.40                 |  |
|                         | coarse                                  | 0.90                       | 0.90                 |  |
| Volume ratio of non-s   | spherical particles in the coarse mode  | 0.97                       | 0.82                 |  |

1. Study in characteristics of aerosol vertical profiles by the ground-based remote sensing



Figure 1. Flowchart diagram of this study.



Figure 2. Two examples showing the determined ABL height (red dots) and the base height of clouds or transported aerosols (blue dots).



Figure 3. Frequency distributions of the extinction coefficient at 532 nm by seasons. The solid horizonal and dashed lines indicate the seasonal means and standard deviations of the ABL height.



Figure 4. Dependencies of the Ångström exponent on the extinction coefficient at 532 nm and the altitude by seasons. The solid and dashed lines indicate the seasonal means and standard deviations of the ABL height.



Figure 5. Dependencies of single-scattering albedo on the extinction coefficient at 532 nm and the altitude by seasons. The solid and dashed lines indicate the seasonal means and standard deviations of the ABL height.



Figure 6. Dependencies of asymmetry factor on the extinction coefficient at 532 nm and the altitude by season. The solid and dashed lines indicate the seasonal means and standard deviations of the ABL height.



**Figure 7.** Optical and physical properties of transported aerosols in the FA: (a) vertical profile of the extinction coefficient at 532 nm with the ABL height (dashed lines), (b) normalized size distribution over the FA, (c) 2-day backward trajectory, and (d) altitude versus time cross section of the backward trajectory. The stars in (a) indicate the start altitude of the backward trajectories shown in (c). The color scale in (c) indicates fire activity from 1 to 9 May 2013, based on MODIS active-fire product data (NEO, 2016). The ochre color indicates desert regions, based on data of the Land Cover Type Climate Modeling Grid product (LP DAAC, 2013).



Figure 8. Results of EXP0 (dashed line) and the difference between EXP1 and EXP0 (solid lines): (a) net downward surface radiation, (b) sensible heat flux, (c) latent heat flux, (d) potential temperature at 12:00 LST, (e) specific humidity at 12:00 LST, and (f) ABL height.



Figure 9. Results of EXP0 (dashed line) and the difference between EXP2 and EXP0 (solid lines): (a) net radiation at the surface, (b) sensible heat flux, (c) latent heat flux, (d) potential temperature at 12:00 LST, (e) specific humidity at 12:00 LST, and (f) ABL height.



**Figure 10.** Dependencies of the daily maximum 2 m temperature (a) and the daily maximum ABL height (b) on the aerosol optical thickness and Ångström exponent in the FA. The "ANG" indicates Ångström exponent. The color of filled circle indicates the value of Ångström exponent, from 0.0 to 0.5 (red), from 0.5 to 1.0 (orange), from 1.0 to 1.5 (green), and from 1.5 to 2.0 (blue). The solid lines are the results of the model simulations for the simplified aerosol vertical profile described in the text.

