

## **Referee's report and detailed comments for transmission for the authors**

Infrasound research has seen a significant resurgence after the opening for signature in 1996 of the Comprehensive nuclear Test Ban Treaty (CTBT). The expansion of the global infrasound International Monitoring System (IMS) and experiment arrays deployed worldwide offers an opportunity to study background noise on a global scale. During the last decade, infrasound has developed into a broad interdisciplinary fields (combining acoustic and seismic technologies, atmospheric science, meteorology and climate) reinvigorating research field in geophysics.

Sparse observations in the middle atmosphere limit the ability to faithfully reproduce the dynamics of the middle atmosphere in numerical weather prediction and climate models. Recent studies have shown the capability of the IMS infrasound network to measure atmospheric perturbations in a wide frequency range, far beyond the frequency band of interest to detect explosions. In particular, the majority of IMS arrays measure pressure fluctuations with frequencies ranging from DC to tens of Hertz, encompassing gravity wave domain and atmospheric tides. Such network provides independent ground-based measurements useful for Numerical Weather Prediction models. The pressure measurements continuously recorded by the worldwide IMS infrasound network therefore constitute a unique set of data that could improve our knowledge on large scale atmospheric perturbations which impact the atmospheric dynamics.

The paper by P. Hupe et al. "Using barometric time series of the IMS infrasound network for a global analysis of thermally induced atmospheric tides" [Paper #amt-2017-465] demonstrates the potential of the IMS infrasound network to better characterize thermally induced atmospheric tides. This paper investigates the seasonal and latitudinal variability of atmospheric tides using historical recordings of the IMS arrays composed of high-quality microbarometers. The temporal and spectral methods presented here allow detailed geographic and temporal variability analyses of the solar and lunar tides' representation.

Compared with MERRA-2 products, these new observations provide additional quantitative measures of solar tide harmonics as well as low-amplitude gravitationally forced lunar tides. It is shown that continuous IMS records represent well the surface pressure fields of solar tides and allow quantifying their seasonal and latitudinal fluctuations. This multi-year and global dataset open doors for further investigations into the source of atmospheric tides and their interaction with planetary waves or gravity waves. Given the accuracy and high temporal resolution of the barometric data from the IMS network, such observations are of high value for continuously calibrating sensors in a wide frequency band using the ambient infrasound noise, refining the knowledge of atmospheric dynamics and data assimilation problems.

Following Marty et al. (2010), this is the first systematic and global study of continuous absolute pressure recordings. The representation of thermal tides in Numerical Weather Prediction models, often limited to the diurnal and the semidiurnal tides, would benefit from this high-resolution and uniform global observations.

Considering the new materials presented in this paper, the analyses carried out, and its revisions compared with an precedent submitted version, I recommend that this paper should be published in Atmospheric Measurement Techniques, as is.

PS: I would suggest to add a time scale on Figure 2 (from January?). It seems that the annual variation is not cyclical. Why?