

**1. Your statements that trace gas retrievals will improve when including snow-covered scenes are too strong. I agree that the high surface reflectivity is potentially benefitting the vertical sensitivity to NO<sub>2</sub> in clear-sky situations. But the reality for all retrievals is that they need highly accurate cloud retrieval schemes that can distinguish between a bright surface and bright, low altitude clouds. I don't think we have those in place yet. I think this line of thought should be reflected in the text.**

Thank you for this suggestion. We have added a paragraph in the conclusion to discuss cloud retrievals as follows (Line 325):

“The potential improvements in NO<sub>2</sub> retrieval performance over snow-covered scenes outlined here were tested for clear-sky conditions. The accuracy of cloud retrieval schemes also impacts the quality of trace gas retrievals. Many cloud retrieval schemes have difficulty distinguishing between a bright surface and bright, low altitude clouds; This may diminish the impact that improved surface snow reflectance can have on observation frequency and sensitivity when clouds are present. However, using accurate surface snow cover information may also lead to corresponding improvements in cloud retrieval accuracy.”

**P7, L14: ...high reflectance is potentially advantageous ...**

This has been changed.

**P8, L36-39: clarify that the number of observations and sensitivity improves strictly in clear-sky situations, in reality snow-covered scenes will often be covered by clouds screening the NO<sub>2</sub> below.**

These lines now read:

“We find that using IMS to identify snow cover and enable inclusion of snow-covered scenes in clear-sky conditions across North America in January...”

**P17, L295-297: “The increase in AMF ... to tropospheric NO<sub>2</sub>.” is a too strong statement. You should clarify that this is only valid if one assumes that a cloud algorithm will handle the snow-covered scenes with 100% reliability. The quality of information will furthermore only be achieved if the NO<sub>2</sub> profile and albedo value are of high quality.**

We now moderate this statement as follows:

“As we assume clear-sky conditions, these are likely upper bounds on potential increases in observation quantity and quality. In practice, the presence of clouds and errors in cloud retrieval algorithms will likely diminish these impacts.”

**2. You test in the paper the presence of snow – but what about the snow albedo value? In Figure 4, these values are listed as 0.81 and 0.83, but based on what? Adding a bit of**

**discussion on the snow albedo values from the various sets and their uncertainties would be useful.**

As stated in Section 2.3, snow reflectivity values used here come from the product described in O’Byrne et al. (2010). We have added information on uncertainty as follows (Line 171):

“This data set is consistent with previous snow reflectivity (e.g. Moody et al., 2007; Tanskanen and Manninen, 2007) over most land types (O’Byrne et al., 2010). Snow-covered reflectivity has an estimated uncertainty of 10-20% in most regions, with higher uncertainties in regions with thin or transient snow.”

**P14, L211: please rephrase considering that snow albedo values may be quite uncertain and that residual clouds may be present affecting this sensitivity – otherwise the readers may wrongfully conclude “snow means better sensitivity”, whereas the message should be that “under clear-sky conditions and with very good knowledge of snow albedo, the presence of snow means better sensitivity”.**

This has been rephrased as follows:

“This shows that satellite observed backscattered radiation in clear-sky conditions is up to five times as sensitive to NO<sub>2</sub> in the boundary layer after accounting for increased reflection by snow, due to the increased absorption by NO<sub>2</sub> in the lower troposphere when the surface reflects more sunlight”

We also added a sentence in the conclusion that mentions the importance of good knowledge of snow albedo (Line 336):

“Accurate knowledge of snow reflectivity is also needed to improve retrievals over snow.”

**Specific comments:**

**P9, L65: please add that ... are often omitted or flagged as unreliable to avoid potential errors**

This has been changed.

**P17, L302: please add ...retrievals over clear-sky reflective snow-free surfaces...**

This has been changed.

**P18, L326: please add ...significant impact on retrieved NO<sub>2</sub> columns and clouds (Vassilkov et al., 2017; Lorente et al., 2018).**

This has been changed.

**Caption Figure 3: please explain how you obtained the snow reflectance value.**

This has been changed. Caption now states:

“Figure 3: (Left) Distribution of Air Mass Factors (AMFs) calculated for OMI NO<sub>2</sub> retrievals over North America for observation geometry of January 2013, using snow-free (Sun et al., 2017) or snow-covered (O’Byrne et al., 2010) surface reflectance. (Right) Maps of AMF for snow-covered and snow-free conditions.”

# Assessing snow extent data sets over North America to inform and improve trace gas retrievals from solar backscatter

Matthew J. Cooper<sup>1</sup>, Randall V. Martin<sup>1,2</sup>, Alexei I. Lyapustin<sup>3</sup>, and Chris A. McLinden<sup>4</sup>

1. Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, Canada.

2. Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA

3. NASA Goddard Space Flight Center, Greenbelt, MD, USA

4. Air Quality Research Division, Environment and Climate Change Canada, Toronto, Ontario, Canada

## Abstract

Accurate representation of surface reflectivity is essential to tropospheric trace gas retrievals from solar backscatter observations. Surface snow cover presents a significant challenge due to its variability and thus snow-covered scenes are often omitted from retrieval data sets; however, the high reflectance of snow is potentially advantageous for trace gas retrievals. We first examine the implications of surface snow on retrievals from the upcoming TEMPO geostationary instrument for North America. We use a radiative transfer model to examine how an increase in surface reflectivity due to snow cover changes the sensitivity of satellite retrievals to NO<sub>2</sub> in the lower troposphere. We find that a substantial fraction (>50%) of the TEMPO field of regard can be snow covered in January, and that the average sensitivity to the tropospheric NO<sub>2</sub> column substantially increases (doubles) when the surface is snow covered.

We then evaluate seven existing satellite-derived or reanalysis snow extent products against ground station observations over North America to assess their capability of informing surface conditions for TEMPO retrievals. The Interactive Multisensor Snow and Ice Mapping System (IMS) had the best agreement with ground observations (accuracy=93%, precision=87%, recall=83%). Multiangle Implementation of Atmospheric Correction (MAIAC) retrievals of MODIS observed radiances had high precision (90% for Aqua and Terra), but underestimated the presence of snow (recall=74% for Aqua, 75% for Terra). MAIAC generally outperforms the standard MODIS products (precision=51%, recall=43% for Aqua; precision=69%, recall=45%

for Terra). The Near-real-time Ice and Snow Extent (NISE) product had good precision (83%) but missed a significant number of snow covered pixels (recall=45%). The Canadian Meteorological Centre (CMC) Daily Snow Depth Analysis Data set had strong performance metrics (accuracy=91%, precision=79%, recall=82%). We use the  $F$  score, which balances precision and recall, to determine overall product performance ( $F = 85\%$ ,  $82(82)\%$ ,  $81\%$ ,  $58\%$ ,  $46(54)\%$  for IMS, MAIAC Aqua(Terra), CMC, NISE, MODIS Aqua(Terra) respectively) for providing snow cover information for TEMPO retrievals from solar backscatter observations. We find that using IMS to identify snow cover and enable inclusion of snow-covered scenes in clear-sky conditions across North America in January can increase both the number of observations by a factor of 2.1 and the average sensitivity to the tropospheric  $\text{NO}_2$  column by a factor of 2.7.

## 1. Introduction

Satellite observations of solar backscatter are widely used as a source of information on atmospheric trace gases (Richter and Wagner, 2011). These observations have provided valuable information on vertical column densities of  $\text{O}_3$ ,  $\text{NO}_2$ ,  $\text{SO}_2$ ,  $\text{CO}$ ,  $\text{HCHO}$ ,  $\text{CH}_4$  and other important trace gases in the troposphere (Fishman et al., 2008). Satellite observations of trace gases have been used to assess air quality (Duncan et al., 2014; Martin, 2008) and to gain insight into atmospheric processes including emissions (Streets et al., 2013), lifetimes (Beirle et al., 2011; Fioletov et al., 2015; de Foy et al., 2015; Valin et al., 2013), and deposition (Geddes and Martin, 2017; Nowlan et al., 2014). The utility of these observations is dependent on their quality, and thus ensuring retrieval accuracy is essential.

Previous studies have found that retrieved  $\text{NO}_2$  vertical column densities are highly sensitive to errors in assumed surface reflectance (Boersma et al., 2004; Lamsal et al., 2017; Martin et al., 2002). Much of this error sensitivity results from observation sensitivity to trace gases in the lower troposphere. The observation sensitivity is accounted for in the air mass factor (AMF) conversion of observed line-of-sight “slant columns” to vertical column densities. Uncertainties in surface reflectance are a significant contributor to AMF uncertainty.

Existing reflectivity climatologies (e.g. Kleipool et al., 2008; Koelemeijer et al., 2003; Liang et al., 2002; Herman and Celarier, 1997) do not represent snow cover well, since the statistical methods to exclude reflective clouds from the climatologies also exclude variable snow cover; Correspondingly, surface snow may be mistaken for cloud, leading to errors in cloud fraction and pressure estimates used in trace gas retrievals (Lin et al., 2015; O’Byrne et al., 2010; Vasilkov et al., 2017). Therefore, snow cover is particularly challenging to retrievals. Misrepresenting surface snow cover can lead to large errors (20-50%) in retrieved NO<sub>2</sub> columns over broad regions with seasonal snow cover (O’Byrne et al., 2010). For this reason, observations over snow are often omitted or flagged as unreliable to avoid potential errors. This limits the ability of satellite retrieved data sets to offer adequate temporal and spatial sampling in winter months. Additionally, over highly reflective surfaces such as snow observation sensitivity to the lower troposphere is larger and has less dependence on *a priori* NO<sub>2</sub> profiles (Lorente et al., 2017; O’Byrne et al., 2010); Thus, omitting snow-covered scenes means omitting the observations with the greatest sensitivity to the lower troposphere. This could be remedied by using a product that would allow for snow cover identification to be done with confidence.

Several data products provide information on snow extent using surface station observations, satellite observed radiances, or visible imagery. Previous evaluations have found it difficult to determine which of these products is definitively the best, partly due to differences in resolution. Most products are more consistent during the winter months when persistent, deep snow is present (Frei et al., 2012; Frei and Lee, 2010). However, disagreements are common during accumulation and melting seasons, over mountains, and under forest canopies. These evaluations have largely focused on local or regional snow cover, or included only cloud-free observations.

The upcoming geostationary Tropospheric Emissions: Monitoring of Pollution (TEMPO) satellite instrument will provide hourly observations of air quality relevant trace gases over North America at an unprecedented spatial and temporal resolution (Zoogman et al., 2017). As is the case for all nadir satellite retrievals, the quality of these observations will depend on the accuracy of the surface reflectance used in the retrieval. As a significant portion of the observed domain experiences snow cover, an accurate representation of snow cover is needed. Current plans to deal with snow cover for TEMPO are to rely on external observations.

In this work, we examine the importance of accurate snow identification by using a radiative transport model to evaluate how the vertical sensitivity of a satellite retrieval is impacted by surface reflectance. We then assess seven snow extent products that are expected to continue to be operational during the TEMPO mission using in situ observations across North America with the intent of determining which product is best suited for providing snow cover information for TEMPO and other future satellite retrievals. Finally, we combine radiative transfer model results with a snow extent product to show how including snow-covered scenes improves both the quantity and quality of information in a retrieval data set.

## **2. Data and Algorithms**

### **2.1. Gridded Snow Products**

#### **2.1.1. IMS**

One of the most widely used sources of snow extent data is the Interactive Multisensor Snow and Ice Mapping System (IMS). IMS provides daily, near-real-time maps of snow and sea ice cover in the Northern Hemisphere at 4km resolution (Helfrich et al., 2007). The maps are produced by a trained analyst using visible imagery from a collection of geostationary (e.g. GOES, MeteoSat) and polar orbiting (e.g. AVHRR, MODIS, SAR) satellite instruments, with additional information from microwave sensors (e.g. DMSP, AMSR, AMSU), surface observations (e.g. SNOTEL), and models (e.g. SNODAS) (Helfrich et al., 2007). By using multiple sources of information with different spatial resolution and temporal sampling, IMS can minimize interference from clouds.

#### **2.1.2. MODIS**

A second commonly used snow and ice product is derived from MODIS satellite observations from the Terra and Aqua satellites (Hall and Riggs, 2007). Terra and Aqua have sun-synchronous, near polar orbits with overpass times of 1030 and 1330 hr respectively. Snow cover is calculated using a Normalized Difference Snow Index (NDSI), which examines the difference between observed radiation at visible wavelengths (where snow is highly reflective) and short IR wavelengths (where there is little reflection from snow). Observations are made at

500 m spatial resolution and aggregated to produce daily snow cover fractions on a 0.05° resolution grid. Past evaluations of the standard MODIS snow product show good agreement in cloud-free conditions but often snow is misidentified as cloud (Hall and Riggs, 2007; Yang et al., 2015).

The Multiangle Implementation of Atmospheric Correction (MAIAC) algorithm is another algorithm processing MODIS observations. MAIAC retrievals uses radiances observed by the MODIS Aqua and Terra satellites to provide atmospheric and surface products including snow detection on a 1 km grid (Lyapustin et al., 2011a, 2011b, 2012). While the NDSI used by the standard MODIS product is also used by MAIAC as one of the criteria, the overall snow and cloud detection in MAIAC are different from the standard MODIS algorithm (Lyapustin et al., 2008).

### **2.1.3. NISE**

The Near-real-time Ice and Snow Extent (NISE) provides daily updated snow cover extent information on a 25x25 km grid (Nolin et al., 2005). NISE uses microwave measurements from the Special Sensor Microwave Imager/Sounder (SSM/I) on a sun-synchronous, quasi-polar orbit to observe how microwave radiation emitted by soil is scattered by snow. Products based on microwave measurements such as NISE are known to miss wet and thin snow, as wet snow emits microwave radiation similar to soil, and thin snow does not provide sufficient scattering.

### **2.1.4. CMC**

The Canadian Meteorological Centre (CMC) Daily Snow Depth Analysis Data is a statistical interpolation of snow depth measurements from 8,000 surface sites across Canada and U.S. interpolated using a snow pack model (Brasnett, 1999). Unlike the aforementioned satellite products that provide snow extent, CMC provides snow depths. Daily snow maps are produced at 25 km resolution. As it a reanalysis product, there is a time delay in availability. The CMC snow depths show good agreement with independent observations over midlatitudes and is considered an improvement over previous snow depth climatologies (Brown et al., 2003).

## **2.2 Surface observations**

These snow identification products are evaluated against surface station observations from the Global Historical Climatology Network-Daily (GHCN-D) database, an amalgamation of daily climate records from over 80,000 surface stations worldwide (Menne et al., 2012a). Most observations over Canada and the United States are collected by government organizations (Environment and Climate Change Canada and NOAA National Climatic Data Center, respectively) with additional measurements from smaller observation networks. While the focus of the database is collecting temperature and precipitation measurements, many stations (1,279 in Canada, 13,932 in United States in 2015 used here) also offer snow depth measurements.

A subset of the surface stations included in GHCN-D may also be used in the CMC reanalysis. It is difficult to definitively know which stations are used, as CMC does not routinely archive this information. However, we estimate that only 5% of the GHCN-D stations used here are located within  $0.1^\circ$  of a possible CMC station, and thus GHCN-D has sufficient independent information sources to evaluate the CMC product.

### 2.3 Radiative transfer calculations

The sensitivity of satellite observations of  $\text{NO}_2$  to its vertical distribution is calculated here using the LIDORT radiative transfer model (Spurr, 2002). The model is used to calculate scattering weights, which quantify the sensitivity of backscattered solar radiation to  $\text{NO}_2$  at different altitudes (Martin et al., 2002; Palmer et al., 2001). The observation sensitivity to lower tropospheric  $\text{NO}_2$  is represented by the air mass factor. Air mass factors for OMI satellite observations in January 2013 are calculated as a useful analog for future TEMPO observations as both instruments are spectrometers observing reflected sunlight at UV to visible wavelengths. AMFs are calculated at 440 nm, at the centre of the  $\text{NO}_2$  retrieval window for OMI and TEMPO where  $\text{NO}_2$  has strong absorption features. Vertical  $\text{NO}_2$  profiles, and other trace gas and aerosol profiles needed for the AMF calculation shown here, are obtained from a simulation of the GEOS-Chem chemical transport model version 11-01 (www.geos-chem.org).

Figure 1 shows maps of snow-free and snow-covered reflectances used here. Snow-free surface reflectance at 470 nm is provided by Nadir BRDF-Adjusted reflectances from the MODIS CMG Gap-Filled Snow-Free Products (Sun et al., 2017). Reflectivities at 354 nm for snow-covered scenes are derived from OMI observations as described by O'Byrne et al. (2010).

This data set is consistent with previous snow reflectivity (e.g. Moody et al., 2007; Tanskanen

and Manninen, 2007) over most land types (O’Byrne et al., 2010). Snow-covered reflectivity has an estimated uncertainty of 10-20% in most regions, with higher uncertainties in regions with thin or transient snow. ~~While this~~ Although the 354 nm wavelength is different than the 440 nm wavelength used to calculate AMFs, snow reflectivity has weak spectral dependence in UV-Visible wavelengths (Feister and Grewe, 1995; O’Byrne et al., 2010). Snow can increase surface reflectance by over a factor of 10 in central North America where short vegetation is readily covered by snow.

### 3. Methods

Here we test daily snow cover products for 2015. Snow products are regridded from their native resolutions to a common 4 km grid (similar to the spatial resolution of TEMPO). A grid box is considered to be snow covered if any observations within that box are snow covered. MAIAC, NISE, and IMS give only a yes/no flag for presence of snow. MODIS products provide a pixel snow fraction, and we consider any pixels with nonzero snow fractions as snow covered. Any CMC grid box with nonzero snow depth is considered snow covered.

GHCN-D surface measurements are used as the ground “truth” for evaluating the satellite and reanalysis snow data products tested here. If measurements from multiple surface data networks exist in the same grid box, the most reliable source is used per the priority order given by GHCN-D (Menne et al., 2012b). If observations from multiple surface stations within the most reliable network within a grid box disagree on the presence of snow on a given day, that day is excluded from the evaluation.

We assess the snow data sets using metrics that are commonly used for evaluating binary data sets (Rittger et al., 2013). These metrics are based on the possible outcomes for identifying snow: true positive (TP), true negative (TN), false positive (FP), and false negative (FN). Accuracy measures the likelihood that a grid box, with snow or without, is correctly classified:

$$Accuracy = \frac{TP + TN}{TP + TN + FP + FN} \quad (1)$$

Precision is the probability that a region identified as snow-covered has snow:

$$Precision = \frac{TP}{TP + FP} \quad (2)$$

Recall is the likelihood that snow cover is detected when present:

$$Recall = \frac{TP}{TP + FN} \quad (3)$$

The  $F$  score balances recall (which accounts for false negatives) and precision (which accounts for false positives) to measure correct classification of snow without the influence of frequent snow-free periods, and therefore is the metric which is most relevant for TEMPO:

$$F = 2 * \frac{precision * recall}{precision + recall} \quad (4)$$

## 4. Results

We first examine the effect of surface reflectivity on retrieval sensitivity by using the LIDORT radiative transfer model to calculate NO<sub>2</sub> air mass factors for both snow-free and snow-covered scenarios using the corresponding snow-free (Sun et al., 2017) or snow-covered (O’Byrne et al., 2010) surface reflectance over North America. We calculate air mass factors over North America in January 2013. We assume cloud-free conditions in all AMF calculations, as the impact of surface reflectance on retrieved cloud fractions is beyond the scope of this paper.

Figure 2 shows the sensitivity of backscattered radiation (scattering weights) over snow-covered and snow-free surfaces for two locations; a midlatitude location (US Midwest, 42°N, 99°W) with a solar zenith angle of 60° and at a high latitude location (Northern Canada, 58°N, 76°W) with a solar zenith angle of 79°. The snow-covered scattering weights are greater than the snow-free scattering weights throughout the troposphere, by factors of 2.0 (2.7) below 5 km, 2.7 (3.7) below 2 km, and 2.6 (5.3) below 1 km at the mid (high) latitude location. This shows that satellite observed backscattered radiation in clear-sky conditions is up to five times as sensitive to NO<sub>2</sub> in the boundary layer in the presence of snow after accounting for increased reflection by snow, due to the increased absorption by NO<sub>2</sub> in the lower troposphere when the surface reflects more sunlight.

Figure 3 shows the distribution of AMF values over North America with and without reflectance from snow. The snow-free AMF distribution is unimodal with a median of 1.2. Allowing for the presence of snow introduces a second mode with a median of 3.2. Mean AMFs increase by a factor 2.0 in the presence of snow, indicating an overall doubling in the sensitivity to tropospheric NO<sub>2</sub> over snow covered surfaces across North America. The impact is larger over

polluted regions, as mean AMFs increase by a factor of 2.2 in regions where NO<sub>2</sub> columns exceed  $1 \times 10^{15}$  molec/cm<sup>2</sup>. Maps of AMF with and without snow cover for January 2013 show that AMF values increase over 69% of the land surface within the TEMPO domain.

We next examine the snow datasets to identify the one most suited for the TEMPO retrieval algorithm. Figure 4 shows the spatial distribution of false positives and false negatives in the data sets. In all data sets, both false positives and negatives are most frequent over mountainous regions, particularly in the Rocky Mountain region, consistent with previous validation studies (Chen et al., 2012, 2014; Frei et al., 2012; Frei and Lee, 2010). These errors are often attributed to differences in representativeness, as snow cover in mountain regions is often spatially inhomogeneous, and thus *in situ* measurements may not be representative of the pixel. A slight increase in the number of false positives in IMS over mid-western and prairie regions may result from crop regions with high snow-free albedos being mistaken for snow in visible imagery (Chen et al., 2012; Yang et al., 2015). NISE, MODIS Aqua, and MODIS Terra have more false negatives overall, especially in the Great Lakes and New England regions. False positives are less frequent than false negatives in all data sets. IMS and CMC have the lowest frequency of false negatives. NISE and MAIAC have the lowest frequency of false positives.

Figure 5 shows the metrics used to evaluate data set performance. Table 1 summarizes these results. All data sets have high accuracy numbers, owing largely to a high number of true negatives during the summer months. MODIS Aqua and Terra have low recall and *F* scores. When only observations with MODIS cloud fractions less than 20% are used, MODIS has better agreement with the ground stations (*F* statistic increases from 0.38 to 0.49 at native resolution for Aqua, 0.43 to 0.63 for Terra), however this reduces the number of usable MODIS observations by up to 60%. NISE has high precision but low recall, indicating that while areas classified as snow-covered by NISE are likely correct, many snow-covered regions are missing in the data set. This is consistent with evaluations by McLinden et al. (2014) and O’Byrne et al. (2010). Although CMC, IMS, and MAIAC products show an increase in frequency of false negatives over the Rocky Mountains, they retain a high precision in this region due to frequent snow cover. While MAIAC Aqua/Terra have high accuracy and precision, lower recall values indicate that they are conservative in identifying the presence of snow. This is possibly a consequence of the method used for identifying cloud, which may incorrectly classify fresh

snowfall as cloud (Lyapustin et al., 2008). Data sets were also evaluated by season with similar results (Appendix Table A1). All data sets have weaker performance metrics during the spring melt season, which has been observed in past evaluations (Frei et al., 2012). IMS has the highest  $F$  score in winter and autumn but is slightly outperformed by MAIAC in spring. Data sets were also evaluated at their native resolutions and at a common 25 km resolution (Appendix Tables A2-3). Results are similar at each resolution with two exceptions: MODIS Aqua and Terra products perform better when regridded from their native  $0.05^\circ$  resolution to a 4 km resolution as it reduces the number of grid boxes missing observations due to cloud, and MAIAC Aqua and Terra perform better at their native resolution than at either 4 km or 25 km as degrading the spatial resolution results in a loss of information.

For all data sets, recall is generally low in two regions: along the Pacific coastline where snow depths are relatively thin, and in the south when snow is rare and generally short lived. Thin snow is likely to be less homogenous across a pixel and more likely to be obscured by forest canopies or tall grasses, and thus is difficult to observe from satellite imagery. Short lived snow in the south is likely to be missed by satellite observations, especially since clouds are often present. However, as IMS uses multiple observations at multiple times of day in addition to incorporating ground station data, it is more likely to find snow in these cases than other satellite products (Hall et al., 2010). Overall, IMS has best agreement with *in situ* observations, with the highest accuracy, recall, and  $F$  statistic and relatively high precision.

While CMC also has strong performance metrics, it is important to consider the information source used to describe snow extent in each product. Products based on satellite observations are advantageous when assessing how surface reflectivity affects backscattered radiation observed from space. For example, thin snow, or snow obscured by tree canopies, may not affect the observed brightness from space, but would be considered snow-covered by a product based on surface observations (e.g. CMC). Also, the reflectivity of a snow-covered surface decreases over time as the snow ages (Warren and Wiscombe, 1980); This effect would not be captured by snow depth measurements. And while snow depth has been used as an indicator of brightness (Arola et al., 2003), it can not account for snow aging or canopy effects. IMS is based on visible satellite imagery and thus determines snow extent based on brightness from space, which is more applicable to satellite retrievals. And while most satellite-based

products rely on observations made at a single overpass time and viewing geometry, IMS has the advantage of incorporating observations from multiple satellites with differing measurement times and geometries, including both geostationary and low Earth orbits. These reasons, in addition to a strong agreement with in situ measurements and near-real-time updates, make IMS best suited for informing TEMPO retrievals.

We next examine the effect on both spatial sampling and sensitivity to the lower troposphere of a retrieval data set if observations with surface snow are included rather than omitted. We use IMS to identify the presence of snow for OMI observations over North America in January 2015. We then use LIDORT to calculate AMFs for these observations using the corresponding snow-free (Sun et al., 2017) or snow-covered (O’Byrne et al., 2010) surface reflectance, and examine the results of either including or omitting snow-covered scenes. Figure 6 shows that including snow-covered scenes results in a significant (factor of 2.1) increase in observation frequency, particularly in the northern US and Canada. Additionally, including snow-covered scenes increases the average AMF by a factor of 2.7 in regions with occasional snow cover. The increase in AMF demonstrates that including snow-covered scenes increases the quality of information about the tropospheric NO<sub>2</sub> column by increasing the observation sensitivity to tropospheric NO<sub>2</sub>. As we assume clear-sky conditions, these are likely upper bounds on potential increases in observation quantity and quality. In practice, the presence of clouds and errors in cloud retrieval algorithms will likely diminish these impacts.

## 5. Conclusion

An accurate representation of snow cover is essential to ensuring satellite retrieval accuracy, including those from TEMPO. Radiative transfer model calculations indicate that clear-sky NO<sub>2</sub> retrievals over reflective snow-covered surfaces are more than twice as sensitive to NO<sub>2</sub> in the boundary layer than over snow-free surfaces. This makes snow an attractive surface over which to observe tropospheric NO<sub>2</sub>. However, the lack of confidence in snow identification has previously led many retrieval procedures to omit observations over snow. We show that increasing this confidence such that these observations could be included not only improves spatial and temporal sampling, but also allows the inclusion of observations with higher quality information on the lower troposphere.

We evaluated seven snow extent data sets to determine their usefulness for informing satellite retrievals of trace gas from solar backscatter observations. All products were more likely to misidentify snow over mountains or where snow cover is thin or short lived. IMS had the best agreement with *in situ* observations ( $F=0.85$ ), and as a satellite based, operational, daily updated product, it is well suited for informing TEMPO satellite retrievals. The low recall value (0.45) for NISE indicated that a significant number of snow covered pixels are missed. The standard MODIS products showed medium precision and low recall owing to cloud contamination. The MAIAC products had the highest precision (0.90 for both Aqua and Terra) of those tested, but is conservative in ascribing the presence of snow (recall=0.74 for Aqua, 0.75 for Terra). CMC had strong performance metrics ( $F=0.81$ ), but as a reanalysis product based on ground observations it may not appropriately represent how a surface snow reflectivity would affect TEMPO observed radiances.

The potential improvements in NO<sub>2</sub> retrieval performance over snow-covered scenes outlined here were tested for clear-sky conditions. The accuracy of cloud retrieval schemes also impacts the quality of trace gas retrievals. Many cloud retrieval schemes have difficulty distinguishing between a bright surface and bright, low altitude clouds; This may diminish the impact that improved surface snow reflectance can have on observation frequency and sensitivity when clouds are present. However, using accurate surface snow cover information may also lead to corresponding improvements in cloud retrieval accuracy.

Future work should investigate snow reflectance products that could be used when snow is detected. This could potentially include Bidirectional Reflectance Distribution Functions (BRDF) that describe reflection at different viewing angles, as this effect has been shown to have significant impact on retrieved NO<sub>2</sub> columns and clouds (Lorente et al., 2018; Vasilkov et al., 2017). Accurate knowledge of snow reflectivity is also needed to improve retrievals over snow. A retrieval algorithm that combines daily snow detection from IMS with a climatology of snow reflectance has the potential to greatly improve upon current methodologies.

## 6. Data Availability

IMS (National Ice Center, 2008), NISE (Brodzik and Stewart, 2016), MODIS Aqua (Hall and Riggs, 2016a), MODIS Terra (Hall and Riggs, 2016b), and CMC (Brown and Brasnett, 2010) data are available from the NASA National Snow and Ice Data Center (<http://nsidc.org>). MAIAC Collection 6 re-processing of MODIS data started in September 2017 and is expected to be completed by the end of year. This study used MAIAC data currently available via ftp at NASA Center for Climate Simulations (NCCS): <ftp://maiac/dataportal.nccs.nasa.gov/DataRelease/>. GHCN-D data are available from the NOAA National Climatic Data Center (Menne et al., 2012b; [www.ncdn.noaa.gov](http://www.ncdn.noaa.gov)). Code for calculating scattering weights and air mass factors, and snow-covered surface reflectances used here, are available at <http://fizz.phys.dal.ca/~atmos>. Snow-free surface reflectances are available at <ftp://rsftp.eeos.umb.edu/data02/Gapfilled/>. The GEOS-Chem chemical transport model used here is available at [www.geos-chem.org](http://www.geos-chem.org).

## 7. References

- Arola, A., Kaurola, J., Koskinen, L., Tanskanen, A., Tikkanen, T., Taalas, P., Herman, J. R., Krotkov, N. and Fioletov, V.: A new approach to estimating the albedo for snow-covered surfaces in the satellite UV method, *J. Geophys. Res.*, 108(D17), 4531, doi:10.1029/2003JD003492, 2003.
- Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G. and Wagner, T.: Megacity emissions and lifetimes of nitrogen oxides probed from space., *Science*, 333(6050), 1737–9, doi:10.1126/science.1207824, 2011.
- Boersma, K. F., Eskes, H. J. and Brinksma, E. J.: Error analysis for tropospheric NO<sub>2</sub> retrieval from space, *J. Geophys. Res. Atmos.*, 109(D4), 2004.
- Brasnett, B.: A Global Analysis of Snow Depth for Numerical Weather Prediction, *J. Appl. Meteorol.*, 38(6), 726–740, doi:10.1175/1520-0450(1999)038<0726:AGAOSD>2.0.CO;2, 1999.
- Brodzik, M. J. and Stewart, J. S.: Near-Real-Time SSM/I-SSMIS EASE-Grid Daily Global Ice Concentration and Snow Extent, Version 5, , doi:<http://dx.doi.org/10.5067/3KB2JPLFPK3R>, 2016.
- Brown, R. D. and Brasnett, B.: Canadian Meteorological Centre (CMC) Daily Snow Depth

370 Analysis Data, Version 1, , doi:<http://dx.doi.org/10.5067/W9FOYWH0EQZ3>, 2010.

371 Brown, R. D., Brasnett, B. and Robinson, D.: Gridded North American monthly snow depth and  
 372 snow water equivalent for GCM evaluation, *Atmosphere-Ocean*, 41(1), 1–14,  
 373 doi:[10.3137/ao.410101](https://doi.org/10.3137/ao.410101), 2003.

374 Chen, C., Lakhankar, T., Romanov, P., Helfrich, S., Powell, A. and Khanbilvardi, R.: Validation  
 375 of NOAA-Interactive Multisensor Snow and Ice Mapping System (IMS) by Comparison with  
 376 Ground-Based Measurements over Continental United States, *Remote Sens.*, 4(12), 1134–1145,  
 377 doi:[10.3390/rs4051134](https://doi.org/10.3390/rs4051134), 2012.

378 Chen, X., Jiang, L., Yang, J. and Pan, J.: Validation of ice mapping system snow cover over  
 379 southern China based on Landsat Enhanced Thematic Mapper Plus imagery, *J. Appl. Remote*  
 380 *Sens.*, 8(1), 84680, doi:[10.1117/1.JRS.8.084680](https://doi.org/10.1117/1.JRS.8.084680), 2014.

381 Duncan, B. N., Prados, A. I., Lamsal, L. N., Liu, Y., Streets, D. G., Gupta, P., Hilsenrath, E.,  
 382 Kahn, R. A., Nielsen, J. E., Beyersdorf, A. J., Burton, S. P., Fiore, A. M., Fishman, J., Henze, D.  
 383 K., Hostetler, C. A., Krotkov, N. A., Lee, P., Lin, M., Pawson, S., Pfister, G., Pickering, K. E.,  
 384 Pierce, R. B., Yoshida, Y. and Ziemba, L. D.: Satellite data of atmospheric pollution for U.S. air  
 385 quality applications: Examples of applications, summary of data end-user resources, answers to  
 386 FAQs, and common mistakes to avoid, *Atmos. Environ.*, 94, 647–662,  
 387 doi:[10.1016/j.atmosenv.2014.05.061](https://doi.org/10.1016/j.atmosenv.2014.05.061), 2014.

388 Feister, U. and Grewe, R.: Spectral albedo measurements in the UV and visible region over  
 389 different types of surfaces, *Photochem. Photobiol.*, 62(4), 736–744, doi:[10.1111/j.1751-](https://doi.org/10.1111/j.1751-1097.1995.tb08723.x)  
 390 [1097.1995.tb08723.x](https://doi.org/10.1111/j.1751-1097.1995.tb08723.x), 1995.

391 Fioletov, V. E., McLinden, C. A., Krotkov, N. and Li, C.: Lifetimes and emissions of SO<sub>2</sub> from  
 392 point sources estimated from OMI, *Geophys. Res. Lett.*, 42(6), 1969–1976,  
 393 doi:[10.1002/2015GL063148](https://doi.org/10.1002/2015GL063148), 2015.

394 Fishman, J., Al-Saadi, J. A., Creilson, J. K., Bowman, K. W., Burrows, J. P., Richter, A.,  
 395 Chance, K. V., Edwards, D. P., Martin, R. V., Morris, G. A., Pierce, R. B., Ziemke, J. R.,  
 396 Schaack, T. K., Thompson, A. M., Fishman, J., Al-Saadi, J. A., Creilson, J. K., Bowman, K. W.,  
 397 Burrows, J. P., Richter, A., Chance, K. V., Edwards, D. P., Martin, R. V., Morris, G. A., Pierce,

398 R. B., Ziemke, J. R., Schaack, T. K. and Thompson, A. M.: Remote Sensing of Tropospheric  
 399 Pollution from Space, *Bull. Am. Meteorol. Soc.*, 89(6), 805–821,  
 400 doi:10.1175/2008BAMS2526.1, 2008.

401 de Foy, B., Lu, Z., Streets, D. G., Lamsal, L. N. and Duncan, B. N.: Estimates of power plant NO  
 402 x emissions and lifetimes from OMI NO<sub>2</sub> satellite retrievals, *Atmos. Environ.*, 116, 1–11, 2015.

403 Frei, A. and Lee, S.: A comparison of optical-band based snow extent products during spring  
 404 over North America, *Remote Sens. Environ.*, 114(9), 1940–1948, doi:10.1016/j.rse.2010.03.015,  
 405 2010.

406 Frei, A., Tedesco, M., Lee, S., Foster, J., Hall, D. K., Kelly, R. and Robinson, D. A.: A review of  
 407 global satellite-derived snow products, *Adv. Sp. Res.*, 50(8), 1007–1029,  
 408 doi:10.1016/j.asr.2011.12.021, 2012.

409 Geddes, J. A. and Martin, R. V.: Global deposition of total reactive nitrogen oxides from 1996 to  
 410 2014 constrained with satellite observations of NO<sub>2</sub> columns, *Atmos. Chem. Phys.*, 17(16),  
 411 10071–10091, doi:10.5194/acp-17-10071-2017, 2017.

412 Hall, D. . and Riggs, G. A.: MODIS/Aqua Snow Cover Daily L3 Global 0.05Deg CMG, Version  
 413 6, , doi:http://dx.doi.org/10.5067/MODIS/MYD10C1.006, 2016a.

414 Hall, D. K. and Riggs, G. A.: Accuracy assessment of the MODIS snow products, *Hydrol.*  
 415 *Process.*, 21(12), 1534–1547, doi:10.1002/hyp.6715, 2007.

416 Hall, D. K. and Riggs, G. A.: MODIS/Terra Snow Cover Daily L3 Global 0.05Deg CMG,  
 417 Version 6, , doi:http://dx.doi.org/10.5067/MODIS/MOD10C1.006, 2016b.

418 Hall, D. K., Fuhrmann, C. M., Perry, L. B., Riggs, G. A., Robinson, D. A. and Foster, J. L.: A  
 419 Comparison of Satellite-Derived Snow Maps with a Focus on Ephemeral Snow in North  
 420 Carolina, in 67th Eastern Snow Conference, Hancock MA USA., 2010.

421 Helfrich, S. R., McNamara, D., Ramsay, B. H., Baldwin, T. and Kasheta, T.: Enhancements to,  
 422 and forthcoming developments in the Interactive Multisensor Snow and Ice Mapping System  
 423 (IMS), *Hydrol. Process.*, 21(12), 1576–1586, doi:10.1002/hyp.6720, 2007.

424 Herman, J. R. and Celarier, E. A.: Earth surface reflectivity climatology at 340–380 nm from

425 TOMS data, *J. Geophys. Res. Atmos.*, 102(D23), 28003–28011, doi:10.1029/97JD02074, 1997.

426 Kleipool, Q. L., Dobber, M. R., de Haan, J. F. and Levelt, P. F.: Earth surface reflectance  
 427 climatology from 3 years of OMI data, *J. Geophys. Res.*, 113(D18), D18308,  
 428 doi:10.1029/2008JD010290, 2008.

429 Koelemeijer, R. B. A., de Haan, J. F. and Stammes, P.: A database of spectral surface reflectivity  
 430 in the range 335–772 nm derived from 5.5 years of GOME observations, *J. Geophys. Res.*,  
 431 108(D2), 4070, doi:10.1029/2002JD002429, 2003.

432 Lamsal, L. N., Janz, S. J., Krotkov, N. A., Pickering, K. E., Spurr, R. J. D., Kowalewski, M. G.,  
 433 Loughner, C. P., Crawford, J. H., Swartz, W. H. and Herman, J. R.: High-resolution NO<sub>2</sub>  
 434 observations from the Airborne Compact Atmospheric Mapper: Retrieval and validation, *J.*  
 435 *Geophys. Res. Atmos.*, 122(3), 1953–1970, doi:10.1002/2016JD025483, 2017.

436 Liang, S., Fang, H., Chen, M., Shuey, C. J., Walthall, C., Daughtry, C., Morissette, J., Schaaf, C.  
 437 and Strahler, A.: Validating MODIS land surface reflectance and albedo products: methods and  
 438 preliminary results, *Remote Sens. Environ.*, 83(1–2), 149–162, doi:10.1016/S0034-  
 439 4257(02)00092-5, 2002.

440 Lin, J.-T., Liu, M.-Y., Xin, J.-Y., Boersma, K. F., Spurr, R., Martin, R. and Zhang, Q.: Influence  
 441 of aerosols and surface reflectance on satellite NO<sub>2</sub> retrieval: seasonal and spatial characteristics  
 442 and implications for NO<sub>x</sub> emission constraints, *Atmos. Chem. Phys.*, 15(19), 11217–11241,  
 443 doi:10.5194/acp-15-11217-2015, 2015.

444 Lorente, A., Folkert Boersma, K., Yu, H., Dörner, S., Hilboll, A., Richter, A., Liu, M., Lamsal,  
 445 L. N., Barkley, M., De Smedt, I., Van Roozendaal, M., Wang, Y., Wagner, T., Beirle, S., Lin, J.-  
 446 T., Krotkov, N., Stammes, P., Wang, P., Eskes, H. J. and Krol, M.: Structural uncertainty in air  
 447 mass factor calculation for NO<sub>2</sub> and HCHO satellite retrievals, *Atmos. Meas. Tech.*, 10(3), 759–  
 448 782, doi:10.5194/amt-10-759-2017, 2017.

449 Lorente, A., Boersma, K. F., Stammes, P., Tilstra, L. G., Richter, A., Yu, H., Kharbouche, S. and  
 450 Muller, J.-P.: The importance of surface reflectance anisotropy for cloud and NO<sub>2</sub> retrievals  
 451 from GOME-2 and OMI, *Atmos. Meas. Tech. Discuss.*, 1–29, doi:10.5194/amt-2018-32, 2018.

452 Lyapustin, A., Wang, Y. and Frey, R.: An automatic cloud mask algorithm based on time series

453 of MODIS measurements, *J. Geophys. Res.*, 113(D16), D16207, doi:10.1029/2007JD009641,  
 454 2008.

455 Lyapustin, A., Martonchik, J., Wang, Y., Laszlo, I. and Korkin, S.: Multiangle implementation of  
 456 atmospheric correction (MAIAC): 1. Radiative transfer basis and look-up tables, *J. Geophys.*  
 457 *Res.*, 116(D3), D03210, doi:10.1029/2010JD014985, 2011a.

458 Lyapustin, A., Wang, Y., Laszlo, I., Kahn, R., Korkin, S., Remer, L., Levy, R. and Reid, J. S.:  
 459 Multiangle implementation of atmospheric correction (MAIAC): 2. Aerosol algorithm, *J.*  
 460 *Geophys. Res.*, 116(D3), D03211, doi:10.1029/2010JD014986, 2011b.

461 Lyapustin, A. I., Wang, Y., Laszlo, I., Hilker, T., G.Hall, F., Sellers, P. J., Tucker, C. J. and  
 462 Korkin, S. V.: Multi-angle implementation of atmospheric correction for MODIS (MAIAC): 3.  
 463 Atmospheric correction, *Remote Sens. Environ.*, 127, 385–393, doi:10.1016/j.rse.2012.09.002,  
 464 2012.

465 Martin, R. V, Chance, K., Jacob, D. J., Kurosu, T. P., Spurr, R. J. D., Bucsela, E., Gleason, J. F.,  
 466 Palmer, P. I., Bey, I. and Fiore, A. M.: An improved retrieval of tropospheric nitrogen dioxide  
 467 from GOME, *J. Geophys. Res. Atmos.*, 107(D20), 2002.

468 Martin, R. V.: Satellite remote sensing of surface air quality, *Atmos. Environ.*, 42(34), 7823–  
 469 7843, doi:10.1016/j.atmosenv.2008.07.018, 2008.

470 McLinden, C. A., Fioletov, V., Boersma, K. F., Kharol, S. K., Krotkov, N., Lamsal, L., Makar,  
 471 P. A., Martin, R. V., Veefkind, J. P. and Yang, K.: Improved satellite retrievals of NO<sub>2</sub> and SO<sub>2</sub>  
 472 over the Canadian oil sands and comparisons with surface measurements, *Atmos. Chem. Phys.*,  
 473 14(7), 3637–3656, doi:10.5194/acp-14-3637-2014, 2014.

474 Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E., Houston, T. G., Menne, M. J., Durre, I.,  
 475 Vose, R. S., Gleason, B. E. and Houston, T. G.: An Overview of the Global Historical  
 476 Climatology Network-Daily Database, *J. Atmos. Ocean. Technol.*, 29(7), 897–910,  
 477 doi:10.1175/JTECH-D-11-00103.1, 2012a.

478 Menne, M. J., Durre, I., Korzeniewski, B., McNeal, S., Thomas, K., Yin, X., Anthony, S., Ray,  
 479 R., Vose, R. S., Gleason, B. E. and Houston, T. G.: Global Historical Climatology Network -  
 480 Daily (GHCN-Daily), Version 3.22, , doi:http://doi.org/10.7289/V5D21VHZ, 2012b.

481 Moody, E. G., King, M. D., Schaaf, C. B., Hall, D. K. and Platnick, S.: Northern Hemisphere  
 482 five-year average (2000–2004) spectral albedos of surfaces in the presence of snow: Statistics  
 483 computed from Terra MODIS land products, *Remote Sens. Environ.*, 111(2), 337–345,  
 484 doi:10.1016/j.rse.2007.03.026, 2007.

485 National Ice Center: IMS Daily Northern Hemisphere Snow and Ice Analysis at 1 km, 4 km, and  
 486 24 km Resolutions, Version 1, , doi:http://dx.doi.org/10.7265/N52R3PMC, 2008.

487 Nolin, A., Armstrong, R. and Maslanik, J.: Near real-time SSM/I EASE-grid daily global ice  
 488 concentration and snow extent, *Digit. Media, Natl. Snow Ice Data Center, Boulder, CO, USA*,  
 489 2005.

490 Nowlan, C. R., Martin, R. V., Philip, S., Lamsal, L. N., Krotkov, N. A., Marais, E. A., Wang, S.  
 491 and Zhang, Q.: Global dry deposition of nitrogen dioxide and sulfur dioxide inferred from space-  
 492 based measurements, *Global Biogeochem. Cycles*, 28(10), 1025–1043,  
 493 doi:10.1002/2014GB004805, 2014.

494 O’Byrne, G., Martin, R. V., van Donkelaar, A., Joiner, J. and Celarier, E. A.: Surface reflectivity  
 495 from the Ozone Monitoring Instrument using the Moderate Resolution Imaging  
 496 Spectroradiometer to eliminate clouds: Effects of snow on ultraviolet and visible trace gas  
 497 retrievals, *J. Geophys. Res.*, 115(D17), D17305, doi:10.1029/2009JD013079, 2010.

498 Palmer, P. I., Jacob, D. J., Chance, K. and Martin, R. V: Air mass factor formulation for  
 499 spectroscopic measurements from satellites’ Application to formaldehyde retrievals from the  
 500 Global Ozone Monitoring Experiment, *J. Geophys. Res.*, 106(D13), 14,514-539,550 [online]  
 501 Available from: <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2000JD900772>, 2001.

502 Richter, A. and Wagner, T.: The Use of UV, Visible and Near IR Solar Back Scattered Radiation  
 503 to Determine Trace Gases, pp. 67–121, Springer, Berlin, Heidelberg., 2011.

504 Rittger, K., Painter, T. H. and Dozier, J.: Assessment of methods for mapping snow cover from  
 505 MODIS, *Adv. Water Resour.*, 51, 367–380, doi:10.1016/j.advwatres.2012.03.002, 2013.

506 Spurr, R. J. D.: Simultaneous derivation of intensities and weighting functions in a general  
 507 pseudo-spherical discrete ordinate radiative transfer treatment, *J. Quant. Spectrosc. Radiat.*  
 508 *Transf.*, 75(75), 129–175 [online] Available from:

509 <https://www.sciencedirect.com/science/article/pii/S002240730100245X> (Accessed 20 July  
510 2017), 2002.

511 Streets, D. G., Canty, T., Carmichael, G. R., de Foy, B., Dickerson, R. R., Duncan, B. N.,  
512 Edwards, D. P., Haynes, J. A., Henze, D. K. and Houyoux, M. R.: Emissions estimation from  
513 satellite retrievals: A review of current capability, *Atmos. Environ.*, 77, 1011–1042, 2013.

514 Sun, Q., Wang, Z., Li, Z., Erb, A. and Schaaf, C. B.: Evaluation of the global MODIS 30 arc-  
515 second spatially and temporally complete snow-free land surface albedo and reflectance  
516 anisotropy dataset, *Int. J. Appl. Earth Obs. Geoinf.*, 58, 36–49, doi:10.1016/j.jag.2017.01.011,  
517 2017.

518 Tanskanen, A. and Manninen, T.: Effective UV surface albedo of seasonally snow-covered  
519 lands, *Atmos. Chem. Phys.*, 7(10), 2759–2764, doi:10.5194/acp-7-2759-2007, 2007.

520 Valin, L. C., Russell, A. R. and Cohen, R. C.: Variations of OH radical in an urban plume  
521 inferred from NO<sub>2</sub> column measurements, *Geophys. Res. Lett.*, 40(9), 1856–1860, 2013.

522 Vasilkov, A., Qin, W., Krotkov, N., Lamsal, L., Spurr, R., Haffner, D., Joiner, J., Yang, E.-S.  
523 and Marchenko, S.: Accounting for the effects of surface BRDF on satellite cloud and trace-gas  
524 retrievals: a new approach based on geometry-dependent Lambertian equivalent  
525 reflectivity applied to OMI algorithms, *Atmos. Meas. Tech.*, 10(1), 333–349, doi:10.5194/amt-  
526 10-333-2017, 2017.

527 Warren, S. G. and Wiscombe, W. J.: A Model for the Spectral Albedo of Snow. II: Snow  
528 Containing Atmospheric Aerosols, *J. Atmos. Sci.*, 37(12), 2734–2745, doi:10.1175/1520-  
529 0469(1980)037<2734:AMFTSA>2.0.CO;2, 1980.


530 Yang, J., Jiang, L., Ménard, C. B., Luo, J., Lemmetyinen, J. and Pulliainen, J.: Evaluation of  
531 snow products over the Tibetan Plateau, *Hydrol. Process.*, 29(15), 3247–3260,  
532 doi:10.1002/hyp.10427, 2015.

533 Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A.,  
534 Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L., Janz, S. J., Andraschko, M. R., Arola,  
535 A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E.,  
536 Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck,

537 J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors,  
538 A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R.,  
539 O'Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D.,  
540 Szykman, J. J., Torres, O., Veefkind, J. P., Veihelmann, B., Wang, H., Wang, J. and Chance, K.:  
541 Tropospheric emissions: Monitoring of pollution (TEMPO), *J. Quant. Spectrosc. Radiat. Transf.*,  
542 186, 17–39, doi:10.1016/j.jqsrt.2016.05.008, 2017.

543


544



545

546 Figure 1: Surface reflectivity at UV-visible wavelengths for snow-covered and snow-free  
547 conditions for January 2013. White space in top panel indicates no snow reflectance information  
548 is available.


549



550

551 Figure 2: Observation sensitivity to NO<sub>2</sub>. Scattering weight profiles calculated for cloud-free  
 552 OMI NO<sub>2</sub> retrievals, with and without surface snow cover, for January 2013 at (Left) 42° N, 99°  
 553 W with a solar zenith angle (ZA) of 60° and (Right) 58° N, 76° W with a solar zenith angle of  
 554 79°.

555



556

557 Figure 3: (Left) Distribution of Air Mass Factors (AMFs) calculated for OMI NO<sub>2</sub> retrievals over  
 558 North America for observation geometry of January 2013, using snow-free (Sun et al., 2017) or  
 559 snow-covered (O'Byrne et al., 2010) surface reflectance~~with and without surface snow cover~~.  
 560 (Right) Maps of AMF for snow-covered and snow-free conditions.





Figure 4: Number of false positive (FP) and false negative (FN) snow attributions by the snow data sets in 2015. All data sets are evaluated at 4 km resolution. Total number of false snow attributions inset. White space indicates no ground stations present.



565

566 Figure 5: Statistical metrics to evaluate snow cover products. All data sets are gridded at 4 km  
 567 resolution. White space indicates no ground stations present.




Figure 6: OMI observation frequency (top) and average AMFs (bottom) over North America in January using IMS to identify surface snow conditions. White space indicates a lack of observations.

573

	Accuracy	Precision	Recall	F
CMC	0.91	0.79	<b>0.83</b>	0.81
IMS	<b>0.93</b>	0.87	<b>0.83</b>	<b>0.85</b>
MAIAC AQUA	0.91	<b>0.90</b>	0.74	0.82
MAIAC TERRA	0.91	<b>0.90</b>	0.75	0.82
MODIS AQUA	0.76	0.51	0.43	0.46
MODIS TERRA	0.82	0.69	0.45	0.54
NISE	0.84	0.83	0.45	0.58

574 Table 1: Evaluation of daily snow extent data set performance for 2015. GHCN-D surface  
575 observations are used as “truth”. All products are regridded to a common 4 km resolution. The  
576 highest value for each metric is shown in bold.

## 577 Appendix

Months	Data Set	Accuracy	Precision	Recall	F
DJF	CMC	0.84	0.84	0.89	0.86
	IMS	<b>0.88</b>	0.90	<b>0.88</b>	<b>0.89</b>
	MAIAC AQUA	0.84	<b>0.93</b>	0.80	0.86
	MAIAC TERRA	0.84	0.92	0.80	0.86
	MODIS AQUA	0.58	0.84	0.34	0.48
	MODIS TERRA	0.60	0.88	0.37	0.52
	NISE	0.63	0.90	0.41	0.57
MAM	CMC	0.90	0.63	0.57	0.59
	IMS	<b>0.93</b>	0.74	<b>0.67</b>	0.70
	MAIAC AQUA	<b>0.93</b>	<b>0.81</b>	0.62	<b>0.71</b>
	MAIAC TERRA	<b>0.93</b>	<b>0.81</b>	0.63	<b>0.71</b>
	MODIS AQUA	0.86	0.43	0.39	0.41
	MODIS TERRA	0.89	0.62	0.40	0.49
	NISE	0.90	0.71	0.34	0.46
SON	CMC	0.91	0.73	<b>0.81</b>	0.76
	IMS	<b>0.92</b>	0.82	0.74	<b>0.78</b>
	MAIAC AQUA	0.91	<b>0.86</b>	0.60	0.71
	MAIAC TERRA	0.90	0.85	0.61	0.71
	MODIS AQUA	0.82	0.51	0.36	0.42
	MODIS TERRA	0.86	0.71	0.39	0.51
	NISE	0.85	0.85	0.25	0.39

578 Table A1: Evaluation of daily snow extent data set performance by season for 2015. GHCN-D  
579 surface observations are used as “truth”. All products are regridded to a common 4 km  
580 resolution. The highest value for each metric/season is shown in bold.

581

582

	Resolution	Accuracy	Precision	Recall	F
CMC	25 km	0.92	0.81	0.81	0.81
IMS	4 km	<b>0.93</b>	0.87	<b>0.83</b>	<b>0.85</b>
MAIAC AQUA	1 km	0.91	<b>0.91</b>	0.71	0.80
MAIAC TERRA	1 km	0.91	0.90	0.71	0.80
MODIS AQUA	0.05°	0.77	0.50	0.30	0.37
MODIS TERRA	0.05°	0.81	0.65	0.32	0.43
NISE	25 km	0.85	0.87	0.37	0.51

583 Table A2: Evaluation of daily snow extent data set performance for 2015. GHCN-D surface  
 584 observations are used as “truth”. The highest value for each metric is shown in bold.

585

	Accuracy	Precision	Recall	F
CMC	0.92	0.81	0.81	0.81
IMS	<b>0.93</b>	0.84	<b>0.85</b>	<b>0.84</b>
MAIAC AQUA	0.87	0.69	0.73	0.71
MAIAC TERRA	0.88	0.68	0.73	0.71
MODIS AQUA	0.78	0.50	0.41	0.45
MODIS TERRA	0.83	0.68	0.43	0.53
NISE	0.85	<b>0.87</b>	0.37	0.52

586 Table A3: Evaluation of daily snow extent data set performance for 2015. GHCN-D surface  
 587 observations are used as “truth”. All products are regridded to a common 25 km resolution. The  
 588 highest value for each metric is shown in bold.