

Response to the comments made by Reviewer # 1 on amt-2018-172

The authors are thanking the anonymous reviewer for offering constructive comments, which have helped us refine the content of our manuscript.

5

Following is the response to each comment and suggestion made by the reviewer.

Reviewers' comments are in BLACK ink

Authors' response is in BLUE ink

10

RC: Both ACAOD and aerosol corrected COD are derived from their retrieval algorithm. However, no validation effort is provided for the derived aerosol corrected COD values. I understand that this is a paper that focuses on ACAOD, but the authors shall at least discuss how the uncertainties in the derived aerosol corrected COD would affect the derived ACAOD and vice versa

15

AR: The near-UV 'color ratio' technique employed in the OMACA algorithm retrieves both ACAOD and aerosol-corrected COD simultaneously. The two pieces of information, i.e., UVAI and reflectance at 388 nm from OMI, allows us retrieving two quantities as reflected in the 2D retrieval diagram shown in Figure 1. While the primary focus of the paper is to highlight the ACAOD product, its spatial-temporal distribution, and initial validation against ORACLES/HSRL-2 observations, we are working with the ORACLES team to perform a detailed validation of the OMACA product, both ACAOD and COD, using airborne in situ and remote sensing measurements. The results of the validation analysis will be covered in a dedicated follow-up publication.

20

25

Section 4 Preliminary Validation is now updated with our intention of a follow-up publication on a detailed validation of the OMACA product.

30

We don't expect to see an interdependency of ACAOD and aerosol-corrected COD retrievals as both quantities are derived simultaneously that explain the two-channel TOA measurements. The inversion philosophy is quite analogous to that adopted in the standard MODIS cloud retrievals where the TOA signal at 0.86 μm is largely responsible for COD magnitudes, and 2.1 μm is for the effective radius. In the present case, the reflectance at 388 nm with an aerosol correction determines the retrieved values of COD and magnitudes of the observed UVAI drives the ACAOD retrievals. So, the assumptions about aerosol and cloud models made in the inversions are the ones that determine the actual uncertainties in both retrievals.

35

5 **RC:** For this product to have a board user community, and especially for modelers, the uncertainties in retrieved properties (such as aerosol corrected COD and ACAOD) shall be provided at the individual retrieval level. Uncertainties are not included in the current data fields as shown in Appendix I. Given the uncertainty in SSA of ± 0.03 , an uncertainty in ACAOD could be introduced on the order of 20-50% (Table 3). I wonder how much the uncertainties in their regional and global time series of ACAOD are attributed to uncertainties in SSA, or are a direct reflection of temporal variations in SSA.

10 **AR:** Reviewer has correctly pointed out here that the uncertainties in the retrieved properties aren't provided at the individual retrieval level. This is a complex task due to the very nature of uncertainty in ACAOD/COD that depends on multiple assumptions made in the inversion. Errors in the ACAOD retrievals resulting from the uncertainty in each major assumption, i.e., SSA, ALH, and AAE, have already been tabulated in Table 3, 4, and 5. The uncertainty in these assumptions can vary on both sides, i.e., underestimation and overestimation, from the assumed state. Moreover, individual uncertainties can also be of opposite signs leading to the partial cancellation of errors in the retrievals. In this situation, it is hard to estimate actual errors in the ACA retrievals especially when the true state of the atmosphere is unknown to the algorithm.

20 Looking at the potential use of the OMACA product in the numerical models, we will consider including the pixel-level retrieval uncertainties in the next upgrade on the basis of the sensitivity of ACAOD/COD to these assumptions individually, such as presented in Table 3, 4, and 5.

25 We assume here that uncertainty in the SSA is random, i.e., distributed on both sides, positive and negative. An OMI-AERONET comparison figure shown to address one of the following comments show that OMI SSAs are spread over both sides of collocated AERONET inversion. Given this assumption, it is expected that the resulting errors in ACAOD/COD retrievals are subjected to the partial cancellation on a monthly regional scale. Moreover, the daily regional time-series of SSA displayed in Figure 2 do not show any apparent long-term trend; rather it demonstrates the known seasonal variations in aerosol absorption properties for both smoke and dust.

30 **RC:** Page 5, lines 32-33, I am not really sure what the authors mean by this sentence "These two components the OMACA algorithm is identical to the ones adopted in the operational cloud-free OMI/OMAERUV two-channel algorithm."

40 **AR:** This sentence refers to the Aerosol Layer Height and Surface Albedo datasets—both are directly adopted from the OMAERUV cloud-free algorithm. Not only that, the aerosol type identification scheme and aerosol models (microphysical and optical properties) are also identical to the ones designed for the OMAERUV.

5 **RC:** Page 5, line 24, The ALH dataset ,which was derived using 30 months of collocated CALIOP and OMI data, is used for aerosol vertical profiles. Was this ALH dataset derived using aerosol above cloud scenes only? If the ALH dataset was derived using cloud free scenes, how representative is the ALH dataset for aerosol above cloud cases?

10 **AR:** The OMI-CALIOP climatology of ALH was derived using mostly clear-sky observations in both datasets with maximum LER in OMI dataset restricted to 0.25. The threshold in LER (0.25) largely avoided ACA scenes that are assigned with the best quality OMACA retrievals (Table 1).

15 The time-series of fractional AOD shown in Figure 9 (right-hand axis in blue ink) of the original manuscript demonstrated that a large fraction of the total column AOD, about 80%-100%, was retrieved as aerosols above the clouds during the seasonal biomass burning and dust episodes. Moreover, the transport of aerosols over the ACA hotspot regions is known to occur above the boundary layer and between the altitudes 3-6 km, as observed from CALIOP lidar. Note that the minimum (maximum) ALH assumed in the OMACA was fixed to 3.0 km (6.0 km) even if the gridded ALH climatology dataset is assigned with lower (higher) layer height. For these reasons, we expect that the OMI-CALIOP ALH database derived from cloud-free observations over the ACA regions is also a representative for scenes with aerosols above cloud. The remaining uncertainty in ALH would translate into the corresponding error in ACAOD already presented in Table 3.

This description has been added in the revision (section 3 Uncertainty Estimates)

25 **RC:** Page 6, lines 29-30, “Observations of aerosols above cloud found outside the boundaries of these 14 pre-selected regions are assigned a fixed SSA of 0.89 and 0.9 for the smoke and dust aerosol types, respectively.” Justifications or references are needed for values mentioned here.

30 **AR:** The SSA values of 0.89 (smoke) and 0.9 (dust) prescribed for regions outside the 14 pre-selected regions are merely our assumption. These values correspond to the aerosol model having a moderate level of absorption for both aerosol types (see Appendix I, aerosol models). We emphasize here that though the OMACA is a global product, it was primarily designed to capture ACA events over major and some minor regions of the world. The selection of regional boundaries was made according to the monthly and seasonal ACA frequency of occurrence maps shown in Figure 5. The pre-selected regions adequately encompass the areas of frequent aerosol-cloud overlaps.

This description has been further clarified in the revised paper (section 2.2.2.3)

40

5 **RC:** Page 8, lines 23-25, “Notice that the current OMACA product does not use the OMMYDCLD product while making above-cloud aerosol retrieval. Instead, we use the information on the geometric cloud fraction derived from OMMYCLD in the post-retrieval analysis.” I wonder why the OMMYDCLD product is not used in the retrieval process. The authors seem to use $LER > 0.2$ to distinguish clear from cloudy scenes. But wouldn't the use of OMMYCLD result in a more accurate estimation of cloud coverage over a given scene? How do the authors deal with partially cloudy scenes?

10 **AR:** As the reviewer has correctly mentioned here, the OMMYDCLD product hasn't been used while deriving the OMACA product. However, we positively consider integrating the OMMYDCLD product into the OMACA algorithm in the near-future upgrade. This will help us to better understand OMI sub-pixel cloud variability and its impact on the retrievals.

15 However, we have used the OMMYDCLD product in the post-retrieval analysis to derive all results presented in this paper. Using the information on the geometric cloud fraction calculated from OMMYDCLD, we adopted a threshold of 0.50 and 0.75 to carry out the frequency of ACA and ACAOD/COD analyses, respectively (section 2.2.3).

20 The minimum LER threshold for detecting cloudy pixels is chosen to be 0.2. The retrieval associated with LER range 0.20-0.25 are flagged as of lower quality due to the chances of encountering partly cloudy-pixels for these retrievals (see Table 1). However, retrievals with $LER > 0.25$ are considered to be more reliable ($QFlag=0$, best quality) owing to increased probability of detecting fully overcast pixels. Currently, the OMACA algorithm treats each identified cloudy pixel as fully overcast scene, even if it is partially cloudy as per the OMMYDCLD product.

30 **RC:** Page 10, lines 20-21, “53% ($AOD > 0.7$, $UVAI > 1.0$) of the total OMAERUV-AERONET SSA (440 nm) retrievals are found to agree within their estimated uncertainties of ± 0.03 .” This means 47% ($AOD > 0.7$, $UVAI > 1.0$) of SSA retrievals are outside of the uncertainty range of ± 0.03 . I am not sure how the authors could come up with this statement “Therefore, we expect that the above-cloud SSA values assigned in the OMACA algorithm over different regions should be accurate within ± 0.03 .”

40 **AR:** The statements referring to the statistics of OMI versus AERONET SSA comparison were derived from Jethva et al. [2014] paper based on the earlier OMAERUV dataset (version 1.4.2 released in 2012). Since then, the OMAERUV algorithm has been upgraded with several major changes, including better treatment of dust particles assuming realistic spheroidal shape

distribution, accounting for angular scattering effects of clouds in the calculation of UV Aerosol Index, use of new minimum surface LER dataset using synergy of multi-year OMI and MODIS observations, and updated cloud screening and retrieval flagging scheme.

5 The regional, daily SSA dataset used in the OMACA product has been derived from this latest
version of the OMAERUV (version 1.8.9.1) product released in 2017. As already described in
the manuscript (section 2.2.2.3), we use UVAI-weighted averages of SSA for smoke and dust
aerosol types using retrievals with measured UVAI>0.8 assuring that only high-quality retrievals
are used in these calculations. The figure shown here compares the SSA derived from the latest
10 OMAERUV product and the AERONET Level 2 dataset over global AERONET sites
(unpublished). The comparative analysis demonstrates that the agreement between the two
independent sets of SSA improves significantly at higher values of UVAI. Quantitatively, about
59% (83%), 65% (88%), and 72% (91%) of the matchups are found to be within the expected
limits of ± 0.03 (± 0.05) difference given the observed range of UVAI>0.8, >1.5, and >2.0,
15 respectively.

The reason for adopting an UVAI-weighted scheme precisely reflects the fact that the agreement
between OMI and AERONET SSA improves at higher aerosol loading/absorption providing
increased confidence in the satellite retrievals.

20 In the revision, we have modified the description of the OMI-AERONET SSA statistics
according to the new comparison results discussed above, since it is consistent with the SSA
dataset used in the OMACA algorithm.

25




Figure. Comparison of OMI-retrieved cloud-free SSA (y-axis) against that of AERONET (x-axis) over globally distributed network for OMI UVAI measurements >0.8 (top), >1.5 (middle), and >2.0 (bottom). Only smoke (red) and dust (blue) aerosols types (relevant to OMACA product) as identified by the OMAERUV algorithm are considered. The statistics of the comparison are shown within each plot.

Response to the comments made by Reviewer # 2 on amt-2018-172

The authors are thanking the anonymous reviewer for offering constructive comments, which have helped us improve the content of our manuscript.

5

Following is the one-to-one response to each comment made by the reviewer.

RC: Referee Comment

AR: Author's Response

10

RC: Introduction: a couple of important papers in the field of above-cloud aerosol studies are missing in the introduction. They should be cited to give the readers a more comprehensive and complete overview of the field. Devasthale and Thomas [2011] is among the first to study the occurrence of above-cloud aerosols.

15

AR: We realized that we missed these citations in the earlier version of the paper. Relevant papers, as suggested by the referee, are now added/cited at appropriate places in the revised paper.

20

- Zhang et al. (2016) cited for direct radiative effects of aerosols above cloud.
- Min and Zhang (2014) cited along with the statement listing several parameters affecting radiative effects.
- Devasthale and Thomas (2011) cited along with the statement “Such situations are commonly observed from satellites over several oceanic and continental regions..”
- Devasthale and Thomas (2011) and Zhang et al. (2016) cited and mentioned in section 5.1 describing the results on frequency occurrence of ACA.
- Lu et al. [2018] now cited along with Wilcox [2012].

25

30

RC: I think a brief overview of the existing above-cloud aerosol retrievals algorithms for the passive sensors will give the readers a “big picture” and understand the relative position of this study. In particular, as the authors are aware, the following algorithms have been developed for POLDER and MODIS

35

AR: A discussion on the existing state of the active and passive-sensor based ACA algorithms are now added to the Introduction (2nd paragraph).

RC: Aerosol type identification (section 2.2.2.1): This part is very important, Because the retrieval algorithm uses different optical properties for different type of aerosols, i.e., dust or smoke. A misclassification can cause retrieval errors and uncertainties. However, the description of this paper is very brief. Some more details need to be added with proper references. For example, it should be mentioned whether and how the identification scheme is validated or evaluated. Has it been compared with CALIOP aerosol subtypes? Why different threshold of CO is used for northern and southern hemispheres?

AR: The aerosol type identification scheme in the OMACA has been directly adopted from the cloud-free OMAERUV algorithm. The scheme uses real-time observation of AIRS CO information in conjunction with OMI UVAI to discern the carbonaceous smoke aerosols from mineral dust, which otherwise not possible to detect using only near-UV measurements. The use of CO measurements also enables the identification of high levels of boundary layer pollution undetectable by near-UV observations alone. Since *Torres et al.* [2013] adequately describes the methodology and implementation of the scheme within OMAERUV, we didn't include a lengthy discussion on this topic in the present manuscript.

The different threshold values of CO in Northern and Southern hemispheres correspond to the average of AIRS CO climatological annual minima over major biomass burning/boreal fire activity regions. These values are 2.2×10^{18} in the Northern Hemisphere (NH) and 1.8×10^{18} for the Southern Hemisphere (SH), based on *Yurganov et al.* [2008, 2010]. The presence of carbonaceous aerosols is assumed if $AI \geq AI$ threshold (0.8) and $CO \geq CO$ threshold (2.2×10^{18} for NH and 1.8×10^{18} for SH) or when CO values larger than 2.8×10^{18} (2.5×10^{18}) are observed in the Northern (Southern) Hemisphere regardless of AI considerations. Conversely, OMI pixels with observed $AI \geq AI$ threshold (0.8) and $CO < CO$ threshold are assigned with the dust aerosol type. Threshold values in AI and CO represent noise and background levels in the respective measurements not necessarily associated with the free troposphere CO burden which is expected to co-exist with the lofted carbonaceous aerosols.

The straightforward way of discerning the absorbing aerosol type works efficiently in most cases, however, may break down under certain situations, i.e., when dust aerosols are present over regions characterized by high CO levels associated with pollution episodes other than the biomass burning smoke for which the scheme would assign absorbing aerosol type as smoke. Note that the aerosol type identification scheme doesn't account for the mixture of aerosols for which either smoke or dust aerosol type is assigned depending upon the threshold values of AI and CO.

A detailed regional-level comparison between CALIOP aerosol sub-type and that of the OMAERUV hasn't been done, but we consider conducting the said analysis in the near future.

A brief description of the aerosol type identification scheme is now provided in the revised paper in section 2.2.2.1.

5

RC: Single scattering albedo (section 2.2.2.1): the SSA is extremely important for ACA retrieval and DRE. I hope the information of aerosol type and the corresponding SSA will be part of the OMIACA so the users can use it in a consistent way with the AOD product

10 **AR:** We fully agree with the referee that the assumption of aerosol SSA above the clouds is extremely crucial for deriving accurate ACAOD retrievals as well as in the quantification of DRE. Section 2.2.2.1 describes how we take advantage of clear-sky SSA retrievals from the OMAERUV product and assign a representative SSA value for each region at daily scale. In this regard, the OMACA stands alone among passive-sensor based ACA algorithms that currently
15 rely on a fixed value of SSA [*Jethva et al., 2013; Myer et al., 2015*].

Looking at its importance, the values of aerosol SSA above-cloud assumed in the OMACA algorithm for the three wavelengths, i.e., 354, 388, and 500 nm, are already reported in the product. The corresponding SDSs are named as “InputSSA354”, “InputSSA388”,
20 “InputSSA500”. Additionally, the aerosol type associated with each valid ACA pixel is also stored in the product as “AerosolType”. Refer to the complete list of SDS stored in each OMACA HDF-EOS file in Appendix I.

25 **RC:** Look-up-tables (section 2.2.24): How is cloud effective radius considered in the LUT? Is it assumed as a constant? Note that the assumption of CER could have impacts on the COD retrieval. Some discussion is needed to clarify this.

30 **AR:** In LUT calculations, clouds are assumed to be liquid in phase and follow the standard C1 size distribution [Deirmendjian, 1969]. The effective radius of C1 water cloud droplet distribution is assumed to be a constant value of 6.0 microns. To answer the reviewer’s concern, we carried out a sensitivity analysis, similar to the ones presented in Table 3, 4, and 5, in which the errors in both ACAOD and aerosol-corrected COD were calculated following the perturbation approach around the assumed CRE value of 12.0 μm . The table shown below lists
35 the errors in aerosol-corrected COD due to the range of uncertainty in the assumed cloud CRE. The analysis was performed assuming reference cloud CRE of 12 μm and for the ACAOD values of 0.5 and 1.0 (388 nm).

| Error in cloud effective radius (μm) | Assumed AOD (388 nm) = 0.5/1.0 Reference cloud effective radius = 12.0 μm % Error in Aerosol-corrected COD (388 nm) | | | |
|---------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|---------------|-------------|---------------|
| | COD=5 | COD=10 | COD=20 | COD=30 |
| -8.0 | -11.31/-11.9 | -10.28/-10.88 | -9.13/-9.83 | -10.11/-10.76 |
| -6.0 | -5.94/-6.54 | -5.38/-5.94 | -4.68/-5.29 | -5.01/-5.62 |
| -4.0 | -2.85/-3.26 | -2.57/-2.94 | -2.18/-2.57 | -2.25/-2.66 |
| 4.0 | 1.17/1.4 | 1.43/1.79 | 1.21/1.62 | 0.95/1.39 |
| 6.0 | 1.34/1.70 | 1.63/2.07 | 1.37/1.87 | 1.05/1.59 |
| 8.0 | 1.51/1.94 | 1.83/2.35 | 1.52/2.11 | 1.16/1.78 |
| 12.0 | 1.85/2.41 | 2.22/2.92 | 1.83/2.59 | 1.36/2.18 |

5 The errors in COD retrievals due to the uncertainty in CRE follow asymmetric behavior to the perturbation around the assumed state. While a large underestimation in CRE of -8 μm produces negative errors of ~10%-11% in the retrieved COD, an overestimation in CRE of +8 to +12 μm yields positive errors of much smaller magnitudes (~1%-2%). The spatial distribution of MODIS
10 monthly cloud CRE over the Southeastern Atlantic Ocean, as shown in Figure 11 of Meyer *et al.* (2015), exhibits spatial variations with smaller droplets (CRE 7-11 μm) concentrated closer to the coast and relatively larger size droplets (11-17 μm) retrieved away from the coast. Given the fixed value of CRE equals 6.0 μm assumed in the OMACA cloud LUTs, the observed variations from MODIS would produce <2% error in the retrieved aerosol-corrected COD.

15 The corresponding errors in ACAOD due to the uncertainty in cloud CRE are found to be marginal. For an ACAOD>0.5, an uncertain assumption in cloud CRE by $\pm 8 \mu\text{m}$ results in ACAOD errors <2% with much smaller magnitudes at higher aerosol loading. This is because at larger ACAODs the aerosol absorption effects dominate over that resulting from varying effective radius of liquid droplets leaving other major algorithmic assumptions, i.e., SSA, ALH, and AAE to determine the resultant uncertainty in ACAOD retrievals.

20 This analysis implies that near-UV wavelengths don't offer a strong sensitivity to the variations in cloud droplet size rather the cloud signal is predominantly driven by the optical thickness. Due to the lack of information on cloud droplet size from OMI, we adopted the standard C1 cloud model validated and used in numerous studies for all cloud LUT calculations.

25 **The description provided above is now added to the section 3 of the revised paper.**

5 **RC:** Partly cloudy pixels: the footprint size of OMI is 13x24 km (338km²). At this scale, there
of the partly cloudy OMI pixels. It seems to me the OMIACA algorithm is applied to both
overcast pixels and partly cloudy pixels, correct? How is the subpixel cloud fraction determined?
How are the partly cloudy pixels treated in the LUT and radiative transfer simulations? How is
the UVAI of clear-sky part of the partly cloudy pixel different from that of the cloudy part, and
what is the meaning of the “observed” UVAI for the partly cloudy pixel? I would strongly
10 recommend the authors to add a separate and dedicated sub-section to discuss the treatment of
partly cloudy pixels in the OMIACA algorithm.

15 **AR:** OMACA algorithm performs retrievals for each pixel of size 13 x 24 km-square at nadir
independently. As the referee has correctly pointed out, there is a possibility of encountering
partly cloudy pixels, especially for measurements with lower reflectivity (388 nm) values. The
algorithm quality flags reported in OMACA product precisely reflect these observed conditions
(Table # 1 QFlag values 0, 1, and 2). Due to the coarser resolution of OMI pixels, there seem to
be is no direct way to infer the sub-pixel cloud variability using only OMI measurements.
Therefore, we have used the OMI-MODIS joint cloud product, OMMYDCLD, post-retrievals for
20 all analyses reported in the manuscript. The OMMYDCLD product, as already explained in the
original manuscript in section 2.2.3, provides the statistics of the MODIS 1-km cloud product
(MYD06) on each collocated OMI footprint.

25 An analysis using the OMMYCLD product over the Southeastern Atlantic Ocean for the period
of Jun-July-Aug 2007 revealed a well-constrained non-linear relationship between the MODIS-
derived COD times the geometric cloud fraction and LER388. A threshold of LER388 of 0.25
adopted for the best quality retrievals (QFlag=0) compares to the COD times geometric cloud
fraction of 3-4. Conversely, given the geometric cloud fraction of unity, the minimum COD
retrieved by OMACA would be in the range 3-4. Retrievals assigned with QFlag=1 further
30 extends the LER388 to 0.20 allowing pixels with relatively lower reflectivity with much stronger
absorption (larger UVAI) above the clouds [Jethva et al., 2013, Figure 6, Aug 12, 2006 case
study].

35 Currently, the OMACA algorithm is designed to perform inversion over fully overcast pixels.
The LUTs are generated assuming fully cloudy conditions and do not explicitly treat partly cloud
pixels. To avoid a large fraction of partly cloudy pixels, therefore, we adopted the geometric
cloud fraction thresholds calculated using OMMYDCLD product of 0.5 and 0.75 for the FOACA
(Figure 5, 6, 7) and AOD/COD analyses (Figure 8, 9, 10, 11).

We will consider using the OMMYDCLD product in the OMACA processing in the next upgrade of the algorithm. We believe that most of the explanation provided here on this concern is already described in the original version of the paper in section 2.2.3 and 2.2.4.

5

RC: Sub-pixel COD variation: A related question is whether and how the algorithm accounts for the subpixel COD variation. What is the physical meaning of the “retrieved COD”? Is it a simple mathematical mean or some kind of weighted mean?

10

AR: As explained in the previous response, each OMACA retrieval corresponds to the respective pixel size derived using a single set of reflectivity and UVAI values. The retrieved values of ACAOD and COD, therefore, represent an overall condition observed in each pixel. It is hard to draw a conclusion that the observed condition is merely a mathematical mean or weighted mean of the sub-pixel cloud variability as the TOA reflectance versus COD relationship exhibits a non-linear behavior especially at higher values of CODs.

15

RC: Spatial distribution of ACA: As mentioned above Zhang et al. (2016) studied the global distribution of different types of ACA. Actually, Figure 5 agrees reasonably well with the Figure 2 of Zhang et al. (2016). In addition, Zhang et al. (2016) also found significant amount of ACA over the north pacific due to the Asian dust and pollution. This study should be cited here and discussed.

20

AR: We referred to Zhang et al. [2016] paper which also shows similar cloudy-sky FOACA results over the global ocean derived using 8-year daytime CALIOP observations. Prior to *Zhang et al.* [2016] and our study, *Devasthale and Thomas* (2011) also conducted FOACA analysis using 4-years of CALIOP data. Both studies are now mentioned and cited in the revised manuscript.

25

30

RC: Figure 5 shows some ACA over the Southern Ocean in January and February. Is this true or some retrieval artifact?

35

AR: Most likely ACA over the Southern Ocean during the winter months inferred from our analysis is an artifact resulting from non-aerosol related enhancement in UVAI observed at certain geometry conditions that are associated with higher solar zenith and viewing zenith angles. Although these artifacts are largely removed from the best quality retrieval group (see Table 1 of the original manuscript) based on thresholds in geometry and are assigned a different

40

quality flag (=3), some residual ACA pixels still reside within the group of good quality retrievals. One of the possible reasons for the non-aerosol related enhancement in the UVAI could be the presence of ice clouds over the Southern Ocean for which the effect of angular scattering is unaccounted for in the calculation of UVAI.

5

RC: It is interesting to see that the FOACA in Figure 5 and the ACA AOD in Figure 8 are highly correlated. Is this a coincidence or there may be some real connection between them? One could argue that the region with high FOACA does not necessarily have a larger ACA AOD. Do you agree?

10

AR: It is assumed here that the reviewer is referring to the high-level of spatial consistency in FOACA and ACAOD maps. This is because of the OMACA algorithm's efficiency to quantitatively retrieve ACAOD and COD when the presence of absorbing aerosols over clouds is identified based on a set of LER and UVAI thresholds. We agree with the reviewer that high frequency occurrence and large ACAOD do not necessarily correlate with each other. FOACA is simply a measure of the temporal occurrence of the aerosol-cloud overlap, whereas ACAOD is a quantitative measure of actual aerosol loading above the clouds.

15

A 12-Year Long Global Record of Optical Depth of Absorbing Aerosols above the Clouds Derived from OMI/OMACA Algorithm

Hiren Jethva^{1,2}, Omar Torres², Changwoo Ahn³

¹Universities Space Research Association, 7178 Columbia Gateway Drive, Columbia, MD 21046, USA

²NASA Goddard Space Flight Center, Earth Science Division, Code 614, Greenbelt, MD 20771, USA

³Science Systems and Applications, Inc., 10210 Greenbelt Rd, Lanham, MD 20706 USA

10 **Correspondence to:** Hiren Jethva (hiren.t.jethva@nasa.gov)

Abstract. Aerosol-cloud interaction continues to be one of the leading uncertain components of the climate models, primarily due to the lack of adequate knowledge of the complex microphysical and radiative processes of the aerosol-cloud system. Situations when the light-absorbing aerosols such as carbonaceous particles and windblown dust overlay low-level cloud decks are commonly found in several regions of the world. Contrary to the known cooling effects of these aerosols in cloud-free scenario over darker surfaces, an overlapping situation of the absorbing aerosols over the cloud can lead to a significant level of atmospheric absorption exerting a positive radiative forcing (warming) at the top-of-atmosphere. We contribute to this topic by introducing a new global product of the above-cloud aerosol optical depth (ACAOD) of absorbing aerosols retrieved from the near-UV observations made by the Ozone Monitoring Instrument (OMI) onboard NASA's Aura platform. Physically based on an unambiguous 'color ratio' effect in the near-UV caused by the aerosol absorption above the cloud, the OMACA (OMI Above-Cloud Aerosols) algorithm simultaneously retrieves the optical depths of aerosols and clouds under a prescribed state of the atmosphere. The OMACA algorithm shares many similarities with the two-channel cloud-free OMAERUV algorithm, including the use of AIRS carbon monoxide for the aerosol type identification, CALIOP-based aerosol layer height dataset, and OMI-based surface albedo database. We present the algorithm architecture, inversion procedure, retrieval quality flags, initial validation results, and results from a 12-year long OMI record (2005-2016) including global climatology of the frequency of occurrence, ACAOD, and aerosol-corrected cloud optical depth. A comparative analysis of the OMACA-retrieved ACAOD collocated with equivalent accurate measurements from the HSRL-2 lidar for the ORACLES phase I operation (August-September 2016) revealed a good agreement ($R=0.77$, $RMSE=0.10$). The long-term OMACA record reveals several important regions of the world, including Southeastern Atlantic Ocean, southern Indian Ocean, South-East Asia, tropical Atlantic Ocean off the coast of western Africa, and northern Arabian sea where the carbonaceous aerosols from the seasonal biomass burning and mineral dust originated over the continents are found to overlie low-level cloud decks with moderate ($0.3 < ACAOD < 0.5$, away from the sources) to higher levels of ACAOD (> 0.8 in the proximity to the sources). No significant long-term trend in the frequency of occurrence of aerosols above the clouds and ACAOD is noticed when OMI observations that are free from the 'row anomaly' throughout the operation are considered. If not

Deleted: coincident and collocated

Deleted: and

5 accounted, the effects of aerosol absorption above the clouds introduce low bias in the retrieval of cloud optical depth with a profound impact at increasing ACAOD and cloud brightness. The OMACA aerosol product from OMI presented in this paper offers a crucial missing piece of information of the aerosol loading above cloud that will help us to quantify the radiative effects of clouds when overlaid with aerosols and its resultant impact on cloud properties and climate.

Keywords: aerosols above clouds, aerosol absorption, color ratio, Ozone Monitoring Instrument, OMACA

1. Introduction

Aerosol-cloud interactions continue to be the most significant source of uncertainty in estimating the role of aerosols and clouds in Earth's changing radiation budget (IPCC, 2014). One of the main hurdles is the lack of adequate knowledge of the complex microphysical and optical processes of an aerosol-cloud system that govern the resultant impact on the regional and global climate. An important aspect of the problem is when the light-absorbing aerosols such as biomass burning generated carbonaceous particles and windblown mineral dust overlay low-level cloud decks. Such situations are commonly observed from satellites over several oceanic and continental regions of the world on daily to seasonal scales (Devasthale and Thomas, 2011; Alfaro-Contreras et al., 2015). The transoceanic transport of elevated fine mode aerosols from biomass burning and coarse mode aerosols from dust storms observed from the ground and satellites is a well-known phenomenon (Prospero et al., 2002; Kaufman et al. 2005; Chand et al., 2008; Torres et al., 2012). Aerosols in the cloud-free scenario over the dark surfaces are known to produce a net cooling effect (negative radiative forcing) on climate. On the contrary, an overlapping situation of the absorbing aerosols over the cloud can lead to a significant level of atmospheric absorption and thus exert a positive radiative forcing at the top-of-atmosphere (TOA) (Kiel and Haywood, 2003; Chand et al.; 2009, Meyer et al., 2013; Feng and Christopher, 2015; Zhang et al., 2016). The magnitude of direct radiative effects of aerosols above cloud depends directly on the amount aerosol loading, microphysical-optical properties of the aerosol layer (Eswaran et al., 2015) and the underlying cloud deck (Meyer et al., 2013), geometric cloud fraction and cloud optical depth (Chand et al., 2009) as well as the diurnal cycle of clouds (Min and Zhang, 2014). The resultant aerosol-driven atmospheric heating can have a significant influence on the atmospheric stability, cloud formation and lifetime, and the hydrological cycle (Wilcox, 2012, Lut et al., 2018). Different climate models treat aerosol-cloud interaction processes differently, which results in significant inter-model discrepancies in aerosol forcing assessments, especially over the absorbing aerosol-cloud overlap regions, i.e., Southeastern Atlantic Ocean and South-East Asia (Schulz et al., 2006). To accurately quantify the direct and semi-direct effects of aerosols in the cloudy skies, it is imperative that a measurement-based global database is made available to the community.

In the past decade, the development of several independent algorithms that quantifies aerosol loading above cloud from satellite-based active as well passive sensors have been a breakthrough. These techniques have shown the potential to retrieve above-cloud aerosol optical depth (ACAOD) using measurements from different A-train sensors (Jethva et al., 2014). The depolarization ratio (Hu et al., 2007) and color ratio methods (Chand et al., 2008) applied to the CALIPSO/CALIOP active lidar measurements of two-way transmittance (Chand et al., 2008) directly provide ACAOD and corresponding Angstrom Exponent. Waquet et al. (2009) make use of polarized radiances measured by PARASOL/POLDER to retrieve ACAOD. An operational version of this algorithm capable of retrieving ACAOD globally has also been developed (Waquet et al., 2013). Taking the advantage of enhanced aerosol absorption sensitivity in the near-UV, Torres et al. (2012) introduced a technique to retrieve ACAOD and underlying aerosol-corrected cloud optical depth (COD), simultaneously. Jethva et al. (2013) have extended the color ratio method to the visible and shortwave IR (SWIR) channel measurements of MODIS to deduce a pair of ACAOD and aerosol-corrected COD. Using an iterative optimal estimation approach applied to the MODIS observations, Meyer et al.

Deleted: as well as

(2015) and Sayer et al. (2016) have developed multi-spectral algorithms to simultaneously retrieve ACAOD, COD, and cloud effective radius of ACA scenes.

5 Satellite-based passive and active sensors can observe the aerosols in the entire atmospheric column above the cloud deck. Conventional satellite retrievals have provided a global distribution of aerosols over only cloud-free regions, leaving the vast cloudy region unmonitored regarding the presence of aerosols. Likewise, field campaigns in the past mainly focused on the measurements and characterization of aerosol properties in cloud-free conditions to evaluate and improve satellite-based retrievals as well as model simulations. The lack of measurements-based aerosol database in the cloudy atmosphere severely limits our understanding of the aerosol effects on cloud radiative forcing and microphysical properties.

10 *Torres et al.* (2012) introduced a remote sensing technique to simultaneously retrieve ACAOD and aerosol-corrected COD, using the near-UV observations made by the Ozone Monitoring Instrument (OMI) onboard NASA's Aura satellite. The method is physically established on an unambiguous 'color ratio' effect produced by the absorbing aerosols above clouds in the two near-UV channels, i.e., 354 nm and 388 nm. Under a prescribed state of the atmosphere, the aerosol-led changes in the cloud radiances measured at the top-of-atmosphere (TOA) relate to a pair of ACAOD and COD. The technique was successfully tested over the case studies of carbonaceous aerosols above the cloud deck found over the Southeastern Atlantic Ocean. Furthermore, ACAOD derived using the near-UV color ratio method was found to be consistent when inter-compared against those inferred from other research-based algorithms applied to the different A-train sensors (MODIS, CALIOP, POLDER) (*Jethva et al.*, 2014).

Deleted: the above-cloud aerosol optical depth (ACAOD)

Deleted: underlying cloud optical depth (COD)

15 In this paper, we apply the near-UV color ratio algorithm on a global scale to produce a Level 2 orbital dataset of ACAOD using OMI observations. First, we describe different components of the algorithm (section 2) followed by the theoretical estimates of the expected uncertainties (section 3). Initial validation results of the OMACA aerosol retrieval using airborne HSRL-2 measurements are presented in section 4. The results of the frequency of occurrence analysis of the aerosols above clouds discussed in section 5. A global 12-year long record of ACAOD and aerosol-corrected COD derived from OMACA along with a quantitative analysis of the impact of aerosol absorption on COD retrievals are presented in section 6. The paper is summarized and concluded in section 7.

2. Description of the OMACA Algorithm

30 2.1 Physical Basis

Light absorbing aerosols such as carbonaceous aerosols (*Kirchstetter et al.*, 2004) and dust particles (*Wagner et al.*, 2012) exhibit strong wavelength dependence in absorption, particularly in the UV region of the spectrum. On the other hand, clouds show minimal to no spectral dependence in reflectance when measured from the space. When absorbing aerosols overlay the cloud deck, the spectral contrast in the UV is further enhanced producing a strong 'color ratio' effect in the two-channel TOA reflectance measurements (*Torres et al.*, 2012). This effect is often referred to as the 'cloud darkening' caused by the aerosol-led spectral absorption. The UV Aerosol Index (UVAI)

measured by OMI is an excellent indicator of the presence of absorbing aerosols in the clear (Torres et al., 1998) as well as the cloudy atmosphere (Torres et al., 2012). Radiative transfer simulation shows that for a prescribed state of the atmosphere, the reduction in the spectral reflectance and UVAI between a pair of wavelengths depends on the optical depth of both aerosols and cloud, single-scattering albedo, and aerosol layer height. Founded on this unambiguous detection of absorbing aerosols above the cloud, Torres et al. (2012) introduced a technique that delivers the simultaneous retrieval of ACAOD and aerosol-corrected COD from OMI's two near-UV observations (354 and 388 nm). Figure 1 shows the two-dimensional domain of simulated reflectance at 388 nm (x-axis) and UVAI (y-axis) for several pairs of ACAOD and COD for the carbonaceous aerosols (left panel) and spheroidal dust particles (right panel). Under a prescribed set of assumptions, i.e., aerosol layer height (ALH), aerosol single-scattering albedo (SSA), surface albedo, and geometry, the two-channel measurements of OMI can be related to a pair of ACAOD and COD.

2.2 Inputs and Ancillary Datasets

2.2.1 Direct Input

We use OMI Level-1b calibrated and geo-located radiance measurements at 354 nm and 388 nm as the primary input to the OMACA algorithm. Foremost, the observed radiances are used to calculate the UVAI (Mie), as described by Torres et al. (2018), which is a residual quantity resulting from the comparison between measured and calculated radiances between 354 nm and 388 nm. Compared to the previously adopted Lambert Equivalent Reflector (LER) based method for calculating UVAI (Herman et al., (1997), Torres et al., (1998)), the new approach offers a better characterization of clouds by accounting for the angular dependence of cloud reflectance (phase function) of liquid water clouds.

2.2.2 Ancillary Datasets

2.2.2.1 AIRS CO for Aerosol Type Identification

The aerosol-type identification scheme for OMACA has directly adopted from the operational cloud-free OMI/OMAERUV (version 1.8.9.1) two-channel algorithm. It uses Aura/OMI UVAI coincident with the [real-time](#) Aqua/AIRS retrievals of Carbon Monoxide (CO) to distinguish three major aerosol types, i.e., carbonaceous aerosols, dust particles, and urban/industrial aerosols. OMACA considers two absorbing aerosol types, i.e., carbonaceous aerosols and desert dust. The aerosol type identification scheme adopts a threshold of 0.8 in UVAI to assign either smoke or dust aerosol type that subsequently depends on the columnar amounts of CO retrieved by AIRS. Since the CO is a primary gaseous component of open-field biomass burning, it constitutes a reliable tracer of carbonaceous aerosol. For the [Northern Hemisphere \(NH\)](#), the threshold in CO is set to 2.0×10^{18} molecules-cm⁻², whereas that for the [Southern Hemisphere \(SH\)](#) is 1.6×10^{18} molecules-cm⁻². [The different threshold values of CO in](#)

Deleted: n
Deleted: h
Deleted: s
Deleted: h

NH and SH correspond to the average of AIRS CO climatological annual minima over major biomass burning/boreal fire activity regions. The presence of carbonaceous aerosols is assumed if UVAI \geq UVAI threshold (0.8) and CO \geq CO threshold or when CO values larger than 2.8×10^{18} (2.5×10^{18}) are observed in the NH (SH) regardless of UVAI considerations. Conversely, OMI pixels with observed UVAI \geq UVAI threshold (0.8) and CO \leq CO threshold are assigned with the dust aerosol type. Threshold values in AI and CO represent noise and background levels in the respective measurements not necessarily associated with the free troposphere CO burden which is expected to co-exist with the lofted carbonaceous aerosols. For the equatorial region bounded by the latitude range 10°S to 10°N, the aerosol type is determined based on a linearly interpolated value of CO threshold between NH and SH.

Deleted: within

Deleted: s

The straightforward way of discerning the absorbing aerosol type works efficiently in most cases, however, may break down under certain situations, i.e., when dust aerosols are present over regions characterized by high CO levels associated with pollution episodes other than the biomass burning smoke for which the scheme would assign absorbing aerosol type as smoke. Note that the aerosol type identification scheme doesn't account for the mixture of aerosols for which either smoke or dust aerosol type is assigned depending upon the threshold values of UVAI and CO. The reader is recommended to refer Torres et al. [2013] for obtaining a detailed description of the scheme and its implementation within the OMAERUV algorithm.

2.2.2.2 Aerosol Layer Height and Surface Albedo

Deleted:

The representation of aerosol vertical profile relies on an averaged aerosol layer height (ALH) dataset derived using the 30-month long combined and co-located measurements of CALIOP vertical backscatter and OMI UVAI (Torres et al., 2013). The aerosol profile follows a quasi-Gaussian distribution around mean ALH given by the CALIOP-OMI dataset. For the surface characterization, we use near-UV surface albedo database derived using the multiyear OMI LER observations. The method adopts a minimum LER approach which ensures minimal or no contamination from the atmosphere, i.e., aerosols and clouds, in the measured reflectivity. Afterward, the minimum LER dataset derived from the OMI observations was scaled in the temporal domain to the seasonality of surface albedo retrieved in the visible wavelengths from MODIS. The dataset contains surface albedo values at 354 and 388 nm at a grid resolution of $0.25^\circ \times 0.25^\circ$. These two components the OMACA algorithm is identical to the ones adopted in the operational cloud-free OMI/OMAERUV two-channel algorithm (Torres et al., 2013).

2.2.2.3 Above-cloud Aerosol Single-scattering Albedo

The aerosol single-scattering albedo (SSA) is one of the most critical assumptions that can govern the accuracy of ACAOD retrieved from OMI (Torres *et al.*, 2012). A perturbation of +0.03 (- 0.03) in SSA yields about +48% (- 25%) error in ACAOD for a reference AOD and COD of 0.5 and 5.0, respectively. The error in ACAOD follows an asymmetric behavior around the perturbed state and is a dynamic function of both aerosol loading (AOD) and underlying cloud brightness (COD). The OMACA algorithm takes advantage of cloud-free standard OMAERUV global retrievals of SSA for characterizing the absorption capacity of aerosols above the cloud. OMI's two-channel OMAERUV algorithm simultaneously retrieves columnar total AOD and SSA at 388 nm in cloud-free conditions on a daily global scale (Torres *et al.*, 2007, 2013). Both retrieved parameters have been evaluated against the ground-based AERONET measurements globally (Ahn *et al.*, 2014; Jethva *et al.*, 2014). Jethva *et al.* (2014) have shown that for carbonaceous and dust aerosol types, which are relevant to the OMACA product, about 52% (77%) of OMI-AERONET matchups agree within the absolute difference of ± 0.03 (± 0.05). Despite the inherent uncertainties in both inversions, a reasonable agreement between the two independent techniques globally and, in fact, a robust comparison over many important sites affected by biomass burning and dust provided the increased confidence and credibility of the OMAERUV aerosol product.

We have used the existing cloud-free OMAERUV SSA product to generate a daily database to prescribe SSA required in OMACA. The whole world is split up into 14 regions based on the patterns of absorbing aerosols above cloud inferred from the frequency of occurrence analysis (given in section 5). Figure 2 (top) shows the geographical boundaries of the selected regions. For each region and each day of OMI observation, the daily, region-specific value of above-cloud SSA was estimated for the carbonaceous and dust aerosol types separately using the respective cloud-free SSA retrieval weighted by the corresponding UVAI observations (>0.8). Since the OMAERUV algorithm assigns a fixed aerosol type, i.e., smoke, dust, or background aerosol, to each valid cloud-free pixels of OMI, it is possible to estimate daily regional SSA value separately for smoke and dust aerosol types. In the case of missing daily regional SSA due to cloud cover or no OMI orbital data, the method relies on the availability of SSA values on nearby days with a sequential preference, i.e., weekly (± 3 days excluding the day in consideration), monthly, or long-term climatology datasets. Observations of aerosols above cloud found outside the boundaries of these 14 pre-selected regions are assigned a fixed SSA of 0.89 and 0.9 for the smoke and dust aerosol types, respectively.

Using ground-based AERONET inversion, Eck *et al.* (2013) showed that the absorption properties of the smoke aerosols over Central/Southern Africa exhibit a seasonal trend where the monthly mean aerosol SSA (440 nm) increases from 0.84 in July to 0.93 in November. An analysis of the OMI-retrieved SSA (388 nm) in the same paper also showed a similar trend as observed by AERONET and suggest that the seasonal change in SSA is widespread over much of southern Africa. The present approach of assigning the above-cloud aerosol SSA captures the daily

variability of aerosol absorption for each region and therefore eliminates the need to assume a time-invariant value of SSA for the retrievals of above-cloud aerosols.

2.2.2.4 Aerosol Models

5 OMACA considers two major partially absorbing aerosol types, i.e., carbonaceous and dust aerosols. The microphysical and optical properties of these aerosol types are adopted from the current operational OMAERUV aerosol models (Torres *et al.*, 2013). Each aerosol type consists of seven distinct sub-models that differ in their imaginary part of the refractive index. The tables included in Appendix list the model properties of both types of aerosol models. Carbonaceous aerosols are assumed as spherical particles with wavelength-dependent imaginary refractive index in the near-UV region (relative spectral dependence of 20%) that accounts for the presence of organics in the biomass burning generated aerosols (Kirchstetter *et al.*, 2004; Jethva and Torres 2011). Dust aerosols are considered spheroidal in shape with a fixed distribution of the spheroidal axis ratio adopted from Dubovik *et al.* (2006) and applied to the near UV observations (Torres *et al.*, 2018).

15 2.2.2.5 Look-up-Tables

OMACA is essentially a look-up-table (LUT) based algorithm. To generate LUT, we employ the vector discrete ordinate radiative transfer model VLIDORT (Spurr, 2006). Clouds are assumed to be liquid in phase and follow the standard C1 size distribution (Deirmendjian, 1969). Aerosol size distribution is assumed to follow a bimodal, log-normal distribution with parameters adopted from the standard OMAERUV aerosol models (see Appendix I). Carbonaceous aerosols are assumed to be spherical with associated scattering phase functions calculated following the Mie theory. Dust particles, on the other hand, are treated as a mixture of randomly oriented spheroids with a fixed distribution of axis ratios (Dubovik *et al.*, 2006). The phase matrix elements of each spheroidal dust aerosol model of OMACA (Appendix I) were calculated using a set of pre-calculated kernels and associated software package made available by Oleg Dubovik (personal communication). The extracted phase matrix elements of each dust model were ingested into VLIDORT to simulate TOA radiances. More details on the treatment of spheroidal dust in the OMI aerosol retrieval framework is given in Torres *et al.*, (2018).

LUTs were generated for carbonaceous and dust aerosol models. Each aerosol type consists of seven discrete aerosol SSA (388 nm) ranging from 0.75 to 1.00, for the 354- and 388-nm wavelengths for a total of seven node values in ACAOD, eight nodes in COD, at different geometries of solar zenith angle, viewing zenith angle, and relative azimuth angle. The simulations were carried out for two surface pressure levels, for four different ALH referenced at respective surface pressure levels, and for five nodes in surface albedo. The node values for each variable are listed in Appendix I. The LUT radiances interpolated at observed geometry, pressure level, ALH, and SSA are matched with the OMI-observed radiance in 2D retrieval domain (Figure 1) to find a pair of ACAOD and COD at 388 nm. The retrieved values at 388 nm are converted to 354 nm and 500 nm wavelengths following the spectral dependence

of extinction associated with the assumed model in the retrieval process. Figure 3 illustrates a general flow diagram of the OMACA algorithm.

2.2.3 Identification of Absorbing Aerosols above Clouds

We adopt a bi-parametric approach to identify the presence of absorbing aerosols above the cloud. The Lambertian Equivalent Reflectivity or LER measured at a near-UV wavelength is proportional to the brightness of the scene. LER represents the reflectivity of the scene when Rayleigh scattering is taken out from the TOA radiance measurements. On the other hand, UVAI is an excellent indicator of the presence of absorbing aerosols in both cloud-free as well overcast skies (Torres *et al.*, 2012) over all surfaces. Radiative transfer simulations show that while LER is directly proportional to COD, the layers of absorbing aerosols above cloud produces higher magnitudes of UVAI that depend on the above-cloud AOD, aerosol model, and cloud brightness (COD). Thus, higher values of LER and UVAI potentially represents scenes of absorbing layers of aerosols over low-level cloud deck.

OMI offers a spatial resolution of $13 \times 24 \text{ km}^2$ at nadir, which intercepts an area of about 338 km^2 on the ground for the VIS part of the instrument (Algorithm Theoretical Document Basis, OMPXCOR). The ground pixel size and associated area increase significantly at the extreme edge of the swath. A new global product called OMMYDCLD processed in-house co-locates the Aqua/MODIS 1-km cloud retrievals (MYD06) with each OMI pixel footprint ($13 \times 24 \text{ km}^2$) (Joanna Joiner and Brad Fisher, personal communication). OMMYDCLD reports statistics of many MODIS cloud parameters for each OMI footprint such as the median COD, histogram of COD, cloud phase information, and many others. In addition to this, the OMMYDCLD also provides the total number of MODIS 1-km pixels (clear and cloudy) as well as the total number of cloudy pixels identified at 1-km spatial resolution. The accessibility of these two parameters allows calculating the geometric cloud fraction as observed by MODIS for each OMI pixel. Notice that the current OMACA product does not use the OMMYDCLD product while making above-cloud aerosol retrieval. Instead, we use the information on the geometric cloud fraction derived from OMMYCLD in the post-retrieval analysis.

2.2.4 Algorithm Quality Flags

Each qualified OMI retrieval of the above-cloud aerosols is assigned with an appropriate algorithm quality flag. Table 1 describes the algorithm quality flags attached to each valid retrieval and their associated observed conditions. Retrievals with the quality flag equal '0' is expected to be the best in quality as they are associated with the larger magnitudes of UVAI (>1.3), and LER388 (>0.25)-both provide high confidence in the detection of absorbing aerosols above the cloud. An analysis using the OMMYCLD product over the southeastern Atlantic Ocean for the period of Jun-July-Aug 2007 revealed a well-constrained non-linear relationship between the MODIS-derived COD times the geometric cloud fraction and LER388. A threshold of LER388 of 0.25 adopted for the

best quality retrievals compares to the COD times the geometric cloud fraction of 3-4. Conversely, given the geometric cloud fraction of unity, the minimum COD retrieved by OMACA would be in the range 3-4.

5 Lower magnitudes in both parameters might result in lesser confidence in the detection of either overcast pixels (0.20<LER<0.25, quality flag=1) or the presence of absorbing aerosols above cloud (0.8<UVAI<1.3, quality flag=2). Lower LER values (0.20-0.25) might pose a risk of identifying broken clouds in the OMI pixels, resulting in a geometric cloud fraction lesser than unity-a condition under which the assumption of fully overcast pixel breaks down. Nevertheless, it is also possible that the increased aerosol loading (AOD>2) with a significant absorption capacity (SSA<0.90) above the fully overcast pixels reduces LER measured at TOA (Figure 6 of *Jethva et al.*,
10 2013). On the other hand, the lower values of UVAI (0.8-1.3) associated with the quality flag '2' may be related to the non-aerosol related artifacts resulting from the inherent uncertainties in the derivation of UVAI. The sources of uncertainties include spectral surface albedo, the unaccounted presence of ice clouds, and viewing geometry of the Sun and satellite. The magnitudes of UVAI depend on several aerosol parameters including ACAOD, COD, SSA, ALH, and spectral dependence of aerosol absorption. Radiative transfer calculations show that for a given value of
15 SSA of 0.90 (388 nm) with an ALH of 3, 4, and 5 km, the UVAI value of 1.3 can be equated to the AOD (388 nm) of 0.30, 0.28, and 0.26, respectively. For a given SSA of 0.84, the values of AOD are 0.22, 0.20, and 0.19. The results of these simulations presented in Table 2 suggest that the minimum value of AOD retrieved using the thresholds in UVAI depends on the actual condition of the scene.

20 Retrievals assigned with the algorithm quality flag '3' are considered to be the lowest in confidence as it represents spurious non-aerosol related enhancement in UVAI (up to 2.0) at certain extreme geometries. However, if the observed UVAI exceeds a value of 2.0, then the retrievals are assigned with the quality flag '0' or '1' or '2' depending upon the observed LER and UVAI. The OMACA algorithm operates over both ocean and land pixels with terrain pressure greater than 800 hPa, which encompasses the majority of the regions of frequent aerosols-cloud
25 overlap (see section 5). Retrievals over oceanic cloud pixels are performed at all Sun glint angles if measured LER exceeds 0.30; for the 0.20<LER<0.30 condition retrievals are performed with pixels having Sun glint angle>20° to avoid glint-related artifacts in the retrievals.

3. Uncertainty Estimates

30 The OMACA algorithm relies on the presumptions about the atmosphere and surface properties. It is, therefore, imperative to estimate the sensitive of the OMACA retrievals to the departure of the actual state of the atmosphere from the one assumed in the inversion. Earlier, Torres et al., (2012) described the theoretical uncertainties in the near-UV based retrieval of ACAOD and aerosol-corrected COD. The analysis was, however, confined to a narrow
35 range of input conditions. Here, we re-perform the same exercise by considering an extended range of perturbation in each assumed parameter.

Deleted: s

Deleted: restrictive

The approach to calculating the uncertainties in the OMACA retrievals follows a perturbation-based method. The errors were calculated by contrasting the retrievals derived assuming a reference state and perturbed state of a particular input parameter. For example, given a fixed set of aerosol size distribution, ALH, and surface albedo, OMACA retrievals are derived assuming a range of aerosol SSA. The retrievals then compared with those derived considering a reference value of SSA, which is the center value of the prescribed range. The errors in the retrievals then can be calculated given both underestimated and overestimated values of an assumed parameter.

Table 3 lists the percent error in ACAOD (388 nm) caused by varying uncertainty in the aerosol SSA for an above-cloud smoke situation. The reference value of SSA (388 nm) was assumed to be 0.89; ALH and surface albedo were referenced at 3.0 km and 0.05, respectively. Errors were calculated for a range of uncertain SSA, i.e., -0.05 (underestimation) to 0.05 (overestimation) in step of 0.01. The optical depth of cloud underneath the aerosol layer was assumed to be 10. The errors in ACAOD are found to behave non-linearly to the perturbations in the assumed SSA. It also depends on the true value of ACAOD. Furthermore, the error conforms to an asymmetric behavior around the reference value of SSA; larger magnitudes of error are associated with the overestimated SSAs, whereas relatively lower errors are obtained when SSA was underestimated. The selection of above-cloud SSA values in actual OMACA retrievals relies on a daily, regional database of cloud-free SSA values retrieved from the standard OMAERUV aerosol product, as described in section 2.2.2.3. The accurateness of assigned above-cloud SSA, therefore, depends on the accuracy of cloud-free OMAERUV SSA retrievals as well as on the validity of the assumption that aerosol absorption properties are invariant between cloud-free and above-cloud aerosols scenes.

Earlier, an intercomparison analysis of OMI-AERONET SSA retrievals, based on previous public version of the OMAERUV product, from over 269 AERONET sites distributed globally showed an agreement within ± 0.03 and ± 0.05 limits for about 51% and 76% of total 5463 collocated matchups (Jethva et al., 2014). When segregated by the range of AOD (440 nm) and UVAI, 49% (AOD $<$ 0.7, UVAI $<$ 1.0) and 53% (AOD $>$ 0.7, UVAI $>$ 1.0) of the total OMAERUV-AERONET SSA (440 nm) retrievals are found to agree within their estimated uncertainties of ± 0.03 . The agreement improved to 74% and 79%, respectively, when the uncertainty limit was relaxed up to ± 0.05 . The statistical comparison was found even better when the matchups were segregated by the aerosol type, i.e., only smoke or dust, over many long-term sites located in the biomass burning and dust dominated regions.

Since its public release in 2012, the OMAERUV algorithm has been upgraded with several major changes, including better treatment of dust particles assuming realistic spheroidal shape distribution, accounting for angular scattering effects of clouds in the calculation of UVAI, use of new minimum surface LER dataset using synergy of multi-year OMI and MODIS observations, and updated cloud screening and retrieval flagging scheme. The regional, daily SSA dataset used in the OMACA product has been derived from this latest version of the OMAERUV (version 1.8.9.1) product released last year in 2017. A new comparative analysis of OMI-AERONET SSAs over global AERONET locations demonstrated that the agreement between the two independent sets of retrievals improved significantly for scenes identified as smoke and dust aerosol type. Quantitatively, about 59% (83%), 65% (88%), and 72% (91%) of

Deleted: A

the matchups are found to be within the expected limits of ± 0.03 (± 0.05) difference given the observed range of $UVAI > 0.8, > 1.5,$ and $> 2.0,$ respectively. The use of UVAI-weighted scheme to estimate daily regional value of SSA precisely reflects the fact that the agreement between OMI and AERONET SSA improves at higher aerosol loading/absorption providing increased confidence in the satellite retrievals.

5

The AERONET-OMI SSA comparative analysis cannot be treated as an actual “validation” of the satellite retrievals as both datasets are essentially derived from the two different platforms using fundamentally different algorithms that rely on a set of assumptions. However, a good-level of agreement between the two SSA datasets at higher UVAI established a consistency and provided an increased level of confidence in the OMAERUV absorption retrievals in cloud-free areas. Table 3 shows that uncertainty in the assumed SSA of $+0.03$ (-0.03) leads to an error in the retrieved ACAOD by $+42\%$ to $+46\%$ (-20% to -25%). The estimated errors are much larger (90%-100%) given the larger uncertainty (± 0.05) in the assumed SSA.

10

Deleted: robust

Deleted: larger

Deleted: AOD (> 0.7) and

Deleted: (> 1.0)

Deleted: Therefore, we expect that the above-cloud SSA values assigned in the OMACA algorithm over different regions should be accurate within ± 0.03 .

Relative to the errors in ACAOD due to the uncertain SSA, departures of ALH from the assumed state results in lower errors (Table 4). For assigning the ALH, OMACA relies on a global, monthly mean dataset derived from the 30-month coincidence and collocated CALIOP and OMI observations (Torres *et al.*, 2013). The expected uncertainty in the derived ALH dataset is about ± 1 km for which the error in ACAOD could vary between -5% to -13% and $+8\%$ to $+21\%$ for an overestimated and underestimated ALH by $+1$ km and -1 km, respectively.

15

The OMACA 2D retrieval domain shown in Figure 1 suggests that the retrieved value of ACAOD is primarily modulated by the magnitudes of UVAI. Several parameters including ACAOD, COD, ALH, SSA, and AAE can influence the magnitudes of UVAI. For instance, given a fixed set of spectral AODs at 354 and 388 nm wavelengths, ALH, and SSA at 388 nm, the magnitude of derived UVAI strongly varies with the assumed value of AAE (Figure 4 of Jethva and Torres, 2011). In other words, for a given value of observed UVAI, different assumptions of AAE would result in different values of the retrieved AOD. The “smoke” and “dust” aerosol models adopted in the OMACA algorithm assume an invariant spectral dependence of the imaginary part of the refractive index. For the carbonaceous and dust aerosol models, the relative spectral dependence in the imaginary index is assumed to be $+20\%$ (Kirchstetter *et al.*, 2004; Jethva and Torres, 2011) and 39% , respectively, between the 354 and 388 nm wavelengths. For a fixed set of size distribution parameters and range of SSA (388 nm), this results in AAE in the range 2.5-3.0 for the carbonaceous aerosol models, and 2.0-4.0 for the dust models (see Appendix I). Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of aerosol AAE (354-388 nm range) is listed in Table 5. The reference value of AAE was assumed to be 2.65 corresponding to the moderately absorbing smoke model, and COD underneath the aerosol layer was assumed to be 10. AAE was perturbed in steps of 0.5 in both directions from the reference value. Similar to the uncertain SSA simulations, errors in ACAOD behaves asymmetrically to the perturbations in AAE with larger (relatively lower) magnitudes of errors are associated with the underestimated (overestimated) AAE.

20

25

30

35

5 The corresponding simulated errors in the retrieval of aerosol-corrected COD resulting from uncertain SSA and ALH are shown in Table 6. The overall errors in COD are found to be much smaller compared those in the ACAOD retrievals. An uncertainty of ± 0.03 in SSA produces an error in COD in the range -2% to 8% depending on the absolute values ACAOD and COD. Similarly, an uncertain assumption of ALH yields an error in COD between -5% to 11%. In addition to the assumptions on aerosol parameters, the LUT calculations also are based on the assumed C1 liquid water cloud model with a fixed value of droplet effective radius (Deirmendjian, 1969). A sensitivity analysis, similar to the ones presented above, was carried out following the perturbation approach around the assumed CRE value of 12.0 μm . Table 7 lists the errors in aerosol-corrected COD due to the range of uncertainty in the assumed cloud effective radius. The analysis was performed assuming a reference effective radius of 12 μm and for the ACAOD of 0.5 and 1.0 (388 nm).

10 The errors in COD retrievals due to the uncertainty in effective radius follow asymmetric behavior to the perturbation around the assumed state. While a large underestimation of -8 μm produces negative errors of $\sim 10\%$ -11% in the retrieved COD, an overestimation of +8 to +12 μm yields positive errors of much smaller magnitudes ($\sim 1\%$ -2%). The spatial distribution of MODIS monthly cloud CRE over the Southeastern Atlantic Ocean, as shown in Figure 11 of Meyer *et al.* (2015), exhibits spatial variations with smaller droplets (CRE 7-11 μm) concentrated closer to the coast and relatively larger size droplets (11-17 μm) retrieved away from the coast. Given the fixed value of CRE equals 6.0 μm assumed in the OMACA cloud LUTs, the observed variations from MODIS would produce $< 2\%$ error in the retrieved aerosol-corrected COD. The corresponding errors in ACAOD due to the uncertainty in cloud effective radius are found to be marginal. For an ACAOD > 0.5 , an uncertainty in cloud effective radius by ± 8 μm results in ACAOD errors $< 2\%$ with much smaller magnitudes at higher aerosol loading. This is because at larger ACAODs the aerosol absorption effects dominate over that produced by varying effective radius of liquid droplets leaving other major algorithmic assumptions, i.e., SSA, ALH, and AAE to determine the resultant uncertainty in ACAOD retrievals.

25 This analysis implies that near-UV wavelengths don't offer a strong sensitivity to the variations in cloud droplet size rather the cloud signal is predominantly driven by the optical depth of cloud layer. Due to the lack of information on cloud droplet size from OMI, we adopted the standard C1 cloud model validated and used in numerous studies for all cloud LUT calculations.

30 The sensitivity of ACAOD and aerosol-corrected retrievals to the uncertainty in three major assumptions made in OMACA considered a broad range of perturbation. However, we anticipate that the prescribed values of SSA, ALH, and AAE are accurate to within ± 0.03 , ± 1 km, and ± 0.5 , respectively, for which the errors in ACAOD can vary from -23% to +46% in the ACAOD range 0.25-1.0. The corresponding errors in aerosol-corrected COD are found to be significantly lower and in the range -5% to +11%. In situations when the uncertainty in the assumed inputs leads to errors of opposite sign, the resulting error in the retrievals is likely to be lower than expected due to the cancellation

Deleted: the

of individual errors. On the other hand, an agglomeration of the errors of the same sign can further amplify the overall uncertainty in the retrievals. Nevertheless, it is practically hard to arrive at the actual uncertainty in the OMACA retrieval for every pixel due to the ill-posed nature of the inversion problem. Evaluating the accuracy of the satellite retrievals requires an independent set of direct measurements of aerosols, in this case above the cloud, discussed in the following section.

4. Preliminary Validation

Unlike the validation exercise of satellite-based aerosol retrievals in cloud-free skies for which the ground-based direct measurements of AOD are amply available from hundreds of sites worldwide such, an assessment of ACAOD retrieved from the satellite is a challenging task due to the lack of such reference aerosol measurements above the clouds. This is because field campaigns in the past mainly focused on the measurements and characterization of aerosol properties in cloud-free conditions leaving vast cloudy areas unmonitored regarding the aerosol measurements. However, the airborne lidar such as High Spectral Resolution Lidar (HSRL) when flying above the top of the aerosol layer can make direct measurements of aerosol extinction and thus provides AOD above the cloud. Also, airborne sunphotometers can make such measurements by flying above the cloud and below the aerosol layer. Using a limited data set of the direct measurements of AOD above the cloud carried out by the NASA Ames Airborne Tracking Sunphotometer (AATS) and Sky-Scanning, Sun-Tracking Atmospheric Research (4STAR) sensors during different field campaigns, *Jethva et al.* (2016) have validated ACAOD retrieved using the ‘color ratio’ method (*Jethva et al.*, 2013), similar to the one presented here, but applied to the TOA visible-near Infrared reflectances measured by the MODIS.

NASA’s ORACLES-Observations of Aerosols above Clouds and their interactions (<https://espo.nasa.gov/oracles>) is an ongoing multi-year field experiment supported by the NASA Earth-Venture Suborbital Program. ORACLES intended to make accurate airborne remote sensing and in situ measurements of aerosols and clouds in the Southeastern Atlantic Ocean. At the time writing this paper, ORACLES has already completed the two phases of its operation, 1st phase conducted in August-September 2016 and the 2nd phase in August 2017. During the 1st phase, HSRL-2 lidar developed by NASA Langley Research Center made extensive measurements of smoke aerosols, including detailed vertical measurements of aerosol backscatter, extinction, and AOD at 355 and 532 nm, above shallow marine clouds while deployed from NASA ER-2 aircraft. HSRL-2 measurements of AOD are routinely compared with that of AERONET and found to agree well ($R=0.98$) with the latter. Taking advantage of highly accurate and valuable dataset of AOD above the cloud, we evaluate the ACAOD retrievals from OMACA for the cases of coincident and collocated OMI-HSRL2 measurements.

Figure 4 shows the comparison of spectral ACAODs measured by HSRL-2 and retrieved from OMI/OMACA for a total of seven ER-2 flights conducted during August-September of 2016 (August 26, September 12, 16, 18, 20, 22, 24). HSRL-2 measurements falling within the boundaries of each OMI pixel, as defined in the OMPICOR product,

were averaged and compared with the ACAOD value of the corresponding OMI pixel. The spatially collocated aircraft-satellite matchups were grouped according to the three different time windows, i.e., no time constraints (in hours), and ΔT of ± 2 hours and ± 1 hours, where ΔT is the absolute time difference (in hours) between the OMI overpass and HSRL-2 measurements. To facilitate the direct comparison, ACAODs from HSRL-2 were interpolated to the OMI wavelengths of 388 nm and 500 nm following the Angstrom Exponent calculated using the 355-532 nm measurements. For the 'no time constraint' group, the collocation procedure yields more than 500 matchup data points with a correlation, RMSE, mean bias of 0.676, 0.23, and -0.11, respectively, at 388 nm. The comparison, however, significantly improves when the matchups are restricted to the narrower time windows of ΔT of ± 2 and ± 1 hour. For the $\Delta T = \pm 1$ -hour matchup group, the comparison yields correlation, RMSE, and mean bias of 0.77, 0.1, and 0.007 with the slope and intercept of 0.6 and 0.19 of the linear regression. Aerosol mass is in constant motion depending on wind speed and direction and therefore, allowing a wider time window between the satellite and aircraft measurements may end up in a mismatch which appears to be the case for relatively poorer comparison when no time limits were imposed on the comparison. Noticeably, the OMI-HSRL2 comparison of ACAOD provides the best agreement at the 500-nm wavelength, where the OMACA does not perform inversion but reports ACAOD based on the spectral dependence of extinction assumed in the aerosol model.

The remaining discrepancies in the comparison could be primarily attributed to the inherent uncertainties associated with both types of measurements, particularly in the satellite retrievals of ACAOD as discussed in the previous section, and spatiotemporal heterogeneity in aerosol fields unresolved by the collocation method. Despite these uncertainties, a reasonable agreement of OMACA-retrieved ACAOD with more accurate measurements from HSRL-2 for the ORACLES campaign provided the credibility and confidence in the product and allowed us to use it for the regional and global analyses presented in the rest of the paper. [While the primary focus of the paper is to highlight the ACAOD product with its spatial-temporal distribution and initial validation against ORACLES/HSRL-2, we are working with the ORACLES team to perform a detailed validation of the OMACA product using airborne in situ and remote sensing measurements. The results of the validation analysis will be covered in a dedicated follow up publication.](#)

5. The frequency of Occurrence of Absorbing Aerosols above Clouds

5.1 Spatial Distribution

The regional and global climate impact of absorbing aerosols above cloud depends on the total aerosol loading above the cloud, the microphysical and optical properties of aerosols and underlying cloud deck as well as the spatial and temporal extent of the aerosol-cloud overlap scenes. We have carried out a global frequency of occurrence of absorbing aerosols above the cloud (FOACA) to identify the regions of frequent aerosols-cloud overlap. We adopt a bi-parametric approach to identify the scenes of absorbing aerosols overlaying the low-level cloud decks, as described in section 2.2.3. FOACA is referenced to the cloudy-sky observations and defined as the ratio of the total number of days with an ACA condition ($LER > 0.25$ & $UVAI > 0.8$) to the total number of days with cloudy-condition ($LER > 0.25$). Additionally, we take advantage of the OMMYDCLD product to calculate the

geometric cloud fraction for each qualifying OMI pixel. For the FOACA analysis, we adopt a less strict threshold of cloud fraction of 0.5. Since the main purpose of this analysis was to identify the presence of absorbing aerosols and cloud in the same atmospheric column and not to quantitatively retrieve ACAOD, a less strict value cloud fraction should adequately represent both aerosols and clouds in the corresponding pixel.

5

Figure 5 shows the monthly averages of FOACA derived from the 12-year record (2005-2016) of OMI following the above-described method. This analysis reveals several important regions of the world where the overlap of absorbing aerosols over clouds are frequently observed. During July through September, carbonaceous aerosols generated from agricultural burning over the central/southern Africa are mobilized over the semi-permanent low-level stratocumulus water clouds in the southeastern Atlantic Ocean (Torres *et al.*, 2012; Alfaro-Contreras *et al.*, 2015; Meyer *et al.*, 2015). With more than 80% of the cloudy-sky observations are identified as aerosols above cloud during the northern hemisphere summer, the southeastern Atlantic Ocean is considered to be one of the prime regions the world and also a natural laboratory to study the aerosols above cloud phenomenon. The springtime biomass burning activities such as burning of forest, savanna/grassland, and crop residue over Southeast Asia (SEA) countries, including Thailand, Myanmar, Laos, Cambodia, and Vietnam release significant amount of trace gases and carbonaceous aerosols in the atmosphere (Elvidge and Baugh, 1996; Streets *et al.*, 2003). Natural color images from satellite show that smoke particles emitted from these activities were mobilized under the influence of winds over the widespread cloud deck over southern China, creating the appearance of the cloud deck darker. FOACA results show that about 20%-40% of the cloudy days are marked with smoke aerosols overlying bright cloud deck. Also, the smoke-cloud overlap seen in the OMI data is not merely confined to over land, but also extended over the western Pacific Ocean albeit less often.

Dust storms originated over the Sahara Desert in North Africa during the summer are often transported across the tropical Atlantic Ocean (Prospero *et al.*, 2002; Kaufman *et al.*, 2005; Huang *et al.*, 2010; Yu *et al.*, 2015). A substantial part of the dust transport occurs over the low-level stratocumulus clouds. The FOACA analysis shows that the presence of mineral dust aerosols above the cloud accounts for 40%-80% of the total cloudy-sky OMI observations. Similarly, mineral dust particles originated from the Arabian Peninsula, Iran and Pakistan often mobilized over the Arabian Sea during the active months (July-Aug) of southwest Indian summer monsoon (Moorthy *et al.*, 2005; Satheesh *et al.*, 2006). The transported dust frequently overlies the low-level clouds such that the dust-cloud overlap accounts for the 40%-60% of the cloud-sky days.

The springtime Asian outflow of air pollutants, both anthropogenic and mineral dust, across the Pacific Ocean is documented in various studies (Liu *et al.*, 2003; Huebert *et al.*, 2003). The FOACA maps for the April and May months reveals that the transport pattern encompasses the entire northern Pacific basin from the eastern coast of Asia to the western coast of North America with 10%-30% of the cloudy-sky scenes identified as aerosols over the clouds. One of the salient features of the FOACA analysis is the smoke transport from southern Africa to the Indian Ocean. Carbonaceous aerosols emitted from the fires in southeastern Africa during the late biomass burning season

35

(September), are often taken away from the continent along the eastward flow and advected above the clouds over the southern Indian Ocean. The FOACA map for the month of September highlights that the transport of smoke over the cloud is confined to within the latitude range 35°S to 20°S with 40%-50% (20%-30%) cloudy scenes marked by the smoke overlaying clouds just off the coast of southeastern Africa (over the western coast of Australia).

Prior to our study, Devsthale and Thomas (2011) Zhang et al. (2016) have studied the global cloud-sky FOACA over ocean using 4 years and 8 years of CALIOP observations, respectively. The results presented in these papers are largely consistent with the ones derived from OMI observations over oceanic regions that are influenced by long-range transport of smoke and dust aerosols from continents as described above.

5.2 Long-term Trends in FOACA

Since mid-2007, OMI observations have been affected by a possible external obstruction that perturbs both the measured solar flux and Earth radiance. This obstruction affecting the quality of radiance at all wavelengths for a particular viewing direction is referred to as “row anomaly” since the viewing geometry is associated with the row numbers on the charge-coupled device detectors. The row anomaly issue was detected first time in mid-2007 for few rows which over the period of operation expanded to other rows in 2008 and later. Figure 6 (a) (top panel) shows the current status of the row anomaly as identified by the anomaly algorithm developed for the NASA OMI total ozone product OMTO3 (Schenkeveld et al., 2017). At present, about half of the total 60 rows across the track are identified and flagged as row anomaly affected positions for which no physical retrievals are being performed.

Earlier, it was learned and concluded that for the derivation of a meaningful trend in the global FOACA only those rows or positions of the OMI instrument should be considered that are mostly unaffected by the row anomaly throughout the OMI operation period (Jethva, H., 2015). This is because UVAI exhibits a dependency on the scan angle in which the westward viewing geometry (left side of the scan, row numbers 1-30) shows higher values than those measured for the eastward-looking geometry (right side of the scan, row number 31-60). The row-averaged UVAI for the OMI operation period shown in the middle panel of Figure 6 (b) reflects the asymmetry in UVAI. OMI lost its most rows on the right side of the scan post-2007 operation due to the row anomaly. The remaining rows on the left side of the scan, where the UVAI values are larger than those on the right side of the scan, introduces a positive shift in the absolute values of UVAI which leads to an overall positive trend in the FOACA. The global trend (%) in FOACA calculated using all rows of OMI, therefore, gives a positive trend (0.178 per year) as shown in the bottom panel of Figure 6 (c). However, when the global trend is calculated using observations from rows which are mostly free of row anomaly throughout the OMI operation period, i.e., row number 1-23, the FOACA shows a negligible trend (0.022 per year). Regionally, we find similar results of positive trends in FOACA when all rows of OMI were used in the calculations (not shown here) against the statistically negligible trend when observations from rows 1-23 were considered, as shown in Figure 7. Based on the present findings and also according to the results published in an interactive comment (Jethva, H., 2015), we strongly recommend the users of

OMACA product to use only those observations that are free of row anomaly throughout the OMI operation (2004 to present) for the trend related analysis.

6. Results from 12-year Long OMACA Record

6.1 Global Distribution of Above-cloud AOD

5 Figure 8 shows the global distribution of ACAOD (388 nm) derived from OMACA algorithm applied to the 12-year long record of OMI observations. The OMACA Level 2 retrievals with algorithm quality flag '0', '1', and '2' were averaged on every $0.5^\circ \times 0.5^\circ$ spatial grid to derive global monthly gridded dataset. Also, a threshold of 0.75 in the geometric cloud fraction calculated using the OMMYDCLD product was used to filter out the L2 pixels with broken cloud fields. Only those grids are considered in the long-term monthly averages for which 1) the total number of
10 days with valid retrievals for a particular month is greater than 3, and 2) number of years greater than 3 out of the 12-year record. The distribution plot reveals moderate to high aerosol loading above the clouds over several regions of the world.

During the northern hemisphere summer, larger magnitudes of ACAOD (>0.7) are retrieved over the Southeastern
15 Atlantic Ocean along the coast of Namibia and Angola. Noticeably, the area coverage of ACAOD expands substantially as the season progresses with retrieval coverage confined to within 1500 km from the western coast of Africa in June to encompassing almost the entire Atlantic Ocean basin (~ 5000 km) in September. Largest magnitudes of ACAOD are observed in August and September when the biomass burning activities also peak in the central/southern Africa. In March and April, biomass burning in Southeast Asia emit large amounts of carbonaceous
20 aerosols, which under the influence of winds are transported above the bright cloud deck over southern China, where OMACA retrieves large values of ACAOD (>0.8) in the vicinity of burning areas. Noticeably the area-coverage of aerosol-cloud overlap extends far from the source burning region to over the East China Sea albeit with a decreasing gradient in the retrieved ACAOD. During the late biomass burning season (September), carbonaceous aerosols emitted from the fires in southeastern Africa often drift away from the continent along the eastward flow (*Garstang
25 et al., 1996*), and advected above the clouds over the southern Indian Ocean. The spatial pattern of retrieved ACAOD encompasses the entire Indian ocean basin stretching from the southeastern coast of Africa with ACAOD in the range 0.4-0.5 to the western coast of Australia with reduced ACAOD in the range 0.2-0.3.

During the northern hemisphere summer months, ACAOD in the range 0.3-0.5 is observed over the tropical Atlantic
30 Ocean where the transport of dust takes place from the Saharan desert to over the oceanic clouds. The area coverage of the retrievals is maximum in July spanning half of the tropical Atlantic Ocean basin with maximum ACAOD (~ 0.5) just off the coast of northern Africa. Also, a gradient in ACAOD is noted over the northern Arabian Sea during summer, owing to the dust transport from the Arabian region to over the low-level clouds over the ocean.

35 OMACA product also captures springtime (April and May), long-range trans-Pacific transport of dust aerosols originated over the Gobi and Taklamakan Deserts possibly mixed with urban pollution and smoke along the

transport pathways and over the clouds. The magnitude of retrieved ACAOD of the above-cloud Asian outflow ranges from 0.4-0.5 near the eastern coast of North-East Asia, reducing to 0.2-0.3 along the transport over the mid-Pacific and up to the western coast of North America.

5 6.2 Regional Time-series of Above-cloud AOD

Figure 9 (left y-axis, color: red) shows the regional, monthly mean time-series of ACAOD (388 nm) for the five prominent regions of the world where the frequent overlap of absorbing aerosols above the cloud is observed. The regional monthly mean ACAODs were calculated in the following way. For each region and month, an averaged value of ACAOD was calculated, and a number of Level 2 observations that went into the averaging was also stored. Subsequently, a set of 12 monthly-averaged values scaled by multiplying them with a fraction which is defined as the ratio Level 2 observations for the individual months to the maximum number of Level 2 observations found over the 12-year period (2005-2016) over the same region. Scaling the monthly-averaged value with the calculated fraction ensures the representativeness of ACAOD statistics over the spatial and temporal domains and therefore facilitates the intercomparison. For instance, a comparison of the two averaged values derived from two different set of statistics, i.e., under-populated and adequately-populated, likely results in the misinterpretation of monthly time-series data. The procedure described above was applied to the OMACA observations for the five prominent aerosol-cloud overlap regions to produce time-series shown in Figure 9.

The temporal evolution of ACAOD (388 nm) over the Southeastern Atlantic Ocean exhibits a repetitive seasonal cycle with monthly mean values reaching up to 0.4 during the dry biomass burning season. Although an inter-annual variation is apparent in the time-series, e.g., lower and higher ACAOD during the burning season of 2012 and 2015, no significant trend is noticed over the 2005-2016 OMI record. Over the tropical Atlantic Ocean, the monthly ACAOD values fall in the range 0.2-0.3 during summer months when dust aerosols from northern Africa are mobilized over the low-level oceanic stratocumulus clouds. The inter-annual variations of ACAOD over the Arabian Sea are found to be significant with the monthly value of 0.4 during the first two years of the record (July and Aug of 2005 and 2006) followed by a drastic reduction in the aerosol loading above the cloud during the middle part of the record. The springtime biomass burning and resulting smoke aerosols above cloud over South-East Asia (4th panel) is evident in the time-series where the peak values of monthly ACAOD vary from 0.2 to 0.5 depending on the year.

It is important to quantify the fraction of the total columnar aerosol loading situated above the cloud for several applications. For instance, the cloud-free aerosol retrieval represents the aerosol in the entire vertical column while avoiding the cloudy-sky scenes. Therefore, the statistics of cloud-free retrievals over the regions with frequent aerosol-cloud overlap becomes restrictive, leading to the partially incomplete representation of aerosol properties on a regional and temporal scales. This kind of scenario affects the calculations of regional climatology, radiative forcing assessments, and aerosol transport. However, the availability of above-cloud (OMACA) and cloud-free total columnar (OMAERUV) AODs from OMI allows us to estimate the fractional aerosol loading above the cloud. The

ratio of monthly mean ACAOD to the total columnar AOD (both at 388 nm) displayed on the right-side y-axis of Figure 9 shows that the fractional aerosol loading over the cloud can be as large as 80%-100% during the peak months of biomass burning and dust transport over the respective regions. A significant fraction of the aerosol column above clouds indicates that the long-range transport of partially absorbing aerosols occurs in the free troposphere and over the low-level clouds. Although the above-cloud and total column AOD are comparable in magnitude during peak aerosol activities,

6.3 Regional Time-series of Aerosol-corrected Cloud Optical Depth

Figure 10 displays the monthly mean evolution of aerosol-corrected COD (left y-axis, color: red) for the five prominent regions of aerosol-cloud overlap. The monthly mean values were calculated following the procedure described in the previous section. The seasonal cycle of COD over the Southeastern Atlantic Ocean exhibits a repetitive behavior with a maximum value of COD reaching up to 8.0 during the peak burning period. On the other hand, the monthly cycle of COD over the tropical Atlantic Ocean (Arabian Sea) during the same season shows more considerable interannual variations with COD in the range 6 to 14 (4 to 10). The magnitudes of aerosol-corrected COD over South-East Asia during the springtime biomass burning season (March-April) are found to be the largest among five regions considered here with values ranging between 16-20 except for the years 2008 and 2013 when COD was lesser than 12.

The right-side y-axis of Figure 10 (color: blue) depicts the concurrent monthly time-series of the percent difference between apparent (non-corrected) and aerosol-corrected CODs the magnitudes of which represent the error in the retrieval of COD at 388-nm wavelength with reference to the corrected COD if the presence of aerosols is ignored in the inversion. Note that only those COD retrievals are considered here for which the absorbing aerosols are identified above the clouds. While there is a significant variation in the magnitudes of the % difference between the two CODs over different regions, generally larger errors are associated with the higher aerosol-corrected COD and ACAODs (shown in Figure 9). More discussion on the impact of aerosol absorption on cloud retrievals is presented in the next section.

6.4 Impact of Aerosol Absorption on Cloud Retrievals

The presence of absorbing aerosols above cloud obstructs the light reflected by the cloud top, and thus reduces cloud-reflected upwelling UV (*Torres et al., 2012*), VIS, and NIR radiation (*Jethva et al., 2013; Meyer et al., 2015*) reaching the TOA. Therefore, cloud retrievals of COD derived from passive sensors such as OMI are expected to be biased low if absorbing aerosols are not accounted for in the inversion. OMACA product reports two sets of COD, one corrected for the presence of absorbing aerosols overlying cloud deck, and one retrieved assuming no aerosols above the cloud which is termed as the apparent COD. The magnitudes of bias in the apparent COD depend on the strength of aerosol absorption and backscattering as well as on the actual value of COD. Note that OMACA does not directly retrieve the aerosol absorption optical depth (AAOD) but retrieves ACAOD assuming an a priori value of SSA (see section 2.2.2.3). Therefore, the AAOD can be readily calculated using these two pieces of information as $AAOD = ACAOD * (1 - SSA)$. Figure 11 shows the percent bias in COD (388 nm), defined as (Apparent COD –

Aerosol- corrected COD)/Aerosol-corrected COD *100, as a function of concurrent AAOD (388 nm). The percent bias was calculated for the distinct range of aerosol-corrected COD and the two aerosol types, i.e., smoke (left panel) and dust (right panel). All OMACA Level 2 orbital data (2005-2016) for the respective regions and the two aerosol types were accumulated separately and subsequently averaged as a function of corresponding AAOD bins of a sampling size of 5000 retrievals.

For both aerosol types, increasing the magnitude of negative bias in the retrieval of apparent COD is related to AAOD suggesting the impact of aerosol absorption on the retrieved COD when the presence of absorbing aerosols is ignored in the inversion. Retrievals identified with the 'smoke' aerosol type, predominantly found over the biomass burning regions of Southeastern Atlantic Ocean and South-East Asia show a much larger range of AAOD and associated bias in COD than that observed with 'dust' aerosols found in the regions of dust transport over the tropical Atlantic Ocean and the Arabian Sea. Noticeably, the magnitudes of negative bias in the cloud retrievals are also co-dependent on the absolute values of COD (here it is aerosol-corrected COD). For instance, for an AAOD of 0.1, the bias in COD is ~ -25% at a lower range of COD (5-10), which gets twofold in magnitude (~ -50%) at the higher range of COD (20-50).

7. Summary and Concluding Remarks

We have developed a global above-cloud aerosol algorithm, formally named as OMACA (OMI Above-cloud Aerosols), to simultaneously derive the columnar optical depth of absorbing aerosols above the cloud and underlying aerosol-corrected cloud optical depth from the near-UV observations made by Aura/OMI. Physically based on the enhanced spectral contrast in the near- UV region (354-388 nm) caused by aerosol absorption above the cloud, OMACA relates the TOA observations in the two channels to a pair of ACAOD and COD under a prescribed set of assumptions. The architect of the OMACA algorithm in terms of the ancillary datasets (CALIOP-OMI based ALH, OMI-based near-UV surface albedo, and use of AIRS CO for the aerosol type identification), aerosol models (smoke and dust), and retrieval approach (two-channel inversion) is analogous to the OMI's two-channel, cloud-free OMAERUV aerosol algorithm. OMACA was applied to the entire record of OMI observations (Oct 2004 to present) to deduce a global research product of AOD above the cloud. Currently, the Level 2 orbital data product is stored on a freely accessible Aura Validation Data Center webpage (<https://avdc.gsfc.nasa.gov/pub/data/satellite/Aura/OMI/V03/L2/OMACA/>). Also, the OMACA product is produced in the forward processing mode with a maximum latency of about three days, which is associated with the availability of AIRS L3 CO data for the aerosol type identification.

An analysis of the frequency of occurrence of the above-cloud absorbing aerosols reveals several important regions of the world where the overlap of absorbing aerosols and cloud are frequently observed on a monthly to seasonal scales. These regions include Southeastern Atlantic Ocean and Southeast Asia where layers of smoke aerosols produced from the seasonal agricultural biomass burning spread thousands of kilometers over the regional low-level stratocumulus cloud deck; the tropical Atlantic Ocean and the Arabian Sea where dust aerosols transported from Sahara and Arabian deserts, respectively, found over low-level clouds; the northern Pacific Ocean where dust

particles originated from Asian deserts possibly mixed with the pollution haze along the eastward transport pathways are found to overlie clouds; and the southern Indian Ocean where the smoke produced from agricultural burning over southeastern Africa drifts along the easterly winds and overlie oceanic cloud deck. Globally as well as on a regional scale, no significant trend in the frequency of ACA was noted when only those observations of OMI instrument (Row # 1 to 23) that are free of row anomaly throughout the OMI operation period (2004 to present) were used in the calculation. The climatology maps of the retrieved ACAOD (388 nm) derived from a 12-year long OMACA record show moderate ($0.3 < \text{ACAOD} < 0.5$, away from the sources) to higher aerosol loading ($\text{ACAOD} > 0.8$ in the proximity to the sources) above the cloud over these prominent aerosols-cloud overlap regions. When compared with the cloud-free, columnar aerosol loading distribution retrieved from the OMI/OMAERUV two-channel algorithm, ACAOD accounts for as large as 60%-100% of the total columnar loading over different regions during peak biomass burning and dust transport seasons.

The aerosol-corrected CODs retrieved at the near-UV wavelength (388 nm) are found to be noticeably higher than those retrieved assuming no aerosols above the cloud. The percent bias in COD with reference to the aerosol-corrected COD is found to strongly correlate with AAOD as well as the magnitude of COD. For instance, carbonaceous aerosols above cloud found over the Southeastern Atlantic Ocean and South-East Asia during respective biomass burning seasons result in a significant negative bias in apparent COD the magnitudes of which increase with increasing aerosol absorption as well as the cloud brightness.

A direct comparison of coincident and collocated ACAODs derived from OMI/OMACA and that measured from airborne HSRL-2 measurements for the ORACLES phase I operation (August-September 2016) over the Southeastern Atlantic Ocean showed a good level of agreement with a correlation and RMSE of 0.7 and 0.1 respectively. We further plan to extend the validation of OMACA using the direct measurements of ACAOD from airborne 4STAR sunphotometer operated from NASA's P3-B aircraft during ORACLES phase I and II. Additionally, a suite of aerosol microphysical and optical measurements from the ORACLES campaign, particularly those characterizing spectral aerosol absorption, will help to verify and improve the region-specific aerosol models employed in OMACA.

Conventional aerosol remote sensing algorithms provide distribution of aerosols in the cloud-free areas leaving behind vast cloudy regions unmonitored regarding the co-presence of aerosols and clouds. OMACA aerosol product offers a quantitative characterization of aerosol loading above cloud on a global scale. Several observational and modeling studies have shown that an overlap of absorbing aerosols above cloud leads to significant atmospheric warming, which can affect cloud lifetime and hydrological cycle. The magnitudes of these effects depend upon the amount of aerosol loading above the cloud, cloud brightness (COD), optical and microphysical properties of aerosols and clouds, and cloud fraction. The OMACA aerosol product from OMI presented in this paper offers a crucial missing piece of information of the aerosol loading above cloud that will help us to quantify the radiative effects of aerosols above the cloud and its resultant impact on clouds and thus climate. A global above-cloud aerosol

product in conjunction with the standard cloud-free aerosol product provides us with an unprecedented all-sky aerosol distribution from space. This can substantially enhance our knowledge of how aerosols affect cloud radiative forcing and microphysical properties, and aerosol transport.

Acknowledgments

Authors acknowledge the support of OMI SIPS team (<https://earthdata.nasa.gov/about/sips/sips-omi>) for processing the OMACA dataset. We thank Dr. P. K. Bhartia at NASA Goddard Space Flight Center for his guidance and helpful suggestions throughout the development of the OMACA product. Peter Leonard at ADNET Systems Inc has extended help in coding the software package. Dr. Oleg Dubovik at the University of Lille1 provided a standalone software package for the simulation of phase matrix elements of spheroidal particle size distribution used to generate the dust aerosol look-up-table. Acknowledgements are due to the HSLR team members for conducting the measurements of aerosols above cloud during ORACLES phase I operation, which helped validate the OMACA product over the Southeastern Atlantic Ocean.

References

- Alfaro-Contreras, R., Zhang, J., Campbell, J. R., and Reid, J. S.: Investigating the frequency and interannual variability in global above-cloud aerosol characteristics with CALIOP and OMI, *Atmos. Chem. Phys.*, 16, 47-69, <https://doi.org/10.5194/acp-16-47-2016>, 2016.
- 5 Liu, H., Jacob, D. J., Bey, I., Yantosca, R. M., Duncan, B. N., and Sachse, G. W.: Transport pathways for Asian pollution outflow over the Pacific: Interannual and seasonal variations, *J. Geophys. Res.*, 108(D20), 8786, doi:10.1029/2002JD003102, 2003.
- Chand, D., Anderson, T. L., Wood, R., Charlson, R. J., Hu, Y., Liu, Z., and Vaughan, M.: Quantifying above-cloud aerosol using spaceborne lidar for improved understanding of cloudy-sky direct climate forcing, *J. Geophys. Res.*, 113, D13206, doi:10.1029/2007JD009433, 2008.
- 10 Chand, D., Wood, R., Anderson, T. L., Satheesh, S. K., and Charlson, R. J.: Satellite-derived direct radiative effect of aerosols dependent on cloud cover, *Nat. Geosci.*, 2, 181–184, doi:10.1038/NGEO437, 2009.
- Deirmendjian, D.: *Electromagnetic scattering on spherical polydispersions*, American Elsevier Publishing Company, Inc., New York, 1969.
- 15 [Devasthale, A. and Thomas, M. A.: A global survey of aerosol-liquid water cloud overlap based on four years of CALIPSO-CALIOP data, *Atmos. Chem. Phys.*, 11, 1143-1154, <https://doi.org/10.5194/acp-11-1143-2011>, 2011.](#)
- Eswaran, K., Satheesh, S. K. and Srinivasan, J.: Dependence of ‘critical cloud fraction’ on aerosol composition. *Atmos. Sci. Lett.*, 16: 380–385. doi:10.1002/asl2.571, 2015.
- Feng, N., and Christopher, S. A.: Measurement-based estimates of direct radiative effects of absorbing aerosols above clouds. *J. Geophys. Res. Atmos.*, 120, 6908–6921. doi: 10.1002/2015JD023252, 2015.
- 20 Garstang, M., Tyson, P. D., Swap, R., Edwards, M., Kallberg, P., and Lindsay, J. A.: Horizontal and vertical transport of air over southern Africa, *J. Geophys. Res.*, 101, 23,721–23,736, 1996
- Herman, J. R., Bhartia, P. K., Torres, O., Hsu, C., Seftor, C., Celarier, E.: Global distribution of UV- absorbing aerosols from Nimbus 7/TOMS data, *J. Geophys. Res.*, 102(D14), 16911–16922, doi:10.1029/96JD03680, 1997.
- 25 [Hu, Y., Vaughan, M., Liu, Z., Powell, K., Rodier S.: Retrieving optical depths and lidar ratios for transparent layers above opaque water clouds from CALIPSO lidar measurements, *IEEE Geosci. Remote Sens. Lett.*, 4\(4\), 523-526, doi:10.1109/LGRS.2007.901085, 2007.](#)
- Huebert, B., Bates, T., Russell, P., Seinfeld, J., Wang, M., Uematsu, M., and Kim, Y. J.: An overview of ACE-Asia: Strategies for quantifying the relationships between Asian aerosols and their climatic impacts, *J. Geophys. Res.*, 108(D23), 8633, doi:10.1029/2003JD003550, 2003
- 30 IPCC, 2013: *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp, doi:10.1017/CBO9781107415324.
- 35 Jethva, H. and Torres, O.: Satellite-based evidence of wavelength-dependent aerosol absorption in biomass burning smoke inferred from Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 11, 10541- 10551, <https://doi.org/10.5194/acp-11-10541-2011>, 2011.
- Jethva, H., Torres, O., Ahn, C.: Global assessment of OMI aerosol single-scattering albedo using ground- based AERONET inversion, *J. Geophys. Res. Atmos.*, 119, 9020-9040, doi:10.1002/2014JD021672, 2014.

- Jethva, H., Torres, O., Remer, L. A., and Bhartia, P. K.: A Color Ratio Method for Simultaneous Retrieval of Aerosol and Cloud Optical Thickness of Above-Cloud Absorbing Aerosols From Passive Sensors: Application to MODIS Measurements, *IEEE Transactions on Geoscience and Remote Sensing*, 51, 3862–3870, [10.1109/TGRS.2012.2230008](https://doi.org/10.1109/TGRS.2012.2230008), 2013.
- 5 Jethva, H., Torres, O., Waquet, F., Chand, D., and Hu, Y.: How do A-train sensors intercompare in the retrieval of above-cloud aerosol optical depth? A case study-based assessment, *Geophys. Res. Lett.*, 41, 186–192, [doi:10.1002/2013GL058405](https://doi.org/10.1002/2013GL058405), 2014.
- Jethva, H.: Interactive comment on “Investigating the frequency and trends in global above-cloud aerosol characteristics with CALIOP and OMI” by R. Alfaro-Contreras et al., *Atmos. Chem. Phys. Discuss.*, 15, C1488–C1496, www.atmos-chem-phys-discuss.net/15/C1488/2015/, 2015.
- 10 Jethva, H., Torres, O., Remer, L., Redemann, J., Livingston, J., Dunagan, S., Shinozuka, Y., Kacenelenbogen, M., Rosenheimer, M. S., and Spurr, R.: Validating MODIS above-cloud aerosol optical depth retrieved from “color ratio” algorithm using direct measurements made by NASA’s airborne AATS and 4STAR sensors, *Atmos. Meas. Tech.*, 9, 5053–5062, <https://doi.org/10.5194/amt-9-5053-2016>, 2016.
- 15 Kaufman, Y. J., Koren, I., Remer, L. A., Tanré, D., Ginoux, P., Fan, S.: Dust transport and deposition observed from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) spacecraft over the Atlantic Ocean, *J. Geophys. Res.*, 110, D10S12, [doi:10.1029/2003JD004436](https://doi.org/10.1029/2003JD004436), 2005.
- Keil, A. and Haywood, J. M.: Solar radiative forcing by biomass burning aerosol particles during SAFARI2000: A case study based on measured aerosol and cloud properties, *J. Geophys. Res.*, 108, 8467, [doi:10.1029/2002JD002315](https://doi.org/10.1029/2002JD002315), 2003.
- 20 Kirchstetter, T. W., Novakov, T., Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *J. Geophys. Res.*, 109, D21208, [doi:10.1029/2004JD004999](https://doi.org/10.1029/2004JD004999), 2004.
- [Lu, Z., Liu, X., Zhang, Z., Zhao, C., Meyer, K., Rajapakse, C., Wu, C., Yang, Z., and Penner, J. E.: Biomass smoke from southern Africa can significantly enhance the brightness of stratocumulus over the southeastern Atlantic Ocean, *Proc. Nat. Acad. Sci.*, 115 \(12\) 2924–2929, <https://doi.org/10.1073/pnas.1713703115>, 2018.](https://doi.org/10.1073/pnas.1713703115)
- 25 Meyer, K., Platnick, S., and Zhang, Z.: Simultaneously inferring above-cloud absorbing aerosol optical thickness and underlying liquid phase cloud optical and microphysical properties using MODIS, *J. Geophys. Res.-Atmos.*, 120, 5524–5547, [doi:10.1002/2015JD023128](https://doi.org/10.1002/2015JD023128), 2015.
- Meyer, K., Platnick, S., Oreopoulos, L., and Lee, D.: Estimating the direct radiative effect of absorbing aerosols overlying marine boundary layer clouds in the southeast Atlantic using MODIS and CALIOP, *J. Geophys. Res.-Atmos.*, 118, 4801–4815, [doi:10.1002/jgrd.50449](https://doi.org/10.1002/jgrd.50449), 2013.
- 30 [Min, M and Zhang, Z.: On the influence of cloud fraction diurnal cycle and sub-grid cloud optical thickness variability on all-sky direct aerosol radiative forcing, *J. Quant. Spectroscopy and Rad. Trans.*, 142, 25–36, <https://doi.org/10.1016/j.jqsrt.2014.03.014>, 2014.](https://doi.org/10.1016/j.jqsrt.2014.03.014)
- 35 Moorthy, K. K., Babu, S. S., and Satheesh, S. K.: Aerosol Characteristics and Radiative Impacts over the Arabian Sea during the Intermonsoon Season: Results from ARMEX Field Campaign. *J. Atmos. Sci.*, 62, 192–206, <https://doi.org/10.1175/JAS-3378.1>, 2005.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental Characterization Of Global Sources Of Atmospheric Soil Dust Identified With The Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) Absorbing Aerosol Product, *Rev. Geophys.*, 40(1), 1002, [doi:10.1029/2000rg000095](https://doi.org/10.1029/2000rg000095), 2002.
- 40

- Satheesh, S., Krishna Moorthy, K., Kaufman, Y., and Takemura, T.: Aerosol optical depth, physical properties and radiative forcing over the Arabian Sea, *Meteorol. Atmos. Phys.*, 91, doi: <https://doi.org/10.1007/s00703-004-0097-4>, 2006.
- 5 Schenkeveld, V. M. E., Jaross, G., Marchenko, S., Haffner, D., Kleipool, Q. L., Rozemeijer, N. C., Veeffkind, J. P., and Levelt, P. F.: In-flight performance of the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 10, 1957–1986, <https://doi.org/10.5194/amt-10-1957-2017>, 2017.
- 10 Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, <https://doi.org/10.5194/acp-6-5225-2006>, 2006.
- Sinha, P., Jaeglé, L., Hobbs, P. V., and Liang, Q.: Transport of biomass burning emissions from southern Africa, *J. Geophys. Res.*, 109, D20204, doi:10.1029/2004JD005044, 2004.
- 15 Spurr, R. J. D.: VLIDORT: A linearized pseudo-spherical vector discrete ordinate radiative transfer code for forward model and retrieval studies in multilayer multiple scattering media, *J. Quant. Spectrosc. Ra.*, 102, 316–342, doi:10.1016/j.jqsrt.2006.05.005, 2006.
- Streets, D. G., Yarber, K. F., Woo, J.-H., and Carmichael, G. R.: Biomass burning in Asia: Annual and seasonal estimates and atmospheric emissions, *Global Biogeochem. Cycles*, 17(4), 1099, doi:10.1029/2003GB002040, 2003.
- 20 Torres, O., Bhartia, P. K., Herman, J. R., Ahmad, Z., and Gleason, J.: Derivation of aerosol properties from satellite measurements of backscattered ultraviolet radiation: Theoretical basis, *J. Geophys. Res.*, 103(D14), 17099–17110, doi:10.1029/98JD00900, 1998.
- Torres, O., Tanskanen, A., Veihelmann, B., Ahn, C., Braak, R., Bhartia, P. K., Veeffkind, P., and Levelt, P.: Aerosols and surface UV products from Ozone Monitoring Instrument observations: An overview, *J. Geophys. Res.*, 112, D24S47, doi:10.1029/2007JD008809, 2007.
- 25 Torres, O., Jethva, H., and Bhartia, P. K.: Retrieval of Aerosol Optical Depth above Clouds from OMI Observations: Sensitivity Analysis and Case Studies, *J. Atmos. Sci.*, 69, 1037–1053, doi:10.1175/JAS-D-11-0130.1, 2012.
- Torres, O., Ahn, C., and Chen, Z.: Improvements to the OMI near-UV aerosol algorithm using A-train CALIOP and AIRS observations, *Atmos. Meas. Tech.*, 6, 3257–3270, <https://doi.org/10.5194/amt-6-3257-2013>, 2013.
- 30 Torres, O., Bhartia, P. K., Jethva, H., and Ahn, C.: Impact of the ozone monitoring instrument row anomaly on the long-term record of aerosol products, *Atmos. Meas. Tech.*, 11, 2701–2715, <https://doi.org/10.5194/amt-11-2701-2018>, 2018.
- Wagner, R., Ajtai, T., Kandler, K., Lieke, K., Linke, C., Müller, T., Schnaiter, M., and Vragel, M.: Complex refractive indices of Saharan dust samples at visible and near UV wavelengths: a laboratory study, *Atmos. Chem. Phys.*, 12, 2491–2512, <https://doi.org/10.5194/acp-12-2491-2012>, 2012.
- 35 [Waquet, F., Riedi, J., Labonnote, L. C., Goloub, P., Cairns, B., Deuzé, J.-L., and Tanré D.: Aerosol remote sensing over clouds using A-train observations, *J. Atmos. Sci.*, 66, 2468–2480, doi:10.1175/2009JAS3026.1, 2009.](#)
- [Waquet, F., Cornet, C., Deuzé, J.-L., Dubovik, O., Ducos, F., Goloub, P., Herman, M., Lapyonok, T., Labonnote, L. C., Riedi, J., Tanré, D., Thieuleux, F., and Vanbauce, C.: Retrieval of aerosol microphysical and optical properties above liquid clouds from POLDER/PARASOL polarization measurements, *Atmos. Meas. Tech.*, 6, 991–1016, <https://doi.org/10.5194/amt-6-991-2013>, 2013.](#)
- 40

Wilcox, E. M.: Direct and semi-direct radiative forcing of smoke aerosols over clouds, *Atmos. Chem. Phys.*, 12, 139-149, <https://doi.org/10.5194/acp-12-139-2012>, 2012.

5 Yu, H., Chin, M., Bian, H., Yuan, T., Prospero, J. M., Omar, A. H., Remer, L. A., Winker, D. M., Yang, Y., Zhang, Y., and Zhang, Z.: Quantification of trans-Atlantic dust transport from seven-year (2007–2013) record of CALIPSO lidar measurements, *Rem. Sens. Environ.*, 159, 232-249, doi: <https://doi.org/10.1016/j.rse.2014.12.010>, 2015.

[Zhang, Z., Meyer, K., Yu, H., Platnick, S., Colarco, P., Liu, Z., and Oreopoulos, L.: Shortwave direct radiative effects of above-cloud aerosols over global oceans derived from 8 years of CALIOP and MODIS observations, *Atmos. Chem. Phys.*, 16, 2877-2900, <https://doi.org/10.5194/acp-16-2877-2016>, 2016.](#)

Deleted: ¶

Appendix I

Optical and microphysical properties of OMACA Dust Aerosol Model

| Model# | Imaginary Index | | | Single-scattering Albedo | | | Absorption Angstrom Exponent | |
|--------|------------------|---------|---------|--------------------------|---------|---------|------------------------------|---------|
| | Wavelength in nm | | | Wavelength in nm | | | | |
| | 354 | 388 | 500 | 354 | 388 | 500 | 354-388 | 354-500 |
| 1 | 0.02303 | 0.01662 | 0.00720 | 0.74982 | 0.77921 | 0.86268 | 1.97901 | 2.34458 |
| 2 | 0.01279 | 0.00923 | 0.00400 | 0.80740 | 0.83778 | 0.91046 | 2.49312 | 2.82591 |
| 3 | 0.00832 | 0.00600 | 0.00260 | 0.84727 | 0.87606 | 0.93640 | 2.90115 | 3.14519 |
| 4 | 0.00561 | 0.00405 | 0.00176 | 0.88062 | 0.90532 | 0.95430 | 3.15286 | 3.38913 |
| 5 | 0.00256 | 0.00185 | 0.00080 | 0.93213 | 0.94886 | 0.97805 | 3.71332 | 3.87787 |
| 6 | 0.00128 | 0.00092 | 0.00040 | 0.96221 | 0.97234 | 0.98620 | 4.03071 | 3.52634 |
| 7 | 0.00000 | 0.00000 | 0.00000 | 1.00000 | 1.00000 | 1.00000 | N/A | N/A |

Real refractive index = 1.55
Mean Radius (fine mode) = 0.052 μm ; Mean Radius (coarse mode) = 0.67 μm
Standard Deviation (fine mode) = 1.697 μm ; Standard Deviation (coarse mode) = 1.806 μm

Minimum Radii (fine mode) = 0.00627012 μm ; Minimum Radii (coarse mode) = 0.0629802 μm
Maximum Radii (fine mode) = 0.431252 μm ; Maximum Radii (coarse mode) = 7.12764 μm

Axis ratio distribution for the spheroidal dust particles (Dubovik et al. 2006)

| Radius | Fractional Weight | Radius | Fractional Weight |
|---------|-------------------|---------|-------------------|
| 0.33490 | 0.0661850 | 1.09540 | 0.0000000 |
| 0.36690 | 0.0650250 | 1.20000 | 0.0000000 |
| 0.40190 | 0.0636350 | 1.31450 | 0.0000000 |
| 0.44030 | 0.0620500 | 1.44000 | 0.0403205 |
| 0.48230 | 0.0587200 | 1.57740 | 0.0429530 |
| 0.52830 | 0.0533500 | 1.72800 | 0.0477625 |
| 0.57870 | 0.0477625 | 1.89290 | 0.0533500 |
| 0.63390 | 0.0429530 | 2.07360 | 0.0587200 |
| 0.69440 | 0.0403205 | 2.27150 | 0.0620500 |
| 0.76070 | 0.0000000 | 2.48832 | 0.0636350 |
| 0.83330 | 0.0000000 | 2.72580 | 0.0650250 |
| 0.91290 | 0.0000000 | 2.98600 | 0.0661850 |
| 1.00000 | 0.0000000 | | |

Optical and microphysical properties of OMACA Carbonaceous Aerosol Model

| Model# | Imaginary Index | | | Single-scattering Albedo | | | Absorption Angstrom Exponent | |
|--------|------------------|--------|--------|--------------------------|--------|--------|------------------------------|---------|
| | Wavelength in nm | | | Wavelength in nm | | | 354-388 | 354-500 |
| | 354 | 388 | 500 | 354 | 388 | 500 | | |
| 1 | 0.0576 | 0.0480 | 0.0288 | 0.7577 | 0.7806 | 0.8265 | 2.4555 | 2.5080 |
| 2 | 0.0480 | 0.0400 | 0.0240 | 0.7876 | 0.8082 | 0.8486 | 2.5124 | 2.5590 |
| 3 | 0.0360 | 0.0300 | 0.0180 | 0.8288 | 0.8549 | 0.8785 | 2.5935 | 2.6238 |
| 4 | 0.0240 | 0.0200 | 0.0120 | 0.8753 | 0.8879 | 0.9117 | 2.6477 | 2.6821 |
| 5 | 0.0120 | 0.0100 | 0.0060 | 0.9346 | 0.9435 | 0.9603 | 2.8481 | 2.9196 |
| 6 | 0.0060 | 0.0050 | 0.0030 | 0.9646 | 0.9696 | 0.9789 | 2.9343 | 2.9955 |
| 7 | 0.0000 | 0.0000 | 0.0000 | 1.0000 | 1.0000 | 1.0000 | N/A | N/A |

Real refractive index = 1.5

Model # 1 to 3

Mean Radius (fine mode) = 0.080132 μm ; Mean Radius (coarse mode) = 0.705495 μm Standard

Deviation (fine mode) = 1.492 μm ; Standard Deviation (coarse mode) = 2.075 μm

Minimum Radii (fine mode) = 0.0161708 μm ; Minimum Radii (coarse mode) = 0.0380559 μm

Maximum Radii (fine mode) = 0.397083 μm ; Maximum Radii (coarse mode) = 13.0788 μm

Model # 4 to 7

Mean Radius (fine mode) = 0.08717 μm ; Mean Radius (coarse mode) = 0.567194 μm Standard

Deviation (fine mode) = 1.537 μm ; Standard Deviation (coarse mode) = 2.203 μm

Minimum Radii (fine mode) = 0.0156197 μm ; Minimum Radii (coarse mode) = 0.0240810 μm

Maximum Radii (fine mode) = 0.486477 μm ; Maximum Radii (coarse mode) = 13.3595 μm

Nodes of OMACA Look-up table

| Node Parameter | Number of Nodes | Node Values |
|--------------------------------|------------------------|---------------------------------------------------------------------|
| ACAOD | 7 | 0.0, 0.1, 0.5, 1.0, 2.5, 4.0, and 6.0 at 500 nm |
| COD | 8 | 2, 5, 10, 20, 30, 40, and 50 (wavelength independent) |
| Solar Zenith Angle | 7 | 0°, 20°, 40°, 60°, 66°, 72°, 80° |
| Viewing Zenith Angle | 14 | 0°, 12°, 18°, 26°, 32°, 36°, 40°, 46°, 50°, 54°, 56°, 60°, 66°, 72° |
| Relative Azimuth Angle | 11 | 0°, 30°, 60°, 90°, 120°, 150°, 160°, 165°, 170°, 175°, 180° |
| Surface Pressure Levels | 2 | 1013.25, 800 hPa |
| Aerosol Layer Height | 4 | 3, 4, 5, 6 km |
| Surface Albedo | 5 | 0.00, 0.05, 0.10, 0.15, 0.20 at 354 and 388 nm |

Content of OMACA HDF-EOS5 Data File

| SDS Name | Dimensions | Description |
|-------------------------------------|-------------------|-----------------------------------------------------------------------------------|
| Geolocation Fields | | |
| <i>Longitude</i> | X,Y | Geodetic longitude of the center part of the pixel (deg) |
| <i>Latitude</i> | X,Y | Geodetic latitude of the center part of the pixel (deg) |
| <i>FoV75Area</i> | X | Mean Area for 75% Field of View Pixels on the WGS-85 Ellipsoid (km ²) |
| <i>FoV75CornerLongitude</i> | X,Y,4 | Corner Latitudes for 75% Field of View Pixels on the WGS-85 Ellipsoid (deg) |
| <i>FoV75CornerLatitude</i> | X,Y,4 | Corner Latitudes for 75% Field of View Pixels on the WGS-85 Ellipsoid (deg) |
| <i>SolarZenithAngle</i> | X,Y | Solar Zenith Angle (deg) |
| <i>ViewingZenithAngle</i> | X,Y | Satellite Viewing Zenith Angle (deg) |
| <i>RelativeAzimuthAngle</i> | X,Y | Relative Azimuth Angle (deg) SolarZenithAngle + 180 – ViewingZenithAngle |
| <i>TerrainPressure</i> | X,Y | Terrain Pressure (mbar) |
| <i>Time</i> | Y | Time at the start of each scan (seconds, TAI93) |
| <i>SecondsInDay</i> | Y | Seconds of day at start of scan |
| <i>XTrackQualityFlags</i> | X,Y | XTrack Quality Flags |
| <i>GroundPixelQualityFlags</i> | X,Y | Groud Pixel Quality Flags |
| Data Fields | | |
| <i>AerosolOpticalDepthOverCloud</i> | X,Y,3 | Aerosol Optical Depth Over Cloud at 354, 388, and 500 nm |
| <i>AerosolCorrCloudOpticalDepth</i> | X,Y,3 | Aerosol-corrected Cloud Optical Depth at 354, 388, and 500 nm |
| <i>ApparentCloudOpticalDepth</i> | X,Y,3 | Apparent (not corrected for aerosols) Cloud Optical Depth at 354, 388, and 500 nm |
| <i>FinalAlgorithmFlags</i> | X,Y | Final Algorithm Flags assigned to each OMACA retrieval |
| <i>FinalAlgorithmFlags_MieAI</i> | X,Y | Final Algorithm Flags associated with UV Aerosol Index calculations |
| <i>CloudOpticalDepth_MieAI</i> | X,Y | Cloud Optical Depth (388 nm) imported from the OMAERUV AI (Mie) algorithm |
| <i>CloudFraction_MieAI</i> | X,Y | Radiative Cloud Fraction (388 nm) imported from the OMAERUV AI (Mie) algorithm |
| <i>InputSSA354</i> | X,Y | Aerosol Single-scattering Albedo at 354 nm assumed in the retrieval |
| <i>InputSSA388</i> | X,Y | Aerosol Single-scattering Albedo at 388 nm assumed in the retrieval |
| <i>InputSSA500</i> | X,Y | Aerosol Single-scattering Albedo at 500 nm assumed in the retrieval |
| <i>UVAerosolIndex</i> | X,Y | UV Aerosol Index (354-388 nm pair) imported from the OMAERUV algorithm |
| <i>NormRadiance</i> | X,Y,3 | Normalized Radiance at 354, 388, and 500 m |
| <i>Reflectivity</i> | X,Y,2 | Reflectivity at 354 and 388 nm |
| <i>Residue</i> | X,Y | Residue (354-388 nm pair) |
| <i>SurfaceAlbedo</i> | X,Y,2 | Surface Albedo at 354 and 388 nm |
| <i>FinalAerosolLayerHeight</i> | X,Y | Final Aerosol Layer Height (km) from the CALIOP-OMI monthly dataset |
| <i>AIRSL3COvalue</i> | XY | AIRS Carbon Monoxide L3 data |
| <i>AerosolType</i> | X,Y | Aerosol Type assigned to each OMACA retrieval |
| <i>Wavelength</i> | 3 | Wavelength of interest (354, 388, 500 nm) |
| <i>PixelQualityFlags</i> | X,Y,3 | Pixel Quality Flags for 354, 388, and 500 nm |

List of Tables

1. A description of the OMACA algorithm retrieval quality flags.
- 5 2. Theoretical simulation of the retrieved ACAOD for given two discrete values of UVAI under different combinations of SSA and ALH assumptions. Threshold values of 0.8 and 1.3 in UVAI correspond to the algorithm quality flags of '0' (best) and '2' (lesser confidence).
- 10 3. Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of SSA at 388 nm. The reference value of SSA (388 nm) assumed in the calculation was 0.89; the error in SSA (leftmost column) represents perturbation from the reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10.
- 15 4. Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of aerosol layer height (ALH). The reference value of ALH assumed in the calculation was 4.0 km; the error in ALH (leftmost column) represents perturbation from the reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10.
- 20 5. Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of aerosol Absorption Angstrom Exponent (AAE). The reference value of AAE (354-388 nm) assumed in the calculation was 2.65; the error in AAE (leftmost column) represents perturbation from this reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10.
- 25 6. Theoretical error (%) in aerosol-corrected COD (388 nm) due to the uncertainty in the assumption of aerosol SSA and layer height (ALH).
7. Theoretical error (%) in aerosol-corrected COD (388 nm) due to the uncertainty in the assumption of cloud effective radius.

Table 1 A description of the OMACA algorithm retrieval quality flags.

| Algorithm Quality Flags | Observation Conditions | Description |
|--------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| 0 | UVAI (Mie) >1.3 & LER388 >0.25 | Best quality retrievals |
| 1 | 1.3 < UVAI (Mie) < 4.3 & 0.20 < LER388 < 0.25 | Less confidence in the detection of total overcast pixels (Use of high-res sensors is recommended) |
| 2 | 0.8 < UVAI (Mie) < 1.3 & LER > 0.25 | Less confidence in the detection of aerosols above cloud |
| 3 | Solar Zenith Angle > 55° & Scattering Angle < 100° & UVAI (Mie) < 2 Solar Zenith Angle > 60° & Scattering Angle < 130° & UVAI (Mie) < 2 Viewing Zenith Angle > 55° & Scattering Angle < 100° & UVAI (Mie) < 2 | Geometry-related artifacts (not recommended for use) |
| 4 | Snow/Ice Contamination | No retrieval |
| 5 | Solar Zenith Angle above threshold (70°) | No retrieval |
| 7 | Terrain Pressure below threshold (800 hPa). | No retrieval |
| 8 | Cross-track anomaly | No retrieval |

Table 2 Theoretical simulations of the retrieved ACAOD for given two discrete values of UVAI under different combinations of SSA and ALH assumptions. Threshold values of 0.8 and 1.3 in UVAI correspond to the algorithm quality flags of '0' (best) and '2' (lesser confidence).

| UVAI (Mie) | Aerosol Single-scattering Albedo (388 nm) | | |
|------------|-------------------------------------------|---------------------|---------------------|
| | SSA388=0.85 | SSA388=0.90 | SSA388=0.94 |
| | <i>ALH 3/4/5 km</i> | <i>ALH 3/4/5 km</i> | <i>ALH 3/4/5 km</i> |
| 0.8 | 0.06/0.055/0.05 | 0.18/0.17/0.16 | 0.35/0.33/0.30 |
| 1.3 | 0.22/0.20/0.19 | 0.30/0.28/0.26 | 0.57/0.52/0.49 |

Table 3 Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of SSA at 388 nm. The reference value of SSA (388 nm) assumed in the calculation was 0.89; the error in SSA (leftmost column) represents perturbation from the reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10.

| Error in SSA (388 nm) | Above-cloud AOD (388 nm) | | | | | |
|--------------------------|--------------------------|--------|--------|--------|--------|--------|
| | 0.25 | 0.50 | 0.75 | 1.0 | 1.5 | 2.0 |
| -0.05 | -29.05 | -30.73 | -32.42 | -32.85 | -36.52 | -37.55 |
| -0.04 | -24.53 | -25.85 | -27.19 | -28.22 | -30.32 | -32.51 |
| -0.03 | -19.76 | -20.73 | -21.72 | -22.95 | -23.92 | N/R |
| -0.02 | -14.21 | -14.84 | -15.48 | -16.31 | -16.84 | N/R |
| -0.01 | -7.71 | -8.02 | -8.33 | -8.73 | -8.96 | N/R |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 0.01 | 11.13 | 11.40 | 11.85 | 13.46 | 15.42 | N/R |
| 0.02 | 24.71 | 25.28 | 26.46 | 29.50 | N/R | N/R |
| 0.03 | 41.60 | 42.80 | 46.25 | N/R | N/R | N/R |
| 0.04 | 63.35 | 66.65 | N/R | N/R | N/R | N/R |
| 0.05 | 90.93 | 98.53 | N/R | N/R | N/R | N/R |

Table 4 Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of aerosol layer height (ALH). The reference value of ALH assumed in the calculation was 4.0 km; the error in ALH (leftmost column) represents perturbation from the reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10.

5

| Error in ALH (km) | Above-cloud AOD (388 nm) | | | | | |
|-------------------|--------------------------|-------|-------|-------|-------|--------|
| | 0.25 | 0.50 | 0.75 | 1.0 | 1.5 | 2.0 |
| -1.0 | 7.74 | 9.45 | 11.40 | 14.93 | 20.78 | N/R |
| -0.5 | 3.43 | 4.24 | 5.17 | 6.83 | 9.91 | N/R |
| 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 0.5 | -2.45 | -3.02 | -3.59 | -4.14 | -4.83 | N/R |
| 1.0 | -4.56 | -5.68 | -6.80 | -7.94 | -9.77 | -12.81 |

Table 5 Theoretical error (%) in ACAOD (388 nm) due to the uncertainty in the assumption of aerosol Absorption Angstrom Exponent (AAE). The reference value of AAE (354-388 nm) assumed in the calculation was 2.65; the error in AAE (leftmost column) represents perturbation from this reference value. The cloud optical depth underneath the aerosol layer was assumed to be 10. Simulations marked with N/R (Not Retrieved) represent the retrieval failure due to the out-of-domain issue.

| Error in AAE (354-388 nm) | Above-cloud AOD (388 nm) | | | | | |
|------------------------------|--------------------------|--------|--------|--------|--------|--------|
| | 0.25 | 0.50 | 0.75 | 1.0 | 1.5 | 2.0 |
| -1.5 | 83.17 | 96.36 | 102.80 | N/R | N/R | N/R |
| -1.0 | 48.42 | 54.12 | 58.50 | N/R | N/R | N/R |
| -0.5 | 20.80 | 22.14 | 24.38 | 26.60 | N/R | N/R |
| 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 0.5 | -11.67 | -12.33 | -12.84 | -13.40 | -13.54 | -14.50 |
| 1.0 | -21.69 | -22.96 | -24.00 | -24.69 | -25.65 | -27.69 |
| 1.5 | -30.69 | -32.54 | -34.12 | -34.68 | -36.89 | -40.09 |

Table 6 Theoretical error (%) in aerosol-corrected COD (388 nm) due to the uncertainty in the assumption of aerosol SSA and layer height (ALH).

| <u>Error in SSA (388 nm)</u> | <u>% Error in Cloud Optical Depth (388 nm) (AOD=0.5/AOD=1.0)</u> | | | |
|----------------------------------|----------------------------------------------------------------------|--------------------|--------------------|--------------------|
| | <u>COD=5</u> | <u>COD=10</u> | <u>COD=20</u> | <u>COD=30</u> |
| <u>-0.05</u> | <u>5.65/10.18</u> | <u>2.48/3.01</u> | <u>0.27/-1.19</u> | <u>-0.41/-2.67</u> |
| <u>-0.03</u> | <u>3.90/6.94</u> | <u>1.69/1.81</u> | <u>0.15/-0.88</u> | <u>-0.33/-1.96</u> |
| <u>0.03</u> | <u>-6.57/-4.5</u> | <u>-1.48/-1.29</u> | <u>0.68/4.76</u> | <u>1.97/8.03</u> |
| <u>0.05</u> | <u>-17.81/0.1</u> | <u>-3.74/3.50</u> | <u>1.21/10.40</u> | <u>3.94/17.48</u> |
| | | | | |
| <u>Error in ALH (km)</u> | <u>% Error in Cloud Optical Depth (388 nm) (AOD=0.5/AOD=1.0)</u> | | | |
| | <u>COD=5</u> | <u>COD=10</u> | <u>COD=20</u> | <u>COD=30</u> |
| <u>-1</u> | <u>1.26/4.57</u> | <u>2.83/9.23</u> | <u>3.07/10.48</u> | <u>3.16/11.32</u> |
| <u>+1</u> | <u>-0.65/-2.19</u> | <u>-1.14/-3.48</u> | <u>-1.23/-3.95</u> | <u>-1.51/-5.03</u> |

Table 7 Theoretical error (%) in aerosol-corrected COD (388 nm) due to the uncertainty in the assumption of cloud effective radius.

| <u>Error in cloud effective radius (μm)</u> | <u>% Error in Cloud Optical Depth (388 nm)</u> | | | |
|-------------------------------------------------------------------|------------------------------------------------|----------------------|--------------------|----------------------|
| | <u>Assumed AOD (388 nm) = 0.5/1.0</u> | | | |
| | <u>COD=5</u> | <u>COD=10</u> | <u>COD=20</u> | <u>COD=30</u> |
| <u>-8.0</u> | <u>-11.31/-11.9</u> | <u>-10.28/-10.88</u> | <u>-9.13/-9.83</u> | <u>-10.11/-10.76</u> |
| <u>-6.0</u> | <u>-5.94/-6.54</u> | <u>-5.38/-5.94</u> | <u>-4.68/-5.29</u> | <u>-5.01/-5.62</u> |
| <u>-4.0</u> | <u>-2.85/-3.26</u> | <u>-2.57/-2.94</u> | <u>-2.18/-2.57</u> | <u>-2.25/-2.66</u> |
| <u>0.0</u> | <u>0.00/0.00</u> | <u>0.00/0.00</u> | <u>0.00/0.00</u> | <u>0.00/0.00</u> |
| <u>4.0</u> | <u>1.17/1.4</u> | <u>1.43/1.79</u> | <u>1.21/1.62</u> | <u>0.95/1.39</u> |
| <u>6.0</u> | <u>1.34/1.70</u> | <u>1.63/2.07</u> | <u>1.37/1.87</u> | <u>1.05/1.59</u> |
| <u>8.0</u> | <u>1.51/1.94</u> | <u>1.83/2.35</u> | <u>1.52/2.11</u> | <u>1.16/1.78</u> |
| <u>12.0</u> | <u>1.85/2.41</u> | <u>2.22/2.92</u> | <u>1.83/2.59</u> | <u>1.36/2.18</u> |

List of Figures

Figure 1 Simulation of UVAI (y-axis) versus Reflectance at 388 nm (x-axis) for the different pairs of ACAOD and COD (both at 388 nm) for the carbonaceous (left) and spheroidal shape dust aerosols (right). Values of ACAOD and COD depicted in the figure correspond to the 388 nm wavelength. The shown 2-D diagram forms the retrieval domain in which the observations from OMI are fitted and related to a pair of ACAOD and COD.

Figure 2 (Top) Geographical boundaries of select 14 regions considered for calculating corresponding regional values of SSA (388 nm). (Remaining panels) Regional mean UVAI weighted cloud-free SSA (388 nm) for carbonaceous, and dust aerosols for the four selected regions (numbered 5, 6, 7, and 8 in the top panel) derived using OMI/OMAERUV operational (version 1.8.9.1) L2 cloud-free SSA retrievals.

Figure 3 A general flowchart of the OMACA algorithm.

Figure 4 Comparison of coincident and collocated spectral ACAODs measured and retrieved from HSRL-2 and OMI/OMACA for a total of seven ER-2 flights operated during ORACLES phase I operation over the Southeastern Atlantic Ocean in August-September of 2016. ΔT represents the absolute time difference (in hours) between the OMI overpass and HSRL-2 measurements.

Figure 5 The monthly mean cloudy-sky frequency of occurrence of above-cloud absorbing aerosols deduced from the 12-year long record (2005-2016) of OMI's near-UV observations.

Figure 6 (a) Chart showing the time evolution of OMI row anomaly and (b) cloudy-sky ($LER_{388} > 0.25$) UVAI. (c) Monthly time-series of the global, cloudy-sky frequency of occurrence (in % with respect to the total cloudy-sky observations) of absorbing aerosols above cloud derived using OMI full scan (Rows 1 to 60) (blue) and anomaly-free observations (Rows 1 to 23) (red). Solid lines represent the linear regression fits to the respective time-series data.


Figure 7 Monthly time-series of the regional cloudy-sky frequency of occurrence (in % with respect to the total cloudy-sky observations) of absorbing aerosols above cloud derived using OMI anomaly-free observations (Rows 1 to 23). Solid lines represent the linear fit to the respective time-series data.

Figure 8 Global distribution of monthly mean above-cloud AOD (388 nm) deduced from the 12-year (2005-2016) OMI observations.


Figure 9 Time-series of regional monthly mean above-cloud AOD at 388 nm (left y-axis, color: red) and the ratio of above-cloud AOD and cloud-free AOD (right y-axis, color: blue), both at 388 nm, derived using OMACA and OMAERUV products, respectively.

5 **Figure 10** Time-series of regional, monthly mean aerosol-corrected (left y-axis, color: red) COD (388 nm) derived using a 12-year long OMACA record. The y-axis on the right side (color: blue) depicts a time-series of the % difference between simultaneous retrievals of apparent/non-corrected and aerosol-corrected COD for the same period and regions. Only anomaly-free observations of OMI (Rows 1 to 23) were used in the calculations.

Figure 11 Percent difference in COD (aerosol-corrected minus non-corrected) as a function of above-cloud absorption AOD (388 nm) for the smoke (left panel) and dust (right panel) dominated regions. Different color codes represent the relationship for a discrete range of aerosol-corrected COD.



5 **Figure 1** Simulation of UVAI (y-axis) versus Reflectance at 388 nm (x-axis) for the different pairs of ACAOD and COD (both at 388 nm) for the carbonaceous (left) and spheroidal shape dust aerosols (right). Values of ACAOD and COD depicted in the figure correspond to the 388 nm wavelength. The shown 2-D diagram forms the retrieval domain in which the observations from OMI are fitted and related to a pair of ACAOD and COD.



5 **Figure 2** (Top) Geographical boundaries of select 14 regions considered for calculating corresponding regional values of SSA (388 nm). (Remaining panels) Regional mean UVA1 weighted cloud-free SSA (388 nm) for carbonaceous, and dust aerosols for the four selected regions (numbered 5, 6, 7, and 8 in the top panel) derived using OMI/OMAERUV operational (version 1.8.9.1) L2 cloud-free SSA retrievals.




Figure 3 A General flowchart of the OMACA algorithm.






Figure 4 Comparison of coincident and collocated spectral ACAODs measured and retrieved from HSRL-2 and OMI/OMACA for a total of seven ER-2 flights operated during ORACLES phase I operation over the Southeastern Atlantic Ocean in August-September of 2016. ΔT represents the absolute time difference (in hours) between the OMI overpass and HSRL-2 measurements.

5



5 **Figure 5** The monthly mean distribution of cloudy-sky frequency of occurrence of above-cloud absorbing aerosols deduced from the 12-year record (2005-2016) of OMI's near-UV observations.



5 **Figure 6** (a) Chart showing the time evolution of OMI row anomaly and (b) cloudy-sky (LER388>0.25) UVAI. (c) Monthly time-series of the global, cloudy-sky frequency of occurrence (in % with respect to the total cloudy-sky observations) of absorbing aerosols above cloud derived using OMI full scan (Rows 1 to 60) (blue) and anomaly-free observations (Rows 1 to 23) (red). Solid lines represent the linear regression fits to the respective time-series data.




Figure 7 Monthly time-series of the regional cloudy-sky frequency of occurrence (in % with respect to the total cloudy-sky observations) of absorbing aerosols above cloud derived using OMI anomaly-free observations (Rows 1 to 23). Solid lines represent the linear fit to the respective time-series data.

5




Figure 8 Global distribution of monthly mean above-cloud AOD (388 nm) deduced from the 12-year (2005-2016)

5 OMI observations.





Figure 9 Time-series of regional monthly mean above-cloud AOD at 388 nm (left y-axis, color: red) and the ratio of above-cloud AOD and cloud-free AOD (right y-axis, color: blue), both at 388 nm, derived using OMACA and

5 OMAERUV products, respectively.



5 **Figure 10** Time-series of regional, monthly mean aerosol-corrected (left y-axis, color: red) COD (388 nm) derived using a 12-year long OMACA record. The y-axis on the right side (color: blue) depicts a time-series of the % difference between simultaneous retrievals of apparent/non-corrected and aerosol-corrected COD for the same period and regions. Only anomaly-free observations of OMI (Rows 1 to 23) were used in the calculations.




Figure 11 Percent difference in COD (aerosol-corrected minus non-corrected) as a function of above-cloud absorption AOD (388 nm) for the smoke (left panel) and dust (right panel) dominated regions. Different color codes represent the relationship for a discrete range of aerosol-corrected COD.

5