Answers to Comments of Reviewer 1

We would like to thank the reviewer for the fruitful comments and suggestions which helped improving the manuscript.

General comment 1

In this paper some interesting analyses are presented about the QBO, ENSO and NAO signal in various long-term ozone data sets. However, I have some problems to find the main aim of the paper. In the abstract it is mentioned that validation is performed for GOME2-A, yet no direct comparison with ground observations has been made. The correlations have been derived for QBO, ENSO and NAO signals, which although interesting as it is, I would not call validation. The term "evaluation" mentioned in the title is a better description. In the title, on the other hand, only GOME-2A is mentioned, while the authors are evaluating SBUV and GTO-ECV in exactly the same way. I suggest to change the title to "The use of QBO, ENSO and NAO perturbations in the evaluation of long-term total ozone satellite measurements." and to use 'evaluation' instead of 'validation' throughout the text.

Answer to general comment 1:

The aim of the paper can be found in the Introduction and reads as follows: "The objective of the present work is to examine the ability of the GOME-2A total ozone data to capture the variability related to dynamical proxies of global and regional importance such as the QBO, ENSO and NAO, in comparison to GB measurements, other satellite data and model calculations. The variability of total ozone from GOME-2A is compared with the variability of total ozone from the other examined data sets during these naturally-occurring fluctuations in order to evaluate the performance of GOME-2A to depict natural perturbations. The analysis is performed in the frame of the validation strategy of GOME-2A data on longer time scales within the project of EUMETSAT, AC SAF. The evaluation of GOME-2A data performed here includes the study of monthly means of total ozone, the annual cycle of total ozone, the annual cycle of total ozone, the annual cycle of the annual cycle [i.e., (max-min)/2], the relation with the QBO (zonal winds at the equator at 30 hPa), the relation with ENSO (correlation with SOI) and the relation with the NAO (correlations with the NAO index in winter (DJF mean))."

The abstract now states "Comparison of GOME-2A total ozone with ground observations shows mean differences of about $-0.7 \pm 1.4\%$ in the tropics (0-30 deg.), about $+0.1 \pm 2.1\%$ in mid-latitudes (30-60 deg.), and about $+2.5 \pm 3.2\%$ and $0.0 \pm 4.3\%$ over the northern and southern high latitudes (60-80 deg.), respectively.". Additional comparisons with ground observations are mentioned in the abstract in different lines as follows: "Differences between deseazonalised GOME-2A and GB total ozone in the tropics are within $\pm 1\%$.", "Differences between GOME-2A and GB measurements at the station of Samoa (American Samoa; 14.25° S, 170.6° W) are within $\pm 1.9\%$.", "We find very good agreement between GOME-2A and GB observations over Canada and Europe as to their NAO-related variability, with mean differences reaching the $\pm 1\%$ levels".

While we analyse other satellite data as well, we give emphasis to GOME-2A. We prefer to keep the title as is.

We now use the term 'evaluation' instead of 'validation' throughout the text.

General comment 2

Throughout the paper correlations are calculated for the comparisons, which I think is very limited. I suggest that the authors provide more information on these comparisons by calculating the regression coefficients.

Answer to general comment 2:

The reviewer asks more information on the comparisons throughout the paper, which is now provided with the regression coefficients as suggested. The regression coefficients for the comparisons are presented in the new Tables 2, 3, 5, 8 (see also answer to comment 8). In addition, in the Supplement of this study we provide global maps of the regression coefficients of QBO, solar cycle, ENSO and NAO, in the Figures S1 (for QBO), S2 (for solar cycle), S3 (for ENSO) and S4 (for NAO), respectively.

Detailed comments:

Comment 1: Line 23: validating => evaluating

Answer to 1: Done

Comment 2: Line 29: Here the GTO-ECV data set is mentioned for the first time. I don't think most readers will have a clear idea what "GOME-type Total Ozone Essential Climate Variable" mean. A short description to describe this data set would be helpful at this point.

Answer to 2: We now write "... GOME-type Total Ozone Essential Climate Variable (GTO-ECV; composed of total ozone observations from GOME (Global Ozone Monitoring Experiment), SCIAMACHY (SCanning Imaging Absorption SpectroMeter for Atmospheric CHartographY), GOME-2A, and OMI (Ozone Monitoring Instrument) combined into one homogeneous time series) ..."

Comment 3: Line 51: Cause & effect are reversed in this sentence. Ozone is considered a greenhouse gas because it warms the Earth's surface not the other way around. In addition, it might be good to mention that not only tropospheric ozone but also stratospheric ozone is a greenhouse gas.

Answer to 3: The line has been revised and now reads as "In addition, ozone is a greenhouse, warming the Earth's surface. In both the stratosphere and the troposphere, ozone absorbs infrared radiation emitted from Earth's surface, trapping heat in the atmosphere. As a result, increases or decreases in stratospheric or tropospheric ozone induce a climate forcing (Hegglin et al., 2015)."

Comment 4: Line 56: "launched in 2018." => "launched end of 2018"

Answer to 4: Changed to "launched on 7 November 2018".

Comment 5: Line 73: Except for the abstract, this is the first time that the SBUV and GTO-ECV datasets are mentioned, therefore, I suggest to add references for both data sets in the text.

Answer to 5: The reference (McPeters et al., 2013) has been added here for SBUV and the references (Coldewey-Egbers et al., 2015; Garane et al., 2018) have been added for GTO-ECV.

Comment 6: Line 89-91: It might be better to refer to more recent papers about the recovery of the ozone layer, for example de Laat et al., Onset of Stratospheric Ozone Recovery in the Antarctic Ozone Hole in Assimilated Daily Total Ozone Columns, JGR, 2017, https://doi.org/10.1002/2016JD025723

Answer to 6: We have added more recent papers about the recovery of the ozone layer, as follows: Solomon et al., 2016; de Laat et al., 2017; Kuttippurath and Nair, 2017; Pazmiño et al., 2018; Stone et al., 2018; Strahan and Douglass, 2018.

The following six papers have been added in list of the references:

Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Neely III, R. R., and Schmidt, A.: Emergence of healing in the Antarctic ozone layer, Science, 30, doi: 10.1126/science.aae0061, 2016.

de Laat, A. T. J., van Weele, M., and van der A., R. J.: Onset of stratospheric ozone recovery in the Antarctic ozone hole in assimilated daily total ozone columns, Journal of Geophysical Research: Atmospheres, 122, 11880-11899, https://doi.org/10.1002/2016JD025723, 2017.

Kuttippurath, J. and Nair, P. J.: The signs of Antarctic ozone hole recovery, Sci. Rep., 7, https://doi.org/10.1038/s41598-017-00722-7, 2017.

Pazmiño, A., Godin-beekmann, S., Hauchecorne, A., Claud, C., Khaykin, S., Goutail, F., Wolfram, E., Salvador, J., and Quel, E.: Multiplesymptoms of total ozone recovery inside the Antarctic vortex during austral spring, Atmos. Chem. Phys, 18, 7557–7572, 2018.

Stone, K. A., Solomon, S., and Kinnison, D. E.: On the identification of ozone recovery, Geophysical Research Letters, 45, 5158-5165, https://doi.org/10.1029/2018GL077955, 2018.

Strahan, S. E. and Douglass, A. R.: Decline in Antarctic Ozone Depletion and Lower Stratospheric Chlorine Determined From Aura Microwave Limb Sounder Observations, Geophys. Res. Lett., 45, 382–390, https://doi.org/10.1002/2017GL074830, 2018.

Comment 7: Line 150-151: When mentioning the various long-term data sets of ozone, also the Multi-Sensor Reanalysis of ozone comes to mind. This data set has also been analysed for QBO, ENSO, NAO and other perturbations in Knibbe et al., ACP, 2014 and therefore is worthwhile to include here and in the discussion at the end of section 3.3.

Answer to 7: We have added the following sentence in response to the comment: "We note here that another long-term data set which has been analysed for QBO, ENSO, NAO and

other perturbations comes from the Multi-Sensor Reanalysis (Knibbe et el., 2014), but is not examined here.". Additionally, the study by Knibbe et al., ACP, 2014 is now included in the discussion at the end of section 3.3, and has been added in the list of references.

Knibbe, J. S., van der A, R. J., and de Laat, A. T. J.: Spatial regression analysis on 32 years of total column ozone data, Atmos. Chem. Phys., 14, 8461-8482, https://doi.org/10.5194/acp-14-8461-2014, 2014.

Comment 8: Line 223: I prefer to see more than only correlation coefficients. The regression parameters could be given here and in the remainder of the analyses.

Answer to 8: The regression parameters for the correlations shown in Figures 1 and 2 are provided in the new Table 2. The regression parameters for the comparisons with the QBO are provided in the new Table 3. The regression parameters for the comparisons with SOI are provided in the new Table 5. The regression parameters for the comparisons with NAO in winter are provided in the new Table 8.

Comment 9: Line 239-240: This sentence seems to saying that the origin of the blue zone (i.e. small amplitude) is attributed to the small amplitude in these parts. Please, give the real origin if this is known.

Answer to 9: The sentence has been corrected and now reads as follows "Interestingly, there is pattern with small amplitude of annual cycle in the southern mid-latitudes with values of about 10-15 DU, seen in Figure 4 as a blue curved line crossing the longitudes around 60 degrees south, which points to small seasonal variations of total ozone in these parts. The seasonal increase in Antarctic is delayed by 2-3 months compared to the north polar region. Only with the breakdown of the polar vortex in late spring, i.e. at a time when the poleward transport over lower latitudes has already ceased, does a strong ozone influx occur in the Antarctic. With this delay the amplitude of the seasonal variation stays much smaller poleward of 55-60° in the south than in the north (Dütsch, 1974)."

The citation (Dütsch, 1974) has been added in the list of references: Dütsch, H. U.: The ozone distribution in the atmosphere, Can. J. Chem, 52, 1491-1504, 1974.

Comment 10: Line 259-265: This analysis was already discussed in section 3.1. Only this time the monthly mean has been subtracted which does not really change the validation. I suggest to remove this or add it in section 3.1.

Answer to 10: We have removed it.

Comment 11: Line 269: A clear phase shift in Figure 5 is mentioned for higher latitudes. Actually for SBUV I see an anti-correlation with the phase for latitudes between -10 and 10, and for GOME2 I see neither phase shift or an anticorrelation. So I would not call this a clear phase shift. A discussion about the clear differences in result of SBUV (pre 2008) and GOME-2 should be added here as well. Answer to 11: For SBUV there is no anti-correlation for latitudes between -10 and 10. The regression coefficients of QBO are all positive in the tropics and negative at higher latitudes as we show in the new Table 5, and display in the Supplement Figure S1.

The part of the text discussing the correlation with the QBO has been revised and now reads as follows:

"The line with dots superimposed on the ozone anomalies in Figure 5 shows the equatorial zonal winds at 30 hPa which were used as a proxy index to study the impact of QBO on total ozone. The general features include a QBO signal in total ozone at latitudes between 10° N and 10° S which almost matches with the phase of QBO in the zonal winds. At higher northern and southern latitudes there is a phase shift in the QBO impact on total ozone. The impact of QBO is more pronounced in the tropics and it is less pronounced in the sub-tropics and mid-latitudes. Strong positive correlations with the QBO are found in the tropics (correlation between GOME-2A and QBO of about +0.77, t-test = 12.91) and weaker (usually of opposite sign) less significant correlations are found at higher latitudes (about -0.15 in the northern and about -0.45 in the southern extra tropics). Similar correlations suggest that the variability that can be attributed to the QBO is about 60% in the tropics, and about 2% and 20% in the northern and the southern extra tropics, respectively.

Table 3 summarizes the correlation and regression coefficients between total ozone and QBO at 30 hPa for the different latitude zones and the different datasets. The numbers speak for themselves: for latitudes between 10° N and 10° S correlations between total ozone from GOME-2A, GTO-ECV, SBUV, GB data and the QBO are all positive. At latitudes between 10° and 30° the correlations turn to negative, in agreement with Knibbe et al. (2014) results, who noted that moving from the tropics towards higher latitudes the regression estimates switch to negative values at approximately 10° N and 10° S. The correlations with the QBO at 30 hPa remain negative up to 60°, a consistent result among all our data sets, something also reported by Knibbe et al. (2014) with the MSR ozone data. The correlation and regression coefficients between GOME-2A and QBO are fairly similar to those found between SBUV and QBO, as well as among all data sets as seen in Table 3, despite the different periods of records."

Comment 12: Line 291-292: The correlations are not removed but the relation between ozone and QBO has been removed. Please, reformulate.

Answer to 12: We have reformulated as follows: "To examine the impact of ENSO on total ozone we first removed variability related to the QBO and the solar cycle, and then performed the correlation analysis with the SOI".

Comment 13: Line 295: If you are using this equation, it would be very interesting to mention also the fitted a0 and a1 instead or in addition to the found correlations.

Answer to 13: The fitted a0 and a1 are provided in addition to the found correlations, as follows: "The QBO-related coefficients *a*0 and *a*1 of Eq. (1) for the deseasonalized GOME-2A, GTO-ECV, TOMS/OMI/OMPS and Oslo CTM3 zonal mean data are presented in Table 3. Additional information for the regression coefficients *a*1 of QBO is provided in the

Supplement Figure S1, which shows the spatial distribution of the regression coefficients in latitude-longitude maps."

Comment 14: Section 3.3, Figure 8 and 9: the GOME2 values in the last 4 year of the Figures 8 and 9 show a much worse comparison than the other years in the time series. Is there any explanation for this? I miss this in the discussion of the results here.

Answer to 14: We have added it in Section 3.3 as follows: "Despite the small differences found, we note here that GOME-2A values in the last 4 years of Figures 8 and 9 slightly deviate from the other data sets, and correlate weaker with SOI than the other years in the time series. For instance, we estimate a drop in the correlation coefficient between GOME-2A and SOI at the station Samoa (+0.58 in the period 2007-2012 and +0.47 in the period 2007-2016), which nevertheless does not alter the statistical significance of the correlation."

Comment 15: Line 367: A discussion of a comparison with the work of Knibbe et al., ACP, 2014 would be useful at this point.

Answer to 15: We have added the following sentence at this point "Our results are also in agreement with Knibbe et al. (2014) who showed negative ozone effects of El Niño between 25° S and 25° N, especially over the Pacific."

Comment 16: Line 370: Here the effects of QBO are removed, but what about the ENSO perturbations? Are these also removed before continuing studying the NAO effects. The two effects have to be separated.

Answer to 16: The effect of ENSO is now removed before continuing studying the NAO effects. The new line now reads "The residuals from Eq. (3), free from seasonal, QBO, solar and ENSO related variations, were later used to study the correlation between total ozone and NAO in winter". Tables and figures 7-12 for ENSO and NAO have been revised accordingly.

Comment 17: Line 293-393: Same as previous remark.

Answer to 17: We now separate the effects using different regressions, one regression to account for the effect of QBO (Eq. 1), a second regression to account for the effect of solar cycle (Eq. 2) and a third regression to account for the effect of ENSO (Eq. 3). Variability related to ENSO is now removed with Eq. (3) before continuing studying the NAO effects. The related text, tables, and figures have been revised accordingly.

Comment 18: Line 469: This is not a real validation because a lot is still unknown about the quantification of the QBO, ENSO and NAO, therefore it is qualitative evaluation not a quantitative validation resulting in uncertainty estimates.

Answer to 18: We have corrected the text to read "to qualitatively evaluate GOME-2A" instead of "validating GOME-2A".

Comment 19: Figure 1: It is very difficult to distinguish the GOME2-A line and the SBUV-line. The legend doesn't seem to be correct either?

Answer to 19: The figures 1 and 2 have been redrawn using a different combination of colors.

Answers to Comments of Reviewer 2

We would like to thank the reviewer for the fruitful comments and suggestions which helped improving the manuscript.

Specific comments

Comment 1: Figures 1 and 2: Maybe you could select a different combination of colors since now it is hard the differences between the different datasets to be distinguished. Alternatively you could plot the monthly differences, instead of the actual total ozone column values.

Answer to 1: The figures 1 and 2 have been redrawn using a different combination of colors.

Comment 2: Page 6 lines 214-215: It is mentioned that the highest differences are found over the southern high latitudes, however from Figures 1 and 2 it is depicted that these are presented over the Northern high latitudes (60 - 80 N) and the highest variability (standard deviation of the mean difference) is observed over the latitude belt (60 - 80 S). In addition, these differences (especially at the high latitudes) can be affected by the fact that you have not used the same days for the construction of the monthly mean values for the different datasets.

Answer to 2: The lines have been revised as suggested, and now read as follows: "In summary, the largest differences between GOME-2A, SBUV (v8.6) and GB measurements are found over the northern high latitudes $(60^{\circ}-80^{\circ} \text{ N})$ and the highest variability (standard deviation of the mean difference) is observed over the latitude belt $(60^{\circ}-80^{\circ} \text{ S})$. In addition, these differences (especially at the high latitudes) can be affected by the fact that not always the same days have been used for the construction of the monthly mean values for the different datasets."

Comment 3: Page 7 lines 220-226: Which statistical test did you use to check the statistical significance?

Answer to 3: We have added this sentence in the text which explains it: "The statistical significance of the correlation coefficients, R, was calculated using the *t*-test formula for R with N-2 degrees of freedom, as used in Zerefos et al. (2018)."

The formula is:

$$t = R \sqrt{\frac{N-2}{1-R^2}}$$

The citation Zerefos et al. (2018) has been added in the list of references:

Zerefos, C. S., Kapsomenakis, J., Eleftheratos, K., Tourpali, K., Petropavlovskikh, I., Hubert, D., Godin-Beekmann, S., Steinbrecht, W., Frith, S., Sofieva, V., and Hassler, B.: Representativeness of single lidar stations for zonally averaged ozone profiles, their trends and attribution to proxies, Atmos. Chem. Phys., 18, 6427-6440, https://doi.org/10.5194/acp-18-6427-2018, 2018.

Comment 4: Page 8 lines 269 – 271: I don't think that you see the amplitude of QBO effect on your total ozone column. The times series are just deseasonalized, but still contain the effect of other signals such as the 11 year solar cycle, ENSO etc and thus not all the variation can be attributed to QBO.

Answer to 4: We agree that not all the variation can be attributed to QBO, and we have revised the part of the text describing the correlation with the QBO as follows:

"The line with dots superimposed on the ozone anomalies in Figure 5 shows the equatorial zonal winds at 30 hPa which were used as a proxy index to study the impact of QBO on total ozone. The general features include a QBO signal in total ozone at latitudes between 10° N and 10° S which almost matches with the phase of QBO in the zonal winds. At higher northern and southern latitudes there is a phase shift in the QBO impact on total ozone. The impact of QBO is more pronounced in the tropics and it is less pronounced in the sub-tropics and mid-latitudes. Strong positive correlations with the QBO are found in the tropics (correlation between GOME-2A and QBO of about +0.77, t-test = 12.91) and weaker (usually of opposite sign) less significant correlations are found at higher latitudes (about -0.15 in the northern and about -0.45 in the southern extra tropics). Similar correlations suggest that the variability that can be attributed to the QBO is about 60% in the tropics, and about 2% and 20% in the northern and the southern extra tropics, respectively.

Table 3 summarizes the correlation and regression coefficients between total ozone and QBO at 30 hPa for the different latitude zones and the different datasets. The numbers speak for themselves: for latitudes between 10° N and 10° S correlations between total ozone from GOME-2A, GTO-ECV, SBUV, GB data and the QBO are all positive. At latitudes between 10° and 30° the correlations turn to negative, in agreement with Knibbe et al. (2014) results, who noted that moving from the tropics towards higher latitudes the regression estimates switch to negative values at approximately 10° N and 10° S. The correlations with the QBO at 30 hPa remain negative up to 60°, a consistent result among all our data sets, something also reported by Knibbe et al. (2014) with the MSR ozone data. The correlation and regression coefficients between GOME-2A and QBO are fairly similar to those found between SBUV and QBO, as well as among all data sets as seen in Table 3, despite the different periods of records."

Comment 5: Figures 5 and 6: You could possible superimpose the QBO proxy on the ozone anomalies.

Answer to 5: The QBO proxy is now superimposed on the ozone anomalies.

Comment 6: Section 3.3: You removed the effect of the annual cycle and QBO, before you correlate your ozone time series with ENSO but the effect of solar cycle could also affect your results.

Answer to 6: We now remove the effect of solar cycle and repeat our calculations. We account for the solar cycle effect in ozone, using the 10.7 cm wavelength solar radio flux

(F10.7) as a proxy, taken from the National Research Council and Natural Resources Canada at <u>ftp://ftp.geolab.nrcan.gc.ca/data/solar_flux/monthly_averages/solflux_monthly_average.txt</u> We used the absolute values of F10.7. The text, tables, and figures 7-12, have been revised accordingly.

Comment 7: Page 9 lines 306-307: Which statistical test did you use for checking the statistical significance?

Answer to 7: We used the *t*-test for *R* with *N*-2 degrees of freedom (see answer to comment 3). We have corrected the sentence as follows: "These correlations were tested as to their statistical significance in the period 2007-2016 using the *t*-test for *R* with *N*-2 degrees of freedom (as in Zerefos et al., 2018), and were found to be statistical significant."

Comment 8: Section 3.4: Here you discuss the correlations between total ozone column and the NAO during winter months, evaluating the known anti-correlation between those two factors. Maybe it would be of interest to look also the correlations during summer, following the study of Osso et al. who reported a reversal in the correlation pattern between NAO and TOC from winter to summer for southern Europe.

Ossó A, Sola Y, Bech J, Lorente J (2011) Evidence for the influence of the North Atlantic Oscillation on the total ozone column at northern low latitudes and midlatitudes during winter and summer seasons. J Geophys Res Atmos 116:D24122. doi: 10.1029/2011JD016539

Answer to 8: We have also looked at the correlations during summer, which appear in the new Figure 13 for southern Europe. The new Figure A2 of Appendix A shows the correlations in global maps. The results are discussed at the end of section 3.4 as follows:

"The anti-correlation between total ozone column and the NAO index during winter also applies to southern Europe and the Mediterranean. Following the study of Ossó et al. (2011) who reported a reversal in the correlation pattern between NAO and total ozone from winter to summer in southern Europe, we have looked at the correlations during summer as well. Figure 13 presents the comparisons for 21 ground-based stations located in the region bounded by latitudes 30° - 47° N and by longitudes 10° W- 40° E. Figure 13a shows results for the summer and Figure 13b shows results for winter. As evident, the anti-correlation between GB total ozone and NAO in winter (R= -0.43, slope= -0.980, t-value= -2.095, p-value= 0.0499, N = 21) reverses sign and becomes positive in the summer (R= +0.60, slope= 0.874, t-value= 3.309, p-value= 0.0037, N= 21), indicating that the NAO explains about 36% of ozone variability in the summer in this region. A similar picture is also seen from GOME-2A, GTO-ECV and SBUV data."

Typos:

Page 5, line 146: 50 -> 5^o Answer: Done

Page 5, line 149: all offsets where -> all offsets were Answer: Done

Page 5, line 179: we made use of the monthly -> we used the monthly Answer: Done

Page 6, line 181: we made use of the monthly -> we used the monthly Answer: Done

Page 6, lines 187 – 190: "Use was made of the principal …" doesn't sound very nice maybe you could change to: "The principal component (PC)-based NAO index (DJF) provided by the … (last access: 15 June 2018) was used (or analyzed). Answer: Changed as suggested.

Page 6, line 190: After dynamical variability add "," Answer: Done

Page 6, line 192: The impact of tropopause variability on -> The impact of the tropopause height variations on Answer: Done

The use of QBO, ENSO and NAO perturbations in the 1 evaluation of GOME-2/MetopA total ozone measurements 2

Kostas Eleftheratos^{1,2}, Christos S. Zerefos^{2,3,4,5}, Dimitris S. Balis⁶, Maria-Elissavet Koukouli⁶, 3

John Kapsomenakis³, Diego G. Loyola⁷, Pieter Valks⁷, Melanie Coldewey-Egbers⁷, Christophe Lerot⁸, Stacey M. Frith⁹, Amund <u>Søvde S.</u> Haslerud¹⁰, Ivar S. A. Isaksen^{10,11}, Seppo Hassinen¹² 4

5

¹Laboratory of Climatology and Atmospheric Environment, Faculty of Geology and Geoenvironment, National and 6

- 7 Kapodistrian University of Athens, Greece
- 8 ²Biomedical Research Foundation of the Academy of Athens, Athens, Greece
- ³Research Centre for Atmospheric Physics and Climatology, Academy of Athens, Athens, Greece 9
- 10 ⁴Mariolopoulos-Kanaginis Foundation for the Environmental Sciences, Athens, Greece
- ⁵Navarino Environmental Observatory (N.E.O.), Messinia, Greece 11
- ⁶Laboratory of Atmospheric Physics, Department of Physics, Aristotle University of Thessaloniki, Greece 12
- 13 ⁷Institut für Methodik der Fernerkundung (IMF), Deutsches Zentrum für Luft- und Raumfahrt (DLR),
- 14 Oberpfaffenhofen, Germany
- 15 ⁸Royal Belgian Institute for Space Aeronomy (BIRA), Brussels, Belgium
- 16 ⁹Science Systems and Applications, Inc., Lanham, MD, USA
- ¹⁰Cicero Center for International Climate Research, Oslo, Norway 17
- ¹¹Department of Geosciences, University of Oslo, Oslo, Norway 18
- ¹²Finnish Meteorological Institute, Helsinki, Finland 19
- 20 Correspondence to: Kostas Eleftheratos (kelef@geol.uoa.gr)

21 Abstract. In this work we present evidence that quasi cyclical perturbations in total ozone (Quasi Biennial 22 Oscillation (QBO), El Nino Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO)) can be used as independent proxies in validating evaluating Global Ozone Monitoring Experiment-2 aboard MetopA (GOME-2A) 23 24 satellite total ozone data, using ground-based measurements, other satellite data and chemical transport model 25 calculations. The analysis is performed in the frame of the validation strategy on longer time scales within the 26 European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT), Satellite Application 27 Facility on Atmospheric Composition Monitoring (AC SAF) project, and covers the period 2007-2016. Comparison 28 of GOME-2A total ozone with ground observations shows mean differences of about $-0.7 \pm 1.4\%$ in the tropics (0-29 30 deg.), about $+0.1 \pm 2.1\%$ in mid-latitudes (30-60 deg.), and about $+2.5 \pm 3.2\%$ and $0.0 \pm 4.3\%$ over the northern 30 and southern high latitudes (60-80 deg.), respectively. In general, we find that GOME-2A total ozone data depict the QBO/ENSO/NAO natural fluctuations in concurrence with co-located Solar Backscatter Ultraviolet Radiometer 31 (SBUV), GOME-type Total Ozone Essential Climate Variable (GTO-ECV: composed of total ozone observations 32 33 from GOME (Global Ozone Monitoring Experiment), SCIAMACHY (SCanning Imaging Absorption SpectroMeter 34 for Atmospheric CHartographY), GOME-2A, and OMI (Ozone Monitoring Instrument) combined into one homogeneous time series) and ground-based (GB) observations. Total ozone from GOME-2A is well correlated with 35 36 the QBO (highest correlation in the tropics of +0.8) in agreement with SBUV, GTO-ECV and GB data which also 37 give the highest correlation in the tropics. The differences between deseazonalised GOME-2A and GB total ozone in

- the tropics are within $\pm 1\%$. These differences were tested further as to their correlations with the QBO. The 38
- 39 differences had practically no QBO signal, providing an independent test of the stability of the long-term variability

40 of the satellite data. Correlations between GOME-2A total ozone and the Southern Oscillation Index (SOI) were 41 studied over the tropical Pacific Ocean after removing seasonal-and, QBO and solar cycle related variability. 42 Correlations between ozone and SOI are the order of $\frac{10.60.5}{10.60.5}$, in consistency with SBUV and GB observations. 43 Differences between GOME-2A and GB measurements at the station of Samoa (American Samoa; 14.25° S, 170.6° 44 W) are within $\pm 1.51.9$ %. We also studied the impact of NAO on total ozone in the northern mid-latitudes in winter. 45 We find very good agreement between GOME-2A and GB observations over Canada and Europe as to their NAO-46 related variability, with mean differences reaching the $\pm 1\%$ levels. The agreement and small differences which were 47 found between the independently produced total ozone data sets as to the influence of QBO, ENSO and NAO show 48 the importance of these climatological proxies as additional tool for monitoring the long-term stability of satellite-49 ground truth biases.

50 1 Introduction

51 Ozone is an important gas of the Earth's atmosphere. In the stratosphere, ozone is considered as good ozone because 52 it absorbs ultraviolet-B radiation from the Sun thus it protects the biosphere from a large part of the Sun's harmful 53 radiation (e.g. Eleftheratos et al., 2012; Hegglin et al., 2015). In the lower atmosphere and near the surface, natural 54 ozone has an equally important beneficial role because it initiates the chemical removal of air pollutants from the 55 atmosphere such as carbon monoxide, nitrogen oxides and methane. Above natural levels however, ozone is 56 considered as bad ozone because it can harm humans, plants and animals. In addition, ozone is a greenhouse gas, 57 warming the Earth's surface. In both the stratosphere and the troposphere, ozone absorbs infrared radiation emitted 58 from Earth's surface, trapping heat in the atmosphere. As a result, increases or decreases in stratospheric or 59 tropospheric ozone induce a climate forcinginereases in tropospheric ozone lead to a warming of the Earth's surface 60 one is a greenhouse gas (Hegglin et al., 2015).

61 Ozone in the atmosphere can be measured by ground-based instruments, by balloons, aircraft and satellites and can 62 be calculated by chemical transport model (CTM) simulations. Measurements by satellites from space provide ozone 63 profiles and column amounts over nearly the entire globe on a daily basis (e.g. WMO, 2014). The three Global 64 Ozone Monitoring Experiment-2 (GOME-2) instruments carried on Metop platforms A, B and C serve this purpose. 65 The first was launched in-on 19 October 2006, the second in-on 19 September 2012 and the last one will be was 66 launched in-on 7 November 2018. The three GOME-2 instruments will provide unique long-term data sets of more 67 than 15 years (2007-2024) related to atmospheric composition and surface ultraviolet radiation using consistent 68 retrieval techniques (Hassinen et al., 2016). The GOME-2 off-line data is set to make a significant contribution 69 towards climate and atmospheric research while providing near real-time data for use in weather forecasting and air 70 quality forecasting applications (Hassinen et al., 2016).

71 Validation of satellite ozone measurements is performed with ground-based (GB) measurements as well as other

72 satellite instruments (Hassinen et al., 2016). Validation of GOME-2A total ozone for the period 2007-2011 was

- 73 performed by Loyola et al. (2011) and Koukouli et al. (2012). It was found that GOME-2 total ozone data agree at
- T4 the $\pm 1\%$ level with GB measurements and other satellite data sets (Hassinen et al., 2016). The consistency between

GOME-2A and GOME-2B total ozone columns, including a validation with GB measurements, was presented by Hao et al. (2014). An updated time series of the differences between GOME-2A and GOME-2B with GB observations can be found in Hassinen et al. (2016). The long-term stability of the two satellite instruments was also noted in that study. Both satellites are consistent over the Northern Hemisphere with negligible latitudinal dependence, while over the Southern Hemisphere there is a systematic difference of 1% between the two satellite instruments (Hassinen et al., 2016).

81 Chiou et al. (2014) compared zonal mean total column ozone inferred from three independent multi-year data records, namely, SBUV (v8.6) total ozone (MCPeters et al., 2013), GOME-type Total Ozone Essential Climate 82 83 Variable (GTO-ECV) (Coldewey-Egbers et al., 2015; Garane et al., 2018), and GB total ozone for the period 1996-84 2011. Their analyses were conducted for the latitudinal zones of 0-30° S, 0-30° N, 50-30° S, and 30-60° N. It was 85 found that, on average, the differences in monthly zonal mean total ozone vary between -0.3 and 0.8% and are well 86 within 1%. In that study it was concluded that despite the differences in the satellite sensors and retrievals methods, 87 the SBUV v8.6 and GTO-ECV data records show very good agreement both in the monthly zonal mean total ozone 88 and the monthly zonal mean anomalies between 60°S and 60°N. The GB zonal means showed larger scatter in the 89 monthly mean data compared to satellite-based records, but the scattering was significantly reduced when seasonal 90 zonal averages were analysed. The differences between SBUV and GB total ozone data presented in Chiou et al. 91 (2014) are well in agreement with Labow et al. (2013), who systematically compared SBUV (v8.6) total ozone data 92 with that measured by Brewer and Dobson instruments at various stations as a function of time, satellite solar zenith 93 angle, and latitude. The comparisons showed good agreement (within $\pm 1\%$) over the past 40 years with very small 94 bias approaching zero over the last decade. Comparisons with ozone sonde data showed good agreement in the 95 integrated column up to 25 hPa with differences not exceeding 5% (Labow et al., 2013).

96 The observed small biases (at the percentage level) between satellite and GB observations of total ozone, as have 97 been documented in the above studies, ensure the provision of accurate satellite ozone measurements. The high 98 accuracy and stability of the satellite instruments is essential for monitoring the expected recovery of the ozone layer 99 resulting from measures adopted by the 1987 Montreal protocol and its amendments (e.g., Zerefos et al., 2009; 100 Loyola et al., 2011; Solomon et al., 2016; de Laat et al., 2017; Kuttippurath and Nair, 2017; Pazmiño et al., 2018; 101 Stone et al., 2018; Strahan and Douglass, 2018). It is known that total ozone varies strongly with latitude and 102 longitude as a result of chemical and transport processes in the atmosphere. Total ozone also varies with season. 103 Seasonal variations are larger over middle and high latitudes and smaller in the tropics (e.g. WMO, 2014). On longer 104 time scales total ozone variability is related to large scale natural oscillations such as the Quasi-Biennial Oscillation (QBO) (e.g. Zerefos et al., 1983; Baldwin et al., 2001), the El Nino Southern Oscillation (ENSO) (e.g. Zerefos et al., 105 106 1992; Oman et al., 2013; Coldewey-Egbers et al., 2014), the North Atlantic Oscillation (NAO) (e.g. Ossó et al., 107 2011; Chehade et al., 2014) and the 11-year solar cycle (e.g. Zerefos et al., 2001; Tourpali et al., 2007; Brönniman et 108 al., 2013). Moreover, volcanic eruptions may also alter the thickness of the ozone layer (Zerefos et al., 1994; 109 Frossard et al., 2013; Rieder et al., 2013; WMO, 2014). These natural perturbations affect the background 110 atmosphere and consequently the distribution of the ozone layer. In this context, the study of the effect of known natural fluctuations in total ozone could serve as additional tool for evaluating the long-term variability of satellitetotal ozone data records.

113 The objective of the present work is to examine the ability of the GOME-2A total ozone data to capture the 114 variability related to dynamical proxies of global and regional importance such as the QBO, ENSO and NAO, in 115 comparison to GB measurements, other satellite data and model calculations. The variability of total ozone from 116 GOME-2A is compared with the variability of total ozone from the other examined data sets during these naturally-117 occurring fluctuations in order to evaluate the performance of GOME-2A to depict natural perturbations. The 118 analysis is performed in the frame of the validation strategy of GOME-2A data on longer time scales within the 119 project of EUMETSAT, AC SAF. The validation evaluation of GOME-2A data performed here includes the study 120 of monthly means of total ozone, the annual cycle of total ozone, the amplitude of the annual cycle [i.e., (max-121 min)/2], the relation with the QBO (zonal winds at the equator at 30 hPa), the relation with ENSO (correlation with 122 SOI) and the relation with the NAO (correlations with the NAO index in winter (DJF mean)).

123 The annual cycle describes regular oscillations in total ozone that occur from month to month within a year. In 124 general, month-to-month variations of total ozone are larger in middle and high latitudes than in the tropics. The 125 QBO dominates the variability of the equatorial stratosphere (~16-50 km) and is easily seen as downward 126 propagating easterly and westerly wind regimes, with a variable period averaging approximately 28 months. 127 Circulation changes induced by the QBO affect temperature and chemistry (Baldwin et al., 2001). ENSO and NAO 128 are naturally-occurring patterns or modes of atmospheric and oceanic variability, which orchestrate large variations 129 in climate over large regions with profound impacts on ecosystems (Hurrell and Deser, 2009). We present the level 130 of agreement between satellite-derived GOME-2A and GB total ozone in depicting natural oscillations like QBO, 131 ENSO and NAO, highlighting the importance of these climatological proxies to be used as additional tool for 132 monitoring the long-term stability of satellite-ground truth biases.

133 2 Data sources

134 The analysis uses GOME-2 satellite total ozone columns for the period 2007-2016. This data forms part of the 135 operational EUMETSAT AC SAF GOME-2/MetopA GDP4.8 data product provided by the German Aerospace 136 Center (DLR). The GOME-2 total ozone data have been monthly averaged on a 1°x1° latitude longitude grid. The 137 overview of the GOME-2A satellite instrument and of the GOME-2 atmospheric data provided by AC SAF can be 138 found in Hassinen et al. (2016).

To examine the natural variability of ozone on longer time scales, we have additionally analysed the GOME/ERS-2, SCIAMACHY/Envisat, GOME-2A, and OMI/Aura merged prototype level 3 harmonized data record (GTO-ECV, 1°x1°) for the period 1995-2016 (Coldewey-Egbers et al., 2015; Garane et al., 2018). This GTO-ECV ozone data product was generated and provided by DLR as part of the European Space Agency Ozone Climate Change Initiative (ESA O3 CCI) project. The ESA O3 CCI merged level-3 record, which is based on GOME/SCIAMACHY/GOME-2A/OMI level-2 data, was obtained using the GODFIT v3.0 retrieval algorithm.

145	More on ESA O3 CCI datasets can be found in the studies by Van Roozendael et al. (2012), Lerot et al. (2014),
146	Koukouli et al. (2015) and Garane et al. (2018).
147	Both datasets are compared with a combined TOMS/OMI/OMPS satellite total ozone data set constructed using data
148	from the Total Ozone Mapping Spectrometer (TOMS) on Nimbus 7 (1979-1993), TOMS on Meteor 3 (1991-1994),
149	TOMS on Earth Probe (1996-2005), the Ozone Monitoring Instrument (OMI) onboard the NASA Earth Observing
150	System (EOS) Aura satellite (2005-present) and data from the next generation Ozone Mapping Profiler Suite
151	(OMPS) nadir profiler instrument, launched in October 2011 on the Suomi National Polar-orbiting Partnership
152	(NPP) satellite (McPeters et al., 2015). The total ozone data are available at $1^{\circ}~x~1.25^{\circ}$ (TOMS) or $1^{\circ}~x~1^{\circ}$
153	(OMI/OMPS) resolution from https://acd-ext.gsfc.nasa.gov/anonftp/toms/ (last access: 15 June 2018). From these
154	data we constructed monthly mean total ozone data on a $5^{\circ} \ge 5^{\circ}$ grid. To account for known biases between the
155	instruments (e.g., Labow et al., 2013) we use the Solar Backscatter Ultraviolet (SBUV) version 8.6 Merged Ozone
156	Data Set (MOD) monthly zonal mean total ozone (https://acd-ext.gsfc.nasa.gov/Data_services/merged/index.html,
157	also see next paragraph; last access: 15 June 2018) as a reference. We adjust each instrument such that the zonal
158	mean in each 5°_{k} band averaged over the instrument lifetime matches the corresponding SBUV MOD zonal mean
159	average. Thus the inherent longitudinal variability is retained from the TOMS/OMI/OMPS measurements but any
160	latitude-dependent bias between the instruments is removed. With the exception of Meteor 3 TOMS in the northern
161	hemisphere, all offsets where were within 2% at low and mid-latitudes. Such a data set should not be used for long-
162	term trends but is sufficient for analyzing periodic variability such as QBO, ENSO and NAO. We used data for the
163	period 1995-2016. We note here that another long-term data set which has been analysed for QBO, ENSO, NAO
164	and other perturbations comes from the Multi-Sensor Reanalysis (Knibbe et el., 2014), but is not examined here.
165	In addition, we compare with satellite SBUV station overpass data from 1995 to 2016. The satellite data are based
166	on measurements from three SBUV-type instruments from April 1970 to the present (continuous data coverage from
167	November 1978). Even though the time series includes different versions of the SBUV instrument, the basic
168	measurement technique remains the same over the advancement of the instrument from the Backscatter Ultraviolet
169	(BUV) to SBUV/2 (Bhartia et al., 2013). Satellite overpass data over various ground-based stations are provided per
170	day from https://acd-ext.gsfc.nasa.gov/anonftp/toms/sbuv/MERGED/ (last access: 15 June 2018). These overpass
171	data are analogous to the SBUV MOD monthly zonal mean data previously mentioned. Both are constructed by first
172	filtering lesser quality measurements and then averaging data from individual satellites when more than one
173	instrument is operating. Monthly averages have been calculated by averaging the daily merged ozone overpass data
174	for stations listed in Supplement Table S1. Details about the data are provided by McPeters et al. (2013) and Frith et
175	al. (2014).
176	We also compare with GB observations of total ozone from a number of stations contributing to the World Ozone

Field Code Changed

Field Code Changed

Formatted: Superscript

Field Code Changed

and Ultraviolet Radiation Data Centre (WOUDC). The WOUDC data centre is one of six World Data Centres which 177 178

are part of the Global Atmosphere Watch programme of the World Meteorological Organization (WMO). The

179 WOUDC data centre is operated by the Meteorological Service of Canada, a branch of Environment Canada. In

180 total, we analysed total ozone daily summaries from 193 ground-based stations operating either Brewer, Dobson,

181	filter, SAOZ or microtops instruments. The GB total ozone measurements are available from the	
182	website https://woudc.org/archive/Summaries/TotalOzone/Daily_Summary/ (last access: 15 June 2018). The various	Field Code Changed
183	stations used in this study are listed in Table S1.	
184	We have also analysed simulations of total ozone from the global 3-D chemical transport model (CTM) Oslo CTM3	
185	(Søvde et al., 2012). The Oslo CTM3 has traditionally been driven by 3-hourly meteorological forecast data from	
186	the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecast System (IFS) model,	
187	whereas in this study we apply the OpenIFS model (<u>https://software.ecmwf.int/wiki/display/OIFS/</u>) (last access: 15	Field Code Changed
188	June 2018), cycle 38r1, which is an improvement from Søvde et al. (2012). Details on the model are given in Søvde	
189	et al. (2012). The Oslo CTM3 comprises both detailed tropospheric and stratospheric chemistry. Photochemistry is	
190	calculated using fast-JX version 6.7c (Prather, 2012), and chemical kinetics from JPL 2010 (Sander et al., 2011).	
191	Total ozone columns compare well with measurements and other model studies (Søvde et al., 2012 and references	
192	therein). The horizontal resolution of the model is 2.25° x 2.25°. We made use of used the global monthly mean total	
193	ozone columns for the period 1995-2016.	
194	To examine the OBO component on total ozone we made use of the monthly mean zonal winds at Singapore at 30	
195	hPa. The zonal wind data at 30 hPa were provided by the Freie Universität Berlin (FU-Berlin) at http://www.geo.fu-	Field Code Changed
196	berlin.de/met/ag/strat/produkte/gbo/gbo.dat (last access: 15 June 2018) (Naujokat, 1986). The impact of ENSO in	
197	the tropics was investigated by using the Southern Oscillation Index (SOI) from the Bureau of Meteorology of the	
198	Australian Government (http://www.bom.gov.au/climate/current/soi2.shtml) (last access: 15 June 2018). The	Field Code Changed
199	correlation between total ozone and the NAO index was mainly computed for the winter-mean (DJF) when the NAO	
200	amplitude is large (e.g. Hurrell and Deser, 2009), but it is also addressed in other seasons. Emphasis is given over	
201	Canada, Europe and the North Atlantic Ocean in winter. Use was made of the The principal component (PC)-based	
202	NAO index (DJF) which was provided by the Climate Analysis Section, NCAR, Boulder, USA (available	
203	at: <u>https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-pc-based</u>) (last access:	Field Code Changed
204	15 June 2018) was used. Total ozone variability is also related to dynamical variability, for example variability in	
205	tropopause height (e.g. Dameris et al., 1995; Hoinka et al., 1996; Steinbrecht et al., 1998). The impact of tropopause	
206	variability height variations on total ozone variability was examined by analyzing the tropopause pressure from the	
207	independently produced NCEP/NCAR (National Centers for Environmental Prediction/National Center for	
208	Atmospheric Research) reanalysis 1 data set computed on a 2.5° grid. The NCEP/NCAR reanalysis data were	
209	provided from the web site at <u>https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.tropopause.html</u> (last	Field Code Changed
210	access: 15 June 2018) (Kalnay et al., 1996).	
211	3 Results and discussion	

212 3.1 Monthly zonal means and annual cycle

213 Figure 1 compares monthly mean total ozone from GOME-2A and SBUV (v8.6) satellite overpass data for stations

shown in Table S1 (Supplement). The GOME-2A data were taken at a spatial resolution of $1^{\circ}x1^{\circ}$ around each of the

215 ground-based monitoring stations listed in Supplement Table S1 and then averaged over the tropics, middle and high 216 latitudes of both Hemispheres in 30° latitudinal zones to provide the large scale monthly zonal means for the 217 GOME-2A data. Accordingly, SBUV satellite overpass data were averaged over each geographical zone to provide 218 the large scale zonal means for the SBUV observations. Mean differences and standard deviations between GOME-219 2A and SBUV total ozone were found to be $+0.1 \pm 0.7\%$ in the tropics (0-30 deg.), about $+0.8 \pm 1.6\%$ in mid-220 latitudes (30-60 deg.), about $+1.3 \pm 2.2\%$ over the northern high latitudes (60-80 deg. N) and about $-0.5 \pm 2.9\%$ over 221 the southern high latitudes (60-80 deg. S). The differences were estimated as [GOME-2A - SBUV] / SBUV (%) 222 from January 2007 to December 2016. Small differences were also found between GOME-2A and GB 223 measurements (Figure 2 and Table 1), where here GB stations data have been averaged over each geographical zone 224 to provide the large scale zonal means for the GB measurements. Mean differences and standard deviations between 225 GOME-2A and GB total ozone were found to be $-0.7 \pm 1.4\%$ in the tropics (0-30 deg.), $+0.1 \pm 2.1\%$ in mid-latitudes 226 (30-60 deg.), $+2.5 \pm 3.2\%$ over the northern high latitudes (60-80 deg. N) and $0.0 \pm 4.3\%$ over the southern high 227 latitudes (60-80 deg. S). We remind that all estimates refer to the period between January 2007 and December 2016.

228 In summary, the largest differences between GOME-2A, SBUV (v8.6) and GB measurements are found over the 229 southern-northern high latitudes (60^o-80^o N) and the highest variability (standard deviation of the mean difference) is 230 observed over the latitude belt (60° - 80° S). In addition, these differences (especially at the high latitudes) can be 231 affected by the fact that not always the same days have been used for the construction of the monthly mean values 232 for the different datasets. In the tropics and mid-latitudes the respective differences are within $\pm 1\%$ or less, and the 233 results are in line with Chiou et al. (2014). Validation results were also presented by Loyola et al. (2011), Koukouli 234 et al. (2012), Coldewey-Egbers et al. (2015), Koukouli et al. (2015), updates of which are included in Hassinen et al. 235 (2016). Our results based on updated to 2017 data largely confirm those studies, pointing to the good performance of 236 GOME-2A when extending the period of record.

237 Next, we have studied the correlation between total ozone from GOME-2A and SBUV satellite data using linear 238 regression analysis for the period 2007-2016. The statistical significance of the correlation coefficients, R, was 239 calculated using the t-test formula for R with N-2 degrees of freedom, as used in Zerefos et al. (2018). The 240 regression model showed statistically significant correlations between the different datasets as follows: R = +0.99 in the tropics, mid-latitudes and the northern high latitudes and R = +0.940.97 in the southern high latitudes. All 241 242 correlation coefficients are highly statically significant (99.9% confidence level). In the long-term, statistically 243 significant correlation coefficients ($R \ge +0.94$) are also found between GOME-2A satellite and GB measurements 244 (Figure 2) despite the different type of instruments used to measure total ozone from the ground. The regression 245 parameters for the correlation coefficients shown in Figures 1 and 2 are provided in Table 2.

A large part of the strong correlations shown in Figures 1 and 2 is attributable to the seasonal variability of total ozone which is presented in Figure 3 for GOME-2A, SBUV and GB data. More specifically, Figure 3 shows the seasonal variations of total ozone from stations mean data, averaged per 10 degree latitude zones north and south. At high latitudes our analysis stops at 80 degrees. There is a very good agreement between the annual cycles of total ozone from the three datasets denoting the consistency of the satellite retrievals with GB observations. Similar 251 annual cycles are also found with the GTO-ECV ozone data (not shown). Similar consistency is also revealed for the 252 amplitudes of the annual cycles, computed as [(maximum value - minimum value)/2] in Dobson Units (DU). Figure 253 4 shows global maps of the amplitude of annual cycle of total ozone for the period 2007-2016 from GOME-2A 254 (upper left panel), GTO-ECV (upper right) and the TOMS/OMI/OMPS (lower left) satellite data. All maps are 255 plotted against the sine of latitude north and south in order to show areas according to their actual size. As can be 256 seen from Figure 4, the amplitude of annual cycle is less than 20 DU in the tropics, increasing as we move towards 257 middle and high latitudes up to 75 DU. Interestingly, there is pattern with small amplitude of annual cycle in the 258 southern mid-latitudes with values of about 10-15 DU, seen in Figure 4 as a blue curved line crossing the longitudes around 60 degrees south, the origin of which is attributed to the small annual variation which points to small seasonal 259 260 variations of total ozone in these parts. The seasonal increase in Antarctic is delayed by 2-3 months compared to the 261 north polar region. Only with the breakdown of the polar vortex in late spring, i.e. at a time when the poleward 262 transport over lower latitudes has already ceased, does a strong ozone influx occur in the Antarctic. With this delay 263 the amplitude of the seasonal variation stays much smaller poleward of 55-60° in the south than in the north (Dütsch, 264 1974). These features are consistent between all examined satellite data sets and are reproduced to a large extend by 265 the Oslo CTM3 model as well, except in the southern mid-latitudes where the model seems to underestimate the 266 observed annual cycle (Figure 4 lower right).

In summary, we find similar annual cycle and amplitude of annual cycle between total ozone from GOME-2A and the other examined total ozone data sets. The mean differences in the annual cycles of GOME-2A and SBUV satellite data are small in the tropics (0-30 deg.: 0.3 ± 2.4 DU), and increase as we move to mid-latitudes (30-60 deg.: 2.4 ± 4.4 DU) and higher latitudes (60-80 deg.: 1.7 ± 4.8 DU). These numbers are consistent with the ones found between GOME-2A and GB measurements (tropics: 1.1 ± 2.3 DU; mid-latitudes: 1.2 ± 5.1 DU; high latitudes: 5.1 ± 7.1 DU). In all latitude zones the correlation coefficients between the annual cycles of GOME-2A – SBUV and GOME-2A – GB data pairs were found to be greater than 0.9.

Before examining correlations with the large scale natural fluctuations QBO, ENSO and NAO, the mean annualcycle has been removed from the ozone data sets as described in the next section.

276 3.2 Correlation with QBO

277 We then studied how changes in dynamics affect the ozone columns in the atmosphere. The time series obtained 278 have been deseasonalised by subtracting the long-term monthly mean from each individual monthly mean value. 279 Ozone column variations for different latitude zones in the Northern and Southern Hemispheres have been 280 compared. Figure 5 compares total ozone deseasonalised anomalies (in % of the mean) from GOME-2A and SBUV 281 satellite retrievals in the tropics $(10^{\circ} \text{ N}-10^{\circ} \text{ S})$, sub-tropics $(10^{\circ}-30^{\circ})$ and mid-latitudes $(30^{\circ}-60^{\circ})$. The right panel of 282 Figure 5 shows the respective anomalies from GTO-ECV data. Mean differences between GOME-2A and SBUV 283 deseasonalised total ozone datamonthly zonal means between 60° N and 60° S are less than $\pm 0.5\%$ (Table 2). As can 284 een from Table 2 and Figure 5, there is a very good agreement between the GOME 2A, GTO ECV and SBUV 285 total ozone anomalies over the entire period of observations. The correlation coefficients between GOME 2A and SBUV are highly significant everywhere (30°-60°-N: +0.94; 10°-30°-N: +0.95; 10°-N-10°-S: +0.98; 10°-30°-S: 286

287 +0.93; 30°-60°-S: +0.87). The same stands when correlating the GTO ECV with SBUV deseasonalised data (30° 288 60°-N: +0.96; 10°-30°-N: +0.97; 10°-N-10°-S: +0.98; 10°-30°-S: +0.96; 30°-60°-S: +0.93).

289 The line with dots superimposed on the ozone anomalies in the middle panel of Figure 5 shows the equatorial zonal 290 winds at 30 hPa which were used as a proxy index to study the impact of QBO on total ozone. The general features 291 include a QBO signal in total ozone at latitudes between 10° N and 10° S which almost matches with the phase of 292 QBO in the zonal winds. At higher northern and southern latitudes there is a clear-phase shift in the QBO impact on total ozone. The impact of QBO is most pronounced in the tropics with amplitudes of +4% to -4% and it is less 293 294 pronounced in the sub-tropics and mid-latitudes. As such, sStrong positive correlations with the QBO are found in 295 the tropics (correlation between GOME-2A and OBO of about +0.77, t-test = 12.91) and weaker (usually of opposite 296 sign) less significant correlations are found at higher latitudes (about -0.15 in the northern and about -0.45 in the 297 southern extra tropics). Similar strong correlations in the tropics and weaker correlations in the extra 298 tropiescorrelation patterns with the QBO are found for the GTO-ECV, SBUV and GB data. These correlations 299 suggest that the variability that can be attributed to the OBO in the tropics is about 60%, and about 2% and 20% in 300 the northern and the southern extra tropics, respectively.

301 Table 3 summarizes the correlation and regression coefficients between total ozone and QBO at 30 hPa for the 302 different latitude zones and the different datasets. The numbers speak for themselves: for latitudes between 10° N 303 and 10º S correlations between total ozone from GOME-2A, GTO-ECV, SBUV, GB data and the QBO are all 304 positive. At latitudes between 10^e and 30^e the correlations turn to negative, in agreement with Knibbe et al. (2014) 305 results, who noted that moving from the tropics towards higher latitudes the regression estimates switch to negative 306 values at approximately 10° N and 10° S. The correlations with the QBO at 30 hPa remain negative up to 60° , a 307 consistent result among all our data sets, something also reported by Knibbe et al. (2014) with the MSR ozone data. 308 The correlation and regression coefficients between GOME-2A and QBO are fairly similar to those found between 309 SBUV and QBO, as well as among all data sets as seen in Table 3, despite the different periods of records.

310 These features are also evident in Figure 6 which compares GOME-2A (and GTO-ECV) satellite total ozone with 311 GB observations with respect to the QBO. Mean differences and standard deviations between GOME-2A and GB 312 and between GTO-ECV and GB deseasonalised total ozone data do not exceed one percent (Table 2). Again, 313 correlation coefficients between deseasonalised GOME-2A and deseasonalised GB data are highly significant in all 314 latitude zones (30°-60° N: +0.91 (slope=0.818, error=0.035, t-value=23.466, N=119); 10°-30° N: +0.91 315 (slope=0.786, error=0.033, t-value=23.529, N=119; 10° N-10° S: +0.94 (slope=0.973, error=0.034, t-value=28.449, 316 <u>N=109;</u> 10°-30° S: +0.87 (slope=0.864, error=0.044, t-value=19.659, N=119; 30°-60° S: +0.88 (slope=0.858, 317 error=0.043, t-value=19.854, N=119). The same stands for the correlations between GTO-ECV and GB data pairs 318 (30°-60° N: +0.94; 10°-30° N: +0.89; 10° N-10° S: +0.94; 10°-30° S: +0.87; 30°-60° S: +0.85). Our results are in 319 line with Eleftheratos et al. (2013) and Isaksen et al. (2014) who compared QBO-related ozone column variations 320 from the chemical transport model Oslo CTM2 with SBUV satellite data for shorter time periods. In summary, it has 321 been shown that GOME-2A depicts the significant effects of QBO on stratospheric ozone in concurrence with 322 SBUV and GB measurements. The instrument captures correctly the variability of ozone in the tropics and the midlatitudes, which is nearly in phase with the QBO in the tropics and out of phase in the northern and the southernmid-latitudes as have been shown by earlier studies (e.g. Zerefos, 1983; Baldwin et al., 2001).

325 3.3 Correlation with ENSO

Apart from the QBO, which affects the variability of total ozone in the tropics, an important mode of natural climate variability in the tropics is ENSO. To examine the impact of ENSO on total ozone in the tropics we first removed correlations with variability related to the QBO and the solar cycle, and then performed the correlation analysis with the SOI. The effect of the QBO was removed from the time series by using a linear regression model for the total ozone variations at each grid box, of the form:

(1)

(3)

331
$$D(t) = a0 + a1 * QBO(t) + residuals(t); 0 < t \le T$$

332 where D(t) is the monthly deseasonalised total ozone and t is the time in months with t=0 corresponding to the initial 333 month and t=T corresponding to the last month. The term a0 is the intercept of the statistical model. To model QBO 334 we made use of the equatorial zonal winds at 30 hPa. The term a_1 is the regression coefficient of QBO. The QBO 335 component was removed from the time series by using a phase lag with maximum correlation of 28 months (month 336 lag -14 to month lag 13). Then, the remainders from Eq. (1) have been analysed to study the correlations between 337 total ozone and SOI at each individual grid box. The QBO-related coefficients $\alpha 0$ and $\alpha 1$ of Eq. (1) for the 338 deseasonalized GOME-2A, GTO-ECV, TOMS/OMI/OMPS and Oslo CTM3 zonal mean data are presented in Table 339 3. Additional information for the regression coefficients $\alpha 1$ of QBO is provided in the Supplement Figure S1, which 340 shows the spatial distribution of the regression coefficients in latitude-longitude maps.

341 The residuals from Eq. (1) were then inserted in a second regression (Eq. 2) to account for the effect of solar cycle
 342 on total ozone, as follows:

343	$O_3(t) = \beta 0 + \beta 1 * F10.7(t) + residuals(t); 0 < t \le T$	(2)
-----	---	-----

344	where $\beta 0$ and $\beta 1$ are now the intercept and regression coefficients of solar cycle, respectively. To model the set	olar
345	cycle we used the 10.7 cm wavelength solar radio flux (F10.7) as a proxy, taken from the National Research Cou	ncil
346	and Natural Resources Can	<u>1ada</u>
347	at ftp://ftp.geolab.nrcan.gc.ca/data/solar flux/monthly averages/solflux monthly average.txt (last access	12
348	December 2018). We used the absolute values of F10.7. Latitude-longitude maps of the regression coefficients β	<u>1 of</u>
349	the solar cycle are presented in the Supplement Figure S2. We note that the global pattern of the regress	sion
350	coefficients of solar cycle from GOME-2A data matches well with what has been shown by Knibbe et al. (20)14)
351	with the reanalysis MSR data.	
352	The remainders from Eq. (2) were used in a third regression (Eq. 3) to study the correlations between total oz	one
353	and SOI at each individual grid box:	

354 $O_3(t) = c0 + c1 * SOI(t) + residuals(t); 0 < t \le T$ _____

where c0 and c1 are now the intercept and regression coefficients of ENSO, accordingly. Estimates of the regression coefficients c1 are shown in the Supplement Figure S3.

357 Figure 7 presents the correlations between SOI and total ozone from GOME-2A (upper left panel), GTO-ECV 358 (upper right) and TOMS/OMI/OMPS satellite data (bottom left), as well as between SOI and the Oslo model 359 simulations (bottom right). All four plots refer to the period 2007-2016. As can be seen from Figure 7 (upper left), 360 correlations of >0.3 between GOME-2A total ozone and SOI are found in the tropical Pacific Ocean at latitudes 361 between 25 deg. north and south. These correlations were tested as to their statistical significance in the period 362 2007-2016 using the t-test for R with N-2 degrees of freedom (as in Zerefos et al., 2018), and were found to be 363 statistical significant. A similar picture of correlation coefficients is also observed by the GTO-ECV and 364 TOMS/OMI/OMPS data. Both data sets show similar results as to the range of correlations (>0.3) in the tropical 365 Pacific for the common period of observations. Nevertheless, the spatial resolution is higher in the GOME-2A and 366 GTO-ECV (1x1 deg.) data than in the TOMS/OMI/OMPS (5x5 deg.) data, so the former data sets perform better 367 when looking at smaller space scales. We have to note here that in both maps there are larger areas with correlation 368 coefficients >0.3 in the southern part of the tropics than in the northern part. However, this was mostly observed 369 during the period 2007-2016. By examining the longer-term data record of the TOMS/OMI/OMPS data which 370 extend back to the 1979, we find symmetry in the pattern of correlations north and south of the equator in the 371 tropical Pacific Ocean (Figure A1 of Appendix A), which indicates that both sides of the tropical Pacific are affected 372 more or less in a similar way by El Niño/La Niña events. Finally, the Oslo CTM3 gives small correlations (<0.3) in 373 the tropical Pacific Ocean around the equator, except over the northern and southern subtropics where the model 374 compares better with the observations.

375 The small rectangle in Figure 7 corresponds to the South Pacific region $(10^{\circ}-20^{\circ} \text{ S}, 180^{\circ}-220^{\circ} \text{ E})$ and the blue cross 376 to the station Samoa (American Samoa; 14.25° S, 189.4° E), in which total ozone has been studied as for the impact 377 of ENSO after removing variability related to the annual cycle and the QBO. Figure 8 shows an example of the 378 ENSO impact on total ozone in the South Pacific Ocean. The upper panel shows the time series of total ozone 379 anomalies from GOME-2A, GTO-ECV and TOMS/OMI/OMPS satellite data together with the SOI (Figure 8a). 380 Comparisons of GOME-2A data with GTO-ECV data, SBUV overpass data and GB measurements at the station 381 Samoa are shown in Figure 8b. The dotted line shows the respective tropopause pressure anomalies from NCEP 382 reanalysis. All data sets point to the strong influence of ENSO on total ozone. Most evident is the strong decrease of 383 about 4% in 1997/98 which was caused by the strongest El Niño event in the examined period. A strong decrease is 384 also observed in the tropopause pressures by NCEP. Notable also is the strong La Niña event in 2010 which caused 385 total ozone to increase by about 4%. We calculate a strong correlation between total ozone from GTO-ECV and SOI of +0.620.66 (99% confidence level), which accounts for about 40% of the variability of total ozone over the 386 387 tropical Pacific Ocean when the annual cycle-and, QBO signal and solar cycle are removed. From the regression 388 with SOI we estimated an ENSO-related term from which we calculated the amplitude of ENSO in total ozone as 389 [maximum ozone - minimum ozone]/2. The amplitude of ENSO in total ozone was estimated to be 8.78.8 DU or 390 3.43.5% of the annual mean. This is comparable to the amplitude of annual cycle (7.7 DU or 3.0% of the mean) and 391 -3 times larger than the amplitude of QBO in this region (2.2 DU or 0.8% of the mean) and the amplitude of solar 392 cycle in this region (4.1 DU or 1.6% of the mean). These results are based on the GTO-ECV total ozone data.
393 Similar results were also found at the station Samoa from GB observations (i.e. correlation with SOI: +0.510.55, amplitude of ENSO: 7.67.7 DU or 3.0% of the mean, amplitude of annual cycle: 6.7 DU or 2.7% of the mean).
395 Statistics of total ozone such as mean, amplitude of annual cycle, amplitude of QBO, amplitude of solar cycle and amplitude of ENSO in total ozone over the selected areas are presented in Table 34. Both satellite, GB and model data show consistent results. It also appears that the station Samoa represents well the greater area in the Southern
398 Pacific as to the impact of ENSO.

399 Differences between GOME-2A and its data pairs in the southern Pacific Ocean are the order of $-0.2 \pm 1.0\%$ 400 between GOME-2A and TOMS/OMI/OMPS data, $-0.3 \pm 0.9\%$ between GOME-2A and GTO-ECV, and $-0.9 \pm 1.8\%$ 401 between GOME-2A and Oslo CTM3. Accordingly, differences at Samoa are: -0.6 ± 1.9% between GOME-2A and 402 GB data, $0.0 \pm 1.4\%$ between GOME-2A and GTO-ECV, and $-0.1 \pm 1.3\%$ between GOME-2A and SBUV. Despite 403 the small differences found, we note here that GOME-2A values in the last 4 years of Figures 8 and 9 slightly 404 deviate from the other data sets, and correlate weaker with SOI than the other years in the time series. For instance, 405 we estimate a drop in the correlation coefficient between GOME-2A and SOI at the station Samoa (+0.58 in the 406 period 2007-2012 and +0.47 in the period 2007-2016), which nevertheless does not alter the statistical significance 407 of the correlation.

408 From Figure 8 it also appears that there are high correlations with the tropopause height. The correlation coefficient 409 between the NCEP tropopause pressure and GOME-2A total ozone over the South Pacific Ocean is of the order of 410 +0.550.59 (Student's t-test statistics results: t-value = 7.115917.946, p-value <0.0001, N = 119). Accordingly, the 411 correlation with GTO-ECV ozone data is the order of +0.590.64 (t-value $= \frac{11.6707713.165}{11.6707713.165}$, p-value < 0.0001, N = 412 $\frac{259252}{259252}$ and with TOMS/OMI/OMPS the order of +0.520.58 (t-value = $\frac{9.4987410.913}{259252}$, p-value < 0.0001, N = 241). 413 The high correlation between the tropopause pressure and total ozone on interannual and longer time scales points to 414 the very strong link between these parameters. These links were already documented in the past (e.g. Steinbrecht et 415 al., 1998, 2001) and are verified with the GOME-2A data. At the same time a strong correlation is also evident 416 between tropopause pressure and SOI, again on interannual and longer time scales (R = +0.66, t-value = 417 14.2503613.825, p-value <0.0001, N = $\frac{264252}{2}$). The above results point to the strong impact of ENSO on the 418 tropical ozone column through the tropical tropopause; warm (El Niño) and cold (La Niña) events affect the 419 variability of the troppopuse which in turn affects the distribution of stratospheric ozone. In the tropics, where total 420 ozone is mainly stratospheric, as the tropopause moves to higher altitudes (lower pressure), the stratosphere is 421 compressed, reducing the amount of stratospheric (total) ozone. This happens during warm (El Niño) episodes. The 422 opposite phenomenon occurs during cold (La Niña) events when the tropopause height decreases (higher pressure) 423 and total ozone is then increased. These events can affect the long-term ozone trends in the tropics when looking at 424 time periods when strong El Niño and La Niña events occur at the beginning and the end of the trend period 425 respectively (Coldewey-Egbers et al., 2014).

Furthermore, in Figure 8 we have marked 7 stations in the greater South Asia region (35°-45° N, 45°-125° E) where
total ozone is anti-correlated with the SOI. Admittedly, these anti-correlations are weak (about -0.3) but we thought

428 worthwhile presenting the time series in these areas as well. Figure 9 shows the variability of total ozone after 429 removing seasonal and QBO related variations, over the South Asia region (upper panel) and over the 7 stations 430 averaged within this region (lower panel). As can be seen from this figure, the explained variance by ENSO is small, 431 not exceeding 9%. All correlations from the comparisons with the SOI are summarized in Table 5. In spite the small correlations with the SOI, the consistency between GOME-2A, GTO-ECV, TOMS/OMI/OMPS and Oslo CTM3 432 433 data anomalies is very high and their differences are within \pm 1%. Differences at the 7 stations in South Asia are as 434 follows: -1.3 \pm 2.4% between GOME-2A and GB data, -0.4 \pm 1.0% between GOME-2A, and GTO-ECV and -0.5 \pm 435 1.0% between GOME-2A and SBUV.

In summary, our findings indicate that GOME-2A captures well the disturbances in total ozone during ENSO events
with respect to satellite SBUV and GB observations. Our findings on the ENSO-related total ozone variations (low
ozone during ENSO warm events, high ozone during ENSO cold events, and magnitude of changes) are in line with
recent studies (e.g. Randel and Thompson, 2011; Oman et al., 2013, Sioris et al., 2014) included in the recent-2014
Ozone Assessment report (Pawson et al., 2014; WMO, 2014). Our results are also in agreement with Knibbe et al.
(2014) who showed negative ozone effects of El Niño between 25^o S and 25^o N, especially over the Pacific.

442 3.4 Correlation with NAO

443 The residuals from Eq. (3), free from seasonal, QBO, solar and ENSO related variations, were later used to study the 444 correlation between total ozone and NAO in winterThe residuals from Eq. (1), free from seasonal and QBO related variations, were also used to study the correlation between total ozone and NAO in winter (DJF mean). The results 445 446 are presented in Figure 10 which shows the correlation coefficients between total ozone and NAO index in winter 447 from the GOME-2A (upper left), GTO-ECV (upper right) and TOMS/OMI/OMPS satellite data (lower left), and the 448 Oslo CTM3 model calculations (lower right). Negative correlations between total ozone and NAO are presented 449 with blue colours while positive correlations with red colours. From Figure 10 (upper left) it appears that total ozone 450 is strongly correlated with NAO in many regions. Strong negative correlation coefficients are observed in the 451 majority of the northern mid-latitudes (R about -0.6) while positive correlations exist in the tropics and some 452 negative correlations in the southern mid-latitudes. These characteristics are observed in both GTO-ECV and 453 TOMS/OMI/OMPS datasets and are reproduced by the Oslo model as well, all for the common period 20082007-454 2016. The regression coefficients on these comparisons are presented in the Supplement Figure S4.

455 We note here that the results of the correlation analysis for the period 20082007-2016 were based on a relative small 456 sample of data from 10 winters(9 winters as DJF means) and therefore many of these correlation coefficients may 457 not be statistically significant. The statistical significance of the correlation coefficients in every grid box was tested only with the TOMS/OMI/OMPS data (Figure A2, Appendix A), which provided us the opportunity to calculate the 458 respective correlations using data for the whole period of record 1979-2016more data (37 winter means). It appears 459 460 that when extending the data back to the 1980's the negative correlations in the southern mid-latitudes in winter 461 disappear while the positive correlations in the tropics become weaker; yet the observed anti-correlation between total ozone and NAO index in the northern mid-latitude zone holds strong. The dotted line in the plot shows areas 462 463 with statistically significant correlation coefficients (99% confidence level). Indeed, in the long-term, statistically

464 significant correlations between total ozone and <u>the NAO index in the long termduring winter</u> are <u>mostly</u> found only
465 over the northern mid-latitudes <u>and the sub-tropics and not elsewhere</u>. <u>A small, statistically significant signal is also</u>
466 seen over Antarctica but it was not analysed further.

467 According to this finding we have restricted the analysis of NAO to the northern mid-latitudes. Rectangles (Figure 468 10, upper left) correspond to two regions in the North Atlantic, i.e., 35°-50° N, 20°-50° W and 15°-27° N, 30°-60° W 469 respectively, which were studied for the impact of NAO on total ozone after removing variability related to the 470 annual cycle and the QBO. In addition we have studied a number of stations in Canada, USA, and Europe 471 contributing ozone data to WOUDC, which are marked by red and green crosses in Figure 10. The red crosses refer 472 to the monitoring stations in Canada and the US, and the green crosses to the stations in Europe. In Figure 11 we present the times series of total ozone anomalies from GOME-2A, GTO-ECV and TOMS/OMI/OMPS satellite data 473 474 along with the NAO index in winter over the North Atlantic. Model calculations are shown as well. The dotted line 475 shows the respective tropopause pressure anomalies from NCEP reanalysis. Comparisons between GOME-2A, 476 GTO-ECV, SBUV (v8.6) overpass data and GB measurements over the various stations in Canada, USA and Europe 477 are shown in Figure 12.

478 The observed anomalies over the North Atlantic Ocean point to the strong influence of NAO on total ozone in 479 winter. Most evident is the strong increase in total ozone in 2010 of more than 8% particularly over 35°-50° N and 480 20°-50° W. This increase was accompanied by a strong increase in tropopause pressures. Both changes (in total 481 ozone and tropopause pressures) occurred under a strong negative phase of NAO, the strongest one in the past 20 482 years. We observe strong anti-correlations among total ozone and NAO index in winter (R = -0.720.74 over 35°-50° 483 N, 20°-50° W), which is statistically significant at the 99% confidence level. This anti-correlation suggests that about 484 50% of the variability of total ozone in winter is explained by NAO when the annual cycle-and QBO signal, QBO, solar cycle and ENSO signals are removed. Differences for GOME-2A and its data pairs are estimated to be $-0.7 \pm$ 485 486 1.1% between GOME-2A and TOMS/OMI/OMPS data, +0.1 ± 1.0% between GOME-2A and GTO-ECV, and -0.2 487 \pm 1.5% between GOME-2A and Oslo CTM3. From the regression with the NAO index we derived a NAO-related 488 term from which we calculated the amplitude of NAO in total ozone as [maximum ozone - minimum ozone]/2. The 489 amplitude of NAO over the North Atlantic region (35°-50° N, 20°-50° W) was estimated to be about 18-16.5 DU or 490 5.85.2% of the annual mean. This is about half of the amplitude of the annual cycle (which is ~37 DU or 11.7% of 491 the mean). These estimates are based on GTO-ECV data. Similar correlation and amplitude were also found with 492 GOME-2A, the combined TOMS/OMI/OMPS satellite data and the Oslo CTM3 model simulations.

A similar but opposite correlation is found over the southern part of the North Atlantic (15°-27° N, 30°-60° W).
Here, we estimate a significant correlation coefficient with NAO of +0.690.60, amplitude of NAO of about 9-7.2
DU (3-22.6% of the annual mean) and amplitude of annual cycle of about 16-15.8 DU (5.7% of the mean). Again, similar estimates are found with the GOME-2A and the TOMS/OMI/OMPS satellite data and reproduced by the model calculations as well. The annual mean total ozone and the amplitudes of annual cycle, QBO, solar cycle and NAO in total ozone over the studied regions in the North Atlantic are summarised in Table 46. Differences between

499 GOME-2A and GTO-ECV data at the southern part of North Atlantic are the order of $-0.6 \pm 0.7\%$. Differences with 500 the TOMS/OMI/OMPS data are estimated to be $-0.9 \pm 0.8\%$, and with the Oslo CTM3 $-0.1 \pm 0.7\%$.

- 501 The time series of total ozone anomalies and of the NAO index for the examined stations in Canada, USA and 502 Europe are presented in Figure 12. Table 5-7 presents the respective statistics. The correlation between total ozone 503 and the NAO index in winter after removing from ozone variability related to the annual cycle, QBO, solar cycle 504 and ENSO and the QBO is -0.440.40 (9590% confidence level). Again, a particular feature was the total ozone 505 increase in 2010 by 6% of the mean associated with the negative NAO phase. Noteworthy on this increase is the 506 consistency with the GB measurements and the satellite SBUV overpassing data, and in general the agreement found 507 between the variability of the tropopause pressures and total ozone. Differences between GOME-2A and GB data 508 are -1.0 \pm 1.8%. Accordingly we estimate differences of about -1.1 \pm 0.5% between GOME-2A and GTO-ECV data 509 and of about -1.3 ± 0.6% between GOME-2A and SBUV data. Table 5 indicates On the basis of GTO-ECV data we estimate that in Canada and USA, the amplitude of NAO in total ozone in winter is about $\frac{10-7}{2}$ DU (or $\frac{32.2}{2}$ % of the 510 511 mean), while it is higher over Europe estimated to be about 16.9 DU (or 52.7% of the mean). These numbers are 512 slightly smaller than the GOME-2A, GB and SBUV estimates, less than about one percent (Table 7).
- 513 The anti-correlation between total ozone column and the NAO index during winter also applies to southern Europe 514 and the Mediterranean. Following the study of Ossó et al. (2011) who reported a reversal in the correlation pattern 515 between NAO and total ozone from winter to summer in southern Europe, we have looked at the correlations during 516 summer as well. Figure 13 presents the comparisons for 21 ground-based stations located in the region bounded by 517 latitudes 30°-47° N and by longitudes 10°W-40°E. Figure 13a shows results for the summer and Figure 13b shows 518 results for winter. As evident, the observed anti-correlation between GB total ozone and NAO in winter (R= -0.43, 519 slope= -0.980, t-value= -2.095, p-value= 0.0499, N = 21) reverses sign and becomes positive in the summer (R= 520 +0.60, slope= 0.874, t-value= 3.309, p-value= 0.0037, N= 21), indicating that the NAO explains about 36% of ozone 521 variability in the summer in this region. A similar picture is also seen from GOME-2A, GTO-ECV and SBUV data.
- 522 In summary, our findings based on GOME-2A, GTO-ECV and SBUV overpass data are in line with those found by 523 Ossó et al. (2011) and Steinbrecht et al. (2011) who analysed TOMS and OMI satellite data, and GB measurements 524 at the station Hohenpeissenberg, respectively. During winter, total ozone variability associated with the NAO is 525 particularly important over northern Europe, the U.S. East Coast, and Canada, explaining up to 30% in total ozone 526 variance for this region (Ossó et al., 2011). Also, both studies found unusually high total ozone columns in 2010 527 over much of the Northern Hemisphere and related them to the negative phase of NAO or AO (the Arctic 528 Oscillation).

529 4 Conclusions

530 We have studied-evaluated the ability of GOME-2/MetopA (GOME-2A) satellite total ozone retrievals to capture

531 known natural oscillations such as the QBO, ENSO and NAO. In general, GOME-2A depicts these natural

oscillations in concurrence with GTO-ECV, TOMS/OMI/OMPS, SBUV (v8.6) satellite overpass data, ground-based
 measurements (Brewer, Dobson, filter and SAOZ) and chemical transport model calculations (Oslo CTM3).

534 Mean differences between GOME-2A and SBUV total ozone were found to be $+0.1 \pm 0.7\%$ in the tropics (0-30

535 deg.), about $+0.8 \pm 1.6\%$ in mid-latitudes (30-60 deg.), about $+1.3 \pm 2.2\%$ over the northern high latitudes (60-80

536 deg. N) and about $-0.5 \pm 2.9\%$ over the southern high latitudes (60-80 deg. S). These differences were estimated as

537 [GOME-2A - SBUV] / SBUV (%) from January 2007 to December 2016. Small differences were also found

538 between GOME-2A and GB measurements, with standard deviations of the differences being \pm 1.4% in the tropics,

539 \pm 2.1% in mid-latitudes, and \pm 3.2% and \pm 4.3% over the northern and the southern high latitudes respectively.

The variability of total ozone from GOME-2A has been compared with the variability of total ozone from other examined data sets as to their agreement to depict natural atmospheric phenomena such as the QBO, ENSO and NAO. First, we studied correlations between total ozone and the QBO after removing from the ozone data sets variability related to the seasonal cycle. Then, we examined correlations between total ozone, and ENSO and NAO, after removing variability related to the QBO and solar cycle, and finally correlations with the NAO after removing variability related to the OBO, solar cycle and ENSO. Our main results are as follows:

QBO: Total ozone from GOME-2A is well correlated with the Quasi-Biennial Oscillation (+0.8 in the tropics) in agreement with GTO-ECV, SBUV and GB data. The amplitude of QBO on total ozone maximizes around the equator and it is estimated to about 42.6% of the mean. Going from low to mid-latitudes there is a elear-phase shift in the QBO impact on total ozone. Correlation coefficients between GOME-2A total ozone and the QBO over 30-60 deg. north and south are -0.1 and -0.5 respectively, in agreement with the correlations between GB total ozone and the QBO (-0.2 and -0.5, accordingly). On the basis of GOME-2A, the amplitude of QBO in total ozone is estimated to be 0.6% of the mean in the northern mid-latitudes and 1.4% of the mean in the southern mid-latitudes.

ENSO: Correlation coefficients among GOME-2A total ozone and SOI in the tropical Pacific Ocean are estimated
to be about +0.6, consistent with GTO-ECV, SBUV and GB observations. It was found that the ENSO signal is
evident and consistent in all examined datasets. The amplitude of the El Nino Southern Oscillation in total ozone is
about 6–9 DU corresponding to about 2.5–3.5% of the annual mean. Differences between GOME-2A, GTO-ECV
and GB measurements during warm (El Niño) and cold (La Niña) events are within ±1.5%. Similar estimates also
result from the Dobson measurements at American Samoa, indicating that Samoa station represents well the greater
area in the Southern Pacific for satellite evaluations as to the impact of ENSO.

- NAO: The respective results as far as the impact of North Atlantic Oscillation over the northern mid-latitudes
 showed a clear NAO signal in winter in all data sets, with amplitudes of about <u>17–2016-19</u> DU (about 5–6% of the
 annual mean) in the North Atlantic, 9-12 DU (3-4% of the mean) over Europe, and 7-10 DU (2-3% of the mean)
 over Canada and the US. Comparison with GB observations over Canada and Europe showed very good agreement
 between GOME-2A, GTO-ECV and GB observations as to the influence by NAO, with differences within ±1%.
- Additionally to the usual validation methods, which compare monthly mean and zonal mean total ozone data and analyse the differences between satellite and GB instruments, we showed here that quasi cyclical perturbations such

as the QBO, ENSO and NAO can serve as independent proxies of spatiotemporal variation in validating_to
gualitatively evaluate_GOME-2A satellite total ozone against ground-based and other satellite total ozone data sets.
The agreement and small differences which were found between the variability of total ozone from GOME-2A and
the variability of total ozone from other satellite retrievals and ground-based measurements during these naturallyoccurring oscillations verify the good quality of GOME-2A satellite total ozone to be used in ozone-climate research
studies.

573 Data availability

574 Satellite SBUV (v8.6) total ozone station overpass data were downloaded from https://acd-575 ext.gsfc.nasa.gov/Data_services/merged/index.html (last access: 15 June 2018) (McPeters et al., 2013; Bhartia et al., 2013). GTO-ECV total ozone data are available at http://www.esa-ozone-cci.org/?q=node/160 (last access: 15 June 576 577 2018) (Coldewey-Egbers et al., 2015; Garane et al., 2018). Ground-based total ozone daily summaries were obtained 578 from the World Ozone and UV Data Centre (WOUDC) at https://woudc.org/archive/Summaries/TotalOzone/Daily_Summary/ (last access: 15 June 2018). The QBO 579 580 component on total ozone was examined by using the monthly mean zonal winds at Singapore at 30 hPa. Zonal 581 wind data at 30 hPa were provided by the Freie Universität Berlin (FU-Berlin) at http://www.geo.fu-582 berlin.de/met/ag/strat/produkte/qbo/qbo.dat (last access: 15 June 2018) (Naujokat, 1986). The Southern Oscillation 583 Index (SOI) was provided by the Bureau of Meteorology of the Australian Government 584 at http://www.bom.gov.au/climate/current/soi2.shtml (Australian Government - Bureau of Meteorology, 2018). The 585 NAO index for December, January and February was provided by the Climate Analysis Section, NCAR, Boulder, 586 USA at https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-pc-based (last 587 access: 15 June 2018) (Hurrell and Deser, 2009). The tropopause pressures from the NCEP/NCAR reanalysis 1 data 588 set were downloaded from https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.tropopause.html (last 589 access: 15 June 2018) (Kalnay et al., 1996).

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

590 Competing interests

591 The authors declare that they have no conflict of interest.

592 Acknowledgements

593 Development of the GOME-2/MetopA total ozone products and their validation has been funded by the AC SAF

594 project with EUMETSAT and national contributions. We further acknowledge the Mariolopoulos-Kanaginis

595 Foundation for the Environmental Sciences, the ESA Ozone CCI project and the NASA Goddard Space Flight

596 Centre. The ground-based data used in this publication were obtained as part of WMO's Global Atmosphere Watch

597 (GAW) and are publicly available via the World Ozone and UV Data Centre (WOUDC). The authors would like to

thank all the investigators that provide quality assured total ozone column data on a timely basis to the WOUDC

599 database. We acknowledge the National Oceanic and Atmospheric Administration (NOAA) for maintaining the

600 American Samoa Dobson station. KE and CS would like to dedicate the study to the memory of Professor Ivar

601 Isaksen (University of Oslo) who passed away on May 16th, 2017.

602 References

Australian Government – Bureau of Meteorology: Southern Oscillation Index (SOI) since 1986, available
 at http://www.bom.gov.au/climate/current/soi2.shtml, last access: 15 June 2018.

Field Code Changed

- 605 Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander,
- 606 M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnersley, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The
- 607 quasi-biennial oscillation, Rev. Geophys., 39, 179-229, doi: 10.1029/1999RG000073, 2001.
- 608 Bhartia, P. K., McPeters, R. D., Flynn, L. E., Taylor, S., Kramarova, N. A., Frith, S., Fisher, B., and DeLand, M.:
- Solar Backscatter UV (SBUV) total ozone and profile algorithm, Atmos. Meas. Tech., 6, 2533–2548, doi:
 10.5194/amt-6-2533-2013, 2013.
- 611 Brönnimann, S., Bhend, J., Franke, J., Flückiger, S., Fischer, A. M., Bleisch, R., Bodeker, G., Hassler, B., Rozanov,
- E., and Schraner, M.: A global historical ozone data set and prominent features of stratospheric variability prior to
- 613 1979, Atmos. Chem. Phys., 13 (18), 9623-9639, doi: 10.5194/acp-13-9623-2013, 2013.
- 614 Chehade, W., Weber, M., and Burrows, J. P.: Total ozone trends and variability during 1979-2012 from merged data
 615 sets of various satellites, Atmos. Chem. Phys., 14, 7059–7074, doi: 10.5194/acp-14-7059-2014, 2014.
- 616 Chiou, E. W., Bhartia, P. K., McPeters, R. D., Loyola, D. G., Coldewey-Egbers, M., Fioletov, V. E., Van
- 617 Roozendael, M., Spurr, R., Lerot, C., and Frith, S. M.: Comparison of profile total ozone from SBUV (v8.6) with
- 618 GOME-type and ground-based total ozone for a 16-year period (1996 to 2011), Atmos. Meas. Tech., 7, 1681–1692,
- 619 doi: 10.5194/amt-7-1681-2014, 2014.
- 620 Coldewey-Egbers, M., Loyola R., D. G., Braesicke, P., Dameris, M., van Roozendael, M., Lerot, C., and W.
 621 Zimmer, W.: A new health check of the ozone layer at global and regional scales, Geophys. Res. Lett., 41, 4363–
 622 4372, doi:10.1002/2014GL060212, 2014.
- 623 Coldewey-Egbers, M., Loyola, D. G., Koukouli, M., Balis, D., Lambert, J.-C., Verhoelst, T., Granville, J., van
- 624 Roozendael, M., Lerot, C., Spurr, R., Frith, S. M., and Zehner, C.: The GOME-type Total Ozone Essential Climate
- 625 Variable (GTO-ECV) data record from the ESA Climate Change Initiative, Atmos. Meas. Tech., 8, 3923–3940, doi:
- **626** 10.5194/amt-8-3923-2015, 2015.
- 627 Dameris, M., Nodorp, D., and Sausen, R.: Correlation between Tropopause Height Pressure and TOMS-Data for the
- 628 EASOE-Winter 1991/1992, Beitr. Phys. Atmosph., 68 (3), 227-232, 1995.

- de Laat, A. T. J., van Weele, M., and van der A., R. J.: Onset of stratospheric ozone recovery in the Antarctic ozone
 hole in assimilated daily total ozone columns, Journal of Geophysical Research: Atmospheres, 122, 11880-
- 631 <u>11899, https://doi.org/10.1002/2016JD025723, 2017.</u>
- 632 Dütsch, H. U.: The ozone distribution in the atmosphere, Can. J. Chem, 52, 1491-1504, 1974.
- Eleftheratos, K., Isaksen, I., Zerefos, C., Nastos, P., Tourpali, K., and Rognerud, B.: Ozone variations derived by a
 chemical transport model, Water, Air and Soil Pollution, 224:1585, doi: 10.1007/s11270-013-1585-2, 2013.
- 635 Eleftheratos, K., Isaksen, I. S. A., Zerefos, C. S., Tourpali, K., and Nastos, P.: Comparison of Ozone Variations from
- 636 Model Calculations (OsloCTM2) and Satellite Retrievals (SBUV), 11th International Conference on Meteorology,
- 637 Climatology and Atmospheric Physics (COMECAP 2012), Athens, Greece, 29 May 1 June 2012, C. G. Helmis
 638 and P. T. Nastos (eds.), Advances in Meteorology, Climatology and Atmospheric Physics, Springer Atmospheric
- 639 Sciences, DOI 10.1007/978-3-642-29172-2_132, © Springer-Verlag Berlin Heidelberg, pp. 945–950, 2012.
- Frith, S. M., Kramarova, N. A., Stolarski, R. S., McPeters, R. D., Bhartia, P. K., and Labow, G. J.: Recent changes
 in total column ozone based on the SBUV Version 8.6 merged ozone data set, J. Geophys. Res., 119, 9735–9751,
 doi: 10.1002/2014JD021889, 2014.
- Frossard, L., Rieder, H. E., Ribatet, M., Staehelin, J., Maeder, J. A., Di Rocco, S., Davison, A. C., and Peter, T.: On
 the relationship between total ozone and atmospheric dynamics and chemistry at mid-latitudes Part 1: Statistical
 models and spatial fingerprints of atmospheric dynamics and chemistry, Atmos. Chem. Phys., 13 (1), 147-164, doi:
 10.5194/acp-13-147-2013, 2013.
- Garane, K., Lerot, C., Coldewey-Egbers, M., Verhoelst, T., Koukouli, M. E., Zyrichidou, I., Balis, D. S., Danckaert,
 T., Goutail, F., Granville, J., Hubert, D., Keppens, A., Lambert, J.-C., Loyola, D., Pommereau, J.-P., Van
 Roozendael, M., and Zehner, C.: Quality assessment of the Ozone_cci Climate Research Data Package (release
 2017) Part 1: Ground-based validation of total ozone column data products, Atmos. Meas. Tech., 11, 1385-1402,
 doi:10.5194/amt-11-1385-2018, 2018.
- Hao, N., Koukouli, M. E., Inness, A., Valks, P., Loyola, D. G., Zimmer, W., Balis, D. S., Zyrichidou, I., Van
 Roozendael, M., Lerot, C., and Spurr, R. J. D.: GOME-2 total ozone columns from MetOp-A/MetOp-B and
 assimilation in the MACC system, Atmos. Meas. Tech., 7, 2937–2951, doi: 10.5194/amt-7-2937-2014, 2014.
- Hassinen, S., Balis, D., Bauer, H., Begoin, M., Delcloo, A., Eleftheratos, K., Gimeno Garcia, S., Granville, J.,
- 656 Grossi, M., Hao, N., Hedelt, P., Hendrick, F., Hess, M., Heue, K.-P., Hovila, J., Jønch-Sørensen, H., Kalakoski, N.,
- 657 Kauppi, A., Kiemle, S., Kins, L., Koukouli, M. E., Kujanpää, J., Lambert, J.-C., Lang, R., Lerot, C., Loyola, D.,
- 658 Pedergnana, M., Pinardi, G., Romahn, F., van Roozendael, M., Lutz, R., De Smedt, I., Stammes, P., Steinbrecht, W.,
- 659 Tamminen, J., Theys, N., Tilstra, L. G., Tuinder, O. N. E., Valks, P., Zerefos, C., Zimmer, W., and Zyrichidou, I.:
- 660 Overview of the O3M SAF GOME-2 operational atmospheric composition and UV radiation data products and data
- 661 availability, Atmos. Meas. Tech., 9, 383–407, doi: 10.5194/amt-9-383-2016, 2016.

- 662 Hegglin, M. I., Fahey, D. W., McFarland, M., Montzka, S. A., and Nash, E. R.: Twenty questions and answers about
- the ozone layer: 2014 update, Scientific Assessment of Ozone Depletion: 2014, 84 pp., World Meteorological
- 664 Organization, Geneva, Switzerland, ISBN: 978-9966-076-02-1, 2015.
- 665 Hoinka, K. P., Claude, H., and Köhler, U.: On the correlation between tropopause pressure and ozone above Central
- 666 Europe, Geophys. Res. Lett., 23 (14), 1753-1756, 1996.
- Hurrell, J. W., and Deser, C.: North Atlantic climate variability: The role of the North Atlantic Oscillation, Journal
 of Marine Systems, 78, 28-41, doi: 10.1016/j.jmarsys.2008.11.026, 2009.
- 669 Isaksen, I. S. A., Berntsen, T. K., Dalsøren, S. B., Eleftheratos, K., Orsolini, Y., Rognerud, B., Stordal, F., Søvde, O.
- A., Zerefos, C., and Holmes, C. D.: Atmospheric ozone and methane in a changing climate, Atmosphere, 5, 518–
 535, doi: 10.3390/atmos5030518, 2014.
- 672 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G.,
- 673 Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J.,
- 674 Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, Bulletin of the
- 675 American Meteorological Society, Vol. 77, No. 3, 437-472, 1996.
- Knibbe, J. S., van der A, R. J., and de Laat, A. T. J.: Spatial regression analysis on 32 years of total column ozone
 data, Atmos. Chem. Phys., 14, 8461-8482, https://doi.org/10.5194/acp-14-8461-2014, 2014.
- Koukouli, M. E., Balis, D. S., Loyola, D., Valks, P., Zimmer, W., Hao, N., Lambert, J.-C., Van Roozendael, M.,
 Lerot, C., and Spurr, R. J. D.: Geophysical validation and long-term consistency between GOME-2/MetOp-A total
 ozone column and measurements from the sensors GOME/ERS-2, SCIAMACHY/ENVISAT and OMI/Aura,
 Atmos. Meas. Tech., 5, 2169–2181, doi: 10.5194/amt-5-2169-2012, 2012.
- Koukouli, M. E., Lerot, C., Granville, J., Goutail, F., Lambert, J-C., Pommereau, J-P., Balis, D., Zyrichidou, I., Van
 Roozendael, M., Coldewey-Egbers, M., Loyola, D., Labow, G., Frith, S., Spurr, R., Zehner, C.: Evaluating a new
 homogeneous total ozone climate data record from GOME/ERS-2, SCIAMACHY/Envisat and GOME-2/MetOp-A,
 J. Geophys. Res. Atmos., 120, doi: 10.1002/2015JD023699, 2015.
- <u>Kuttippurath, J. and Nair, P. J.: The signs of Antarctic ozone hole recovery, Sci. Rep., 7,</u>
 <u>https://doi.org/10.1038/s41598-017-00722-7, 2017.</u>
- Labow, G. J., McPeters, R. D., Bhartia, P. K., and Kramarova, N.: A comparison of 40 years of SBUV
 measurements of column ozone with data from the Dobson/Brewer network, J. Geophys. Res. Atmos., 118, 73707378, doi:10.1002/jgrd.50503, 2013.
- 691 Lerot, C., Van Roozendael, M., Spurr, R., Loyola, D., Coldewey-Egbers, M. Kochenova, S., van Gent, J., Koukouli,
- 692 M., Balis, D., Lambert, J.-C., Granville, J., and Zehner, C.: Homogenized total ozone data records from the
- 693 European sensors GOME/ERS-2, SCIAMACHY/Envisat, and GOME-2/MetOp-A, J. Geophys. Res. Atmos., 119,
- 694 1639–1662, doi: 10.1002/2013JD020831, 2014.

- 695 Loyola, D. G., Koukouli, M. E., Valks, P., Balis, D. S., Hao, N., Van Roozendael, M., Spurr, R. J. D., Zimmer, W.,
- 696 Kiemle, S., Lerot, C., and Lambert, J.-C.: The GOME-2 total column ozone product: retrieval algorithm and ground-
- 697 based validation, J. Geophys. Res., 116, D07302, doi: 10.1029/2010JD014675, 2011.
- 698 McPeters, R. D., Bhartia, P. K., Haffner, D., Labow, G. J., and Flynn, L.: The version 8.6 SBUV ozone data record:
- 699 An overview, J. Geophys. Res., 118, 8032–8039, doi: 10.1002/jgrd.50597, 2013.
- 700 McPeters, R. D., Frith, S, and Labow, G. J.: OMI total column ozone: extending the long-term data record, Atmos.
- 701 Meas. Tech., 8, 4845-4850, soi:10.5194/amt-8-4845-2015, 2015.
- Naujokat, B., 1986: An update of the observed quasi-biennial oscillation of the stratospheric winds over the tropics,
 J. Atmos. Sci., 43, 1873-1877.
- Oman, L., Douglass, A., Ziemke, J., Rodriguez, J., Waugh, D., and Nielsen, J.: The ozone response to ENSO in
 Aura satellite measurements and a chemistry-climate simulation, J. Geophys. Res., 118 (2), 965-976, doi:
 10.1029/2012JD018546, 2013.
- 707 Ossó, A., Sola, Y., Bech, J., and Lorente, J.: Evidence for the influence of the North Atlantic Oscillation on the total
 708 ozone column at northern low latitudes and midlatitudes during winter and summer seasons, J. Geophys. Res., 116
 709 (D24), D24122, doi: 10.1029/2011JD016539, 2011.
- 710 Pawson, S., and Steinbrecht, W. (Lead Authors), Charlton-Perez, A. J., Fujiwara, M., Karpechko, A. Yu.,
- 711 Petropavlovskikh, I., Urban, J., and Weber, M.: Update on global ozone: Past, present, and future, V. E. Violetov
- 712 and U. Langematz (Eds), Chapter 2 in Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and
- 713 Monitoring Project Report No. 55, World Meteorological Organization, Geneva, Switzerland, 2014.
- Pazmiño, A., Godin-beekmann, S., Hauchecorne, A., Claud, C., Khaykin, S., Goutail, F., Wolfram, E., Salvador, J.,
 and Quel, E.: Multiplesymptoms of total ozone recovery inside the Antarctic vortex during austral spring, Atmos.
 Chem. Phys, 18, 7557–7572, 2018.
- Prather, M.: Fast-JX version 6.7c, available at: <u>ftp://halo.ess.uci.edu/public/prather/Fast-J/</u> (last access: 15 June 2018), 2012.
- Randel, W. J., and Thompson, A. M.: Interannual variability and trends in tropical ozone derived from SAGE II
 satellite data and SHADOZ ozonesondes, J. Geophys. Res., 116 (D7), D07303, doi: 10.1029/2010JD015195, 2011.
- 721 Rieder, H. E., Frossard, L., Ribatet, M., Staehelin, J., Maeder, J. A., Di Rocco, S., Davison, A. C., Peter, T., Weihs,
- 722 P., and Holawe, F.: On the relationship between total ozone and atmospheric dynamics and chemistry at
- 723 midlatitudes Part 2: The effects of the El Niño/Southern Oscillation, volcanic eruptions and contributions of
- atmospheric dynamics and chemistry to long-term total ozone changes, Atmos. Chem. Phys., 13 (1), 165-179, doi:
- 725 10.5194/acp-13-165-2013, 2013.
- 726 Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E.,
- 727 Kurylo, M. J., Moortgat, G. K., Orkin V. L., and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in

Field Code Changed

Atmospheric Studies, Evaluation No. 17, JPL Publication 10-6, Jet Propulsion Laboratory, Pasadena, 2011,
 (http://jpldataeval.jpl.nasa.gov; last access 15 June 2018), 2011.

Field Code Changed

- 730 Sioris, C. E., McLinden, C. A., Fioletov, V. E., Adams, C., Zawodny, J. M.m Bourassa, A. E., Roth, C. Z., and
- 731 Degenstein, D. A.: Trend and variability in ozone in the tropical lower stratosphere over 2.5 solar cycles observed
- 732 by SAGE II and OSIRIS, Atmos. Chem. Phys., 14, 3479-3496, doi: 10.5194/acp-14-3479-2014, 2014.
- 733 Søvde, O. A., Prather, M. J., Isaksen, I. S. A., Berntsen, T. K., Stordal, F., Zhu, X., Holmes, C. D., and Hsu, J.: The
- chemical transport model Oslo CTM3, Geosci. Model Dev., 5, 1441–1469, doi: 10.5194/gmd-5-1441-2012, 2012.
- Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Neely III, R. R., and Schmidt, A.: Emergence of healing in the
 Antarctic ozone layer, Science, 30, doi: 10.1126/science.aae0061, 2016.
- 737 Steinbrecht, W., Claude, H., Köhler, U., and Hoinka, K. P.: Correlations between tropopause height and total ozone:
- **738**Implications for long-term changes, J. Geophys. Res., 103 (D15), 19183-19192, 1998.
- Steinbrecht, W., Claude, H., Köhler, U., and Winkler, P.: Interannual changes of total ozone and Northern
 Hemisphere circulation patterns, Geophys. Res. Lett., 28, 1191-1194, 2001.
- Steinbrecht, W., Köhler, U., Claude, H., Weber, M., Burrows, J. P., and van der A, R. J.: Very high ozone columns
 at northern mid-latitudes in 2010, Geophys. Res. Lett., 38 (6), L06803, doi: 10.1029/2010GL046634, 2011.
- 743 <u>Stone, K. A., Solomon, S., and Kinnison, D. E.: On the identification of ozone recovery, Geophysical Research</u>
 744 <u>Letters, 45, 5158-5165, https://doi.org/10.1029/2018GL077955, 2018.</u>
- Strahan, S. E. and Douglass, A. R.: Decline in Antarctic Ozone Depletion and Lower Stratospheric Chlorine
 Determined From Aura Microwave Limb Sounder Observations, Geophys. Res. Lett., 45, 382–390,
 https://doi.org/10.1002/2017GL074830, 2018.
- Tourpali, K., Zerefos, C. S., Balis, D. S., and Bais, A. F.: The 11-year solar cycle in stratospheric ozone:
 Comparison between Umkehr and SBUVv8 and effects on surface erythemal irradiance, J. Geophys. Res., 112
 (D12), D12306, doi: 10.1029/2006JD007760, 2007.
- Van Roozendael, M., Spurr, R. J. D., Loyola, D., Lerot, C., Balis, D. S., Lambert, J.C., Zimmer, W., van Gent, J.,
 van Geffen, J., Koukouli, M., Doicu, A., and Zehner, C.: Sixteen years of GOME/ERS-2 total ozone data: The new
 direct-fitting GOME Data Processor (GDP) version 5 Algorithm description, J. Geophys Res., 117, D03305, doi:
 10.1029/2011JD016471, 2012.
- WMO (World Meteorological Organization), *Scientific Assessment of Ozone Depletion: 2014*, Global Ozone
 Research and Monitoring Project–Report No. 55, 416 pp., Geneva, Switzerland, 2014.
- Zerefos, C. S.: On the quasi-biennial oscillation in stratospheric temperatures and total ozone, Advances in Space
 Research, 2, 177–181, 1983.

- Zerefos, C. S., Bais, A. F., and Ziomas, I. C.: On the Relative Importance of Quasi-Biennial Oscillation and El
 Nino/Southern Oscillation in the Revised Dobson Total Ozone Records, J. Geophys. Res., 97 (D9), 10135–10144,
 1992.
- Zerefos, C., Contopoulos, G., and G. Skalkeas G. (Eds.): Twenty Years of Ozone Decline, Proceedings of the
 Symposium for the 20th Anniversary of the Montreal Protocol, Springer, Netherlands, Part of Springer Science +
 Decline, M. E. D. V. 170, and 101, 2160, 2000
- 764 Business Media B. V, 470 pp., ISBN: 978-90-481-2468-8, 2009.
- 765 Zerefos, C. S., Kapsomenakis, J., Eleftheratos, K., Tourpali, K., Petropavlovskikh, I., Hubert, D., Godin-Beekmann,
- 766 S., Steinbrecht, W., Frith, S., Sofieva, V., and Hassler, B.: Representativeness of single lidar stations for zonally
 767 averaged ozone profiles, their trends and attribution to proxies, Atmos. Chem. Phys., 18, 6427-6440,
- 768
 https://doi.org/10.5194/acp-18-6427-2018, 2018.
- Zerefos, C. S., Tourpali, K, and Bais, A. F.: Further studies on possible volcanic signal to the ozone layer, J.
 Geophys. Res., 99 (D12), 25741–25746, 1994.
- 771 Zerefos, C. S., Tourpali, K, Isaksen, I. S. A., and Schuurmans, C. J. E.: Long term solar induced variation in total
- ozone, stratospheric temperatures and the tropopause, Adv. Space Res., 27 (12), 1943–1948, 2001.
- 773







780 Figure A1. Map of correlation coefficients between total ozone from TOMS/OMI/OMPS satellite data and 781 SOI for the whole period 1979-2016, after removing variability related to the seasonal cycle-and-the, QBO 782 and solar cycle. The dotted line bounds the regions where the correlation coefficients are statistically 783 significant at the 99% confidence level (t-test). Only correlation coefficients above/below ±0.2 are shown. 784 Ozone data for the period 1991-1993 after the Mt Pinatubo eruption were not used in the correlation analysis to avoid any data contamination by the volcanic aerosols.



⁷⁸⁹

790 Figure A2. Map of correlation coefficients between total ozone from TOMS/OMI/OMPS satellite data and the 791 NAO index in-during winter (DJF meanDecember, January, February (D-J-F); upper left), spring (March, 792 April, May (M-A-M); upper right), summer (June, July, August (J-J-A); lower left) and autumn (September, 793 October, November (S-O-N); lower right) for the whole period 19801979-2016, after removing variability

794	related to the seasonal cycle-and the, QBO, solar cycle and ENSO. The dotted line bounds the regions where
795	the correlation coefficients are statistically significant at the 99% confidence level (t-test). Only correlation
796	coefficients above/below ±0.2 are shown. Ozone data for the period 1991-1993 after the Mt Pinatubo eruption
797	were not used in the correlation analysis to avoid any data contamination by the volcanic aerosols.

801

802 Table 1. Mean differences and their standard deviations in percent between total ozone from GOME-2A,

803 SBUV (v8.6) satellite overpass data and ground-based observations over different latitude zones, as shown in

804 Figures 1 and 2.

805

	[GOME-2A – SBUV] / SBUV (%)	[GOME-2A - GROUND] / GROUND (%)
	Stations mean data	Stations mean data
60°-80° N	$+1.3 \pm 2.2$	$+2.5 \pm 3.2$
30°-60° N	$+0.8 \pm 1.6$	$+0.1 \pm 1.9$
0°-30° N	0.0 ± 0.7	-0.5 ± 1.2
0°-30° S	$+0.1 \pm 0.7$	-0.9 ± 1.6
30°-60° S	$+0.9 \pm 1.6$	0.0 ± 2.4
60°-80° S	-0.5 ± 2.9	0.0 ± 4.3

806 807

808 Table 2. Statistics of the correlations shown in Figures 1 and 2 between total ozone from a) GOME-2A data 809

28.304

<u>21.575</u> _4.198

2.944

810

Tuble I Dutibueb 0	i the correlatio		igures i unu	- between t	otul ozone n		
and SBUV (v8.6) ov	erpass data, an	d b) GOME-2.	A data and g	round-based	measuremen	<u>nts.</u>	
(a) GOME-2A and	Correlation	Intercept	Slope*	Error	t-value	p-value	Ν
<u>SBUV (v8.6)</u>		(DU)					_
<u>60º-80º N</u>	+0.987	4.925	<u>0.999</u>	0.015	<u>65.224</u>	<u><0.0001</u>	<u>117</u>
<u>30º-60º N</u>	+0.984	<u>5.002</u>	<u>0.993</u>	0.017	<u>59.784</u>	<u><0.0001</u>	<u>118</u>

<u>0.894</u>

0.919

<u>1.023</u> 0.984

0.012

0.017

0.021

72.404

53.874 49.123 39.985

< 0.0001

< 0.0001

< 0.0001

< 0.0001

<u>118</u>

118

<u>118</u> <u>88</u>

811

<u>0°-30° N</u>

30º

<u>60°-80°</u>

-60

(b) GOME-2A and	Correlation	Intercept	Slope*	Error	t-value	p-value	N
Ground-based		<u>(DU)</u>					
<u>60°-80° N</u>	+0.973	<u>7.651</u>	<u>1.002</u>	0.022	<u>45.155</u>	<u><0.0001</u>	<u>118</u>
<u>30º-60º N</u>	<u>+0.977</u>	<u>15.772</u>	<u>0.952</u>	<u>0.019</u>	<u>49.671</u>	<u><0.0001</u>	<u>119</u>
<u>0°-30° N</u>	+0.982	<u>49.534</u>	<u>0.810</u>	<u>0.014</u>	<u>56.951</u>	<u><0.0001</u>	<u>119</u>
<u>0°-30° S</u>	+0.916	<u>56.520</u>	<u>0.778</u>	0.032	<u>24.655</u>	<0.0001	<u>119</u>
<u>30°-60° S</u>	+0.946	12.423	0.958	0.030	31.612	<0.0001	<u>119</u>
<u>60°-80° S</u>	+0.939	0.405	0.999	0.039	25.439	< 0.0001	<u>89</u>

812 * Error, t-value and p-value refer to slope.

+0.989

+0.981

+0.977

+0.974

- 813
- 814
- 815

818 Table 2. Mean differences and their standard deviations in percent between deseasonalised total ozone data
 819 from GOME-2A, SBUV (v8.6) satellite overpass data and ground based observations over different latitude
 820 zones, as shown in Figures 5 and 6.

	[GOME 2A SBUV] (%)	[GOME 2A GROUND] (%)
	Stations mean deseasonalized data	Stations mean deseasonalized data
30°-60°-N	-0.1 ± 0.7	-0.1 ± 0.9
10°-30°-N	-0.3 ± 0.5	-0.8 ± 0.8
10° N 10° S	$+0.1 \pm 0.6$	$+0.1 \pm 1.0$
10°-30°-S	0.0 ± 0.7	<u>-0.1 ± 0.9</u>
30°-60°- \$	-0.1 ± 1.0	-0.4 ± 1.0

2A data, b) GTO-E	<u>CV data, c) SBL</u>	JV (v8.6) over	pass data, an	d d) ground-	based measu	rements.	
(a) GOME-2A and	Correlation	Intercept	Slope*	Error	t-value	p-value	1
OBO		(%)					-
30°-60° N	-0.073	-0.045	-0.008	0.010	-0.791	0.4307	1
10º-30º N	-0.099	-0.048	-0.008	0.008	-1.077	0.2835	1
10° N-10° S	+0.767	0.654	0.114	0.009	12.910	< 0.0001	1
10º-30º S	-0.472	-0.273	-0.048	0.008	-5.799	< 0.0001	1
$30^{\circ}-60^{\circ}$ S	-0.424	-0.262	-0.046	0.009	-5.063	<0.0001	1
(b) GTO-ECV and	Correlation	Intercept	Slope*	Error	t-value	<u>p-value</u>	l
<u>QBO</u>		<u>(%)</u>					
<u>30°-60° N</u>	<u>-0.116</u>	<u>-0.090</u>	<u>-0.012</u>	<u>0.007</u>	<u>-1.869</u>	<u>0.0628</u>	2
<u>10º-30º N</u>	<u>-0.142</u>	<u>-0.100</u>	<u>-0.014</u>	<u>0.006</u>	<u>-2.293</u>	<u>0.0226</u>	2
<u>10º N-10º S</u>	<u>+0.779</u>	<u>0.705</u>	<u>0.109</u>	<u>0.005</u>	<u>19.949</u>	<u><0.0001</u>	2
<u>10º-30º S</u>	<u>-0.484</u>	<u>-0.306</u>	<u>-0.046</u>	<u>0.005</u>	<u>-8.873</u>	<u><0.0001</u>	2
<u>30°-60° S</u>	<u>-0.417</u>	<u>-0.312</u>	<u>-0.048</u>	<u>0.007</u>	<u>-7.345</u>	<u><0.0001</u>	2
<u>(b) SBUV v(8.6)</u>	Correlation	Intercept	<u>Slope*</u>	Error	<u>t-value</u>	<u>p-value</u>	<u>l</u>
and QBO		<u>(%)</u>					
<u>30º-60º N</u>	<u>-0.165</u>	<u>-0.112</u>	<u>-0.018</u>	<u>0.007</u>	<u>-2.694</u>	<u>0.0075</u>	2
<u>10º-30º N</u>	<u>-0.177</u>	<u>-0.114</u>	<u>-0.018</u>	<u>0.006</u>	<u>-2.901</u>	<u>0.0040</u>	2
<u>10º N-10º S</u>	<u>+0.748</u>	<u>0.648</u>	<u>0.104</u>	<u>0.006</u>	<u>18.223</u>	<u><0.0001</u>	2
<u>10^e-30^e S</u>	<u>-0.488</u>	<u>-0.287</u>	<u>-0.046</u>	<u>0.005</u>	<u>–9.037</u>	<u><0.0001</u>	2
<u>30°-60° S</u>	<u>-0.458</u>	<u>-0.328</u>	<u>-0.051</u>	<u>0.006</u>	<u>–8.333</u>	<u><0.0001</u>	2
(b) Ground-based	Correlation	Intercept	<u>Slope*</u>	Error	<u>t-value</u>	<u>p-value</u>	
and QBO	0.150	<u>(%)</u>	0.017	0.007	2.504	0.0100	-
<u>3060- N</u>	<u>-0.158</u>	<u>-0.123</u>	<u>-0.017</u>	0.007	<u>-2.594</u>	0.0100	<u></u>
$10^{2} - 30^{2}$ N	<u>-0.142</u>	<u>-0.083</u>	<u>-0.016</u>	0.007	<u>-2.317</u>	0.0213	2
10^{-} N-10 ⁻ S 10 ⁰ 20 ⁰ S	<u>+0.695</u>	0.253	0.095	0.006	<u>15.327</u>	<u><0.0001</u>	2
<u>10-30-5</u>	<u>-0.490</u>	<u>-0.208</u>	<u>-0.040</u>	0.005	<u>-9.091</u>	<u><0.0001</u>	2

Table <u>34</u>. Annual mean total ozone, amplitude of annual cycle, amplitude of QBO<u>, amplitude of solar cycle</u> and amplitude of ENSO in the period 1995 2016 from <u>GOME-2A</u>, GTO-ECV, the combined TOMS/OMI/OMPS satellite data and Oslo CTM3 model calculations over the South Pacific region
 (10°-20° S, 180°-220° E) and at station Samoa (14.25° S, 189.4° E) located within this region.

		South	Pacific Ocean			statio	n Samoa	
	<u>GOME-</u> <u>2A*</u>	GTO-ECV	TOMS/OMI/OMPS	Oslo CTM3	GOME-2A*	GTO-ECV	GROUND	SBUV <u>(v8.6)</u>
Annual mean	<u>255.3 DU</u>	254.7 DU	253.0 DU	259.5 DU	<u>252.7 DU</u>	252.2 DU	249.2 DU	251.9 DU
Amplitude of annual cycle	<u>7.4 DU</u> (2.9%)	7.7 DU (3.0%)	7.3 DU (2.9%)	5.2 DU (2.0%)	<u>7.1 DU</u> (2.8%)	6.7 DU (2.7%)	6.7 DU (2.7%)	7.3 DU (2.9%)
Amplitude of QBO	<u>2.7 DU</u> (1.0%)	2.2 DU (0.9%)	2.4 DU (0.9%)	2.3 DU (0.9%)	<u>3.0 DU</u> (1.2%)	2.2 DU (0.9%)	2.7 DU (1.1%)	2.0 DU (0.8%)
<u>Amplitude of</u> solar cycle	<u>2.1 DU</u> (0.8%)	<u>4.1 DU (1.6%)</u>	<u>4.6 DU (1.8%)</u>	<u>1.8 DU (0.7%)</u>	<u>2.0 DU</u> (0.8%)	<u>4.5 DU (1.8%)</u>	<u>1.6 DU (0.6%)</u>	<u>4.5 DU</u> (1.8%)
Amplitude of ENSO	<u>6.2 DU</u> (2.4%)	8.7<u>8.8</u> DU (3.4<u>3.5</u>%)	6.0 DU (2.4%)	8.9 <u>8.8</u> DU (3.4%)	<u>5.6 DU</u> (2.2%)	7.6<u>7.7</u> DU (3.0%)	5.7<u>5.5</u> DU (2.3<u>2.2</u>%)	7.6<u>7.5</u> DU (3.0%)

845 <u>*period 2007-2016</u>

(<u>10^o-20^o S, 180^o-220^o</u>	E), b) station	Samoa (14.25	^o S, 189.4 ^o E)	, c) South A	<u>sia (35º-45º l</u>	N, 45º-125º E)	and d)
stations in South Asi	<u>ia.</u>						
(a) South Pacific	Correlation	Intercept	Slope*	Error	t-value	p-value	Ν
<u></u>	with SOI	(%)					_
GOME-2A	+0.56	-0.238	<u>0.118</u>	0.016	7.236	<u><0.0001</u>	<u>119</u>
GTO-ECV	+0.66	-0.069	0.145	0.010	14.014	<0.0001	252
TOMS/OMI/OMPS	+0.62	-0.139	0.134	0.011	12.285	<0.0001	241
Oslo CTM3	+0.55	-0.064	0.144	0.014	10.501	<0.0001	252
Tropopause	<u>+0.66</u>	<u>-0.761</u>	<u>0.241</u>	<u>0.017</u>	<u>13.825</u>	<u><0.0001</u>	252
(b) Samoa	Correlation	Intercept	Slope*	Error	t-value	p-value	N
	with SOI	<u>(%)</u>	_			_	
GOME-2A	<u>+0.47</u>	<u>-0.217</u>	<u>0.108</u>	<u>0.018</u>	<u>5.823</u>	<u><0.0001</u>	<u>119</u>
GTO-ECV	+0.55	<u>-0.100</u>	<u>0.127</u>	0.012	<u>10.366</u>	< 0.0001	252
SBUV overpass	<u>+0.59</u>	<u>-0.114</u>	<u>0.127</u>	<u>0.011</u>	<u>11.398</u>	<u><0.0001</u>	<u>251</u>
<u>GB (WOUDC)</u>	+0.42	<u>-0.058</u>	<u>0.106</u>	<u>0.017</u>	<u>6.194</u>	<u><0.0001</u>	<u>178</u>
Tropopause	+0.65	<u>-0.799</u>	0.223	<u>0.017</u>	<u>13.405</u>	<u><0.0001</u>	252
(c) South Asia	Correlation	Intercept	Slope*	Error	<u>t-value</u>	<u>p-value</u>	<u>N</u>
	with SOI	<u>(%)</u>					
GOME-2A	<u>-0.23</u>	<u>0.090</u>	<u>-0.044</u>	<u>0.018</u>	<u>-2.525</u>	<u>0.0129</u>	<u>119</u>
GTO-ECV	<u>-0.30</u>	<u>0.073</u>	<u>-0.074</u>	<u>0.015</u>	<u>-5.047</u>	<u><0.0001</u>	<u>252</u>
TOMS/OMI/OMPS	<u>-0.28</u>	<u>-0.212</u>	<u>-0.073</u>	<u>0.016</u>	<u>-4.553</u>	<u><0.0001</u>	<u>241</u>
Oslo CTM3	<u>-0.18</u>	<u>0.140</u>	<u>-0.040</u>	<u>0.014</u>	<u>-2.877</u>	<u>0.0044</u>	<u>252</u>
Tropopause	<u>-0.27</u>	<u>-0.188</u>	<u>-0.129</u>	<u>0.029</u>	<u>-4.476</u>	<u><0.0001</u>	<u>252</u>
(d) South Asia (7	Correlation	Intercept	Slope*	<u>Error</u>	<u>t-value</u>	<u>p-value</u>	<u>N</u>
stations mean)	with SOI	<u>(%)</u>					
GOME-2A	<u>-0.23</u>	<u>0.090</u>	<u>-0.043</u>	<u>0.017</u>	<u>-2.518</u>	<u>0.0132</u>	<u>119</u>
GTO-ECV	<u>-0.30</u>	<u>0.067</u>	<u>-0.072</u>	<u>0.014</u>	<u>-5.040</u>	<u><0.0001</u>	252
SBUV overpass	<u>-0.27</u>	<u>0.086</u>	<u>-0.066</u>	<u>0.015</u>	<u>-4.464</u>	<u><0.0001</u>	251
CD (WOUDC)	0.26	0.427	_0.103	0.017	-5.912	<0.0001	240

Table 46. Annual mean total ozone, amplitude of annual cycle, amplitude of QBO, amplitude of solar cycle and amplitude of NAO in the period 1995-2016 from <u>GOME-2A</u>, GTO-ECV, the combined TOMS/OMI/OMPS satellite data and Oslo CTM3 model calculations over the North Atlantic Ocean:
(a) region 35°-50° N, 20°-50° W, and (b) region 15°-27° N, 30°-60° W.

	North Atlantic Ocean									
		(a) 35°	50° N, 20°-50° W		(b) 15°-27° N, 30°-60° W					
	<u>GOME-</u> <u>2A*</u>	GTO-ECV	TOMS/OMI/OMPS	Oslo CTM3	GOME-2A*	GTO-ECV	TOMS/OMI/OMPS	Oslo CTM3		
Annual mean	<u>319.7 DU</u>	315.9 DU	317.3 DU	311.2 DU	<u>276.6 DU</u>	276.4 DU	274.4 DU	282.6 DU		
Amplitude of annual cycle	<u>37.4 DU</u> (11.7%)	37.0 DU (11.7%)	36.9 DU (11.6%)	32.0 DU (10.3%)	<u>12.7 DU</u> (4.6%)	15.8 DU (5.7%)	15.1 DU (5.5%)	15.5 DU (5.5%)		
Amplitude of QBO	<u>2.5 DU</u> (0.8%)	2.3 DU (0.7%)	2.6 DU (0.8%)	3.2 DU (1.0%)	<u>3.0 DU</u> (1.1%)	2.8 DU (1.0%)	3.9 DU (1.4%)	4.3 DU (1.5%)		
Amplitude of solar cycle	<u>0.4 DU</u> (0.1%)	<u>0.3 DU (0.1%)</u>	<u>2.2 DU (0.7%)</u>	<u>2.3 DU (0.7%)</u>	<u>3.5 DU</u> (1.3%)	<u>2.7 DU</u> (1.0%)	<u>3.3 DU (1.2%)</u>	<u>1.0 DU</u> (0.3%)		
Amplitude of NAO (winter)	<u>18.3 DU</u> (5.7%)	18.3<u>16.5</u> DU (5.8<u>5.2</u>%)	17.5<u>18.4</u> DU (5.5<u>5.8</u>%)	20.3 <u>18.3</u> DU (6.5<u>5.9</u>%)	<u>4.2 DU</u> (1.5%)	8.8<u>7.2</u> DU (<u>3.22.6</u>%)	7.2<u>5.0</u> DU (<u>2.61.8</u>%)	8.0 DU (2.8%)		

867 <u>*period 2007-2016</u>

Table <u>57</u>. Annual mean total ozone, amplitude of annual cycle, amplitude of QBO, <u>amplitude of solar cycle</u> and amplitude of NAO in the period 1995-2016 from <u>GOME-2A</u>, GTO-ECV satellite data, ground-based observations and SBUV (v8.6) satellite overpass data over: (a) Canada and USA (11

873

stations mean), and (b) Europe (41 stations mean).

	(a) Canada and USA				(b) Europe				
		30°-50° N, 60°-110	^o W (11 stations me	ean)	$35^{\circ}-55^{\circ}$ N, 10° W- 40° E (41 stations mean)				
	GOME-2A*	GTO-ECV	GROUND	OUND SBUV <u>(v8.6)</u> GOME-2A		GTO-ECV	GROUND	SBUV <u>(v8.6)</u>	
Annual mean	<u>324.2 DU</u>	320.6 DU	322.5 DU	320.9 DU	<u>329.9 DU</u>	325.7 DU	326.9 DU	326.8 DU	
Amplitude of annual cycle	<u>38.1 DU</u> (11.7%)	34.1 DU (10.6%)	33.2 DU (10.3%)	34.0 DU (10.6%)	<u>39.3 (11.9%)</u>	40.5 DU (12.4%)	39.2 DU (12.0%)	40.7 DU (12.4%)	
Amplitude of QBO	<u>2.1 DU</u> (0.6%)	2.5 DU (0.8%)	3.5 DU (1.1%)	2.6 DU (0.8%)	<u>2.7 DU</u> (0.8%)	1.9 DU (0.6%)	2.8 DU (0.8%)	2.2 DU (0.7%)	
Amplitude of solar cycle	<u>0.3 DU</u> (0.1%)	<u>0.5 DU (0.2%)</u>	<u>1.4 DU (0.4%)</u>	<u>0.5 DU (0.2%)</u>	<u>2.1 DU</u> (0.6%)	<u>0.8 DU (0.2%)</u>	<u>1.0 DU (0.3%)</u>	<u>0.3 DU (0.1%)</u>	
Amplitude of NAO (winter)	<u>9.8 DU</u> (3.0%)	9.5<u>6.9</u> DU (<u>3.02.2</u>%)	10.2<u>8.7</u> DU (3.2<u>2.7</u>%)	11.1<u>9.3</u> DU (3.5<u>2.9</u>%)	<u>9.8 DU</u> (3.0%)	12.7<u>8.9</u> DU (3.9<u>2.7</u>%)	16.5<u>11.8</u> DU (<u>5.1<u>3.6</u>%)</u>	14.7<u>9.9</u> DU (4<u>.53.0</u>%)	

*period 2007-2016

<u>Table 8. Statistics of the comparisons between total ozone, tropopause pressures and NAO index in winter</u> (DJF mean) for a) the northern part of North Atlantic $(35^{\circ}-50^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its southern part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its part $(15^{\circ}-27^{\circ} \text{ N}, 20^{\circ}-50^{\circ} \text{ W})$, b) its part $(15^{\circ}-27^{\circ}-50^{\circ} \text{ W})$, b) its part $(15^{\circ}-27^{\circ}-50^{\circ} \text{ W})$. 883

<u>30°-60° W), c) 11 stations in Canada and USA, and d) 41 stations in Europe.</u>

(a) Northern part of	Correlation	Intercept	Slope*	Error	<u>t-value</u>	<u>p-value</u>	<u>N</u>
North Atlantic	with NAO in	<u>(%)</u>					
	winter						
GOME-2A	<u>-0.85</u>	<u>0.035</u>	<u>-2.474</u>	<u>0.568</u>	<u>-4.355</u>	<u>0.0033</u>	<u>9</u>
GTO-ECV	<u>-0.74</u>	<u>0.412</u>	<u>-2.188</u>	<u>0.453</u>	<u>-4.827</u>	<u>0.0001</u>	<u>21</u>
TOMS/OMI/OMPS	<u>-0.74</u>	<u>0.734</u>	<u>-2.386</u>	<u>0.538</u>	<u>-4.436</u>	<u>0.0004</u>	<u>18</u>
Oslo CTM3	<u>-0.75</u>	<u>0.639</u>	<u>-2.457</u>	<u>0.498</u>	<u>-4.937</u>	<u><0.0001</u>	<u>21</u>
Tropopause	<u>-0.83</u>	<u>0.665</u>	-3.112	<u>0.480</u>	<u>-6.478</u>	<u><0.0001</u>	<u>21</u>

$\Lambda O in \qquad (\%)$					
$\frac{1}{\sqrt{0}}$					
ter					
<u>-0.132</u>	<u>0.661</u>	<u>0.386</u>	<u>1.712</u>	<u>0.1306</u>	<u>9</u>
<u>60</u> <u>-0.202</u>	<u>1.097</u>	<u>0.333</u>	<u>3.291</u>	<u>0.0038</u>	<u>21</u>
<u>-0.334</u>	<u>1.138</u>	0.402	<u>2.832</u>	<u>0.0120</u>	<u>18</u>
<u>65</u> <u>-0.077</u>	<u>1.188</u>	<u>0.316</u>	<u>3.761</u>	<u>0.0013</u>	<u>21</u>
<u>-0.702</u>	<u>1.547</u>	<u>0.482</u>	<u>3.207</u>	<u>0.0046</u>	<u>21</u>
	$\begin{array}{c c} \underline{\text{ter}} \\ \hline 54 \\ \hline 50 \\ \hline -0.202 \\ \hline 58 \\ \hline -0.334 \\ \hline 55 \\ \hline -0.077 \\ \hline 59 \\ \hline -0.702 \\ \hline \end{array}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

(a) CA/USA (11	Correlation	Intercept	<u>Slope*</u>	Error	<u>t-value</u>	<u>p-value</u>	<u>N</u>
stations mean)	with NAO in	<u>(%)</u>					
	winter						
GOME-2A	<u>-0.71</u>	<u>-0.042</u>	<u>-1.305</u>	<u>0.493</u>	-2.647	<u>0.0331</u>	<u>9</u>
GTO-ECV	<u>-0.40</u>	<u>0.308</u>	<u>-0.904</u>	<u>0.479</u>	<u>-1.886</u>	<u>0.0746</u>	<u>21</u>
SBUV overpass	<u>-0.50</u>	<u>0.318</u>	<u>-1.209</u>	<u>0.476</u>	<u>-2.541</u>	<u>0.0199</u>	<u>21</u>
GB (WOUDC)	<u>-0.46</u>	<u>0.268</u>	<u>-1.046</u>	<u>0.477</u>	<u>-2.190</u>	<u>0.0419</u>	<u>20</u>
Tropopause	<u>-0.41</u>	0.268	<u>-0.739</u>	<u>0.377</u>	<u>-1.959</u>	<u>0.0650</u>	21

(b) Europe (41 stations mean)	Correlation with NAO in winter	Intercept (%)	<u>Slope*</u>	<u>Error</u>	<u>t-value</u>	<u>p-value</u>	<u>N</u>
GOME-2A	<u>-0.46</u>	0.089	-1.282	<u>0.897</u>	-1.428	0.1963	<u>9</u>
GTO-ECV	<u>-0.42</u>	0.315	<u>-1.141</u>	<u>0.573</u>	<u>-1.992</u>	0.0609	<u>21</u>
SBUV overpass	<u>-0.47</u>	<u>0.389</u>	-1.264	<u>0.543</u>	-2.329	<u>0.0311</u>	<u>21</u>
GB (WOUDC)	-0.48	0.625	-1.327	0.560	-2.368	0.0287	21
Tropopause	<u>-0.40</u>	0.048	<u>-0.989</u>	0.523	-1.891	<u>0.0739</u>	21

* The slope is in % per unit change of the explanatory variable. Error, t-value and p-value refer to slope,





896Figure 1. Monthly mean total ozone from GOME-2A as compared with monthly mean total ozone from897SBUV (v8.6) satellite overpass data for the period 2007-2016 over the Northern and the Southern Hemisphere898based on stations mean data. R is the correlation coefficient between the two lines. Error bars show the899standard deviation of each monthly mean. Mean differences $\pm \sigma$ are given as [GOME-2A – SBUV] / SBUV900(%).





907Figure 2. Same as in Figure 1 but for GOME-2A and GB observations. R is the correlation coefficient908between the two lines. Error bars show the standard deviation of each monthly mean. Mean differences $\pm \sigma$ 909are given as [GOME-2A – GROUND] / GROUND (%).



Figure 3. Comparison of the annual cycle of total ozone from GOME-2A with that from SBUV (v8.6) satellite overpass data and GB observations in the period 2007-2016 based on stations data averaged per 10 degree latitude zones. The annual cycle is distorted above 60 deg. S due to the Antarctic ozone hole.



Figure 4. Comparison of the amplitude [i.e., (max-min)/2] of the annual cycle of total ozone from GOME-2A
 (upper left) with the amplitude of the annual cycle of total ozone from GTO-ECV (upper right), the combined

925 TOMS/OMI/OMPS satellite data (lower left) and Oslo CTM3 model simulations (lower right).



- 930 Figure 5. (Left panel) Time series of deseasonalised total ozone from GOME-2A and SBUV (v8.6) satellite
- 931 overpasses over different latitude zones along with the equatorial zonal winds at 30 hPa as an index of the
- 932 QBO; (Right panel) same as in left panel but for GTO-ECV and SBUV. Values with red colour refer to the 933 mean differences $\pm \sigma$ (in %) between GOME-2A and SBUV deseasonalised data averaged over various
- WOUDC stations (150 stations in the northern mid-latitudes ($30^{\circ}-60^{\circ}$ N), 21 stations in the northern
- subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ} \text{ S}-10^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ} \text{ S}-10^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ} \text{ S}-10^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ} \text{ S}-10^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 8 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in the tropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics $(10^{\circ}-30^{\circ} \text{ N})$, 10 stations in southern subtropics
- 936 12 stations in the southern mid-latitudes (30°-60° S)). The QBO proxy is superimposed on the ozone
- 937 <u>anomalies.</u>
- 938



943 Figure 6. Same as in Figure 5 but for GOME-2A and GB observations (left panel), and for GTO-ECV and GB observations (right panel). The QBO proxy is superimposed on the ozone anomalies.







Figure 7. Map of correlation coefficients between total ozone and SOI for GOME-2A (upper left), GTO-ECV
(upper right), TOMS/OMI/OMPS satellite data (lower left) and Oslo CTM3 model simulations (lower right).
Rectangles correspond to the South Pacific region (10-20 °S, 180-220 °E) and South Asia region (35-45 °N, 45125 °E), blue cross to the station Samoa (14.25 °S, 189.4 °E) and red triangles to stations in South Asia, in
which total ozone has been studied as for the impact of ENSO after removing variability related to the annual
cycle and the, QBO and solar cycle. Positive correlations are shown by red colours while negative correlations
by blue colours. Only correlation coefficients above/below ±0.2 are shown.





958Figure 8. (a) Example of regional time series of total ozone (%) over the South Pacific region $(10^{\circ}-20^{\circ} \text{ NS},$ 959180°-220° E) along with SOI. The dotted line shows the respective tropopause pressure variability from960NCEP. R is the correlation coefficient between GTO-ECV total ozone and SOI (statistical significance of R is951given in parentheses). The difference refers to the mean difference $\pm \sigma$ (in %) between GTO-ECV and the962combined TOMS/OMI/OMPS satellite data. (b) Same as in (a) but for SBUV overpass and GB data at the963station Samoa. The difference refers to the mean difference $\pm \sigma$ (in %) between GTO-ECV and GB data.





967 Figure 9. (a) Example of regional time series of total ozone (%) over South Asia ($35^{\circ}-45^{\circ}$ N, $45^{\circ}-125^{\circ}$ E) along 968 with SOI. The dotted line shows the respective tropopause pressure variability from NCEP. *R* is the 969 correlation coefficient between GTO-ECV total ozone and SOI (statistical significance of *R* is given in 970 parentheses). The difference refers to the mean difference $\pm \sigma$ (in %) between GTO-ECV and the combined 971 TOMS/OMI/OMPS satellite data. (b) Same as in (a) but with SBUV overpass and GB data averaged at 7 972 stations in South Asia. The difference refers to the mean difference $\pm \sigma$ (in %) between GTO-ECV and GB 973 data.









978 Figure 10. Map of correlation coefficients between total ozone and the NAO index in winter (DJF 979 mean)during winter (December, January, February; D-J-F) for GOME-2A (upper left), GTO-ECV (upper 980 right), TOMS/OMI/OMPS satellite data (lower left) and Oslo CTM3 model simulations (lower right). Rectangles correspond to regions in the North Atlantic (35°-50° N, 20°-50° W; 15°-27° N, 30°-60° W), and red 981 982 and green crosses to stations in Canada/USA and Europe, in which total ozone has been studied as for the 983 impact of NAO after removing variability related to the annual cycle and the, QBO, solar cycle and ENSO. 984 Positive correlations are shown by red colours while negative correlations by blue colours. Only correlation 985 coefficients above/below ±0.2 are shown.





990Figure 11. Example of regional time series of total ozone (%) over the North Atlantic regions (a) $35^{\circ}-50^{\circ}$ N,991 $20^{\circ}-50^{\circ}$ W and (b) $15^{\circ}-27^{\circ}$ N, $30^{\circ}-60^{\circ}$ W in winter (DJF mean) along with the NAO index. The dotted line992shows the respective tropopause pressure variability from NCEP reanalysis. *R* is the correlation coefficient993between GTO-ECV total ozone and the NAO index. The differences refer to the mean differences $\pm \sigma$ (in %)994between GTO-ECV and the combined TOMS/OMI/OMPS satellite data.





999Figure 12. Comparison with GB observations over: (a) Canada and USA and (b) Europe in winter (DJF1000mean). R is the correlation coefficient between GTO-ECV total ozone and the NAO index. The differences1001refer to the mean differences $\pm \sigma$ (in %) between GTO-ECV and GB data.



