

Interactive comment on “A shape model of internally mixed soot particles derived from artificial surface tension” by H. Ishimoto et al.

Anonymous Referee #2

Received and published: 30 September 2018

This manuscript presents optical modeling of internally mixed soot particles, a subject that is of interest for atmospheric remote sensing. The modeling approach and computational techniques are sound and reasonable. The results are representative and semi-extensive, and thus should be useful in remote sensing analysis and relevant optical interpretation. I suggest the following revisions for the authors' consideration:

- 1) Is Eq. (1) necessary? Remove it, if it is unnecessary.
- 2) The discussion on the efficiency of computational methods only focuses on the shape aspect. Actually, the refractive index has large impact on the efficiency comparison between FDTD and DDA.
- 3) It might be better to have a table of the refractive indices at the 10 wavelengths.

Printer-friendly version

Discussion paper



4) A reference is required for the Maxwell-Garnett mixing rule.

5) The results are presented for single particles. It is unclear to obtain the size-averaged results from the simulated results. More discussion is required on the comparison between simulations and observations/measurements.

6) In summary, it might be better to summarize the new knowledge gained from the present modeling study.

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2018-249, 2018.

[Printer-friendly version](#)

[Discussion paper](#)

