

Interactive comments on “Technical note: Absorption aerosol optical depth components from AERONET observations of mixed dust plumes” by Sung-Kyun Shin et al.

Referee comments are noted in black. Our replies are given in blue.

We would like to thank all Referees for their constructive comments. Please find our point-by-point replies below. We have also attached a revised version of the manuscript with all changes marked.

Anonymous Referee #1

Manuscript touches important problem: separation the components of aerosol mixture characterized by different absorbing properties basing on AERONET measurements. Manuscript is clearly written, provides new useful results and is suitable for publishing in AMT.

Referees #3 and #4 provided extended comments, so I have just several notes.

For separation of dust and non-dust components authors use depolarization ratio recalculated from AERONET inversions. So I wonder if such separation can be done directly from spheroids volume fraction. I think authors should explain necessity of using lidar related characteristics.

We agree with the reviewer that new methods have to be compared with existing ones. However, we think that the suggested comparison of percentage of spherical particles (sphericity) to the contributions of dust/non-dust as derived using the methodology presented in our paper is not useful and would rather add confusion than clarification. Two points support our reasoning:

1. Sphericity is a retrieval parameter of the AERONET inversion rather than a physically meaningful quantity. As such, it is specific to the AERONET retrieval. In contrast to PLDR, it cannot be compared to independent measurements.
2. Sphericity is no longer included as an output parameter of the AERONET version 3 retrieval. For the data points considered in our study, we have collated version 3 PLDRs with version 2 sphericity to investigate a possible connection between the two parameters. Figure 1 shows that while low values of sphericity are generally connected to higher PLDRs, there is not clear relationship of decreasing PLDRs with increasing sphericity. In contrast, there is a clear link between PLDR and the contribution of non-spherical particles as documented in the literature.

The advantage of the new method is that it relies on physically meaningful parameters that can be observed through independent measurements. It is based solely on AERONET products, and thus, does not require co-located lidar measurements.




Figure 1: Correlation between sphericity (from AERONET version 2) and 1020-nm PLDR (from AERONET version 3) for level 2.0 data from Beijing, Cape Verde, and Dakar.

p.2 ln.10. BC as emitted from incomplete anthropogenic combustion or biomass burning is generally considered the main light absorber among atmospheric aerosols” Actually contribution of brown carbon to absorption can be also significant. I think authors should comment it.

In this study we use the terminology BC for carbon particles with the morphological and chemical properties typical of soot particles from combustion including a black, blackish or brown substance formed by combustion. In this sense we determined BC as more likely to be a primary source (without mixing). We acknowledge the potential contribution of brown carbon to aerosol absorption. However, we believe that the use of one absorbing aerosol type is sufficient to present the general idea of our methodology. We have revised our statement to:

“In dust-free conditions, BC as emitted from incomplete combustion involved in anthropogenic activities or biomass burning is generally considered the main light absorber among atmospheric aerosols (Bond and Bergstrom, 2006; Bond et al., 2013; Russell et al., 2010), and thus, the main contributor to non-dust AAOD. The term BC refers to carbon particles with the morphological and chemical properties typical of soot particles from combustion including a black, blackish or brown substance formed by combustion (Andreae and Gelencsér, 2006). We point out that the contribution of brown carbon (BrC) to aerosol absorption can also be significant. However, we opt for a single absorbing aerosol component as it allows us to present the general idea of our new methodology in a straightforward manner.”

Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131-3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.

p.4 ln.12. Authors assume depolarization of non-dust particles to be of 0.02. Actually depolarization ratio of smoke varies in a wide range and can exceed 10% (Butron et al, 2012). How such variations may influence results?

We thank the authors for this comment. We are aware of detection of PLDRs that are larger than 0.02. To obtain the value used in our study, we have repeated the investigation of Shin et al. (2018) for AERONET stations dominated by biomass-burning smoke. For clarification, we have added the following text:

“The latter value has been obtained from the analysis of δ derived at AERONET stations dominated by biomass-burning aerosols, analogous to the dust-focused study of Shin et al. (2018).”

Eq(5). Authors introduce the layer thickness “h” but looks like never use it later.

Correct. We have introduced this parameter for the sole purpose of resolving the connection between AOD (the columnar parameter provided by sun photometer) and extinction coefficient (the height-resolved parameter provided by aerosol lidar).

p.4.ln.28. “We take the values of 44 sr and 54 sr for Asian and Saharan dust, respectively”
Lidar ratios even for pure dust can vary in a wide range. This should be commented.

We thank the Referee for this comment. We are referring to the study of *Shin et al. (2018)* in which statistics of the lidar ratios at the different dust sources are presented. Nevertheless, we have revised our statement to:

“It varies according to the desert source and can cover a wide range even for pure dust. We take the mean values of 44 sr and 54 sr for Asian and Saharan dust, respectively from the AERONET-based study of Shin et al. (2018).”

Technical note: Absorption aerosol optical depth components from AERONET observations of mixed dust plumes

Sung-Kyun Shin¹, Matthias Tesche^{1,2}, Detlef Müller¹, and Youngmin Noh³

¹School of Physics, Astronomy and Mathematics, University of Hertfordshire, Hatfield, Hertfordshire, United Kingdom

²Leipzig Institute for Meteorology, Leipzig University, Leipzig, Germany

³Department of Environmental Engineering, Pukyong National University, Busan, Republic of Korea

Correspondence: Detlef Müller (d.mueller@herts.ac.uk)

Abstract. Absorption aerosol optical depth (AAOD) as obtained from sun/sky photometer measurements provides a measure of the light-absorbing properties of the columnar aerosol loading. However, it is not an unambiguous, aerosol-type specific parameter, particularly if several types of absorbing aerosols, for instance black carbon (BC) and mineral dust, are present in a mixed aerosol plume. The contribution of mineral dust to total aerosol light-absorption is particularly important at UV wavelengths. In this study we refine a lidar-based technique ~~for~~ **applied to** the separation of dust and non-dust aerosol types for the use with Aerosol Robotic Network (AERONET) direct sun and inversion products. We extend the methodology to retrieve AAOD related to non-dust aerosol (AAOD_{nd}) and BC (AAOD_{BC}). We test the method at selected AERONET sites that are frequently affected by aerosol plumes that contain a mixture of Saharan or Asian mineral dust and biomass-burning smoke or anthropogenic pollution, respectively. We find that aerosol optical depth (AOD) related to mineral dust as obtained with our methodology is frequently smaller than coarse-mode AOD. This suggests that the latter is not an ideal proxy for estimating the contribution of mineral dust to mixed dust plumes. We present the results of the AAOD_{BC} retrieval for the selected AERONET sites and compare them to coincident values provided in the Copernicus Atmospheric Monitoring System aerosol re-analysis. We find that modelled and AERONET AAOD_{BC} are most consistent for Asian sites or at Saharan sites with strong local anthropogenic sources.

1 Introduction

Atmospheric aerosols have a strong impact on the Earth's radiation budget and climate (Stocker *et al.*, 2013). The main interactions between atmospheric particles and the climate system are through scattering and absorption of radiation (direct effect) and through modification of the microphysical properties of clouds (indirect effect). Estimates of the aerosol radiative forcing, i.e. of the perturbation of radiant fluxes ~~due to~~ **by** aerosol particles, require information on aerosol loading as well as on the aerosol's optical and microphysical properties (Bellouin *et al.*, 2013). Aerosol optical depth (AOD) is the height integral of the aerosol extinction coefficient. It provides a measure of the columnar aerosol loading and is routinely obtained from ground-based and spaceborne remote-sensing observations. Despite ~~our~~ **the** unprecedented global coverage of atmospheric aerosol information, it is still challenging to assess the aerosol radiative effect accurately. Not only are the sources of aerosols, their lifetime and the processes that affect their optical and microphysical characteristics highly inhomogeneous in space and time

(Stocker *et al.*, 2013). Aerosol particles from different natural and anthropogenic sources also often mix **with each other** and undergo aging processes, which reflects in the optical and microphysical properties of the bulk aerosol. Better estimates of the aerosol radiative forcing require an improved consideration of the properties and contributions of the different aerosol types in mixed aerosol plumes.

5 Remote sensing measurements are an important way to obtain insight into optical and microphysical aerosol properties. For instance, ground-based AErosol RObotic NETwork (AERONET, Holben *et al.* 1998, 2001) sun/sky radiometers provide long-term observations of aerosol products including spectral AOD, particle size distribution, and complex refractive index for the atmospheric column even at remote locations. AERONET also provides absorption aerosol optical depth (AAOD) which is a measure of the column aerosol loading of light-absorbing particles such as black carbon (BC), carbonaceous aerosols or
10 mineral dust. However, AAOD becomes ambiguous if several types of absorbing aerosols are present in a mixed aerosol plume.

In dust-free conditions, BC as emitted from incomplete anthropogenic combustion **involved in anthropogenic activities** or biomass burning is generally considered the main light absorber among atmospheric aerosols (Bond and Bergstrom, 2006; Bond *et al.*, 2013; Russell *et al.*, 2010), and thus, the main contributor to non-dust AAOD. **The term BC refers to carbon particles with the morphological and chemical properties typical of soot particles from combustion including a black,**
15 **blackish or brown substance formed by combustion (Andreae and Gelencsér, 2006). We point out that the contribution of brown carbon (BrC) to aerosol absorption can also be significant. However, we opt for a single absorbing aerosol component as it allows us to present the general idea of our new methodology in a straightforward manner.**

Schuster *et al.* (2005) inferred columnar BC concentrations based on the Maxwell Garnett effective medium approximation with AERONET-retrieved complex refractive indices. Koven and Fung (2006) separated the absorption properties of BC from
20 the absorption of dust by exploiting the spectral absorption properties **that can be** inferred from the AERONET inversion. Russell *et al.* (2010) utilized AERONET-retrieved SSA, AAOD, and absorption Ångström exponent (AAE) as indicator to separate **classify observations with respect to** the contributions of BC, organic matter (OM), and mineral dust to the absorbing aerosol fraction.

Passive remote-sensing techniques can only provide the properties of the total aerosol mixture. Determining the optical
25 properties of a certain aerosol type in a mixed aerosol plume requires additional information. For instance, the Ångström exponent (AE or α , Ångström 1964) as inferred from spectral AOD measurements gives qualitative information on aerosol size that can be used for aerosol-type classification: **values greater than 2 indicate small particles such as biomass-burning smoke while values smaller than 1 indicate large particles like sea salt and mineral dust.** ~~and to infer the fine or coarse mode fraction in the aerosol size distribution.~~ Schuster *et al.* (2006) **found that the variation of the Ångström exponent is**
30 **associated with bimodal aerosol size distributions. The authors focused on the fine or coarse fraction of aerosols.** More detailed and quantitative information can be obtained from active aerosol remote sensing with lidar. In particular, the particle linear depolarization ratio (PLDR or δ) is an intensive parameter that is very sensitive to particle shape. It can be used to obtain the contribution of dust and non-dust particles to **the optical properties of** a mixed aerosol plume under the assumption that this plume consists of only those two aerosol types in an external mixture (Shimizu *et al.*, 2004; Tesche *et al.*, 2009b).
35 Burton *et al.* (2014) developed a generalised version of the methodology to separate contributions to mixtures of two aerosol

types while *Mamouri and Ansmann* (2014) further refined it to also separate between the contribution of fine and coarse dust particles.

In this study, we use AERONET version 3 level 2 products to refine the lidar-based aerosol-type separation methodology to resolve the contributions of dust and non-dust aerosol to the total and absorbing fractions of AOD. This is most useful over
5 and downwind of deserts where mineral dust can contribute significantly to AAOD – particularly at short wavelengths. We also propose a method to obtain the fraction of BC-related absorption to the non-dust AAOD. We describe our methodology in Section 2. In section 3, we present and discuss our results. We summarise our findings and provide concluding remarks in Section 4.

2 Data and methodology

10 2.1 AERONET sun/sky radiometer observations

AERONET (<http://aeronet.gsfc.gov>, *Holben et al.* 1998, 2001) operates automatic sun/sky radiometers for direct sun and sky radiation observation at sites all over the globe. AERONET instruments measure AOD at several wavelengths from 340 nm to 1640 nm **always including observations at 440, 670, 870, and 1020 nm**. The AOD uncertainty is estimated as 0.01 to 0.02 depending on wavelength in the absence of cloud contamination. The calibrated sky radiance measurements typically
15 have uncertainties below 5%. The Ångström exponent and the fine-mode fraction (FMF, *O'Neill et al.* 2003) are obtained from the spectral AOD measurements. **The level 2 product available from the AERONET portal includes inversion results**
~~The AERONET inversion is performed~~ for measurements with a 440-nm AOD larger than 0.4 (*Dubovik et al.*, 2006). **The AERONET inversion** It uses direct-sun and sky-radiance measurements at 440, 675, 870, and 1020 nm to infer columnar particle properties such as the volume size distribution, the complex refractive index, and the single-scattering albedo (SSA or ω). The uncertainty in SSA is expected to be of the order of 0.03 (*Dubovik et al.*, 2000). Knowledge of SSA ~~allows~~ **is used** to determine the fraction of AOD related to light absorption, referred to as absorption aerosol optical depth (AAOD) as:

$$AAOD = (1 - \omega) \times AOD. \quad (1)$$

Detailed descriptions of the instrumentation, calibration, methodology, data processing, and data quality assurance are provided in *Holben et al.* (1998, 2001), *Dubovik et al.* (2002, 2006), *Eck et al.* (2005) and *Giles et al.* (2018). The recently released
25 version 3 of the AERONET aerosol retrieval added spectral PLDRs and lidar ratios (S) to the list of inversion products. The representativeness of these values for pure mineral dust conditions has recently been discussed by *Shin et al.* (2018). *Noh et al.* (2017) investigated the reliability of the PLDR retrieved from AERONET sun/sky radiometer observations and found the strongest correlation between the 1020-nm PLDR inferred from AERONET data and the 532-nm PLDR from lidar observations. In this contribution we use AERONET version 3 level 2.0 inversion products inferred from observations of
30 mineral dust downwind of the Saharan and Asian deserts.

2.2 AOD and AAOD components in mixed dust plumes

In order to retrieve the AOD and AAOD for non-dust aerosols in mixed dust plumes, the optical properties of the mixture need to be separated according to the contributions of dust and non-dust particles, respectively. This is possible by using lidar measurements of the PLDR δ which depends mainly on the shape of the particles and their size with respect to the measurement wavelength. The PLDR is zero for spheres and increases with increasing particle non-sphericity. *Tesche et al.* (2009b) present a method to separate mixtures of Saharan dust and biomass burning particles while *Shimizu et al.* (2004) retrieved the contribution of dust and non-dust particles in plumes of Asian dust mixed with spherical particles. *Noh* (2014) expanded these methods to retrieve the fractional contribution of the different aerosol types in the mixture to the bulk measurements of SSA, as well as the SSA for dust (ω_d) and non-dust (ω_{nd}) particles.

While δ is measured directly with lidar, it can also be computed from AERONET data and has been included as a standard product in version 3 of the AERONET retrieval. For an external aerosol mixture, **this parameter** is used to calculate the contribution of dust (R_d) and non-dust (R_{nd}) to the particle backscatter coefficient following *Shimizu et al.* (2004) and *Tesche et al.* (2009b) as:

$$R_d = \frac{(\delta - \delta_{nd})(1 + \delta_d)}{(\delta_d - \delta_{nd})(1 + \delta)} \quad (2)$$

and

$$R_{nd} = 1 - R_d. \quad (3)$$

Here, δ_d and δ_{nd} indicate δ of dust and non-dust particles, respectively. Their values can be determined from lidar measurements (*Burton et al.*, 2014; *Freudenthaler et al.*, 2009) or from AERONET observations representative for pure mineral dust (*Shin et al.*, 2018). At the standard lidar wavelength of 532 nm, typical values are $\delta_d = 0.33$ and $\delta_{nd} = 0.02$ (*Freudenthaler et al.*, 2009; *Burton et al.*, 2014). *Shin et al.* (2018) recently discussed AERONET-derived δ_d for mineral dust from different source regions. They **authors** conclude that in general, values of δ at 870 and 1020 nm from the AERONET version 3 inversion product seem to be most reliable ~~when compared to the literature~~. **Their finding is based on values found in literature that reports** on lidar observations of mineral dust. We consequently apply the aerosol-type separation procedure to AERONET measurements at 1020 nm. ~~using~~ **We used** values of $\delta_d = 0.30$ ($\delta_d = 0.31$) for mixed Asian (Saharan) dust plumes (*Shin et al.*, 2018) and $\delta_{nd} = 0.02$. **The latter value has been obtained from the analysis of δ derived at AERONET stations dominated by biomass-burning aerosols, analogous to the dust-focused study of *Shin et al.* (2018).** When δ was lower than δ_{nd} or higher than δ_d , R_d was set to 0 or 1, respectively.

The ratios R_d and R_{nd} obtained from using δ refer to the lidar measurements in the backscatter direction (i.e. the scattering angle of 180°) and allow for inferring the dust-related backscatter coefficient β_d as:

$$\beta_d = \beta R_d. \quad (4)$$

This approach needs to be refined so that it can be also applied to sun/sky photometer measurements which provide information on total light extinction, i.e. AOD is the height integral of the extinction coefficient α , rather than the backscatter

coefficient. For a single aerosol layer of depth h , it can be expressed as $AOD = \alpha h$. The extinction coefficient is connected to β through the lidar ratio $S = \alpha/\beta$. Consequently, dust AOD can be expressed as:

$$AOD_d = S_d \beta_d h. \quad (5)$$

The use of Eq. (5) for the total aerosol and the dust fraction together with Eq. (4) leads to the dust and non-dust AOD as:

$$5 \quad AOD_d = AOD \times R_d \times \frac{S_d}{S} \quad (6)$$

and

$$AOD_{nd} = AOD - AOD_d. \quad (7)$$

AOD and S are the total AOD and lidar ratio of the aerosol mixture as provided by AERONET, respectively. The S_d is the AERONET-derived lidar ratio of pure dust particles. **It The lidar ratio varies according to the desert source and can cover a**
 10 **wide range even for pure dust.** We take the **mean** values of 44 sr and 54 sr for Asian and Saharan dust, respectively from **the**
AERONET-based study of Shin et al. (2018). As before, values at 1020 nm are used in the calculation.

To convert the 1020-nm AOD to other wavelengths λ , we use the Ångström exponent $\hat{a}_d = 0.06$ for pure Saharan dust (Tesche et al., 2009a). We obtain:

$$AOD_{d,\lambda} = AOD_{d,1020} \times \left(\frac{1020 \text{ nm}}{\lambda} \right)^{\hat{a}_d} \quad (8)$$

15 and

$$AOD_{nd,\lambda} = AOD_\lambda - AOD_{d,\lambda}. \quad (9)$$

The contributions of dust and non-dust aerosols to the total AOD can now be described by the extinction-related dust ratio χ as:

$$\chi_{d,\lambda} = \frac{AOD_{d,\lambda}}{AOD_\lambda} = R_d \frac{S_d}{S} \quad (10)$$

20 and

$$\chi_{nd,\lambda} = \frac{AOD_{nd,\lambda}}{AOD_\lambda} = 1 - R_d \frac{S_d}{S}. \quad (11)$$

This means that the contribution of mineral dust to the extinction coefficient decreases (increases) with respect to the contribution to the backscatter coefficient (i.e. to R_d) if the second aerosol type in the mixture has a lidar ratio larger (smaller) than that of mineral dust. Mixtures with absorbing aerosols will show total lidar ratios larger than that of pure dust, which means
 25 that in the cases considered here, χ_d is generally smaller than R_d . The total SSA of the mixed dust/pollution plume as **that is**
 provided by individual AERONET measurements is ~~now considered to be the result of mixing the SSA of dust and non-dust particles~~
can be calculated according to the following the mixing rule:

$$\omega_\lambda = \chi_{d,\lambda} \omega_{d,\lambda} + \chi_{nd,\lambda} \omega_{nd,\lambda}. \quad (12)$$

Re-arranging Eq. (12) gives the SSA related to non-dust particles

$$\omega_{nd,\lambda} = \frac{\omega_{\lambda} - \chi_{d,\lambda}\omega_{d,\lambda}}{\chi_{nd,\lambda}}. \quad (13)$$

The spectral SSA for pure dust particles is taken from the literature (see Table 1). The non-dust fraction to AAOD can now be derived as

$$AAOD_{nd,\lambda} = (1 - \omega_{nd,\lambda})AOD_{nd,\lambda}. \quad (14)$$

We can assume that the light-absorbing features of the non-dust part of the aerosol plume are caused primarily by BC. As it has been shown that BC is not an ideal light absorber, i.e., $\omega_{BC,\lambda} \neq 0$, (Bond and Bergstrom, 2006; Bond et al., 2013). Thus, we need to account for the SSA of BC to obtain the BC-related AAOD as:

$$AAOD_{BC,\lambda} = AOD_{nd,\lambda}(1 - \omega_{nd,\lambda})(1 - \omega_{BC,\lambda}) = AAOD_{nd,\lambda}(1 - \omega_{BC,\lambda}). \quad (15)$$

Bond and Bergstrom (2006) report on single-scattering albedos of 0.10 to 0.28 for fresh BC. Similar values for fresh BC have also been reported by Khalizov et al. (2009) and Cross et al. (2010). Here, we use values of $\omega_{BC,\lambda}$ from Haywood and Ramaswamy (1998). These values are provided together with the other input parameters in Table 1.

2.3 Connection between AAOD, AAOD_{nd} and AAOD_{BC}

Substituting Eq. (12) in Eq. (1) leads to the equation for the AAOD (of dusty mixtures) that accounts for the contribution of the different components as:

$$AAOD = (1 - (\chi_{d,\lambda}\omega_{d,\lambda} + \chi_{nd,\lambda}\omega_{nd,\lambda}))AOD. \quad (16)$$

The connection between total and non-dust AAOD for non-dust components with different values of ω_{nd} between 0.90 and 0.96, an ω_d of 0.98, and a total AOD of unity is presented in Figure 1. In case of $\chi_d = 1$, all absorption is due to caused by mineral dust. As the contribution of dust to the mixture decreases, the overall AAOD increases as a result of the stronger absorption of the non-dust particles. The ratio between AAOD_{nd} and total AAOD in Figure 1 changes linearly with χ_d in case of equal values of ω_{nd} and ω_d . The relation becomes increasingly non-linear with increasing difference in the absorbing properties of the dust and non-dust particles. This means that total AAOD as provided by AERONET for dusty mixtures is likely to represent the non-dust component at larger wavelengths, where dust is less absorbing, while its interpretation is less ambiguous at shorter wavelengths.

The approach described above assumes that BC is the major absorber in mixtures of non-dust aerosols. Because ω_{BC} is not zero, it is obvious from Eq. (15) that AAOD_{BC} is always smaller than AAOD and vanishes as AAOD_{nd} disappears, i.e. for $\chi_d = 1$.

2.4 CAMS aerosol re-analysis

We use the European Centre for Medium-range Weather Forecast's (ECMWF) Copernicus Atmospheric Monitoring Service (CAMS) aerosol re-analysis data (Inness et al., 2013) to assess the results of the AAOD_{BC} retrieval methodology. The CAMS

re-analysis assimilates satellite data into a data assimilation system and global model to correct for model departures from observational data (Bellouin *et al.*, 2013; Inness *et al.*, 2013). The re-analysis data provides not only total AOD at 469, 550, 670, 865, and 1240 nm but also the AOD of five aerosol species: mineral dust, sea salt, sulphate, BC, and OM at 550 nm. Mineral dust and sea salt are being separated into three different size classes each, and BC and OM are distinguishable by
5 according to their hydrophilic and hydrophobic properties (Bellouin *et al.*, 2013).

3 Results

3.1 AERONET statistics

For this study, we have selected AERONET sites downwind of the major dust sources in Africa and Asia. We will refer to the two regions as Saharan and Asian for the remainder of this work. Details on the stations are provided in Table 2. An
10 overview of the mean AOD and PLDR at 1020 nm as well as the FMF for the two regions are provided in the histograms in Figure 2 and in Table 2. While both regions show comparably similar features in the histograms of AOD (with larger mean values for Saharan stations), there is a clear difference in the distribution and mean values of PLDRs: Saharan stations most of the time show values above 0.25 while values below 0.15 form the majority of observations at Asian stations. The latter also show a considerable number of cases (30%) with $\delta_{1020} < 0.02$, for which we assume that dust is completely absent. The
15 distribution of δ_{1020} is directly related to the contribution of mineral dust at the respective sites which is also reflected in the FMF. Most observations at Saharan sites show $FMF < 0.2$ with highest values of 0.4 while the observations at Asian sites show a broad distribution across all possible values with peaks at 0.3 and 0.5. Overall, the two regions allow for assessing the methodology proposed here in situations dominated by mineral dust (Saharan) as well as in dusty mixtures with a broad range of dust/non-dust mixing ratios **Saharan and Asian**.

20 ~~FFigure 3 shows the effect of the different dust contributions is also apparent~~ in the histograms of extinction and absorption Ångström exponents for the two regions in Figure 3. An absorption Ångström exponent close to unity is the theoretical value for black carbon (Bergstrom, 1973; Bohren and Huffman, 1983) while higher values of 1.5 have been associated with biomass burning and those exceeding 2.0 represent an increasing contribution of mineral dust (Bond *et al.*, 2013). Due to the dominance of mineral dust, Saharan observations show a weak spectral dependence of AOD while a broad range of values between 1 and
25 4 is found for the absorbing Ångström exponent in , see Figure 3b. Similar values between 1.5 and 3.5 have been reported by Russell *et al.* (2010) for Arabian and Saharan dust. The large absorbing Ångström exponents result from the strong spectral dependence of the absorbing properties of mineral dust (Müller *et al.*, 2009; Petzold *et al.*, 2009). This effect is also reflected in the spectral variation of the single-scattering albedo (not shown). The observations at Asian sites show a higher extinction Ångström exponent **that peaking** at 1.0 to 1.25 and a lower absorption Ångström exponent with a maximum between 1.0 and
30 1.5. Consequently, this leads to a less pronounced spectral dependence of the single-scattering albedo (not shown). Figure 3 confirms the first impression provided by Figure 2 regarding the different contribution of mineral dust to the total AOD in the two regions.

The dust ratio χ_d as derived using Eq. (10) for the observations in the two regions is presented in Figure 4. The general shape of the histograms of χ_d resembles that of δ_{1020} in Figure 2b. The crucial difference is that PLDR marks a proxy of the contribution of mineral dust to the **lidar** measurement of the backscatter coefficient ~~as performed with lidar~~ while χ_d quantifies the contribution of mineral dust to the AERONET sun/sky photometer measurement of columnar AOD. The large occurrence rate of χ_d of zero and unity refers to observations of δ_{1020} below and above the thresholds for non-dust and dust particles, respectively. Figure 4 reveals an occurrence rate of 47% and 4% for pure dust conditions for Saharan and Asian sites, respectively, when considering cases with $\chi_d > 0.9$ as pure dust. It also shows that situations with dust contributions below 50% are rare for the Saharan stations while they are most common for the Asian sites. This suggests that the selected data set includes a wide spread of situations for testing the methodology proposed here.

A closer view on the relationship between δ_{1020} and χ_d is provided in Figure 5. The figure shows the spread of χ_d that is introduced when transforming the simple theoretical relationship of Eq. (2) for lidar backscatter measurements (*Shimizu et al.*, 2004; *Tesche et al.*, 2009b) to extinction data by means of Eq. (10). Depending on the value of the total lidar ratio for the aerosol mixture with respect to the reference value for pure dust conditions (Table 1), χ_d is either increased or decreased with respect to R_d . Figure 5 shows that χ_d is almost exclusively larger than R_d for observations at Asian sites as the majority of AERONET-derived values of S is smaller than the reference value for Asian dust ~~presented by~~, see *Shin et al.* (2018) (not shown). The same is the case for the Saharan observations with $\delta_{1020} < 0.2$ while above that value, χ_d is spread evenly to both sides of R_d . The latter **feature** is related to the fact that the frequency distribution of S for the Saharan observations peaks around the value for pure Saharan dust of 54 sr (not shown) and the generally larger occurrence rate of pure-dust cases used to define the reference value in *Shin et al.* (2018). When considering the effect of FMF (not shown), we find that low values of FMF are generally linked to higher values of δ_{1020} for both Asian and Saharan sites. However, there are occasional cases for which low FMF can be found for low values of δ_{1020} ~~which~~. **Such cases** might introduce artifacts when using FMF as a means for separating dusty from dust-free aerosol conditions.

3.2 Coarse-mode AOD versus dust AOD

A comparison of the coarse-mode AOD as provided by AERONET to the dust AOD obtained using Eqs. (6) and (7), respectively, is presented in Figure 6. Unsurprisingly, we find that lower coarse-mode and dust AODs are related to lower coarse-mode volume concentrations (not shown). For the Asian stations, we find that coarse-mode AOD tends to overestimate the contribution of mineral dust to AOD. The effect is particularly pronounced at AODs below 0.5 at 1020 nm and coarse-mode volume concentrations below 0.5. This means that other coarse particles, such as marine aerosols, are likely to be present under these conditions. As a consequence, fine-mode AOD, if used as proxy for non-dust aerosols, would lead to a systematic underestimation of the contribution of non-dust aerosol to total AOD. For the Saharan stations, coarse-mode AOD is found to be a suitable proxy for dust AOD. However, coarse-mode AOD shows few values below 0.1 while dust AOD can be as low as zero. Because the concentration of fine-mode aerosol is generally small at the selected Saharan sites, any comparison to non-dust AOD is inconclusive. In contrast to the Asian sites, the AOD related to fine-mode or non-dust particles is generally much lower to that of coarse-mode or dust particles, respectively (not shown).

We conclude that that coarse-mode AOD and dust AOD cannot necessarily be considered as synonymous. This needs to be kept in mind when using AERONET observations in the calibration/validation of spaceborne remote-sensing observations and aerosol transport modelling - particularly for locations with a high occurrence rate of complex aerosol mixtures.

3.3 AERONET-derived $AAOD_{BC}$ and model assessment

- Figure 7 presents the connection between $AAOD$ and $AAOD_{BC}$ at the standard AERONET wavelengths for observations at the Asian and Saharan sites. $AAOD_{BC}$ has been obtained from the non-dust $AAOD$ following Eq. (15). ~~We found that~~ absolute values of $AAOD$ are generally larger for Asian compared to Saharan sites and ~~that~~ the contribution of mineral dust to aerosol absorption at all wavelengths is generally larger at Saharan compared to Asian sites. A majority of $AAOD_{BC}$ values at Asian sites follows the theoretical curve for dust-free situations (i.e. with $\chi_{dust} = 0$) and the connection between $AAOD$ and $AAOD_{BC}$ is almost linear – particularly at longer wavelengths and larger $AAOD_{BC}$. For the same $AAOD$, a larger dust ratio χ_{dust} leads to a smaller $AAOD_{BC}$ and ~~a~~ **its** corresponding observation is located further away from the solid line (not shown). The abundance of pure dust conditions at the Saharan sites therefore leads to the larger spread of $AAOD_{BC}$ in Figure 7, ~~that~~ **and this feature** is particularly pronounced at 440 nm.

- To evaluate the quality of the methodology **that is used** for retrieving $AAOD_{BC}$, we ~~have~~ compared AERONET-derived values to the ones provided by CAMS aerosol reanalysis data for the sites considered in this study. We ~~have~~ investigated cases in which total AOD from AERONET and CAMS agree within 30%, 10%, and 5% of each other. We used these thresholds as a crude measure **that allowed us to introduce levels** of consistency between the two data sets and to assure that we consider cases in which the modelled aerosol situation is most likely resembling observations. The plots in Figure 8 show a very different situation for the Asian and Saharan sites: the former show correlated results and slopes of the linear fit that are reasonably close to the 1:1 line (particularly when requiring less than 5% difference in measured and modelled AOD), while the latter suggest that the CAMS $AAOD_{BC}$ is strongly underestimating the contribution of BC to light absorption in mixed Saharan dust plumes. The best model resemblance of $AAOD_{BC}$ is found for Dakar, where local pollution has a much stronger effect on aerosol composition than at the other Saharan sites (*Petzold et al.*, 2011). This suggests that $AAOD_{BC}$ as derived here from AERONET observations is more likely to describe aerosol absorption in anthropogenic pollution than in biomass-burning.

- We have presented a very selective analysis of AERONET observations as a proof of concept of the proposed methodology. More conclusive findings will require a thorough investigation of observations at a much larger set of AERONET sites.

4 Summary and conclusions

- We have presented a methodology to separate the contribution of dust and non-dust aerosol to total AOD measured with AERONET instruments based on lidar parameters provided in the version 3 level 2.0 inversion product. We showed how to derive the $AAOD$ related to the non-dust component as well as to the BC fraction. We have analysed AERONET time series at six sites that are frequently affected by Asian or Saharan dust, respectively. We found that coarse- and fine mode AOD cannot always be considered as synonymous with the AOD related to dust and non-dust aerosol, respectively. We note

that our methodology is the first **attempt** to enable such a differentiation solely on products provided by AERONET. We have compared retrieved values of $AAOD_{BC}$ to collocated model results provided by the CAMS aerosol reanalysis. This comparison has been restricted to only those AERONET-CAMS matches, for which total AOD agrees within 30% or better. We find that our methodology for obtaining $AAOD_{BC}$ from AERONET provides values that resemble CAMS aerosol modelling for Asian sites.

- 5 Little correlation was found for Saharan sites that are not frequently affected by a considerable contribution of anthropogenic pollution. This suggests that $AAOD_{BC}$ as derived here is less useful for observations of biomass-burning smoke – though the currently investigated data set has been far too small to draw a robust conclusion.

We consider the presented methodology as a useful tool for a more detailed calibration and validation of spaceborne remote-sensing observations and aerosol dispersion modelling with AERONET measurements. It will be particularly valuable at locations ~~with that show~~ a frequent occurrence of complex mixtures of mineral dust and anthropogenic pollution, e.g. east Asia or southern Europe but also individual highly polluted big cities downwind from major deserts.

- 10

Data availability. The data used in this work are freely available through the AERONET portal at <http://aeronet.gsfc.gov/>.

Author contributions. SKS, MT, and DM had the idea for this study. SKS and MT performed the data analysis and prepared the figures. All authors contributed to the discussion of the findings and the preparation of the manuscript.

- 15 *Competing interests.* The authors declare that no competing interests are present.

Acknowledgements. We thank the PIs of the AERONET site used in this study for maintaining their instruments and providing their data to the community. We also would like to thank AERONET for their continuous efforts in providing high-quality measurements and derivative products. All data used in this work can be accessed through the AERONET web page <http://aeronet.gsfc.gov/>. This work was supported by the Korean Meteorological Administration Research and Development program under Grant KMI (2018-04010).

References

- Ångström, A.: The parameters of atmospheric turbidity, *Tellus* 16, 64-75, <http://doi.org/10.1111/j.2153-3490.1964.tb00144.x>, 1964.
- Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131-3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.
- 5 Bellouin, N., Quaas, J., Morcrette, J.-J., and Boucher, O.: Estimates of aerosol radiative forcing from the MACC re-analysis, *Atmos. Chem. Phys.*, 13, 2045-2062, <http://doi.org/10.5194/acp-13-2045-2013>, 2013.
- Bergstrom, R. W.: Extinction and absorption coefficients of the atmospheric aerosol as a function of particle size, *Contr. Atmos. Phys.*, 46, 223-234, 1973.
- Bohren, C. and Huffman, D.: Absorbing and scattering of light by small particles. Wiley, <https://doi.org/10.1002/9783527618156>, 1983.
- 10 Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Sci. Tech.*, 40, 27-67, <https://doi.org/10.1080/02786820500421521>, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.*, 118, 5380-5552, doi:10.1002/jgrd.50171, 2013.
- 15 Burton, S. P., Vaughan, M. A., Ferrare, R. A., and Hostetler, C. A.: Separating mixtures of aerosol types in airborne High Spectral Resolution Lidar data, *Atmos. Meas. Tech.*, 7, <https://doi.org/10.5194/amt-7-419-2014>, 2014.
- Cross, E. S., Onasch, T. B., Ahern, A., Wrobel, W., Slowik, J. G., Olfert, J., Lack, D. A., Massoli, P., Cappa, C. D., Schwarz, J. P., and Spackman, J. R.: Soot particle studies—instrument inter-comparison—project overview, *Aerosol Sci. Tech.*, 44, 592-611, <https://doi.org/10.1080/02786826.2010.482113>, 2010.
- 20 Dubovik, O., Smirnov, A., Holben, B., King, M. D., Kaufman, Y. J., Eck, T. F., and Slutsker, I.: Accuracy assessments of aerosol optical properties retrieved from Aerosol Robotic Network (AERONET) Sun and sky radiance measurements, *J. Geophys. Res.—Atmos.*, 105, 9791-9806, <https://doi.org/10.1029/2000JD900040>, 2000.
- 25 Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.: Variability of absorption and optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59, 590-608, [http://doi.org/10.1175/1520-0469\(2002\)059<0590:VOAAOP>2.0.CO;2](http://doi.org/10.1175/1520-0469(2002)059<0590:VOAAOP>2.0.CO;2), 2002.
- Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten, H., Muñoz, O., and Veihelmann, B.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, *J. Geophys. Res.—Atmos.*, 111, <https://doi.org/10.1029/2005JD006619>, 2006.
- 30 Eck, T., Holben, B., Dubovik, O., Smirnov, A., Goloub, P., Chen, H., Chatenet, B., Gomes, L., Zhang, X.Y., Tsay, S. C.: Columnar aerosol optical properties at AERONET sites in central eastern Asia and aerosol transport to the tropical mid-Pacific, *J. Geophys. Res.—Atmos.*, 110, <https://doi.org/10.1029/2004JD005274>, 2005.
- Freudenthaler, V., Esselborn, M., Wiegner, M., Heese, B., Tesche, M., Ansmann, A., Müller, D., Althausen, D., Wirth, M., and Fix, A.: Depolarization ratio profiling at several wavelengths in pure Saharan dust during SAMUM 2006, *Tellus B*, 61, 165-179, <https://doi.org/10.1111/j.1600-0889.2008.00396.x>, 2009.

- Giles, D. M., Sinyuk, A., Sorokin, M. S., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J., Campbell, J., Welton, E. J., Korkin, S., and Lyapustin, A.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 Database – Automated Near Real-Time Quality Control Algorithm with Improved Cloud Screening for Sun Photometer Aerosol Optical Depth (AOD) Measurements, *Atmos. Meas. Tech. Discuss.*, <https://doi.org/10.5194/amt-2018-272>, in review, 2018.
- 5 Haywood, J.M. and Ramaswamy, V.: Global sensitivity studies of the direct radiative forcing due to anthropogenic sulfate and black carbon aerosols, *J. Geophys. Res.–Atmos.*, 103, 6043-6058, <https://doi.org/10.1029/97JD03426>, 1998.
- Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J., Setzer, A., Vermote, E., Reagan, J., Kaufman, Y. J., and Nakajima, T.: AERONET—A federated instrument network and data archive for aerosol characterization, *Rem. Sens. Environ.*, 66, 1-16. [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
- 10 Holben, B., Tanré, D., Smirnov, A., Eck, T., Slutsker, I., Abuhassan, N., Newcomb, W., Schafer, J., Chatenet, B., and Lavenue, F.: An emerging ground-based aerosol climatology: Aerosol optical depth from AERONET, *J. Geophys. Res.–Atmos.*, 106, 12067-12097, <https://doi.org/10.1029/2001JD900014>, 2001.
- Inness, A., Baier, F., Benedetti, A., Bouarar, I., Chabrillat, S., Clark, H., Clerbaux, C., Coheur, P., Engelen, R., and Errera, Q.: The MACC reanalysis: an 8 yr data set of atmospheric composition, *Atmos. Chem. Phys.*, 13, 4073-4109, <https://doi.org/10.5194/acp-13-4073-2013>, 15 2013.
- Khalizov, A. F., Xue, H. X., Wang, L., Zheng, J., and Zhang, R. Y.: Enhanced light absorption and scattering by carbon soot aerosol internally mixed with sulfuric acid, *J. Phys. Chem. A*, 113, 1066–1074, doi:10.1021/jp807531n, 2009.
- Koven, C.D. and Fung, I.: Inferring dust composition from wavelength-dependent absorption in Aerosol Robotic Network (AERONET) data, *J. Geophys. Res.–Atmos.*, 111, <https://doi.org/10.1029/2005JD006678>, 2006
- 20 Mamouri, R.-E. and Ansmann, A.: Fine and coarse dust separation with polarization lidar, *Atmos. Meas. Tech.*, 7, 3717-3735, <https://doi.org/10.5194/amt-7-3717-2014>, 2014.
- Müller, T., Schladitz, A., Massling, A., Kaaden, N., Kandler, K. and Wiedensohler, A.: Spectral absorption coefficients and imaginary parts of refractive indices of Saharan dust during SAMUM-1, *Tellus*, 61B, 79-95, <https://doi.org/10.1111/j.1600-0889.2008.00399.x>, 2009.
- Noh, Y.M.: Single-scattering albedo profiling of mixed Asian dust plumes with multiwavelength Raman lidar, *Atmos. Environ.*, 95, 305-317, 25 <https://doi.org/10.1016/j.atmosenv.2014.06.028>, 2014.
- Noh, Y., Müller, D., Lee, K., Kim, K., Lee, K., Shimizu, A., Sano, I., and Park, C. B.: Depolarization ratios retrieved by AERONET sun–sky radiometer data and comparison to depolarization ratios measured with lidar, *Atmos. Chem. Phys.*, 17, 6271-6290, <https://doi.org/10.5194/acp-17-6271-2017>, 2017.
- O'Neill, N., Eck, T., Smirnov, A., Holben, B., and Thulasiraman, S.: Spectral discrimination of coarse and fine mode optical depth, *J. Geophys. Res.–Atmos.*, 108, <https://doi.org/10.1029/2002JD002975>, 2003.
- 30 Petzold, A., Rasp, K., Weinzierl, B., Esselborn, M., Hamburger, T., Dörnbrack, A., Kandler, K., Schütz, L., Knippertz, P., Fiebig, M. and Virkkula, A.: Saharan dust absorption and refractive index from aircraft-based observations during SAMUM 2006, *Tellus*, 61B, 118-130, <https://doi.org/10.1111/j.1600-0889.2008.00383.x>, 2009.
- Petzold, A., Veira, A., Mund, S., Esselborn, M., Kiemle, C., Weinzierl, B., Hamburger, T., Ehret, G., Lieke, K. and Kandler, K.: Mixing of mineral dust with urban pollution aerosol over Dakar (Senegal): impact on dust physico-chemical and radiative properties, *Tellus*, 63B, 619-634, <https://doi.org/10.1111/j.1600-0889.2011.00547.x>, 2011.

- Russell, P., Bergstrom, R., Shinozuka, Y., Clarke, A., DeCarlo, P., Jimenez, J., Livingston, J., Redemann, J., Dubovik, O., and Strawa, A.: Absorption Angstrom exponent in AERONET and related data as an indicator of aerosol composition, *Atmos. Chem. Phys.*, 10, 1155-116, <https://doi.org/10.5194/acp-10-1155-2010>, 2010.
- Schuster, G.L., Dubovik, O., Holben, B.N., and Clothiaux, E.E.: Inferring black carbon content and specific absorption from Aerosol Robotic Network (AERONET) aerosol retrievals, *J. Geophys. Res.–Atmos.*, 110, <https://doi.org/10.1029/2004JD004548>, 2005.
- Schuster, G. L., Dubovik, O., and Holben, B. N.: Angstrom exponent and bimodal aerosol size distributions, *J. Geophys. Res.–Atmos.*, 111, <https://doi.org/10.1029/2005JD006328>, 2006.
- Shimizu, A., Sugimoto, N., Matsui, I., Arao, K., Uno, I., Murayama, T., Kagawa, N., Aoki, K., Uchiyama, A., and Yamazaki, A.: Continuous observations of Asian dust and other aerosols by polarization lidars in China and Japan during ACE-Asia, *J. Geophys. Res.–Atmos.*, 109, <https://doi.org/10.1029/2002JD003253>, 2004.
- Shin, S.-K., Tesche, M., Kim, K., Kezoudi, M., Tatarov, B., Müller, D., and Noh, Y.: On the spectral depolarisation and lidar ratio of mineral dust provided in the AERONET version 3 inversion product, *Atmos. Chem. Phys.*, 18, 12735-12746, <https://doi.org/10.5194/acp-18-12735-2018>, 2018.
- Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M., 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, 1535 pp. Cambridge Univ. Press, Cambridge, UK, and New York.
- Tesche, M., Ansmann, A., Müller, D., Althausen, D., Mattis, I., Heese, B., Freudenthaler, V., Wiegner, M., Esselborn, M., Pisani, G., and Knippertz, P.: Vertical profiling of Saharan dust with Raman lidars and airborne HSRL in southern Morocco during SAMUM, *Tellus B*, 61, 144-164, <https://doi.org/10.1111/j.1600-0889.2008.00390.x>, 2009a.
- Tesche, M., Ansmann, A., Müller, D., Althausen, D., Engelmann, R., Freudenthaler, V., and Groß, S.: Vertically resolved separation of dust and smoke over Cape Verde using multiwavelength Raman and polarization lidars during Saharan Mineral Dust Experiment 2008, *J. Geophys. Res.–Atmos.*, 114, D13202, <https://doi.org/10.1029/2009JD011862>, 2009b.
- Yu, X., Cheng, T., Chen, J., and Liu, Y.: A comparison of dust properties between China continent and Korea, Japan in East Asia, *Atmos. Environ.*, 40, 5787-5797, <https://doi.org/10.1016/j.atmosenv.2006.05.013>, 2006.




Figure 1. Change in $AAOD$ (red), $AAOD_d$ (black), and $AAOD_{nd}$ (magenta) with dust ratio χ_d for an aerosol mixture of dust ($\omega_d = 0.98$) and non-dust ($\omega_{nd} = 0.90 - 0.96$) and an AOD of unity. The blue lines mark the contribution of non-dust aerosol to $AAOD$ for different values of ω_{nd} .




Figure 2. Histograms of the 1020-nm AOD (a), 1020-nm $PLDR$ (b), and FMF (c) for the considered AERONET stations affected by Saharan (blue) and Asian dust (red). Coloured number provide the value for the bar that exceeds the scale. Details on the considered AERONET stations and mean values are provided in Table 2.




Figure 3. Histograms of the 440-870-nm extinction (a) and absorption (b) Ångström exponents for the considered AERONET stations affected by Saharan (blue) and Asian dust (red). Coloured number provide the value for the bar that exceeds the scale. Details on the considered AERONET stations are presented in Table 2.




Figure 4. Histograms of χ_d for the considered AERONET stations affected by Saharan (blue) and Asian dust (red). Coloured numbers provide the values for bars that exceed the scale.




Figure 5. 2d histograms of 1020-nm PLDR and χ_d for the considered AERONET stations affected by Asian (left) and Saharan dust (right). The black lines refer to the backscatter-related dust ratio R_d .




Figure 6. 2d histograms of 1020-nm coarse-mode *AOD* and dust-related 1020-nm *AOD* for the considered AERONET stations affected by Asian (upper panel) and Saharan dust (lower panel).




Figure 7. 2d histograms of $AAOD$ and $AAOD_{BC}$ at the four AERONET standard wavelengths for the Asian (**left upper panel**) and Saharan (**right lower panel**) stations considered in this study. Solid lines refer to the theoretical values of $AAOD_{BC}$ (using Eq. (15) and the values in Table 1) in the absence of mineral dust. Dashed lines mark the 1:1 line. Observations would follow this slope only if BC was a perfect absorber, i.e. if all absorption in the non-dust fraction would be due to were caused by BC or $\omega_{BC} = 0$.




Figure 8. Comparison of AERONET-derived $AAOD_{BC}$ with CAMS model estimates for the sites listed in Table 2 for cases in which the total AOD from CAMS and AERONET agrees within 30% (black circles, thin lines), 10% (red circles, medium lines), and 5% (red dots, bold lines). Numbers in the plots refer to the total number of collocated points and the number of matches with the given AOD agreement. Dashed lines mark the 1:1 line. Solid lines are linear fits of the data. Numbers in the plots refer to the number of collocations and squared correlation coefficients for all cases (solid black, no R^2 given) and those with an AOD agreement within 30% (open black), 10% (open red), and 5% (solid red).

Table 1. List of input parameters used for the retrieval of $AAOD_{\text{nd}}$ and $AAOD_{\text{BC}}$ in this study. Values of δ at 1020 nm are used for the separation of optical properties of dust and non-dust particles. The dust-related Ångström exponent is needed to transform findings at 1020 nm to other wavelengths. The values of ω_{d} and ω_{BC} are used to retrieve $AAOD_{\text{nd}}$ and $AAOD_{\text{BC}}$, respectively.

Parameter	Symbol	Value				Reference
		440 nm	675 nm	870 nm	1020 nm	
total AOD	AOD					
total PLDR	δ	from individual AERONET				
total lidar ratio	S	version 3 level 2.0 measurements				
total SSA	ω					
non-dust PLDR	δ_{nd}	-	-	-	0.02 ± 0.01	<i>Shimizu et al. (2004)</i>
dust PLDR (Asian)	δ_{d}	-	-	-	0.30 ± 0.04	<i>Shin et al. (2018)</i>
dust PLDR (Saharan)	δ_{d}	-	-	-	0.31 ± 0.03	<i>Shin et al. (2018)</i>
dust lidar ratio (Asian)	S_{d}	-	-	-	44 ± 6 sr	<i>Shin et al. (2018)</i>
dust lidar ratio (Saharan)	S_{d}	-	-	-	54 ± 9 sr	<i>Shin et al. (2018)</i>
dust Ångström exponent	α_{d}		0.06 ± 0.21			<i>Tesche et al. (2009a)</i>
dust SSA	ω_{d}	0.94	0.98	0.99	0.99	<i>Eck et al. (2005); Yu et al. (2006)</i>
BC SSA	ω_{BC}	0.25 ± 0.13	0.17 ± 0.01	0.13 ± 0.03	0.07 ± 0.02	<i>Haywood and Ramaswamy (1998)</i>

Table 2. Overview of the AERONET sites included in this study in terms of location, length of time series and number of available version 3 level 2.0 data points. The last three columns refer to mean values and standard deviation of AOD_{1020} , δ_{1020} , and FMF for the respective sites and regions. The figures in this work refer to the combined Asian and Saharan data sets.

Station	Location	Period	N	AOD_{1020}	δ_{1020}	FMF
Beijing	39.98 °N, 116.38 °E	2001–2018	2713	0.45 ± 0.29	0.06 ± 0.07	0.42 ± 0.17
Gwangju_GIST	35.23 °N, 126.84 °E	2004–2018	956	0.25 ± 0.12	0.06 ± 0.07	0.51 ± 0.19
XiangHe	39.75 °N, 116.96 °E	2001–2018	4300	0.41 ± 0.25	0.06 ± 0.07	0.44 ± 0.18
combined Asian		2001–2018	7969	0.41 ± 0.26	0.06 ± 0.07	0.44 ± 0.18
Banizoumbou	13.55 °N, 2.67 °E	1995–2018	4217	0.60 ± 0.31	0.29 ± 0.05	0.11 ± 0.08
Capo_Verde	16.73 °N, 22.94 °W	1994–2018	1689	0.55 ± 0.25	0.30 ± 0.05	0.09 ± 0.04
Dakar	14.39 °N, 16.96 °W	1996–2018	4118	0.54 ± 0.28	0.28 ± 0.06	0.12 ± 0.08
combined Saharan		1994–2018	10024	0.57 ± 0.29	0.29 ± 0.05	0.11 ± 0.07