

Interactive comment on “Albedo-Ice regression method for determining ice water content of Polar Mesospheric Clouds using ultraviolet observations from space”

by Gary E. Thomas et al.

Anonymous Referee #1

Received and published: 23 December 2018

General comments:

This manuscript deals with an alternative approach to retrieve the ice water content (IWC) of PMCs from satellite-borne UV-backscatter observations with the CIPS instrument on the AIM satellite. Due to orbital drifts CIPS is since 2016 operated in a different way and the original approach to retrieve IWC is not possible any more for a large part of the measurements. The novel approach estimates IWC from backscatter measurements at a single scattering angle. Overall, the approach works well. The paper is generally well written and is suitable for publication in AMT. I ask the authors to consider the comments listed below. In my opinion the paper should be accepted, once the (mainly minor) issues listed below were addressed.

Specific comments and responses:

Our responses are in italics. When we respond “OK”, it means that we are following your recommendation. We also include a revised pdf document with our tracked changes, where all additions are in red type. Line numbers are referred to “new” if they refer to the new document, which has been reformatted.

Line 16: "265nm" -> "265 nm" **OK**

Line 114: “..it is necessary to derive scattered radiances from the same algorithm used by CIPS.”

I don't really understand this statement “derive scattered radiances from the same algorithm used by CIPS.” CIPS measures scattered radiances and the algorithms are used to infer physical properties of PMCs, right? I guess I'm missing a point?

No, we were not clear. We replaced the sentences (lines 113-117), (now 91-94) with the following: “Since the SOFIE technique uses near-IR solar extinction in ice-water absorption bands, the primary measurement is ice water content. As shown in Sec 2.2, we reversed the process, to derive radiances from IWC, and then applied the same regression method to the results.”

[I suggest an alternative wording for the last sentence (and note the change from section 2.2 to 2.3)]: “As shown in Sec 2.3, we inverted the retrieved SOFIE IWC to derive the equivalent 265-nm albedo, and then applied the regression method described above to the results.”

Line 138: “the monochromatic scattering cross-section” It actually is the “differential” scattering cross section. Perhaps this can/should be added. **OK.**

Line 140: “number density of ice particles”
Perhaps better “column density”?

in new line 107 we changed “number density” to “column density”.

Equation (2): I suggest using a slightly different symbol for the albedo than in equation (1). The left hand sides of equation (1) and (2) are the same, but the right hand sides differ.

Agreed. We placed the letter “m” as a superscript, and identified it as such in the previous sentence (new line 117).

Line 167: add space before “denotes”. OK

Line 209: I don’t fully understand the meaning of this sentence. Is the microphysical model a 1-D model?

No. It is fully 3-D.

“(2) radiance and IWC may be calculated accurately, so that effects of cloud inhomogeneity are absent;”

We added the following sentences after (new) line 172.” With regard to the latter point, we describe in more detail the model calculations. The model grid is 4° in latitude, 5° in longitude, and variable in the vertical. Ice particles of varying sizes fill many of these cells, but the density of particles within each cell is, by definition, constant. For a given model cloud, the integration is made through a vertical ‘stack’ of all ice-filled cells generated in a given computer run, and within each particle size grid. The total radiance is the sum of contributions from the size range 20 to 150 nm. The observation angles are always assumed to be zero, in other 175 words, the integration is performed in the vertical only. Thus cloud ‘boundaries’ in the horizontal plane are not an issue. This contrasts with real heterogeneous clouds where these approximations would not hold.” As you will note in the response to Reviewer 2, we added two more clarifying statements.

Line 216: “thick.).“ -> “thick).“

Same line: I suggest adding “resolution“ to read “The model height grid resolution is variable, being highest. ”, otherwise “being highest” doesn’t make sense.

We changed the sentence to read (now line 186) “The model height grid is variable, so that the smallest layer thickness is 0.26 km, which resolves the narrow ice layers”

• • •

Line 220 and Figure 1: I’m wondering how the linear regressions are actually done. Is it a single linear fit of y vs. x, or do you fit both y vs. x and x vs. y and determine an average slope and offset? Alternatively, there are routines taking both differences in x and y direction into account, when minimizing chi-square. There may be a large difference between fitting y vs. x or x vs. y.

This is an interesting point. However it is largely a mathematical issue, irrelevant to the desired objective. The albedo is the measured quantity, and IWC is the derived quantity - the albedo is the natural independent variable. Certainly, there are circumstances where

the opposite is true, when IWC is the measured variable (as in the case of SOFIE) and one desires to know the albedo- (however one would also have to specify the scattering angle in that case). As the reviewer notes, in this case the fitting process would have to be x vs y, but our regression results (y vs. x) apply to the question being asked.

Line 227: “In fact, we found that the linear relationship breaks down for very small albedo,”
The reason is probably, that particle populations with really small particles (< 10 nm) have a non-zero IWC, but the albedo is essentially zero, right?

Yes, we would agree with that. In line (new) 199, We replaced this phrase with “below the detection threshold of CIPS and SBUV, and are a result of the very faint small particles.”

Figure 1: Both the IWC and the albedo values have lower limits. Perhaps I missed it, but what is the reason for this lower limit. For IWC the limit seems to be 20 g/km².

“The lower limit for the albedo (SA=90°) is 1G, which is the detection limit of the CIPS experiment.” We added this comment to the figure caption.

Figure 2: The caption should clarify that the displayed error is a relative error given in Line 216:

We added the word “relative” to the word ‘error’ in figure caption 2. We added further clarification by adding : “The errors are relative to the model values”

Closing parenthesis missing after “Rapp and Thomas (2006)”. OK

Line new 114: “DIFFERENTIAL scattering cross-section”? OK

Line 289: “The mean difference is -13%. Looking at the Figure, the mean difference seems to be larger.

-13% is an unweighted mean over all bins. If we had weighted the error according to the number of points in each bin, the error would be larger, and positive.

We added the sentence (now line 243)” Assuming the microphysical model is accurate, the accuracy of the CIPS UV measurements ranges from over +100% for very small albedo to -60% for high albedos. We emphasize that this is not an AIR result, but is an attempt to assess how particles that are too small to be visible to UV measurements affect the accuracy of the CIPS IWC results.”

Also, what is the reason for the characteristic variation of the difference with increasing albedo? Is there a simple explanation?

We elaborate further in the Conclusion section (lines 353 -356): “Assuming the microphysical model is accurate, the accuracy of UV measurements ranges from over +100% for very small albedo to -60% for high albedos. The overall accuracy of IWC (averaging over all

albedo bins) is $-13 \pm 17\%$. The CIPS algorithm overestimates the small-particle population (20-30 nm) as a result of the Gaussian [this change should also be made in the paper] approximation when the mean particle size is small, and the opposite is true when the mean size is large. Distinct from the more fundamental errors due to the invisibility of very small ice particles and the Gaussian approximation, we also estimated the errors in the AIR approximation, relative to the AIM SOFIE data which apply to larger values of IWC than the model. For the dimmer and more frequent clouds, Fig. 2 shows that the error in ensemble averages is of the order of 5%.

Line 289: “.. subset .. have” -> “.. subset .. has” OK

Line 303: “As previously shown, the AIR data applies to the ice mass of ‘UV-visible’ clouds, not to their total ice mass.” I don’t really understand this statement. What is the meaning of “UV-visible clouds”? Was this shown in the current paper? If not, please provide a reference.

In line 112 we added the sentence “As shown by Rapp and Thomas (2006), particles with sizes < 20 nm are not detectable by UV measurements because of their small cross-section values – hence we refer to ‘UV-visible clouds’.”

Caption Figure 4, line 3: “SA = 40 deg” should read “SA = 50 deg”. OK.

Line 366: “AIR overestimates IWC by up to 15% AIR may also underestimate IWC, right? I also think that the overestimation may be much larger than 15%.

See the above additions where we address these issues in more detail (line 243):
“Assuming the microphysical model is accurate, the accuracy of the CIPS UV measurements ranges from over +100% for very small albedo to -60% for high albedos. We emphasize that this is not an AIR result, but is an attempt to assess how particles that are too small to be visible to UV measurements affect the accuracy of the CIPS IWC results.”

Line 367: “SA = 100 deg” -> “SA = 110 deg” OK

Line 375: “The accuracy of the average IWC results was estimated by removing half the data from an entire season and comparing the two results. “ I’m not sure, I fully understand this statement. AIR is applied to SBUV data and then you split the data set in half and compare the results. How does this allow you to estimate the accuracy of the “average IWC results”? I guess I’m missing the point here?

We added (line 101) the following sentence to emphasize that the errors refer to systematic differences between several divided data sets; “For a highly-populated region (more than 1000 clouds per season at latitudes higher than 70°), the differences in IWC ranged between ± 3 –5 g-km⁻², thus can be considered typical systematic errors. For a less-populated region (50°–64° latitude) where there were many fewer clouds (<50), the differences were larger, ± 5 –10 g-km⁻².”

Next sentence: You report on “changes” of $\pm 3 - 5 \text{ g km}^2$ etc., but were there any systematic differences? These values were the systematic differences between pairs of divided data sets. We added the word “systematic” to $\pm 3 - 5 \text{ g / km}^2$. We specifically added the phrase “systematic” errors. See above.

Interactive comment on “Albedo-Ice regression method for determining ice water content of Polar Mesospheric Clouds using ultraviolet observations from space” by Gary E. Thomas et al.

Anonymous Referee #2

Received and published: 14 January 2019

This manuscript describes the retrieval of the ice water content (IWC) of mesospheric clouds based on measuring cloud albedo. This retrieval method has been developed with particular focus on AIM/CIPS since changes in orbit have made the original IWC retrieval based on phase function analysis impossible. However, the method is also applicable to other mesospheric datasets like SBUV. This gives the method potential importance for inter-comparisons of mesospheric datasets and for the analysis of long- term variability. The approach is straight-forward, and the results are convincing. I recommend the manuscript for publication after some minor revisions. I mainly would like to see a number of clarifications.

New and old text are in “”.

One issue I would like the authors to discuss more: Why is the AIR method not even better than described in the manuscript? The AIR method is based on finding a linear relationship between ice water content and cloud albedo. For typical mesospheric clouds, one can argue for such a relationship even on a theoretical basis, as long as scattering coefficients depend on the particle size to a power in the vicinity of 3. The authors show that the AIR method works well on a statistical basis. However, the AIR results presented in the manuscript show much scatter in the relationship between albedo and IWC, and the authors point out in several places that we cannot expect AIR to work for individual IWC retrievals. I would like to see more discussion on why the method is "not better than this". A reference is e.g. Hultgren and Gumbel (J. Geophys. Res., 119, 14129-14143, 2014). That paper shows many examples of a close relationship between cloud scattering coefficient and ice mass density, which works well even for the altitude-dependent quantities, not only for the column-integrated quantities considered in the current paper. Can I dare the authors to make a more quantitative statement: Can we take the AIR results for real and apply the method to individual retrievals, by providing a suitable statement about the error bar of such individual AIR IWC retrievals? How large (in percent) would such an error bar be?

Apart from the question of why one would be interested in only one cloud measurement, we have responded by calculating the overall percent error in single "measurements" (model simulations) of albedo, given the scattering angle. We looked at all albedos > 1G and SA=90°. The distribution of AIR errors is given below. The std deviation of the Gaussian fit is 19%. For most applications this is probably too large an error to be useful. The dispersion of particle sizes leads to a distribution that is quasi-random. This means that averaging will improve the mean by the square root of the number n of measurements. We separated the data into sets of 100 and 500 randomly chosen iwc-albedo pairs, and calculated the % errors. The errors in the means of the AIR distributions [σ/\sqrt{n}] were 2.16% and 0.9%; these are equivalent to a standard error of ~19% for a random data set. These numbers will vary with scattering angle, and with specific albedo values, but this provides an example of how AIR would work with real data, on a time interval when only a few hundred measurements are available, e.g. one month.

We added the following sentence in line 225 (now line 225): "For the conditions in Fig. 1(c), the mean error of AIR for a single model simulation is 19%. The error can be reduced substantially by averaging. For example, for 100 measurements, the AIR error is only 2%. Figure 1 also shows...." We also added the reference Hultgren and Gumbel to line 130.

See Supplemental Figure 1: "singlemeasurementerrors.jpeg"

Section 2 (Theoretical basis): The major result of this study is that IWC is linearly related to cloud albedo. Therefore, the description in line 156-160 is confusing. In line 156-157, the authors refer to "the results of this study that IWC is linearly related to the column density of ice particles". The fact that IWC is linearly related to the column density of ice particles is somehow trivial (although dependent on the details of the particle population). In line 158-159, it is also stated that "As pointed out by Englert and Stevens (2007), such a relationship exists for certain SA values..." However, the relevant finding by Englert and Stevens is about the relationship between IWC and albedo. I therefore suspect that this paragraph should be

about the relationship between IWC and albedo, not between IWC and column density. Please clarify and reformulate.

We agree that we should have stated albedo rather than column density. We changed the sentence from "Anticipating the results of this study that IWC is linearly related to the column density of ice particles," to "Anticipating the results of this study that IWC is linearly related to cloud albedo, .."

Section 2.1 (Model results): Please describe in more detailed the processes included in the model simulations and the resulting variability in mesospheric clouds. Is gravity wave activity included in the simulations? Does the cloud database include multiple layer clouds or other conditions that may lead to clouds deviating from a straight growth/sedimentation scenario?

In response to reviewer 1, we added more detail on the model calculations (lines 171-onward). In response to your specific request, we added the sentence (now line 177): "The model contains variability due to waves of various sorts, including tides and gravity waves. However, it does not capture all known details of PMC, such as double layers. Since we are dealing with integrated quantities, this should not be an important issue. Furthermore, we don't place full reliance on the model, which is why we also use two independent data sets."

Section 2.2 (AIR results from CIPS): Figure 8 shows the AIR method applied to the CIPS from the Northern Hemisphere 2011. When it comes to demonstrating that the AIR method works, this choice of season is unfortunate. CIPS data from the years 2010-2013 has been used in the regression analysis to "train" the model. When subsequently investigating the ability of the model to retrieve IWC, a season should be chosen that has not already been used to train the model. I suggest to choose another season.

Since we 'trained' the AIR method to literally hundreds of thousands of individual cloud data, showing how it works for a single day does not detract from illustrating its usefulness..

Some other details: Line 84: The notation "meteor 'smoke' nucleation" may be misleading. It is better to write "nucleation on meteoric 'smoke'".

Agreed: We replace this phrase by "The processes treated by the model include nucleation on meteor 'smoke' particles,.."

Line 274: To avoid confusion, please clarify what is meant by "mean ice particle volume evaluated at r_m ", i.e. make clear that you refer to an integration over the Gaussian particle size distribution.

Agreed. We replaced the sentence in (now) line 231 with "V denotes the ice particle volume, averaged over the Gaussian distribution with a mean particle radius value r_m ."

Line 287: Clarify that by "simulated CIPS retrieved IWC" you mean the AIR result.

We apologize for this misconception. We added the following sentence (line 243): “We emphasize that this is not an AIR result, but is an attempt to assess how particles that are too small to be visible to UV measurements affect the accuracy of the CIPS IWC results.”

Line 394: The authors refer to $n = 3-5$ as typical exponents for the size-dependence ($\sigma \sim r^n$) of the scattering cross section for typical mesospheric clouds. Can this be motivated better? Otherwise a larger range may be appropriate from $n = 2$ (geometrical optics limit) to $n = 6$ (Rayleigh limit). The reference to Hultgren and Gumbel (2014) has also been mentioned above. This reference is interesting even here as it discusses ideas underlying the relationship between cloud brightness and ice (including e.g. r^n dependence of the scattering coefficient, dependence on particle size distribution) that are also discussed in the current paper.

See below for our response to why AIR works as well as it does.

Figure 1: In order to better understand the behavior of the model data, it would be instructive for the reader to see the data points all the way down to the zero point (small albedo, small IWC). I do not see a reason not to show these points.

For your curiosity, this plot shows the results down to very small albedo. This information is not relevant to the CIPS data, which has a detection limit of 1G. The behavior of the relationship is different, with a different slope and more dispersion. We believe that showing this behavior detracts from our message. The blue dots show results from effective particle sizes < 20nm. Red dots show results for reff between 20 & 30 nm and the black dots show the contributions from larger sizes. The light blue line is the AIR result, which is curved because of the log-log scale.

See Supplemental Figure 2, “IWCvsfaintalbedo.jpeg”

Figure 2: To avoid confusion, I suggest to mention the units of the contour lines (g km^{-2}) in the figure caption.

Again, we apologize for the misconception. We added the sentence: “Contour lines are labelled as percent errors relative to the accurate model values.”

Figure 11: In the two plots, there is an obvious lower limit to the data points (in terms of a straight line nearly parallel to the red line). I (and possibly other readers) do not understand why there is such a well-defined lower limit. Please add an explanation.

This was a very helpful suggestion, and we looked further into this matter.

The ratio of albedo to IWC shows a clearly defined lower limit to this ratio, which is nearly independent of the effective particle size at least for the larger particles.. This behavior is due to the fact that the ratio of IWC/albedo (which is independent of column density) asymptotes to a straight line for the larger values of particle size ($30 < \text{reff} < 50$ nm) which are those responsible for PMC. Below is a plot of this ratio versus effective radius, showing this behavior, for $SA=90$

deg. The constant ratio at large r_{eff} is due to the fact that the r -dependence of the cross-section in this range is r^3 , and cancels the IWC dependence on r^3 . The same behavior occurs for the other scattering angles, but the asymptotes are different. The near-constancy of the ratios means that there will be distinct lower limits to the regressions of IWC vs Albedo, which slant upwards in the plots due to the linear variation of albedo on column density. This is evident in Figure 1 where the lower boundary of the scatter is quite linear. These points refer to the largest particles in the population. The other reason for the sharpness of the lower boundary of the ratio is that the largest particles are the ones responsible, and these have a very steep fall-off in the size distributions. $r^3 / \sigma_{\lambda}(r, \Phi)$

See Supplemental Fig. 3, “RatioIWCtoAlb.jpeg

We added the discussion of this issue in Sec. 3, Effects of Mean Particle Size, beginning on line 329,

“The AIR approximation is based on the notion that particle size effects can be ignored in retrieving IWC from albedo measurements. That is, they contribute in a sense to the ‘noise’ of the measurement, which can be minimized by averaging. In fact, the particle size (or more accurately, the term r^3) is a principal ‘driver’ of $\langle IWC \rangle$ itself, so it is not obvious that particle size effects play a minor role in deriving IWC. The dependence of albedo on column density adequately captures this part of the variability (albedo is strictly linear in column density). The AIR slope term is $\sim r^3 / \sigma_{\lambda}(r, \phi)$ averaged over a distribution of particle sizes, r . The size dependence of the cross-section varies as a power of r , within two limits, the geometric-optics limit, r^2 , and the small-particle (Rayleigh) limit, r^6 . In the intermediate and realistic conditions of PMC, the exponent has an intermediate value. Fortunately, there is a “sweet spot” (or better, a ‘sweet region’ of the r -domain) in which the r -dependence of σ_{λ} is $\sim r^3$, so that the slope term is constant (for fixed SA). This behavior occurs for all relevant values of SA, and for the albedo values typical of CIPS. It accounts mainly for the effectiveness of the AIR method. The other aspect favorable to AIR is the steep fall-off of the particle size distribution at the largest sizes, which contributes to the sharpness of the lower boundaries in the spread of points in Fig. 1.”

Albedo-Ice regression method for determining ice water content of Polar Mesospheric Clouds using ultraviolet observations from space

Gary E. Thomas¹, Jerry Lumpe², Charles Bardeen³ and Cora E. Randall^{1,4}

¹Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, Colorado, USA

² Computational Physics, Inc., Boulder, Colorado, USA

³ National Center for Atmospheric Research, Boulder, Colorado, USA

Correspondence to: Gary E. Thomas (thomas@lasp.colorado.edu)

Abstract. High spatial-resolution images of Polar Mesospheric Clouds (PMC) from a camera array onboard the Aeronomy of Ice in the Mesosphere Satellite have been obtained since 2007. The Cloud Imaging and Particle Size Experiment (CIPS) detects scattered ultraviolet (UV) radiance at a variety of scattering angles, allowing the scattering phase function to be measured for every image pixel. With well-established scattering theory, the mean particle size and ice water content (IWC) are derived. In the nominal mode of operation, approximately seven scattering angles are measured per cloud pixel. However, because of a change in the orbital geometry in 2016, a new mode of operation was implemented such that one, or at most two, scattering angles per pixel are now available. Thus particle size and IWC can no longer be derived from the standard CIPS algorithm. The Albedo-Ice Regression (AIR) method was devised to overcome this obstacle. Using data from both a microphysical model and from CIPS in its normal mode, we show that the AIR method provides sufficiently accurate average IWC so that PMC IWC can be retrieved from CIPS data into the future, even when albedo is not measured at multiple scattering angles. We also show from the model that 265nm UV scattering is sensitive only to ice particle sizes greater than about 20-25 nm in (effective) radius, and that the operational CIPS algorithm has an average error in retrieving IWC of $-13\pm 17\%$.

1 Introduction

Polar Mesospheric Clouds (known as noctilucent clouds in the ground-based literature) have been studied for over a century from high-latitude ground observations, but only since the space age have we understood their physical nature, as water-ice particles occurring in the extremely cold summertime mesopause region. Their seasonal and latitudinal variations have now been well documented (DeLand et al., 2006). Interest in these clouds ‘at the edge of space’ has been stimulated by suggestions that they are sensitive to global change in the mesosphere (Thomas et al., 1989). This expectation has been supported recently by a time series analysis of Solar Backscattered Ultraviolet measurements of PMC (Hervig et al., 2016) and by model calculations (Lübken et al., 2018).

The Aeronomy of Ice in the Mesosphere satellite (AIM) (Russell et al., 2009) was designed to provide a deeper understanding of the basic processes affecting PMC, through remote sensing of both the clouds and their physical environment (temperature, water vapor, and meteor smoke density, among other constituents). One of the two active experiments on board AIM is a camera array, the Cloud Imaging and Particle Size (CIPS) experiment, which provides high spatial resolution images of PMC (McClintock et al., 2009). CIPS measures scattered ultraviolet (UV) sunlight in the nadir in a spectral region centered at 265 nm, where ozone absorption allows the optically-thin ice particles to be visible above the Rayleigh scattering background issuing from the ~50-km region (Rusch et al., 2009; Bailey et al.,

2009). Because of its wide field of view and 43-second image cadence, CIPS views a cloud element multiple times in its sun-synchronous orbital passage over the polar region, thus providing consecutive measurements of the same location at multiple (typically seven) scattering angles (SA). Together with scattering theory, the brightness of the cloud (albedo) at multiple angles provides constraints needed to estimate the mean ice particle size (Lumpe et al., 2013). From the particle size and albedo measurements, the ice water content is calculated for each cloud element (7.5 x 7.5 km² in the most recent CIPS retrieval algorithm). However, over time, the AIM orbit plane has drifted from its nominal noon-midnight orientation to the point where the satellite is currently operating in a terminator orbit. Responding to this altered geometry and the desire to broaden the scope of AIM, new measurement sequences were implemented to provide observations of the entire sunlit hemisphere, rather than just the summertime high-latitude region. Because the total number of images per orbit is fixed by data storage limitations, a new mode (the ‘continuous imaging mode’) of observations, with a reduced three-minute image cadence, was implemented in February 2016. The present sampling in a single Level 2 pixel contains many fewer scattering angles (often only one). To maintain consistency in the study of inter-annual variations of PMC, this necessitates a revised method of retrieving ice water content (IWC) where only a single albedo measurement is available. IWC is a valuable measure of the physical properties of PMC since it largely removes the effects of scattering-angle geometry, is a convenient PMC climate variable when averaged over season, and can be used in comparing with contemporaneous measurements of PMC that use different observational techniques.

The Albedo-Ice Regression (AIR) method was developed to fill the need to retrieve PMC IWC with only a single measurement of albedo. Based on the simple notions that both albedo and IWC depend linearly upon the ice-particle column density, multiple linear relationships are established between IWC and cloud directional albedo, depending upon scattering angle. The regressions are derived from three data sources: (1) the Specified Dynamics version of the Whole Atmosphere Community Climate Model (SD-WACCM) combined with the Community Aerosol and Radiation Model for Atmospheres (CARMA); (2) CIPS data for the years 2007-2013, when multiple scattering angles were available to derive IWC; and (3) Solar Occultation For Ice Experiment (SOFIE), which provides IWC and particle sizes. These three sources provide many thousands of albedo-IWC-particle size combinations, from which the AIR regressions are derived. Although the AIR method may be inaccurate for a single retrieval of IWC, averages over many observations result in close agreement as the number of data points increases. The utility of AIR thus depends upon the availability of large data sets that apply to roughly the same atmospheric conditions. For example, we will show CIPS results for July and January averages for ascending and descending portions of the orbit.

In this paper we first describe the theoretical framework relating the scattered radiance to mesospheric ice particles. It is desirable to use model data to test the AIR method, without the complications of cloud heterogeneity and viewing geometry. We utilized a first-principles microphysical model that accurately simulates large numbers of cloud properties (number density and particle size distribution). The processes treated by the model include **nucleation on meteor ‘smoke’ particles**, growth, and sedimentation, occurring in a saturated environment at density and temperature conditions provided by the main global climate model (Bardeen et al., 2010). Several runs for one-day and multiple-day periods during summer solstice conditions for solar conditions applying to 1995 were analyzed. Cloud radiances

(albedos) at 265 nm were calculated for the SA range encountered by the CIPS experiment. We chose a set of cloud simulations to derive a single set of two AIR coefficients through linear regression. The accuracy of the AIR approximation was then tested on the same data, and on other model runs, using averages as a function of SA, and increasing IWC threshold values. Thresholding is necessary to account for the fact that different measurement techniques have different detection sensitivities. This is not a signal/noise issue, rather the ability to discriminate PMC against a background that is usually larger than the PMC signal itself. We show in particular how seasonal means of IWC can be derived from Solar Backscatter Ultraviolet Spectrometer (SBUV) radiance data, without the need to derive particle size.

1 Having tested the technique for model data, we use the same approach with real-life PMC data collected from CIPS in the normal pre-2016 operating mode. This mode provided scattering angles needed to define an ice scattering phase function, from which mean particle size was derived based on assumed properties of the underlying size distribution (Lumpe et al., 2013). The regressions were run for a period of 40 days in each of four seasons, each comprising millions of separate cloud measurements, and from both summertime hemispheres. The results were combined into a single set of AIR coefficients, and again the AIR technique was tested on monthly averages. These averages were constructed over all years of nominal spacecraft operations (2007-2013 in the northern hemisphere, and 2007-8 through 2013-14 in the southern hemisphere). Note that testing the accuracy of the AIR technique during the nominal mission period allows the method to be used even during the ‘continuous imaging mode’ of CIPS operation.

2 We then employed highly-accurate data from SOFIE for ice column density and mean particle size. **Since the SOFIE technique uses near-IR solar extinction in ice-water absorption bands, the primary measurement is ice water content. As shown in Sec 2.3, we inverted the retrieved SOFIE IWC to derive the equivalent 265-nm albedo, and then applied the regression method described above to the results.**

3 After describing the AIR method, we discuss briefly the application of the method to a third contemporaneous experiment, the SBUV satellite experiment, which has in common the same limitations as CIPS in its continuous-imaging mode, namely that measurements of nadir albedo are made at a single scattering angle. This has already resulted in a publication (DeLand and Thomas, 2015) where we provided a time series of PMC IWC from the AIR method extending back to the first SBUV experiment in 1979.

2 Theoretical Basis

Here we provide a brief overview of the theoretical basis of the IWC retrieval technique, referring to previous publications for more detail (Thomas and McKay, 1985; Rusch et al., 2009; Bailey et al., 2009, Lumpe et al., 2013). The basic measurement is PMC cloud radiance $I(\Phi, \theta)$ where Φ is the scattering angle (angle between the sun and observation vectors) and θ is the view angle, which is the angle subtended by the nadir and observation direction, measured from the point of scattering. Since the ice layer is optically thin, and secondary scattering is negligible, the albedo is described by first-order scattering. The ratio of scattered (detected) radiance to the incoming solar irradiance F_λ is the albedo A_λ , where

$$A_{\lambda}(\Phi, \theta) = I_{\lambda}(\Phi, \theta) / F_{\lambda} = \sec \theta \int_{z_t}^{z_b} dz' \int_{r_{\min}}^{r_{\max}} dr' \sigma_{\lambda}(r, \Phi) n(r', z'). \quad (1)$$

Here $n(r', z') dr' dz'$ is the **column** density of ice particles (cm^{-2}) in the ranges $r', r'+dr'$ and $z', z'+dz'$. For CIPS measurements, each camera has a finite bandpass, centered at 265 nm, and is characterized by a function R_{λ} with an effective width of 10 nm (McClintock et al., 2009). The albedo A_{λ}^m derived from this instrument is given by

Here z' and r' are the height and particle radius variables, and z_b and z_t define the height limits of the ice layer, with the majority of the integrand extending between 83 and 85 km. r_{\min} and r_{\max} are particle radii which span the particle size regime responsible for scattering (from ~ 20 nm to ~ 150 nm). **As shown by Rapp and Thomas (2006), particles with sizes < 20 nm are not detectable by UV measurements because of their small cross-section values – hence we refer to ‘UV-visible clouds’.** σ_{λ} is the **differential** scattering cross-section ($\text{cm}^2\text{-sr}^{-1}$) at wavelength λ and scattering angle Φ . $n(r', z') dr' dz'$ is the number density of ice particles (cm^{-2}) in the ranges $r', r'+dr'$ and $z', z'+dz'$. For CIPS measurements, each camera has a finite bandpass, centered at 265 nm, and is characterized by a function R_{λ} with an effective width of 10 nm (McClintock et al., 2009). The albedo A_{λ}^m derived from this instrument is given by

$$A_{\lambda}^m(\Phi, \theta) = \sec \theta \int d\lambda' R_{\lambda'} \int_{z_t}^{z_b} dz' \int_{r_{\min}}^{r_{\max}} dr' \sigma_{\lambda}(r, \Phi) n(r', z'). \quad (2)$$

In the model, the ice particles are assumed spherical, but the scattering theory should take account of the non-spherical nature of ice crystals. The best agreement of theory with near-IR mesospheric ice extinction occurs for a randomly rotating oblate- spheroid shape, of axial ratio two (Hervig and Gordley, 2010). This shape is assumed in the calculation of the cross-section, which is accomplished through a generalization of Mie-Debye scattering theory, the T-matrix method (Mishchenko and Travis, 1998). The radius in the T-matrix approach is defined as the radius of the volume-equivalent sphere. In the model calculations, we will ignore the view angle effect. In the reported CIPS data, the $\sec \theta$ factor is applied to the reported albedos, so that A always refers to the nadir albedo ($\theta = 0^\circ$).

The ice water content (IWC) is the integrated mass of ice particles over a vertical column through the layer. Its definition is

$$IWC = \rho \int_{z_b}^{z_t} dz' \int_{r_{\min}}^{r_{\max}} dr' (4\pi / 3) r'^3 n(r', z'). \quad (3)$$

ρ denotes the density of water-ice at low temperature (0.92 g-cm^{-3}). Anticipating the results of this study that IWC is linearly related to the column density of ice particles,

$N = \int dr' \int dz' n(r, z') dz'$, we explore the physical basis of this result. As pointed out by Englert and Stevens (2007) **and Hultgren and Gumbel (2014)** such a relationship exists for certain SA

values, for which $\sigma \sim r^3$, in which case it is easily seen that Eq. (2) is proportional to IWC. However, we find that a linear approximation is valid for a much wider range of scattering angles. To understand this result, we imagine that all particles have the same radius, so that $n = n_c \delta(r - r_c)$, where δ is the Dirac δ -function. Then Eqs. (1) and (3) ‘collapse’ to a simpler result,

$$A_\lambda(\Phi, 0) = \sigma_\lambda(r_c, \Phi)N(r_c), IWC(r_c) = \rho V(r_c)N(r_c). \quad (4)$$

Here $N(r_c) = n_c \Delta z$ where Δz is the effective vertical layer thickness. Eliminating the column density $N(r_c)$, IWC is written

$$IWC(r_c) = \rho V(r_c)A(\Phi, 0) / \sigma_\lambda(r_c, \Phi). \quad (5)$$

$V(r_c)$ denotes the particle volume. Thus in this special case, $IWC(r_c) \sim A_\lambda(\Phi, 0)$. A superposition of the effects of all participating particle sizes will exhibit a similar proportionality. When $IWC(r)$ is integrated over all r , the contributions from each size are straight lines, each having different intercepts and slopes.

As previously discussed, the value of the AIR method is in evaluating *average* IWC (denoted by $\langle IWC \rangle$) over many albedo observations made at numerous scattering angles. The accuracy of the method should be assessed primarily on this basis, not on how well an individual albedo measurement yields the correct value of IWC. **However we also address the error of using individual albedo measurements in estimating IWC.** An additional issue is the differing detection thresholds for IWC among the various experiments. In the case of the scattered-light experiments, the detection threshold depends upon how well the cloud radiance data can be separated from the bright Rayleigh-scattered background. The CIPS experiment retrieval method relies upon high spatial resolution over a large field of view, and the differing scattering-angle dependence of PMC and the Rayleigh-scattering background (Lumpe et al., 2013). The SBUV retrieval relies upon differing wavelength-dependence of PMC and background, but primarily on the PMC radiance residuals being higher (2 sigma) than fluctuations from a smoothly-varying sky background (Thomas et al., 1991; DeLand and Thomas, 2015). The AIM SOFIE method is very different, being a near-IR solar extinction measurement in multiple wavelength bands. SOFIE can detect much weaker clouds with smaller effective sizes than either CIPS or SBUV. Particle radii values as small as 10 nm can be retrieved from the SOFIE data (Hervig et al., 2009). To compare the various experiments, it is necessary to ‘threshold’ the data from more sensitive experiments with a cutoff value of IWC.

In the next three sections, we present the AIR results from the model, CIPS and SOFIE, using averages over many cloud occurrences. It is not our intention to compare the different thresholded data sets to one another (this task will be relegated to a separate publication), but to illustrate how even measurements made at a single scattering angle (e.g., SBUV) can yield averaged IWC values that are sufficiently accurate to assess variations in daily and seasonal averages. These variations are of crucial value to determining solar cycle and long-term trends

in the atmospheric variables (mainly temperature and water vapor) that control ice properties in the cold summertime PMC region. We examine the accuracy of AIR through simulations of scattered radiance from the model, and from CIPS and SOFIE data. Since these data sources yield particle radii, they can provide both the ‘actual’ and approximate values of IWC from the regression formulas. Hervig and Stevens (2014) used the spectral content of the SBUV data to provide limited information on particle size. Together with the albedos themselves, they used this information to derive seasonally-averaged ice water content. They showed that the variation of mean particle size over the 1979-2013 time period was relatively low (standard deviation of ± 1 nm). They also found a very small systematic increase with time, as discussed in Sec. 3.

2. 1 Model Results

Using a microphysical model as a reference source of IWC ‘data’ is useful, in the following ways: 1) in contrast to the CIPS and SOFIE retrieval algorithms, no artificial assumptions are needed concerning the size distribution of ice particles; (2) limitations due to background removal are absent; (3) radiance and IWC may be calculated accurately, so that effects of cloud inhomogeneity are absent. **With regard to the latter point, we describe in more detail the model calculations. The model grid is 4° in latitude, 5° in longitude and variable in the vertical. Ice particles of varying sizes fill many of these cells, but the density of particles within each cell is, by definition, constant. For a given model cloud, the integration is made through a vertical ‘stack’ of all ice-filled cells generated in a given computer run, and within each particle size grid. The total radiance is the sum of contributions from the size range 20 to 150 nm. The observation angles are always assumed to be zero, in other words, the integration is performed in the vertical only. Thus cloud ‘boundaries’ in the horizontal plane are not an issue. This contrasts with real heterogeneous clouds where these approximations would not hold. The model contains variability due to waves of various sorts, including tides and gravity waves. However, it does not capture all known details of PMC, such as double layers. Since we are dealing with integrated quantities, this should not be an important issue. Furthermore, we don’t place full reliance on the model, which is why we also use two independent data sets.**

To gain insight into the accuracy of the AIR approach, it is sufficient to work with monochromatic radiance at the central wavelength of the various passbands. The integrations of Eqs. (1) and (3) were approximated by sums over **variable** increments of radius, and over all sub-layers within the model ice cloud (a typical ice layer is several km thick.). **The model height grid is variable, so that the smallest layer thickness is 0.26 km, which resolves the narrow ice layers** (see Bardeen et al., 2010 for more details). We then performed the linear regression for SA values over which CIPS observations are made.

Figure 1 displays the regressions for six scattering angles, and 2514 individual model clouds. The units of IWC are $\text{g}\cdot\text{km}^{-2}$, or equivalently $\mu\text{g}\cdot\text{m}^{-2}$, which are commonly used in the literature. Each plot is divided into two groups according to the effective radii r_{eff} for each cloud. r_{eff} is defined in the literature (Hansen and Travis (1974) as

$$r_{eff} = \int dr' n(r') r'^3 / \int dr' n(r') r'^2. \quad (6)$$

Figure 1 clearly illustrates that particle size contributes to the ‘scatter’ from the linear fits. For the conditions in Fig. 1(c), the mean error of AIR for a *single* model simulation is 19%. The error can be reduced substantially by averaging. For example, for 100 measurements, the AIR error in the average IWC is only 2%. Figure 1 also shows the existence of a non-zero intercept of IWC vs albedo. The non-zero intercept was at first surprising since we expected that for an albedo of zero, IWC should also be zero. In fact, we found that the linear relationship breaks down for very small albedo, and the points in the plot narrow down in this limit (not shown). In albedo units of 10^{-6} sr^{-1} (hereafter referred to as 1 G) this departure from linearity occurs for $A < 1 \text{ G}$ and $IWC < 10 \text{ g-km}^{-2}$, conditions which fortunately are below the detection threshold of CIPS and SBUV, and are a result of the very faint small particles. For more sensitive detection techniques, this limitation must be kept in mind. A limitation of the present model (not necessarily all models) is that it does not simulate the largest particles in PMC and the largest values of IWC, as seen in both AIM SOFIE and CIPS experiments. The largest model IWC value is 180 g-km^{-2} and the largest effective radius is 66 nm, whereas CIPS and SOFIE find particle radii up to 100 nm and IWC up to 300 g-km^{-2} . This limitation is irrelevant for the AIR CIPS results (to be discussed), but could limit the application of the AIR technique to SBUV data. In Sec. 3 we will return to the issue of the AIR accuracy, as applied to SBUV data.

We chose to use averages for the entire model run, which includes different latitudes, longitudes, and UT, but the data can be **divided** in many different ways. It is certainly preferable in data sets to choose a small time and space interval over which temperature and water vapor are not expected to vary, but this is not necessary for the model. All that we ask of the model is whether the AIR results provide an accurate estimate of $\langle IWC \rangle$, taken over the ensemble of model cloud albedos calculated at a variety of scattering angles.

As discussed above, we are also interested in the accuracy of AIR in the thresholded data, that is, how AIR represents $\langle IWC \rangle$ in comparisons of data sets with varying detection sensitivities to PMC. Figure 2 displays the error in the ensemble-average (2488 model clouds) as a function of the IWC threshold and scattering angle. Despite the large data scatter from the linear fit shown in Fig. 1, the averaging removes almost all the influence of the ‘random error’. In this case, the overall error is less than 3%. The influence of particle size is of course not a random error, but acts like one in the averaging process. However, the AIR coefficients also depend weakly upon the mean effective radius, defined in Eq. (6) for a single cloud, which varies from one latitude to another and from year to year. The effect of variable r_{eff} on the AIR error is discussed in Sec. 3.

2.2 AIR Results from CIPS

A detailed description of the Version 4.20 CIPS algorithm, together with an error analysis of individual cloud observations, was presented in Lumpe et al. (2013). Here we describe only what is necessary to understand how IWC is derived from the data. Even though an accurate determination of the scattering-angle dependence of radiance (often called the scattering

phase function) is obtained by seven independent measurements, this does not fully define the distribution of particle sizes. Instead, additional constraints need to be introduced to derive the mean particle size. The particles are assumed to be the same oblate-spheroidal shape as defined for the model calculations, and to have a Gaussian size distribution (see eq. 11 in Rapp and Thomas, 2006). A relationship between the Gaussian width s and the mean particle radius r_m is derived from that found in vertically-integrated lidar data (Baumgarten et al., 2010). The net result is that two parameters, the mean particle size and the Gaussian width, are retrieved from a given scattering phase function. However, there is only one independent variable, since the two are related by $s(r_m)$. Thus Eq. (3) simplifies to

$$IWC = \rho V(r_m) A(\Phi = 90^\circ, 0) / \sigma_\lambda(r_m, \Phi). \quad (7)$$

V denotes the ice particle volume, averaged over the Gaussian distribution with a mean particle radius value r_m . A refers to the retrieved albedo, corrected to view angle $\theta = 0^\circ$ and interpolated to scattering angle $\Phi = 90^\circ$. Note the resemblance of Eq. (7) to Eq. (5). $A(\Phi = 90^\circ, 0)$, along with r_m and IWC are products reported in the CIPS PMC data base, found at (<http://lasp.colorado.edu/aim/>). $\sigma_\lambda(r_m, \Phi = 90^\circ)$ is the mean scattering cross-section, integrated over the assumed Gaussian distribution with mean radius r_m and distribution width s .

Before discussing the AIR results, we first apply the CIPS algorithm to the model data to test how well it works on a set of realistic particle sizes. As mentioned earlier, UV measurements of ice particles are not sensitive to particle radii < 20-25 nm. We applied the CIPS algorithm to 6672 model clouds, using seven scattering-angle points, spanning the range 50°-150° (the results are insensitive to the values chosen). We then calculated the % difference between the exact model calculation of IWC and the simulated CIPS retrieved IWC for every model cloud. Figure 3 shows the result as a function of $A(\Phi=90^\circ)$. **Assuming the microphysical model is accurate, the accuracy of the CIPS UV measurements ranges from over +100% for very small albedo to -60% for high albedos. We emphasize that this is not an AIR result, but is an attempt to assess how particles that are too small to be visible to UV measurements affect the accuracy of the CIPS IWC results.** The mean difference and standard deviation for the (albedo) bin-averages for two model days is $-13 \pm 17\%$. With the caveat that not all ice is retrieved, a large subset of CIPS IWC data thus has an acceptable accuracy (an average of 84% of the modelled ice mass is contained in particles with radii exceeding 23 nm). We note that IWC in the model used to derive the AIR approximation refers to *all* particle sizes.

The procedure for deriving AIR coefficients from the CIPS data is as follows: (1) Regression coefficients were derived from data pertaining to 0-40 days from summer solstice (day from solstice, DFS=0 to 40) on every third orbit. This meant that ~200 orbits per season were used. The regression analysis was performed on four years of data (2010-2013). The data were binned in 5-degree SA bins and only the best quality pixels with six or more points in the phase function were used; (2) data from each northern and southern summer season were treated separately. The coefficients and standard deviations of the fit were then interpolated to a finer

SA grid from 22° to 180° in increments of 1°; (3) The coefficients from each hemisphere were averaged, and these coefficients were then used to create an AIR IWC data base, to accompany the normal CIPS products. As previously shown, the AIR data applies to the ice mass of ‘UV-visible’ clouds, not to their total ice mass.

We emphasize that using the AIR data is unnecessary for seasons prior to the northern summer season of 2016 – however the AIR data have great importance since that time because the observing mode was changed, resulting in measured phase functions that contain many fewer (and often only one) scattering angles. As illustrated in Fig. 4, it is trivial to infer both IWC and $A(90^\circ)$ from a single measurement of albedo. This alternative 90-deg albedo value, ALB_AIR, is now included along with IWC AIR in the CIPS Level 2 data files. Fig. 5 shows the AIR results for monthly-averaged IWC (July and January) compared to the same averages of the more accurate results from the operational (OP) retrieval described in Lumpe et al. (2013). The data have been separated into different hemispheres, and into ascending and descending nodes of the sun-synchronous orbit, and apply to the years of the nominal operating mode. The ALB_AIR results are systematically higher than the operationally retrieved 90-deg albedo, whereas there is no consistent bias in the IWC (AIR) value compared to the operational product. However, for both quantities the interannual changes between the AIR and OP results agree very well. This is reflected in the very high correlation coefficients of the two sets of values. A more stringent test of the AIR method comes from daily values of CIPS IWC. Shown in Figs. 6 and 7 are polar projections of IWC (AIR) and the more accurate operational IWC data product. These ‘daily daisies’ are taken from overlapping orbit strips pertaining to 28 June of two different years. Figure 6 shows data from 2012, when CIPS was still in normal mode. The AIR result shows remarkable agreement with the operational IWC data. By 2016 (see Figure 7) CIPS is in continuous imaging mode and the standard IWC retrieval is limited due to the scarcity of pixels with three or more scattering angles. Here the AIR approach is clearly superior and does a good job of filling in the polar region where CIPS detects high-albedo clouds. The differences in patterns are due primarily to variations of particle size, rather than errors in the AIR method.

AIR accuracy can also be tested in the study of latitudinal variations. Figure 8 compares daily-averaged IWC from the CIPS Level 3C data, for both the standard and AIR algorithms, for the Northern Hemisphere 2011 season. It is clear that AIR is adequate even for 24-h averages. For example, it is capable of defining the beginning and ending of the PMC season, a metric that has valuable scientific value (e.g., Benze et al., 2012)

2.3 AIR Results from SOFIE

A third independent data set of IWC and particle size is available from the AIM SOFIE experiment. SOFIE provides very accurate values of IWC, through precise near-IR extinction measurements, independent of particle size. It assumes the same Gaussian distribution of particle sizes as CIPS, so that the reported value of mean particle radius r_m is consistently defined. SOFIE data are useful to investigate the extent to which the AIR approximation can be applied to an independent data set. To do so, it is necessary to calculate 265-nm albedo at various SA values, given the values of r_m , ice column density N from the data base, and the mean cross-section $\sigma_\lambda(r_m, \Phi)$. The latter quantity is averaged over the assumed Gaussian distribution. The equation for the albedo is

$$A_{\lambda}(\Phi, 0) = \sigma_{\lambda}(r_m, \Phi)N. \quad (8)$$

Given $A_{\lambda}(\Phi, 0)$ and IWC for each PMC measurement (one occultation per orbit), we can once again perform regressions and find AIR coefficients for the SOFIE data set. The comparison of AIR results from all three data sets is shown in Fig. 9, where the constant term C is the y-intercept and S is the slope in the AIR regression,

$$IWC(AIR) = C(\Phi) + S(\Phi) * A(\Phi, 0). \quad (9)$$

Figure 10 displays the results from the three data sets, expressed as contour plots of AIR-derived IWC as functions of SA and Albedo. This comparison shows that the three sets of IWC resemble one another far better than would be anticipated from the AIR coefficients in Fig. 9, where the constant coefficient differs significantly between data sets. Since the result of the regression in yielding IWC is more significant than the coefficients themselves, the comparisons of Fig. 10 are the more appropriate diagnostic. The fact that the IWC derived from AIR is more accurate than would be expected from the differing coefficients is due to the fact that the errors of the constant and slope coefficients are anti-correlated. The agreement between the three results will be even better when taken over a large data set with variable SA and albedo. The comparisons of IWC from different satellite experiments as a function of year and hemisphere will be the subject of a separate publication.

Figure 11 shows that the regressions with AIM SOFIE data obey a linear relationship between IWC and albedo for IWC <220 g-km⁻², but for SA values <90°, AIR overestimates IWC by up to 15%, depending upon the SA. For SA=110° the regressions are still linear up to 300 g-km⁻², values above which are seldom encountered in the data.

2.4 SBUV Data

The AIR coefficients from the model have been used by DeLand and Thomas (2015) to derive mean IWC from SBUV data, which spans the largest time interval of any satellite data set (1979-present). The 273 nm wavelength used in the SBUV Version 3 analysis is sufficiently close to the effective wavelength of the broader passband of the CIPS cameras (Benze et al., 2009) that the same coefficients may be applied to both data sets. The accuracy of the average IWC results was estimated by removing half the data from an entire season and comparing the two results. **For a highly-populated region (more than 1000 clouds per season at latitudes higher than 70°), the differences in IWC ranged between ±3–5 g-km⁻², thus can be considered typical systematic errors. For a less-populated region (50°–64° latitude) where there were many fewer clouds (<50), the differences were larger, ±5–10 g-km⁻².** Even the larger errors are sufficiently small for intercomparison of SBUV and contemporaneous PMC measurements. Figure 12 shows a comparison of SBUV IWC, using the model AIR coefficients, to the results of a more accurate

determination of IWC derived from particle size determinations using SBUV spectral information (Hervig and Stevens, 2014). The comparison is for data residuals from July averages over the time series 1979-2017. Given the different assumptions underlying the two data sets, the agreement is very good (with an rms difference of 3% for the residuals, and 5% for the actual values of $\langle IWC \rangle$).

3. Effects of Mean Particle Size

The AIR approximation is based on the notion that particle size effects can be ignored in retrieving IWC from albedo measurements, that is they contribute in a sense to the ‘noise’ of the measurement, which can be minimized by averaging. In fact, the particle size (or more accurately, the term r^3) is a principal ‘driver’ of $\langle IWC \rangle$ itself, so it is not obvious that particle size effects play a minor role in deriving IWC. The dependence of albedo on column density adequately captures this part of the variability (albedo is strictly linear in column density). The AIR slope term is $\sim r^3 / \sigma_\lambda(r, \Phi)$ averaged over a distribution of particle sizes, r . The size dependence of the cross-section varies as a power of r , within two limits, the geometric-optics limit, r^2 , and the small-particle (Rayleigh) limit, r^6 . In the intermediate and realistic conditions of PMC, the exponent has an intermediate value. Fortunately, there is a ‘sweet spot’ (or better, a ‘sweet region’ of the r -domain) in which the r -dependence of σ_λ is $\sim r^3$, so that the slope term is constant (for fixed SA). This behavior occurs for all relevant values of SA, and for the albedo values typical of CIPS. It accounts mainly for the effectiveness of the AIR method. The other aspect favorable to AIR is the steep fall-off of the particle size distribution at the largest sizes, which contributes to the sharpness of the lower boundaries in the spread of points in Fig. 1. Averaging over many values of r results in the AIR slope term that, in the limit of large number, the term depends predominantly on Φ . This is an example of ‘regression to the mean’, and illustrates how the approximation is designed to work for large numbers of clouds. In a fictitious case where the mean cloud particle size is larger in one year than another, but the cloud column number remains the same, the mean albedo would increase according to Eq. (8), resulting in an increase of $\langle IWC \rangle$. We might expect that the slope term would be different in the two cases. Our study with three different data sets shows that the regression slope itself remains almost the same among the three data sets, despite their differing in mean particle size.

In fact, Hervig and Stevens (2014) found from SBUV spectral data a small long-term trend in $\langle IWC \rangle$ and in addition a trend in the mean particle size ($+0.23 \pm 0.16$ nm/decade). This contributed an additional 20% to the overall long-term trend in $\langle IWC \rangle$. The ignored dependence on mean particle size using the AIR method thus adds a systematic uncertainty in derived $\langle IWC \rangle$ trends, which can be as large as 20%, according to their analysis. This error undoubtedly varies inversely with the number of clouds in the averaging process. For example, the number of CIPS observations per PMC season greatly exceeds that of SBUV, so that the error in $\langle IWC \rangle$ should be correspondingly smaller.

4. Conclusions

We have described the theoretical basis and accuracy for an approximation for retrieving the average ice water content (IWC) of Polar Mesospheric Clouds (PMC) from measurements of UV

albedo at a single scattering angle. This approach provides a continuous set of consistent CIPS measurements of IWC from year to year, regardless of the number of scattering angles for which albedo at a single location is measured. The consistent AIR IWC data base enables robust IWC comparisons throughout the AIM mission, from 2007 to the present. A comparison of IWC derived from the microphysical model and from the CIPS algorithm suggests that CIPS is capable of measuring 84% of the total ice content of PMC (for particle sizes exceeding ~ 23 nm).

Assuming the microphysical model is accurate, the accuracy of the CIPS UV measurements ranges from over +100% for very small albedo to -60% for high albedos. The overall accuracy of IWC (averaging over all albedo bins) is $-13\pm 17\%$. The CIPS algorithm overestimates the small-particle population (20-30 nm) as a result of the Gauss approximation when the mean particle size is small, and the opposite is true when the mean size is large. These errors are a result of the CIPS approximations and the invisibility of small particles, and are irrelevant to the AIR approximation.

Distinct from the more fundamental errors due to the invisibility of very small ice particles and the Gaussian approximation, we also estimated the errors in the AIR approximation, relative to the AIM SOFIE data which apply to larger values of IWC than the model. AIR is less accurate for high IWC (>220 g-km⁻²), but very-high mass clouds (IWC > 300 g-km⁻²) are infrequent and do not influence seasonal averages of IWC. For the dimmer and more frequent clouds, Fig. 2 shows that the error in ensemble averages is of the order of 3%. The accuracy of the AIR results for ensemble averages has a small systematic dependence on mean particle size- the error depends inversely on the size of the ensemble. The inter-annual and hemispheric variations of IWC derived from CIPS and SBUV measurements throughout an entire 11-year period (2007-2018) will provide detailed information on PMC variability over the recent solar cycle 24.

Figure Captions

Figure 1. Linear regressions of model PMC albedo versus model PMC ice water content. The black points represent model clouds with $r_{\text{eff}} < 40$ nm. The red points apply to $r_{\text{eff}} > 40$ nm. The blue line is the linear least-squares fit to all points. (a) through (f) are for different scattering angles. The lower limit for the albedo (SA=90°) is 1G, which is the detection limit of the CIPS experiment.

Figure 2. Relative errors of ensemble averages, $\langle \text{IWC} \rangle$ using the AIR approximation, taken over all cloud model simulations for conditions of summer solstice. $\langle \text{IWC} \rangle$ is 'thresholded' by the variable IWC in the vertical axis, so that $\langle \text{IWC} \rangle$ applies to all values above IWC. Contour lines are labelled as percent errors relative to the accurate model values.

Figure 3. Differences of IWC derived from the model cloud 'data' and the accurate IWC from the model, plotted against the 265-nm albedo (in G units, see text), evaluated at SA=90°. The error bars are the standard deviations in intervals of 2G.

Figure 4. Illustration showing how IWC=98 g-km⁻² (horizontal arrow) and A(90°) = 16 G (thick downward arrow) are derived from the AIR method from a single measurement of cloud albedo at 60 G and SA=50° (upward arrow). Each straight-line plot is calculated from Eq. (9).

Figure 5. Comparison of CIPS $A(90^\circ)$ (top) and $\langle IWC \rangle$ (bottom) calculated from the operational (OP) and AIR algorithms. Data points correspond to July northern hemisphere (NH) and January southern hemisphere (SH) averages in a 5-degree latitude bin centered at 70° . Left and right panels are for ascending and descending legs data, respectively.

Figure 6. Polar projection map of IWC from CIPS, Day 180 (28 June 2012). Left and right panels show the operational IWC product and the AIR result, respectively.

Figure 7. Same as Fig. 6 except for 28 June 2016.

Figure 8. Filled circles and dotted line: IWC (AIR) averaged over 1-deg latitude bins centered on 70° (green) and 80° (blue), and over 15 orbits (from which daily averages are derived). Solid line: standard L3C IWC averaged in the same way.

Figure 9. AIR coefficients for three different sources of IWC and particle size: Model (solid line with open circles), CIPS (solid line), and SOFIE (dashed line).

Figure 10. Contour plots of the AIR approximations for IWC versus cloud albedo (G) for the three data sources: (a) model, (b) SOFIE, and (c) CIPS

Figure 11. Examples of SOFIE AIR regressions for two (specified) scattering angles, (a) 80° and (b) 110° .

Figure 12. Comparison of annually-averaged northern hemisphere July-averaged residuals ($\langle IWC \rangle$ -long-term mean) derived by two independent methods from SBUV 273 nm albedo data. Black curve: $\langle IWC \rangle$ derived from the AIR approximation. Blue curve: $\langle IWC \rangle$ derived from the same SBUV albedo data, but including mean particle size variations (see text). A three-year smoothing has also been applied.

Data Availability

The CIPS operational PMC data, along with the AIR data, can be found at <http://lasp.colorado.edu/aim/>.

Author Contribution

Author G.T. formulated the AIR approximation, and derived the AIR coefficients from the microphysical model (provided by author C.B.), and from the AIM SOFIE data (<http://sofie.gats-inc.com/sofie/index.php>). Authors J.L. and C.R. calculated the AIR coefficients from the CIPS data.

Disclaimer

The authors declare that they have no conflict of interest.

Acknowledgements

We thank M. DeLand and M. Hervig for providing us with the data used in Fig. 12. We gratefully acknowledge the tremendous effort of the engineering, mission operation and data systems

teams whose dedication and skill resulted in the success of the CIPS instrument. **The contributions of two anonymous reviewers greatly enhanced the clarity of the manuscript.** AIM is funded by NASA's Small Explorers Program under contract NAS5-03132.

References

Bardeen, C. G., O. B. Toon, E. J. Jensen, M. E. Hervig, C. E. Randall, S. Benze, D. R. Marsh, and A. Merkel: Numerical simulations of the three-dimensional distribution of polar mesospheric clouds and comparisons with Cloud Imaging and Particle Size (CIPS) experiment and the Solar Occultation For Ice Experiment (SOFIE) observations, *J. Geophys. Res.*, 115, D10204, doi:10.1029/2009JD012451, 2010.

Bailey, S.M., Thomas, G.E., Rusch, D.W., Merkel, A.W., Jeppesen, C., Carstens, J. N., Randall, C.E., McClintock, W.E., Russell III, J.M.: Phase functions of polar mesospheric cloud ice as observed by the CIPS instrument on the AIM satellite, *J. Atmos., Sol.-Terr. Phys.*, 71, 373-380, doi:10.1016/j.jastp.2008.09.039, 2009.

Benze, S., Randall, C. E., DeLand, M. T., Thomas, G.E., Rusch, D. W., Bailey, S.M., Russell III, J. M., McClintock, W., Merkel, A. W. and Jeppesen, C.: Comparison of polar mesospheric cloud measurements from the Cloud Imaging and Particle Size experiment and the Solar Backscatter Ultraviolet instrument in 2007, *J. Atmos. Solar-Terr. Phys.*, 71, 365-372, 2009.

Benze, S., Randall, C.E., Karlsson, B., Harvey, V.L., DeLand, M.T., Thomas, G.E., Shettle, E.P.: On the onset of polar mesospheric cloud seasons as observed by SBUV, *J. Geophys. Res.* 117 <http://dx.doi.org/10.1029/2011JD017350>. D07104, 2012.

DeLand, M. T., Shettle, E. P., Thomas, G. E. and Olivero, J. J.: A quarter-century of satellite PMC observations, *J. Atmos. Sol. Terr. Phys.*, 68, 9–29, 2006.

DeLand, M. T., and Thomas, G. E.: Updated PMC trends derived from SBUV data, *J. Geophys. Res. Atmos.*, 120, doi:10.1002/2014JD022253, 2015.

Englert, C. R. and Stevens, M. H.: Polar mesospheric cloud mass and the ice budget: 1. Quantitative interpretation of mid-UV cloud brightness observations, *J. Geophys. Res.*, 112, D08204, doi:10.1029/2006JD007533, 2007.

Gordley, L. L., Hervig, M. E., Fish, C., Russell III, J. M., Bailey, S. M., Cook, J., Hansen, J., Shumway, A., Paxton, G., Deaver, L., Marshall, T., Burton, J., Magill, B., Brown, C., Thompson, E., and Kemp, J.: The solar occultation for ice experiment, *J. Atmos. Sol. Terr. Phys.*, 71, 300-315, 2009.

Hansen, J. E. and Travis, L.D.: Light scattering in planetary atmospheres, *Space Sci. Rev.* 16, 527-610, 1974.

Hervig, J. E., Gordley, L.L., Stevens, M. H., Russell III, J. M., Bailey, S. M., and Baumgarten: Interpretation of SOFIE PMC measurements: Cloud identification and derivation of mass density, particle shape, and particle size, *J. Atmos. Sol. Terr. Phys.*, 71, 316-330, 2009.

Hervig, M. E., and Gordley, L. L.: Temperature, shape, and phase of mesospheric ice from Solar Occultation for Ice Experiment observations, *J. Geophys. Res.*, 115, D15208, doi:10.1029/2010JD013918, 2010.

Hervig, M. E., and Stevens, M. H.: Interpreting the 35-year SBUV PMC record with SOFIE observations, *J. Geophys. Res. Atmos.*, 119,12,689–12,705, doi:10.1002/2014JD021923,2014.

Hervig, M. E., Berger, U., & Siskind, D. E.: Decadal variability in PMCs and implications for changing temperature and water vapor in the upper mesosphere, *J. Geophys Res. Atmos.*, 121, 2383–2392, <https://doi.org/10.1002/2015JD024439>, 2016.

Hultgren, K. and Gumbel, J.: Tomographic and spectral views on the lifecycle of polar mesospheric clouds from ODIN/OSIRIS, *J. Geophys Res. Atmos.*, 119, 14,129-14,143, doi:10.1002/ 2014JD022435.

Lübken, F.-J., Berger, U., and Baumgarten, G.: On the anthropogenic impact on long-term evolution of noctilucent clouds, *Geophys. Res. Lett.*, 45, <https://doi.org/10.1029/2018GL077719>, 2016.

Lumpe, J.D., Bailey,S.M.,Carstens,J.N.,Randall,C.E.,Rusch,D.W.,Thomas,G.E.,Nielsen,K., Jeppesen, C., McClintock, W.E.,Merkel,A.W.,Riesberg,L.,Templeman, B., Baumgarten,G.,Russell III, J.M.: Retrieval of polar mesospheric cloud properties from CIPS: algorithm description, error analysis and cloud detection sensitivity, *J.Atmos.Sol.- Terr.Phys.*,104,167–196, [http://dx.doi.org/ 10.1016/j.jastp.2013.06.007](http://dx.doi.org/10.1016/j.jastp.2013.06.007), 2013.

McClintock, W.E., Rusch, D.W., Thomas, G.E., Merkel, A.W, Lankton, M.R., Drake, V.A., Bailey, S.M. and Russell III J.M.: The cloud imaging and particle size experiment on the Aeronomy of Ice in the mesosphere mission: Instrument concept, design, calibration, and on-orbit performance, *JASTP* 71, 340-355, doi:10.1016/j.jastp.2008.10.011, 2009.

Mishchenko, M.I., Travis, L.D.: Capabilities and limitations of a current Fortran implementation of the T-matrix method for randomly oriented, rotationally symmetric scatterers. *J. Quant. Spectrosc. Radiat. Transf.* 60, 309–324, 1998.

Rapp, M. and G. E. Thomas: Modeling the microphysics of mesospheric ice particles: Assessment of current capabilities and basic sensitivities, *J. Atmos. Sol.-Terr. Phys.*, 68, 715- 744, 2006.

Russell, J. M. III,, Bailey, S.M., Gordley, L. L., Rusch, D.W., Hora'nyi, M, Hervig, M. E., Thomas, G. E.,Randall, C. E, Siskind, D. E., Stevens, M.H., Summers, M. E., Taylor, M. J., Englert, C. R., Espy,

P. J., McClintock, W. E. and Merkel, A.W.: The Aeronomy of Ice in the Mesosphere (AIM) mission: Overview and early science results, *J. Atmos. Sol.-Terr. Phys*, 289 –299, 2009.

Rusch, D. W., Thomas, G.E., McClintock, W., Merkel, A.W., Bailey, S.M., Russell III, J.M., Randall, C.E., Jeppesen, C. and Callan, M., The cloud imaging and particle size experiment on the aeronomy of ice in the mesosphere mission: Cloud morphology for the northern 2007 season, *J. Atmos. Sol.-Terr. Phys*, 356–364, 2009.

Thomas, G.E. and McKay, C.P.: On the mean particle size and water content of polar mesospheric clouds. *Planetary Space Science* 33 (10), 1209–1224, 1985.

Thomas, G. E., Olivero, J. J., Jensen, E. J., Schröder, W., and Toon, O. B.: Relation between increasing methane and the presence of ice clouds at the mesopause. *Nature*, 338, 490–492, 1989.

Thomas, G. E., McPeters, R. D. and Jensen, E. J.: Satellite observations of polar mesospheric clouds by the Solar Backscattered Ultraviolet radiometer: Evidence of a solar cycle dependence, *J. Geophys. Res.*, 96, 927–939, .

Albedo-Ice Regression, SD-WACCM/CARMA microphysical model




Figure 1

AIR Error in $\langle IWC \rangle$: Model




Figure 2




Figure 3




Figure 4

Comparison of OP retrieval to AIR: DFS=[10,40] ; Lat = 70°
 ASC DSC




Figure 5




Figure 6




Figure 7

CIPS Level 3C IWC - NH 2011




Figure 8




Figure 9






Figure 10





Figure 11




Figure 12