

We thank the reviewer for his/her careful reading of the manuscript and offering many constructive feedbacks and helpful suggestions. We have incorporated most of the suggestions and believe the revised paper is substantially improved. In order for the reviewers and the editor to more readily identify our changes, we've submitted two versions of the revised paper, one with "track changes" and the other with same changes incorporated.

Because we made substantial revisions, we include below a list of main changes, after which we provide a detailed response to the reviewer's comments. The original comments by the reviewer are in black font, our replies in blue.

Major changes in the revised manuscript:

- We added an appendix to investigate the potential retrieval error in ALH due to several assumptions made in the retrieval algorithm, including the surface reflectance, smoke single scattering albedo, aerosol optical depth, and half width of the assumed quasi-Gaussian aerosol vertical profile.
- In the validation for EPIC-retrieved ALH, we now use two sets of CALIOP-based ALH (updated Figures 9 and 10): one with a background aerosol amount added for undetected CALIOP aerosol layers, the other one without. While there is a mean difference of 0.36 km between these two sets of CALIOP ALHs, we found EPIC ALH retrievals are in general consistent with the both sets with a RMSE of 0.57-0.58 km.
- The language of the manuscript has been substantially improved by incorporating reviewers' comments and authors' further proof reading.

Anonymous Referee #1

This paper demonstrates nicely that aerosol layer height information can be retrieved from EPIC/DSCOVER data. This is especially of interest since hourly information can be

retrieved – a unique contribution indeed. The paper should be published following minor revisions which are mostly due to suggested minor phrasing corrections.

We thank the reviewer's positive comments to the significance of this article.

Page 7, lines 20-23. These two sentences are awkward. The sentence "Besides, cloud mask thresholds" leaves the reader in a state of uncertainty. The phrase "might need.." is inconclusive. It is suggested to delete the sentence "Besides, cloud mask thresholds". Perhaps one can replace this sentence with "This is a topic of further investigation."

Following the reviewer's suggestion, we removed the sentence "Besides, cloud mask thresholds ... operational applications."

Page 10, lines 10-15. The sentence "To compensate for this bias,.. " is not clear. I am having difficulty in accepting the methodology used to account for undetected aerosol. How can one impose an exponentially-decaying background aerosol amount to an undetected aerosol layer if you don't know if the undetected aerosol is there or not? To assume that undetected aerosol is everywhere is problematic. The amount of undetected aerosol likely varies from place to place. Furthermore, the summertime Arctic aerosols do not correspond to conditions elsewhere. I think it would be best to estimate the bias in ALHCALIOP due to the undetected aerosol for a number of observations, state the uncertainty in the paper, and then calculate ALHCALIOP without adding undetected AOD amounts anywhere.

Thank you for this suggestion. Through a close inspection, we found that the background aerosol profile was not added in the CALIOP ALH shown in Figures 9 and 10a. So, in the revised manuscript, we keep those figures based on your suggestion. At the same time, in the validation analysis we also included the CALIOP ALH by adding undetected background aerosol. We found the mean difference between these two sets of CALIOP ALH was 0.36 km, and our EPIC ALH retrievals are consistent with both sets

of CALIOP ALH with a RMSE of 0.57-0.58 km (see the revised Figures 9 and 10 in the revised manuscript).

Page 11, lines 28-29. What are typical ALH uncertainties due to MODIS surface products uncertainties and GOME-2 LER uncertainties?

To answer this question, we added an error analysis and discussion (in the article Appendix) for ALH retrievals due to uncertainties in surface reflectance and other parameters. We found a surface reflectance error of 0.01 may lead to ALH retrieval errors from 0.1 km to 0.6 km for both water and vegetation surface types, depending on the aerosol loading and vertical allocation.

Technical corrections:

Page 1, line 3: change to “from the EPIC/DSCOVR”

Corrected.

Page 1, line 38: change to “temperature, influence the measured aerosol extinction profiles”

Corrected.

Page 3, line 4: change to “aerosol extinction profiles measured”

Corrected.

Page 3, line 20: change to “ALH utilizing the O₂”

We changed to “ALH using the O₂”.

Page 6, line 2: change to “where $C(\lambda)$ is the EPIC”

Corrected.

Page 6, line 29: change to “from analyzing USGS (United States Geological Survey)”

Corrected.

Page 7, line 30: change to “constructed with the UNL-VRTM model.”

We changed to “constructed using the UNL-VRTM model.”

Page 8, line 1: change to “It also incorporates HITRAN spectroscopic gaseous absorption with up to 22 trace gases”

Corrected.

Page 8, line 27: change to “satellite instrument, separate over-land”

Corrected.

Page 10, line 6: change to “(2013), the CALIOP day time aerosol extinction threshold is 0.01 – 0.03 km⁻¹ for 80-km horizontal resolution and up to 0.07 km⁻¹ for 5-km horizontal resolution.”

We changed to “(2013), aerosol extinction threshold in a daytime CALIOP scan is 0.01 – 0.03 km⁻¹ for 80-km horizontal averaging resolution and increases to 0.07 km⁻¹ for 5-km horizontal averaging resolution.”

Page 10, line 19: change to “65% of the ALH retrievals are within an uncertainty envelope of”

Corrected.

Page 10, line 22: change to “The collocation method follows Ichoku et al. (2002), but was”

Corrected.

Page 10, line 33: change to “smoke by using the UV aerosol”

Corrected.

Page 11, line 9: change to “satellite, since both perform hyperspectral measurements from the UV to the NIR and both cover the O2 A and B bands”

We changed to “satellite, both of which obtain hyperspectral measurements from the UV to the NIR covering the O2 A and B bands”

Page 11, line 21: change to “dust ALH from the EPIC experiment (Xu”

We changed to “dust ALH from the EPIC experiments”

Page 11, line 28: change to “information. Surface reflectance values are specified using MODIS”

Corrected.

Page 12, line 5: change to “The three years of data recorded” Page 12, line 18: change to “group at the University of Iowa”

Corrected.

Page 12, line 20: change to “acknowledge the AERONET program”

Corrected.

Page 20, line 6: change to “includes all ten EPIC bands,”

Corrected.

Page 21, line 8: change to “resulting in less absorption by O₂ and “ Page 22, line 4: change to “and surface reflectance (A_s) values”.

Corrected.

We thank Dr. Feng Xu for his careful reading of the manuscript and offering many constructive feedbacks and helpful suggestions. We have incorporated most of the suggestions and believe the revised paper is substantially improved. In order for the reviewers and the editor to more readily identify our changes, we've submitted two versions of the revised paper, one with "track changes" and the other with same changes incorporated.

Because we made substantial revisions, we include below a list of main changes, after which we provide a detailed response to the reviewer's comments. The original comments by the reviewer are in black font, our replies in blue.

Major changes in the revised manuscript:

- We added an appendix to investigate the potential retrieval error in ALH due to several assumptions made in the retrieval algorithm, including the surface reflectance, smoke single scattering albedo, aerosol optical depth, and half width of the assumed quasi-Gaussian aerosol vertical profile.
- In the validation for EPIC-retrieved ALH, we now use two sets of CALIOP-based ALH (updated Figures 9 and 10): one with a background aerosol amount added for undetected CALIOP aerosol layers, the other one without. While there is a mean difference of 0.36 km between these two sets of CALIOP ALHs, we found EPIC ALH retrievals are in general consistent with the both sets with a RMSE of 0.57-0.58 km.
- The language of the manuscript has been substantially improved by incorporating reviewers' comments and authors' further proof reading.

Dr. Feng Xu (Referee #2)

In this paper, Xu et al. extends their earlier work on aerosol layer height (ALH) and aerosol optical depth (AOD) retrieval from ocean to land - using EPIC observations. Comparison to CALIOP retrieval of ALH and AERONET product of AOD show an error

of 0.58 km in retrieved ALH and an error of 0.05 in retrieved AOD, respectively. This work was very well organized. I don't have questions on the general technical routines presented in this work as the previous publications by the authors along this line of research have laid down a solid basis for proceeding with this work.

We thank the reviewer's positive comments to the significance of this article.

My comments below are for the authors to somehow clarify their approach that readers can better digest the ideas behind it:

1. What is source of oxygen profile adopted in EPIC ALH and AOD retrievals ? Is there any impact from temperature on oxygen profile and then on ALH retrieval ?

The O₂ mixing ratio profile is obtained from the mid-latitude standard atmosphere (McClatchey et al., 1972), with surface pressure enlisted from MERRA-2 data. O₂ is well mixed in the atmosphere. O₂ absorption cross sections are calculated from the HITRAN spectroscopic line parameters in a very high spectral resolution (see Table 2). The high-spectral radiance data then convolved to EPIC bands. Although temperature and pressure have influence to the line position and line width of spectral O₂ absorption. Such influence tends to be negligible in the convolved spectral radiance. So, we don't think there is any impact from temperature on oxygen profile and ALH retrieval.

We have clarified this in section 3.3: "... Here, we enlisted surface pressure information from the Modern-Era Retrospective analysis for Research and Applications Version 2 (MERRA-2) datasets (Gelaro et al., 2017). MERRA-2's 1-hourly surface pressure at 0.5° by 0.675° grids were interpolated to the location and scan time of each EPIC pixel. In addition, the atmospheric temperature-pressure profile also impacts the width and strength of O₂ absorption lines. However, such influence on the radiative transfer is negligible for EPIC's 1-to-2-nm-wide bands. In this study, our algorithm employs a

standard temperature-pressure profile representing the mid-latitude-summer atmosphere (McClatchey et al., 1972).”

2. The sensitivity of O2-A and B bands to the profiles of smoke aerosols are clear from Figure 3. I’m curious how much the sensitivity will change if aerosol absorption changes gradually from being strongly absorbing to being weakly absorbing?

We performed sensitivity and error analysis to answer this question in the Appendix. We found the DAOS ratios used for ALH retrieval are sensitive to SSA, especially for large AOD values (see Figure A1e). However, SSA only has marginal impact to the ALH retrieval error (see Figure A2).

3. I believe the authors have published it elsewhere, but it would be helpful to some readers if some comments from authors’ side can be made on the sensitivity of O2-A and B bands to the width of aerosol layer.

In our retrieval algorithm we fixed the profile half-width as 1 km, because EPIC measurements are not able to resolve both the ALH and half-width at the same time. In the added appendix, we also performed sensitivity and error analysis to answer this question. We found the DAOS ratios used for ALH retrieval are sensitive to the profile half-width, especially for elevated aerosols (see Figure A1c). From the error analysis, we found a change of 0.5 km in the half-width may cause up to 0.3 km error in the retrieved ALH (Figure A2).


4. To give people a better idea about the “real” sensitivity that measurements have to AOD and ALH, it will be nicer if the authors can describe in the caption of Figure 3 EPIC’s measurement errors in the two O2 bands. Probably it would be clear if Z-score

(ratio of difference of signals in A/B band signals as normalized by measurement errors) is plotted as its axis.

We thank the reviewer for this suggestion. We added that uncertainty in EPIC DOAS ratios are within 3%. In addition, we added Jacobian of DOAS ratio to ALH, as well as the ALH retrieval error due to the EPIC measurements error, in the appendix. We hope those can give readers a better idea of about the real sensitivity.

5. Page 5, though there is a reason (geolocation) on the aggregation of pixels into a box of 3 x 3 pixels, do the authors think the price to pay (reduction of spatial resolution from 24 km to 8 km) too high. What if retrieval is directly implemented on 1 by 1 grid to retain EPIC's original 8 km resolution?

We indeed have tried the retrieval with native EPIC resolution, which is shown in the figure below for the same smoke case on August 25 in the article. The only reason we do 3x3 pixel aggregation is the EPIC geolocation. I understand EPIC version 3 data with better geolocation and calibration will be released soon. We have a plan to implement the native-resolution retrievals with the new EPIC data.



6. Page 8, the smoke particle properties are described for retrieving AOD and ALH. It might helpful if some comments can be given regarding if the pre-determined aerosol model have certain errors and its potential impact on ALH and AOD retrieval accuracy.

Sure. In the added appendix, we included the analysis and discussion about the sensitivity of our retrieval algorithm and the impacts for ALH retrievals from uncertainties in AOD and in the assumed aerosol model (such as SSA and the Gaussian profile width).

Some suggested editorial changes:

1. The reference for Lewis [1994] is not complete. Second author's last name is missing. Correct the citation of this reference in line 24 of Page 6 as well.

The reference is corrected as:

Lewis, P., and Barnsley, M. J.: Influence of the sky radiance distribution on various formulations of the earth surface albedo, in: Proc. Conf. Phys. Meas. Sign. Remote Sens, Val d'Isere, France, 707-715, 1994.

And the citation for this reference is changed to "Lewis and Barnsley (1994)"

2. Delete the redundant set of words "over land" in line 26 of Page 8.

Corrected.

3. Line 27 of Page 8: should "separated" be "separate"?

Corrected.

We thank the reviewer for his/her careful reading of the manuscript and offering many constructive feedbacks and helpful suggestions. We have incorporated most of the suggestions and believe the revised paper is substantially improved. In order for the reviewers and the editor to more readily identify our changes, we've submitted two versions of the revised paper, one with "track changes" and the other with same changes incorporated.

Because we made substantial revisions, we include below a list of main changes, after which we provide a detailed response to the reviewer's comments. The original comments by the reviewer are in black font, our replies in blue.

Major changes in the revised manuscript:

- We added an appendix to investigate the potential retrieval error in ALH due to several assumptions made in the retrieval algorithm, including the surface reflectance, smoke single scattering albedo, aerosol optical depth, and half width of the assumed quasi-Gaussian aerosol vertical profile.
- In the validation for EPIC-retrieved ALH, we now use two sets of CALIOP-based ALH (updated Figures 9 and 10): one with a background aerosol amount added for undetected CALIOP aerosol layers, the other one without. While there is a mean difference of 0.36 km between these two sets of CALIOP ALHs, we found EPIC ALH retrievals are in general consistent with the both sets with a RMSE of 0.57-0.58 km.
- The language of the manuscript has been substantially improved by incorporating reviewers' comments and authors' further proof reading.

Anonymous Referee #3

General Comments:

This work builds upon the authors' prior work on retrieving aerosol layer height (ALH) over ocean by extending the algorithm to work over land surfaces. One of the most important differences between the ocean and land studies is the characterization of surface reflectance, which is much more variable and complex than that over ocean. Further, their algorithm has a new smoke aerosol model for retrieving biomass-burning smoke ALH. The ALH retrievals are critical for obtaining accurate aerosol products in the UV. They also employ new data aggregation and spectral fitting strategies. Overall, the work is sound and relevant, and should be published.

We thank the reviewer's positive comments to the significance of this article.

Specific Comments:

Since the work focuses on land surfaces, what are the effects of non-Lambertian surface reflection (BRDF) on the retrievals?

By neglecting BRDF effects in the surface reflectance, the ALH retrieval could be biased. Further studies are needed to examine the detailed impacts. On the other hand, since EPIC measurements are mostly in the back-scattering direction, the impacts from BRDF could be limited. This will be one of our future efforts to improve the ALH retrieval accuracy.

So we added following text in section 3.3 to clarify this: "It should be noted that the effects of non-Lambertian surface reflection may bias the ALH retrieval, although this type of impact is limited by EPIC's backscattering direction only observation pattern. Further studies are needed to examine the detailed impacts, which will be one of our future efforts."

Is there a difference in ALH retrieval sensitivity as a function of single scattering albedo?

We performed sensitivity and error analysis to answer this question in the Appendix. We found the DAOS ratios used for ALH retrieval are sensitive to SSA, especially for large

AOD values (see Figure A1e). However, SSA only has marginal impact to the ALH retrieval error (see Figure A2).

What are the uncertainties in ALH retrievals due to uncertainties in the surface albedo and pressure climatologies used in the algorithm?

The uncertainties in ALH retrievals due to uncertainties in surface albedo are analyzed and discussed in the added Appendix. We found a surface reflectance error of 0.01 may lead to ALH retrieval errors from 0.1 km to 0.6 km for both water and vegetation surface types, depending on the aerosol loading and vertical allocations.

We indeed use surface pressure from MERRA-2 reanalysis data, applied to the high-latitude summer standard atmosphere for atmospheric pressure levels. Since, O₂ is well mixed the atmosphere, we believe the use of MERRA-2 surface pressure can well capture the amount of O₂, and thus has negligible uncertainty due to the atmospheric pressure.

There are a lot of grammatical and typographical errors (way more than acceptable for a manuscript published in this journal) in the manuscript, which I have tried to capture in the technical comments section. These must be corrected before the manuscript can be published.

We really thankful to the reviewer for his/her careful reading of the manuscript and providing very detailed technical comments. We have corrected all of these comments as listed below.

Technical Comments:

Line 18, page 1: in visible -> in the visible

Corrected.

Line 19, page 1: flexible spectral fittings that account for -> flexible spectral fitting that accounts for

Corrected.

Line 21, page 1: to derive the ALH that represents an optical centroid altitude -> to derive ALH, which represents an optical centroid altitude

Corrected.

Line 22, page 1: the measurements -> measurements Line 23, page 1: United State -> the United States

Corrected.

Lines 23-24, page 1: the algorithm can well capture -> that the algorithm can be used to obtain

Corrected.

Lines 24-25, page 1: Validations are performed against aerosol extinction profile -> Validation is performed against aerosol extinction profiles

Corrected.

Line 26, page 1: and AOD -> , and against AOD Line 26, page 1: in average -> on average

Corrected.

Line 29, page 1: the EPIC's -> EPIC's

Corrected.

Line 32, page 1: earth's -> Earth's

Corrected.

Line 34, page 1: arrange the authors alphabetically

Done.

Lines 37-38, page 1: The thermal signature of dust in particular can likewise influence the earth longwave budget and through the interference of retrievals of water vapor and temperature, thus influencing measure atmospheric state -> The thermal signature of dust, in particular, can likewise influence the Earth's longwave budget, and through interference with retrievals of water vapor and temperature, influence measurement of the atmospheric state

Done.

Line 39, page 1-line 3, page 2: Additionally, the knowledge of ALH is essential in retrieving aerosol absorption properties . . . , in retrieving aerosol microphysical properties . . . , and in the atmospheric correction for ocean color remote sensing -> Additionally, knowledge of ALH is essential for retrieving aerosol absorption properties . . . , aerosol microphysical properties ..., and for atmospheric correction for ocean color remote sensing

Done.

Line 5, page 2: arrange the authors alphabetically; leave a space between the two citations

Done.

Line 18, page 2: from UV to near-infrared-> from the UV to the near-infrared

Corrected.

Line 19, page 2: Figure 1b-c -> Figures 1b-c

Corrected.

Line 20, page 2: the spectral -> spectral

Corrected.

Line 22, page 2: present -> presented

Corrected.

Line 23, page 2: the EPIC -> EPIC

Corrected.

Line 24, page 2: demonstrate -> demonstrated

Corrected.

Line 26, page 2: in determining -> for determining

Corrected.

Line 28, page 2: robust strategies in the -> robust strategies for the

Corrected.

Line 33, page 2: to retrieving -> for retrieving

Corrected.

Line 33, page 2-line 1, page 3: implicating our algorithm development limited to water and vegetated land surface -> limiting our algorithm development to water and vegetated land surfaces

Done.

Line 1, page 3: the assumptions -> assumptions

Corrected.

Lines 2-3, page 3: The ALH retrievals are demonstrated in Section 4, that were applied to smoke events over Canada and the United States in August 2017. -> ALH retrievals of smoke events over Canada and the United States in August 2017 are demonstrated in Section 4.

Corrected.

Line 3, page 3: ALH and AOD from EPIC with -> ALH and AOD from EPIC against

Corrected.

Line 5, page 3: remove "In conclusion,"

Corrected.

Line 11, page 3: Figure 1b-c -> Figures 1b-c

Corrected.

Line 13, page 3: the scattering of presented aerosol particles interact with -> scattered light from aerosol particles interacts with

Corrected.

Line 16, page 3: estimate -> estimated

Corrected.

Line 18, page 3: ' among those are -> ; among those are

Corrected.

Line 20, page 3: with O2 absorption -> using the O2

Corrected.

Line 22, page 3: thus reduce the chance of a photon -> thus reducing the chance of that photon

Corrected.

Line 23, page 3: TOA -> Top Of the Atmosphere (TOA)

Corrected.

Line 28, page 3: leave a space between citations

Done.

Lines 29-30, page 3: a state-of-the-art -> the state-of-the-art

Corrected.

Line 31, page 3: formulated by -> for

Corrected.

Line 33, page 3: at the geometry of -> for the geometry

Corrected.

Line 5, page 4: contribution from surface -> contribution from the surface

Corrected.

Line 7, page 4: findings -> the findings; the O2 A and B band -> O2 A and B band

Corrected.

Line 8, page 4: leave a space between citations

Done.

Line 11, page 4: the retrieval accuracy -> a retrieval accuracy

Corrected.

Line 13, page 4: earth -> Earth

Corrected.

Line 18, page 4: earth -> Earth

Corrected.

Line 20, page 4: remove the comma

Done.

Line 31, page 4: at six EPIC -> in six EPIC

Corrected.

Line 4, page 5: at EPIC bands -> in EPIC bands

Corrected.

Line 5, page 5: EPIC original pixels -> original EPIC pixels

Corrected.

Line 9, page 5: of available pixels -> for the available pixels

Corrected.

Line 11, page 5: specific surface type -> the specific surface type

Corrected.

Lines 12-13, page 5: While the retrieval procedure is based up on our algorithm . . . from the EPIC (Xu et al., 2017), it was upgraded in a few aspects -> While the retrieval procedure is based on our algorithm . . . from EPIC measurements (Xu et al., 2017), it was upgraded in several ways.

Done.

Line 13, page 5: algorithm extends -> algorithm is extended

Corrected.

Line 14, page 5: O2 -> O₂

Corrected.

Line 24, page 5: Obtain -> Obtaining

Corrected.

Lines 29-30, page 5: The two O₂ absorption channels (688 nm and 764 nm) were calibrated by lunar surface reflectivity with EPIC lunar observations at the time of full moon as seen from the earth -> The two O₂ absorption channels (688 nm and 764 nm) were calibrated using lunar surface reflectivity from EPIC lunar observations at the time of full moon as seen from Earth

Corrected.

Line 32, page 5: by calibration factors derived by above studies -> using calibration factors from previous studies

Corrected.

Line 34, page 5: top-of-the-atmosphere (TOA) -> TOA

Corrected.

Line 2, page 6: where $C(\lambda)$ is EPIC measured signal in the units of -> where $C(\lambda)$ is the EPIC measured signal in units of

Corrected.

Line 6, page 6: Determine -> Determining

Corrected.

Line 8, page 6: leave a space between the citations

Done.

Line 12, page 6: leave a space between the citations

Done.

Line 16, page 6: in MODIS's first seven channels -> in the first seven MODIS channels;
leave a space between the citations

Corrected.

Line 24, page 6: leave a space between the citations

Done.

Lines 25-26, page 6: Lambertian surface albedo at MODIS bands of 469, 555, 645, and
858 nm -> Lambertian surface albedo in the 469, 555, 645, and 858 nm MODIS bands

Corrected.

Line 28, page 6: EPIC-bands -> EPIC bands; in the forms -> in the form

Corrected.

Line 31, page 6: spectral locations -> the spectral locations

Corrected.

Line 32, page 6: at each EPIC band -> in each EPIC band; Figure 6c-h -> Figures 6c-h

Corrected.

Line 12, page 7: Mask -> Masking

Corrected.

Line 15, page 7: the land and water -> land and water

Corrected.

Lines 21-22, page 7: higher-resolution geostationary sensors' cloud mask information -> higher resolution cloud mask information from geostationary sensors

Corrected.

Lines 22-23, page 7: if applied operationally -> for operational applications

We removed this sentence "Besides, cloud mask thresholds used in this work might need to be adjusted if applied operationally."

Line 26, page 7: with MODIS land surface -> using MODIS land surface

Corrected.

Line 30, page 7: constructed with -> constructed using the

Corrected.

Line 3, page 8: of the current retrieval -> for the current retrieval

Corrected.

Line 4, page 8: circumstances -> scenarios

Corrected.

Line 5, page 8: simulated by -> simulated using

Corrected.

Line 6, page 8: at the selected 6 bands -> for the selected six bands

Corrected.

Line 9, page 8: leave a space between the citations

Done.

Line 10, page 8: by following -> following

Corrected.

Line 15, page 8: leave a space between the citations

Done.

Lines 17-18, page 8: total AOD at the wavelength of 680 nm -> the total AOD at 680 nm

Corrected.

Line 19, page 8: fittings -> fitting

Corrected.

Line 20, page 8: both the water -> both water

Corrected.

Line 21, page 8: fittings -> fitting; account for specifics of surface reflectivity -> accounts for the specifics of surface reflectivity

Corrected.

Lines 21-22, page 8: First, TOA reflectance in EPIC's "atmospheric window" channels are matched with LUTs to determine AOD, because at these channels the TOA reflectance is independent of ALH. -> First, the TOA reflectance in the EPIC "atmospheric window" channels are matched with LUTs to determine AOD, since the TOA reflectance does not depend on ALH in these channels.

Corrected.

Line 26, page 8: because over land the satellite signal tends to be dominated by surface contributions over land -> since the satellite signal tends to be dominated by surface contributions over land

Corrected.

Line 27, page 8: separated -> separate

Corrected.

Line 28, page 8: in characterizing -> for characterizing

Corrected.

Line 30, page 8: the surface type -> surface type

Corrected.

Line 33, page 8: In contrast, the band of 780 nm is excluded for the spectral fitting -> In contrast, the 780 nm band is excluded for spectral fitting

Corrected.

Lines 1-2, page 9: weights to ratios in the O2 A and B bands are given differently for different surfaces -> different weights are given for the ratios in the O2 A and B bands for different surfaces

Corrected.

Line 5, page 9: Demonstration -> demonstration

Corrected.

Line 9, page 9: shown in EPIC RGB images -> shown in the EPIC RGB images

Corrected.

Line 10, page 9: plumes emitted from wildfires in western Canada and, crossing -> plumes emitted from wildfires in western Canada and crossing

We changed to “plumes emitted from wildfires in western Canada and transported”

Lines 11-12, page 9: The retrieved smoke ALH are shown in Figure 7b and 8b; and retrieved 680-nm AOD in Figure 7c and 8c. -> The retrieved smoke ALH is shown in Figure 7b and 8b, and retrieved 680-nm AOD in Figure 7c and 8c.

Corrected.

Line 13, page 9: and ALH retrievals -> ALH retrievals

Corrected.

Line 18, page 9: towards southeast -> southeast

Corrected.

Line 23, page 9: validations -> validation

Corrected.

Line 25, page 9: observation -> observations

Corrected.

Line 29, page 9: in 532 nm -> at 532 nm

Corrected.

Line 31, page 9: defined in our EPIC algorithm -> as defined in our EPIC algorithm

Corrected.

Line 1, page 10: with the layers where aerosols are detected -> for the layers where aerosols are detected

Corrected.

Line 4, page 10: backscattering ratio that depends -> backscattering ratio, which depends

Corrected.

Line 6, page 10: daytime CALIOP scan -> a daytime CALIOP scan

Corrected.

Line 7, page 10: reaches up to -> increases to

Corrected.

Line 9, page 10: predominately -> predominantly

Corrected.

Lines 10-11, page 10: To compensate for this bias, we use a exponentially-decayed background aerosol extinction profile for substitute of aerosol extinction coefficients of these undetected aerosol layers within troposphere. -> To compensate for this bias, we use an exponentially-decaying background aerosol extinction profile to provide a proxy for aerosol extinction coefficients of these undetected aerosol layers within the troposphere.

Corrected.

Line 13, page 10: summertime atmosphere of the Arctic -> summertime Arctic atmosphere

Corrected.

Line 15, page 10: bias of ALHCALIOP -> bias in ALHCALIOP

Corrected.

Lines 16-17, page 10: Quantitatively, 67

Lines 18-20, page 10: Considering all EPIC- CALIOP ALH pairs, 65

Corrected.

Line 21, page 10: observations of 675 nm AOD -> 675 nm AOD observations

Corrected.

Line 22, page 10: (Ichoku et al., 2002) -> Ichoku et al. (2002)

Corrected.

Lines 22-24, page 10: “but was modified to associate a subset of satellite retrievals within a 3 X 3 AOD subset centered at each site to a subset of 1-hour AERONET observations around EPIC scan time.” It is not clear what the authors mean by 3 X 3 AOD subset. The sentence needs to be revised for clarity.

We revised the sentence into “but was modified to associate compare a subset of satellite retrievals within EPIC AOD retrievals over a 3x3 AOD subset pixels centered at each the AERONET sites to a subset of with 1-hour AERONET AOD observations around the EPIC scan time.”.

Line 24, page 10: EPIC scan time -> the EPIC scan time

Corrected.

Lines 24-25, page 10: Comparison of EPIC AOD and AERONET are shown in Figure 10b. -> A comparison of EPIC and AERONET AODs is shown in Figure 10b.

Corrected.

Lines 25-26, page 10: The collocated AOD pairs, though with limited data samplings, have over 77

Corrected.

Line 26, page 10: EPIC AOD -> The EPIC AOD

Corrected.

Line 34, page 10: UV aerosol index -> the UV aerosol index

Corrected.

Line 1, page 11: because -> since

Corrected.

Line 9, page 11: both perform -> both of which obtain

Corrected.

Line 10, page 11: leave a space between the citations; Is it Omar et al or Torres et al?

Done. Double checked, and it is Omar et al.

Line 14, page 11: which are in contrast to clouds which-> which are in contrast to clouds that

Corrected.

Line 15, page 11: Because -> Since

Corrected.

Line 16, page 11: correlation -> the correlation

Corrected.

Lines 17-18, page 11: may results in a value UVAI from less than 1 to about 4 -> may result in UVAI values ranging from less than 1 to about 4

Corrected.

Line 19, page 11: EPIC's O2 bands -> the EPIC O2 bands

Corrected.

Lines 21-22, page 11: Based on our previous efforts in retrieving over-water dust ALH from the EPIC (Xu et al., 2017), we extend the retrieval algorithm to biomass burning

smoke aerosols over both the water and vegetated land surfaces. -> We extend our retrieval algorithm for retrieving over-water dust ALH from EPIC (Xu et al., 2017) to biomass burning smoke aerosols over both water and vegetated land surfaces.

Corrected.

Line 23, page 11: flexible spectral fittings that account for specifics of-> flexible spectral fitting that accounts for the specifics of

Corrected.

Line 25, page 11: then uses -> and then uses

Corrected.

Line 28, page 11: And, surface reflectance -> Surface reflectance

Corrected.

Lines 31-32, page 11: We found the algorithm captures AOD and ALH multiple times daily over both the water and vegetated land surfaces. -> The algorithm is able to retrieve AOD and ALH multiple times daily over both water and vegetated land surfaces.

Corrected.

Lines 33-34, page 11: , showing EPIC retrieved ALH has a rmse of 0.58 km and captures 52

Corrected.

Line 1, page 12: mrse -> rmse

Corrected.

Lines 1-2, page 12: and over 77

Corrected.

Line 2, page 12: What does an error envelope of +/- (0.05 + 10

We changed the “an uncertainty envelope of $\pm (0.05 + 0.1AOD)$ ”.

Line 4, page 12: the EPIC’s UV bands -> the EPIC UV bands

Corrected.

Line 9, page 12: dust or smoke -> (dust or smoke)

Corrected.

Line 15, page 12: NASA the DSCOVR Earth Science Algorithms Program -> the NASA DSCOVR Earth Science Algorithms Program

Corrected.

Line 16, page 12: Office of Naval Research (ONR’s) -> the Office of Naval Research (ONR)

Corrected.

Line 17, page 12: under the award -> under award

Corrected.

Line 20, page 12: NASA’s -> the NASA; AERONET program -> the AERONET program

Corrected.

Line 21: the AOD data -> AOD data

Corrected.

Table 2 caption: in constructing the LUTs -> for constructing the LUTs

Corrected.

Figure 1 caption: Change to: EPIC instrument filter response function (blue) and atmospheric spectral transmission 5 (orange). Panel (a) includes all ten EPIC bands, whereas panels (b) and (c) show zoom-ins for the 688-nm channel in the O2 B-band and the 764-nm channel in the O2 A-band, respectively. Here, the atmospheric transmission is simulated by the UNL-VRM model, with a spectral step size and a spectral full width at half maximum of 0.02 nm.

Done.

Line 5, page 21: physical principal for -> physical principle of

Corrected.

Line 6, page 21: scattering of aerosol -> scattering by aerosol

Corrected.

Line 7, page 21: path way -> pathlength

Corrected.

Line 8, page 21: than in the lower-altitude aerosol -> than those scattered by the lower-altitude aerosol; less chance -> lower chance

Corrected.

Line 5, page 22: at the geometry of -> for the geometry

Corrected.

Figure 4 legend: Change "Green vegetations" to "Green vegetation surfaces"

Corrected.

Figure 5: What does 3X3 aggregation mean? Do you aggregate 9 pixels at a time? Why? Some explanation is needed in the text and better wording in the Figure.

In the figure, we changed “3X3 aggregation” to “3x3 pixel aggregation”, and changed “3x3 averaging” to “averaging over the 3x3 pixels”

Line 5, page 25: the statistics -> statistics

Corrected.

Line 6, page 25: red dot line -> red dotted lines; their respective -> the respective

Corrected.

Lines 7-8, page 25: reflectance at each EPIC band versus reflectance at corresponding MODIS bands -> reflectance in each EPIC band versus reflectance in the corresponding MODIS bands

Corrected.

Line 4, page 26: UTC time -> UTC times

Corrected.

Line 6, page 26: CALIOP sub-orbital track with an overpass time 19:05 UTC -> the CALIOP sub-orbital track with an overpass time of 19:05 UTC

Corrected.

Line 8, page 26: EPIC scan time -> the EPIC scan time

Corrected.

Line 5, page 27: UTC time -> UTC times

Corrected.

Line 6, page 27: CALIOP overpass time was at 18:15. -> The CALIOP overpass was at 18:15 UTC.

Corrected.

Line 4, page 28: Comparison of ALH retrieved from EPIC and the ALH derived from CALIOP level-2 aerosol extinction profile -> Comparison of ALH retrieved from EPIC and CALIOP level-2 aerosol extinction profile

Corrected.

Line 5-6, page 28: CALIOP orbital tracks are marked on EPIC RGB images in Figure 7–8. -> The CALIOP orbital tracks are marked on EPIC RGB images in Figures 7–8.

Corrected.

Line 7-8, page 28: Error bar of EPIC ALH represents standard deviation for an array of 3x3 24-km retrieval pixels, while the error bar of CALIOP ALH represents standard deviation of over 5 adjacent CALIOP 5-km footprints. -> The error bar for EPIC ALH represents the standard deviation for an array of 3x3 24-km retrieval pixels, while that for CALIOP ALH represents the standard deviation of over 5 adjacent CALIOP 5-km footprints.

Done.

Line 4, page 29: counterparts from -> corresponding Line 5, page 29: Color of -> The color of

Corrected.

Line 6, page 29: scatter -> scatter point; EPIC 680-nm AOD value of -> the EPIC 680-nm value for

Corrected.

Line 7, page 29: Dotted lines -> The dotted lines Line 8, page 29: one-by-one -> the one-to-one

Corrected.

Line 9, page 29: regression fitting -> regression fit; scatters -> scatter points; the linear -
> linear

Corrected.

Line 10, page 29: scatters -> scatter points

Corrected.

Line 4, page 30: UVAI were -> UVAI was

Corrected.

Line 4, page 31: linear regression fitting -> the linear regression fit

Corrected.

Line 5, page 31: scatters -> scatter points

Corrected.

Detecting layer height of smoke aerosols over vegetated land and water surfaces via oxygen absorption bands: Hourly results from the EPIC/DSCOVR in deep space

Xiaoguang Xu^{1,2}, Jun Wang¹, Yi Wang¹, Jing Zeng¹, Omar Torres³, Jeffrey S. Reid⁴, Steven D. Miller⁵, J. Vanderlei Martins², Lorraine A. Remer²

¹Department of Chemical and Biochemical Engineering, Center for Global and Regional Environmental Research, and Informatics Initiative, The University of Iowa, Iowa City, Iowa, 52241, USA

²Joint Center for Earth Systems Technology and Department of Physics, University of Maryland Baltimore County, Baltimore, Maryland, 21250, USA

³Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, 20770, USA

⁴Marine Meteorology Division, Naval Research Laboratory, Monterey, California, 93943, USA

⁵Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, Colorado, 80523, USA

15 Correspondence to: Jun Wang (jun-wang-1@uiowa.edu); Xiaoguang Xu (Xiaoguang-xu@uiowa.edu)

Abstract. We present an algorithm for retrieving aerosol layer height (ALH) and aerosol optical depth (AOD) for smoke over vegetated land and water surfaces from measurements of the Earth Polychromatic Imaging Camera (EPIC) onboard the Deep Space Climate Observatory (DSCOVR). The algorithm uses Earth-reflected radiances in six EPIC bands in the visible and near-infrared and incorporates flexible spectral fitting that accounts for specifics of land and water surface reflectivity. The fitting procedure first determines AOD using EPIC atmospheric window bands (443 nm, 551 nm, 680nm, and 780 nm), then uses oxygen (O₂) A and B bands (688 nm and 764 nm) to derive ALH, which represents an optical centroid altitude. ALH retrieval over vegetated surface primarily takes advantage of measurements in the O₂ B band. We applied the algorithm to EPIC observations of several biomass burning events over the United States and Canada in August 2017. We found that the algorithm can be used to obtain AOD and ALH multiple times daily over water and vegetated land surface. Validation is performed against aerosol extinction profiles detected by the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) and against AOD observed at nine Aerosol Robotic Network (AERONET) sites, showing, on average, an error of 0.58 km and a bias of -0.13 km in retrieved ALH and an error of 0.05 and a bias of 0.03 in retrieved AOD. Additionally, we show that the aerosol height information retrieved by the present algorithm can potentially benefit the retrieval of aerosol properties from EPIC's ultraviolet (UV) bands.

30 1 Introduction

Aerosol vertical distribution is an important but poorly constrained variable that strongly influences how aerosol particles affect earth's energy budget. In particular, absorption of solar radiation by smoke and dust particles can result in diabatic heating, alter atmospheric stability, and affect cloud formation and life cycle. These effects depend critically on the altitude of aerosol layers (Babu et al., 2011; Ge et al., 2014; Koch and Del Genio, 2010; Satheesh et al., 2008; Wendisch et al., 2008). Consequently, aerosol profile factors into the magnitude and even the sign of aerosol direct and indirect effects. An accurate representation of aerosol altitude is thus essential for model prediction of weather and climate (Choi and Chung, 2014; Samset et al., 2013). The thermal signature of dust, in particular, can likewise influence the Earth's longwave budget, and through the interference of retrievals of water vapor and temperature, influence measurement of atmospheric state

Deleted: satellite

Deleted: earth

Deleted: s

Deleted: the

Deleted: that

Deleted: the

Deleted: well capture

Deleted: s

Deleted: are

Deleted: in

Deleted: the

Deleted: ;Ge et al., 2014

Deleted: earth

Deleted: thus influencing

Deleted: measure

(Maddy et al., 2012). Additionally, the knowledge of ALH is essential ~~for~~ retrieving aerosol absorption properties in the ultraviolet (UV) channels (Torres et al., 1998), ~~aerosol microphysical properties from multi-angular photopolarimetric measurements (Chowdhary et al., 2005; Waquet et al., 2009), and~~ ~~for~~ atmospheric correction for ocean color remote sensing (Duforêt et al., 2007).

Deleted: in

Deleted: in retrieving

Deleted: ;Chowdhary et al., 2005

Deleted: in the

5 Despite the importance of aerosol vertical distribution, the simulation of aerosol layer height (ALH) in current climate models is subject to large inter-model variation and uncertainty (Kipling et al., 2016; Koffi et al., 2012). Although the assimilation of space-borne Lidar (e.g., CALIOP, or the Cloud-Aerosol Lidar with Orthogonal Polarization) observations can improve the prediction of vertical allocation (Zhang et al., 2011), a significant challenge remains due to the sparsity of lidar observations over space and time. Therefore, frequent satellite observations of global aerosol vertical distribution based on
10 more abundant observation sources are critically needed.

Deleted: ;Kipling et al., 2016

One such source, the recently-launched Deep Space Climate Observatory (DSCOVR) mission, has introduced an unprecedented opportunity to acquire ALH information multiple times daily. Launched on 11 February 2015, the DSCOVR spacecraft flies a Lissajous orbit around the Sun-Earth L₁ Lagrangian point about 1.5 million km from Earth in the direction of the Sun. While its primary payload is the Plasma-Magnetometer that measures solar wind, DSCOVR also carries the Earth
15 Polychromatic Imaging Camera (EPIC) to image the sunlit disk of Earth every 60 – 100 minutes from its stable L₁ vantage point. EPIC captures its imagery at a 2048-by-2018-pixel resolution, resolving about 8-km pixel resolution at the disk's center. EPIC was equipped with a double wheel filter to sequentially measure Earth-reflected solar radiance in ten narrow bands spanning from ~~the~~ UV to ~~the~~ near-infrared (NIR) spectrum (Figure 1a), with central wavelengths at 318 nm, 325 nm, 340 nm, 388 nm, 443 nm, 551 nm, 680 nm, 688 nm, 764 nm, and 780 nm (Marshak et al., 2018).

Deleted: Figure

20 As shown in ~~Figures~~ 1b-c, two of EPIC's bands are located within the oxygen (O₂) "A" and "B" bands (764 nm and 688 nm), each associated with a reference continuum band at 780 nm and 680 nm, respectively. These four bands, offering
~~spectral contrasts between absorption bands and continuum bands (known as the Differential Optical Absorption Spectroscopic, or DOAS, ratios), were originally designed for determining cloud height (Yang et al., 2013). Recently, Xu et al. (2017) presented~~ an algorithm to simultaneously retrieve aerosol optical depth (AOD) and ALH using the EPIC
25 measurements via these four bands and, for the first time, ~~demonstrated~~ EPIC's promising application for determining dust plume height over ocean surfaces during the daytime hours.

Deleted: the

Deleted: present

Deleted:

Deleted: demonstrate

The present study, building upon the development of Xu et al. (2017) ~~for~~ determining dust ALH over ocean, extends the algorithm to retrieve ALH from EPIC measurements over land surfaces as well. The new developments in this study includes implementing smoke aerosol optical properties, land surface characterization, and more robust strategies ~~for~~ the
30 procedures of pixel selection and spectral fitting. The augmentation of the Xu et al. (2017) algorithm takes an important additional step towards our goal of providing more frequent global ALH and AOD information for multi-species global aerosol.

Deleted: in

Deleted: in

The paper is outlined as follows. Section 2 briefly reviews the physical principle for remote sensing of ALH in the O₂ absorption bands and explains the challenges for retrieving ALH over land surfaces, limiting our algorithm development to water and vegetated land surface. Section 3 describes the key elements, procedures, and assumptions of the updated ALH retrieval algorithm. ALH retrievals smoke events over Canada and the United States in August 2017 are demonstrated in Section 4. Section 5 evaluates the retrieved smoke ALH and AOD from EPIC against aerosol extinction profiles measured by CALIOP and AOD measured at AERONET sites. Section 6 summarizes our findings and discusses potential broader applications of this new ALH information.

2 Remote sensing principle and challenges

The ALH retrieval from EPIC takes advantage of the absorption features of molecular O₂ in the A band at 759 – 771 nm and the B band at 686 – 695 nm. The electronic transitions from the ground state to two excited states of an O₂ molecule, coupled with vibrational-rotational transitions, produce O₂ absorption lines in the A and B bands with distinct spectral variability (Figures 1b-c). As a result, the large variability of atmospheric opacity within the O₂ A and B bands leads to a wide range of penetration depths of solar radiation (Ding et al., 2016). The spectroscopic characteristics in the O₂ bands are related to how scattered light from aerosol particles interacts with O₂ absorption at different altitudes. Furthermore, since O₂ molecules are well mixed in the atmospheric column, the altitude-dependence of O₂ absorption can provide information on the path length of light scattered by aerosol particles and thus is related to the height of aerosol layers. The premise of this retrieval dated back to Hanel (1961) and Yamamoto and Wark (1961), who estimated cloud top pressure based on the amount of absorption by molecular CO₂ and O₂ above the cloud layer. More recently, a number of satellite sensors have been designed to capture the O₂ absorption feature to provide cloud top height and ALH, among those are GOME, SCIAMACHY, MERIS, POLDER, DSCOVR, and most recently TROPOMI (Xu et al., 2018a and references therein).

Figure 2 illustrates the physical principle for sensing of ALH using the O₂ absorption spectroscopic approach, which relies on the fact that a scattering aerosol layer can scatter sun light back to space, shortening the path length of a photon traveling in the atmosphere, and reducing the chance of that photon being absorbed by O₂ molecules. As a result, an elevated scattering layer enhances the Top Of the Atmosphere (TOA) reflectance within the O₂ absorption bands as detected by a satellite. In other words, for a given aerosol layer of fixed AOD placed at different altitudes, the higher the altitude, the larger the TOA reflectance, i.e., $I_H > I_L$ in Figure 2. This relationship contrasts to that of the reference continuum band, where the TOA reflectance is not sensitive to vertical location of the aerosol layer but depends only on the column AOD, i.e., $I_H = I_L$ for a given AOD. Based on this principle, the ratio of TOA reflectance between in-band and continuum band, or the DOAS ratio, provides a practical way to infer the ALH (e.g., Dubuisson et al., 2009b; Duforêt et al., 2007; Xu et al., 2017).

To build the links between DOAS ratios and ALH, we simulate TOA reflectance as observed by EPIC measurements with the state-of-the-art Unified Linearized Vector Radiative Transfer Model (UNL-VRM; described in section 3.5). Figure 3

Deleted: to

Deleted: implicating

Deleted: limited

Deleted: the

Deleted: The

Deleted: , that were applied to smoke events over Canada and the United States in August 2017. Section 5 evaluates the retrieved

Deleted: with

Deleted: In conclusion, s

Deleted: S

Deleted: P

Deleted: C

Deleted: Figure

Deleted: the scattering of presented

Deleted: dates

Deleted: estimate

Deleted: ,

Deleted: with

Deleted: thus

Deleted: e

Deleted: a

Deleted: ;Duforêt et al., 2007

Deleted: a

shows simulations of the relationship between smoke ALH (y axis) and DOAS ratios (x axis) for EPIC channels in O₂ A and B bands and for various AOD values and surface reflectance (A_s , indicated by different colors). Those simulations were performed for a typical biomass-burning aerosol model as observed by EPIC for the geometry of $[\theta_0, \theta, \Delta\phi] = [42^\circ, 37^\circ, 165^\circ]$, where θ_0 and θ are solar and viewing zenith angles, and $\Delta\phi$ the relative azimuth angle between the sun and the viewer. As seen from Figure 3, the DOAS ratios in general increase with the rise of ALH. Meanwhile, the relationship is strongly dependent on the aerosol loading and surface reflectivity.

Deleted: formulated by

Deleted: at

The sensitivity of DOAS to ALH is enhanced for lower surface reflectance and larger AOD. Conversely, the sensitivity decreases rapidly for smaller AOD (e.g., AOD = 0.1 in Figure 3a, d) or as A_s increases, a condition where it is difficult to discriminate between the aerosol scattering contribution to TOA reflectance and contribution from the surface. Therefore, it is challenging to obtain ALH information for low aerosol loading conditions and over bright surfaces. This is consistent with findings of many previous information content studies for determining ALH from O₂ A and B band measurements (Ding et al., 2016; Dubuisson et al., 2009b). Duforêt et al. (2007) showed that the centroid altitude of a single aerosol layer over a dark surface can be retrieved from Polarization and Directionality of the Earth's Reflectances (POLDER) measurements with an error of less than 1 km when AOD is over 0.2. Similarly, Dubuisson et al. (2009b) suggested AOD > 0.3 and $A_s < 0.06$ in order to achieve a retrieval accuracy of 0.5 km from POLDER.

Deleted: the

Deleted: the

Deleted: ;Ding et al., 2016

Deleted: the

With the above retrieval principles and challenges in mind, for our algorithm development we need to determine over what underlying Earth surfaces EPIC measurements have sufficient information to allow a practical retrieval of ALH. First and foremost, the surface reflectance must be low enough in one of the O₂ A and B bands to provide unambiguous signal from the aerosol. According to Figure 3, the DOAS-ALH sensitivity becomes substantial when $A_s < 0.1$ in moderate aerosol loading conditions ($\tau_{680} = 0.4$). Hence, $A_s = 0.1$ is set as the upper threshold, above which an ALH retrieval will not be attempted, despite the sensitivity increases in high aerosol loading ($\tau_{680} = 1.0$) for higher A_s .

Deleted: earth

Figure 4 shows the magnitude of surface reflectance from the ASTER spectral library for typical Earth surface types (Baldrige et al., 2009). The water surface is dark, with reflectance lower than 0.03 in both the O₂ A and B bands. Over land, green vegetation is the only surface type that offers a reflectance below 0.1 in the O₂ B band ($A_s = 0.1$ is indicated by the red dotted line). In fact, A_s in the O₂ A-band is considerably above 0.1 for all land types considered in Figure 4. These findings provide important guidance on the ALH algorithm design. Specifically, EPIC can retrieve ALH only over water and vegetated surfaces. In particular, ALH retrieval over vegetated surface would enlist DOAS ratios in the O₂ B-band. Additionally, retrieval of ALH requires a sufficiently high aerosol loading. We set AOD thresholds for retrieving ALH over both the water and vegetation per the algorithm overview to follow.

Deleted: earth

Deleted: ,

Deleted: in the O₂ B band.

Deleted: A

Deleted: L

Deleted: H

Deleted: R

Deleted: A

3. EPIC aerosol layer height retrieval algorithm

3.1 Algorithm overview

Figure 5 illustrates the processing of EPIC data and the ALH retrieval procedure. Briefly, the retrieval algorithm entails the following steps:

1. Calculate TOA reflectance ~~in~~ six EPIC visible and NIR bands (443, 551, 680, 688, 764, and 780 nm) from the calibrated EPIC level 1B digital data.
2. Identify EPIC pixels that are suitable for aerosol height retrieval. Through various tests, this step screens out pixels having clouds, over-water sun glints, and bright land surfaces, which are performed separately for water and land pixels. Surface pressure comes from MERRA-2 reanalysis data and we determine surface reflectance ~~in~~ EPIC bands using GOME-2 and MODIS surface products.
3. Aggregate the ~~original~~ EPIC pixels into a box of 3×3 individual pixels, an area with size of about 24 km at nadir. In many cases, not all pixels within a box are suitable for retrieval (i.e., cloud, glint, and bright land). If the number of available pixels within a box is not less than 4 (of the total of 9), calculate mean values of TOA reflectance, satellite geometries, and surface reflectance ~~for the~~ available pixels. Otherwise, do not conduct an aerosol retrieval for the box.
4. Invert the aggregated EPIC observations using pre-calculated lookup tables to obtain smoke ALH and AOD. The inversion uses a flexible spectral fitting strategy that considers ~~the~~ specific surface type.

While the retrieval procedure is based ~~on our~~ algorithm of retrieving dust ALH over ocean from EPIC measurements (Xu et al., 2017), it was upgraded in ~~several ways~~. First, the algorithm ~~is extended~~ to retrieve ALH over vegetated land surface. The capability of inferring ALH over vegetation is predicated on the O₂ B band, where the surface reflectance is sufficiently low. Accordingly, the new algorithm uses methods separated for land and water surfaces in determining surface reflectance and screening cloud. Second, the algorithm incorporates a smoke aerosol optical model in order to retrieve biomass-burning smoke ALH. Third, rather than aggregating EPIC data from satellite pixels to regular latitude-longitude grids, the new algorithm retrieves over aggregated boxes, each consisting of an array of 3×3 EPIC pixels. This change is based on the consideration that geolocation of EPIC data has made significant improvements in the new version of level 1B data (Geogdzhayev and Marshak, 2018). Lastly, to obtain AOD and ALH the new algorithm employs a flexible spectral fitting strategy by considering the underlying surface reflectivity. In the following procedural subsections, we describe these changes in full detail.

3.2 Obtaining EPIC TOA reflectance

For this algorithm development, we used the EPIC Level 1B (L1B) Version 02 imagery data, available from NASA's Atmospheric Science Data Center (ASDC) at <https://eosweb.larc.nasa.gov> (last accessed Nov 26, 2018). Various pre- and post-launch calibrations were applied to the L1B EPIC data. EPIC visible and near-IR (NIR) channels at 443, 551, 680, and 780 nm were cross-calibrated with independent LEO satellite instruments, including the MODIS onboard the Terra and Aqua satellites (Geogdzhayev and Marshak, 2018). The two O₂ absorption channels (688 nm and 764 nm) were calibrated ~~using~~ lunar surface reflectivity with EPIC lunar observations at the time of full moon as seen from ~~Earth~~ (Ohtake et al.,

Deleted: at

Deleted: at

Deleted: original

Deleted: of

Deleted: up

Deleted: the

Deleted: a few aspects

Deleted: extends

Formatted: Subscript

Deleted: by

Deleted: the e

2010). The image data in EPIC L1B products is in digital units of counts/second, that is converted into reflectance for each visible and NIR channel using calibration factors provided at ASDC online: https://eosweb.larc.nasa.gov/project/dscovr/DSCOVr_EPIC_Calibration_Factors_V02.pdf (last accessed Nov 26, 2018).

The TOA reflectance values are calculated as:

$$R_{\lambda} = \frac{K(\lambda)C(\lambda)}{\mu_0}, \quad (1)$$

where $C(\lambda)$ is the EPIC measured signal in units of counts/second at the wavelength of λ , $K(\lambda)$ is a calibration factor, and μ_0 is the cosine of solar zenith angle, θ_0 . Additionally, pixels that are far from nadir and strongly distorted by the Earth ellipsoid (e.g., view zenith angle $\theta > 70^\circ$ or solar zenith angle $\theta_0 > 70^\circ$) are excluded.

10 3.3 Determining surface albedo and pressure

Past studies show that an accurate ALH retrieval critically depends on the appropriate assumption of surface reflectance (Corradini and Cervino, 2006; Ding et al., 2016; Dubuisson et al., 2009a). As discussed in the preceding section, our retrievals are confined to water and dark vegetated surfaces. Land and water mask information for each EPIC pixel was determined using the GSHHG coastline database (Wessel and Smith, 1996). Following Xu et al. (2017), surface reflectance over water surface was obtained from the GOME-2 surface Lambert-equivalent reflectivity (LER) database (Koelemeijer et al., 2003; Tilstra et al., 2017). GOME-2 LER products contain spectral LER albedo at 21 one-nanometer-wide channels, available globally for each month of the year at $0.25^\circ \times 0.25^\circ$ grid spacing ($\sim 25 \times 25$ km cells). From these data, we selected LER albedos at 440, 555, 670, 758, and 780 nm to represent surface reflectance for the nearest EPIC bands.

The Moderate-resolution Imaging Spectroradiometer (MODIS) BRDF/Albedo product (MCD43) provides parameters that can be used to determine surface reflectance over land in the first seven MODIS channels (Lucht et al., 2000; Schaaf et al., 2002). Those parameters, including surface bidirectional reflectance distribution function (BRDF) and albedos, were inverted from the atmospherically-corrected (i.e., Rayleigh and aerosol components removed) MODIS reflectance observations from both the Terra and Aqua satellites over a 16-day period. We used black-sky albedo and white-sky albedo compiled in the Level-3 daily Climate Modeling Grid (CMG) Albedo Product (MCD43C3) at a spatial resolution of $0.1^\circ \times 0.1^\circ$. Here, black-sky albedo is the directional hemispherical reflectance if the illumination comes only from the sun at solar noon. White-sky albedo is the bi-hemispherical reflectance under isotropic sky-light illumination. The actual bi-hemispherical reflectance can be computed from white-sky and black-sky albedos via a weighting coefficient that depends primarily on solar angle and columnar optical depth (Kokhanovsky et al., 2005; Lewis and Barnsley, 1994; Schaaf et al., 2002). Following Kokhanovsky et al. (2005), we calculated the Lambertian surface albedo in the 469, 555, 645, and 858 nm MODIS bands by weighting the contributions of white-sky and black-sky albedos. It should be noted that the effects of non-Lambertian surface reflection may bias the ALH retrieval, although this type of impact is limited by EPIC's backscattering direction only observation pattern. Further studies are needed to examine the detailed impacts, which will be one of our future efforts.

Deleted: f

Deleted: by

Deleted: derived by above studies. The calibration factors for all EPIC channels are provided at ASDC online:

Deleted: top-of-the-atmosphere (TOA)

Deleted: the

Deleted: Determine

Deleted: x

Deleted: 's

Deleted: first seven

Deleted: ;Lucht et al., 2000

Deleted: J.

Deleted: at MODIS bands of

The differences in spectral position and width of corresponding EPIC and MODIS channels may result in discrepancies. To compensate, we adjust MODIS reflectance values to equivalent EPIC bands. The adjustment factors, in the form of linear regression coefficients, were determined from analyzing USGS (United States Geological Survey) hyperspectral data for green vegetation. Figure 6a shows the selected reflectance spectra of 47 vegetation samples from the USGS Spectral Library Version 7 (Kokaly et al., 2017) and the spectral locations of EPIC (blue) and MODIS (red-dotted) bands. Linear regression of reflectance values in each EPIC band and corresponding MODIS band is illustrated as scatterplots in Figures 6c-h. The regression slope and offset, as well as coefficient of determination (R^2), are listed in Figure 6b. As shown, the reflectance between EPIC bands and their closest MODIS bands are highly correlated (R^2 from 0.96 to 1.0), and therefore, best-fitting equations in Figure 6b can be used to empirically derive EPIC surface reflectance from MODIS surface reflectance products.

The retrieval also requires auxiliary meteorological information to realistically characterize O_2 absorption properties. Surface pressure strongly affects the reflectance in O_2 absorption bands as observed from space, because it is a direct proxy for the columnar loading of air (and thus oxygen) molecules. Here, we enlisted surface pressure information from the Modern-Era Retrospective analysis for Research and Applications Version 2 (MERRA-2) datasets (Gelaro et al., 2017). MERRA-2's 1-hourly surface pressure at 0.5° by 0.675° grids were interpolated to the location and scan time of each EPIC pixel. In addition, the atmospheric temperature-pressure profile also impacts the width and strength of O_2 absorption lines. However, such influence on the radiative transfer is negligible for EPIC's 1-to-2-nm-wide bands. In this study, our algorithm employs a standard temperature-pressure profile representing the mid-latitude-summer atmosphere (McClatchey et al., 1972).

3.4 Masking clouds, sun glints, and bright land surface

After acquiring EPIC TOA reflectance and surface reflectance, the algorithm conducts a masking exercise to select clear-sky pixels suitable for aerosol retrieval. This includes the screenings of clouds, sun glint over water surfaces, and highly reflective land surfaces. Over both land and water, the cloud mask combines a set of brightness and homogeneity tests following Martins et al. (2002). The brightness tests screen out cloud pixels where EPIC TOA reflectance in two or three visible bands (443 and 680 nm over land; 443, 680, and 780 nm over water) exceed prescribed thresholds. The homogeneity test, on the other hand, identifies cloud pixels by evaluating the standard deviation of TOA reflectance at 443 and 551 nm within 3×3 neighboring pixels. Still, thin and small subpixel clouds can be missed in the relatively coarse resolution 8 km EPIC pixels, leading to an overestimation of AOD and an underestimation of ALH. Therefore, contamination by small-scale clouds is one of the known issues of retrieval quality and coupling higher resolution cloud mask information from geostationary sensors may help to overcome this issue. Besides, cloud mask thresholds used in this work might need to be adjusted for operational applications.

Deleted: -

Deleted: s

Deleted: the

Deleted: at

Deleted: the

Deleted: -

Deleted: '

Formatted: cloud mask information

Formatted: Strikethrough

Deleted: if applied operationally

Formatted: Strikethrough

In addition, EPIC pixels affected by sun glint and highly-reflective land surface are also removed from consideration by the mask. We identify the over-water glint area as those pixels having a glint angle smaller than 30° (Levy et al., 2013). Highly reflective land-surface pixels are identified ~~using~~ MODIS land surface products. Any EPIC pixel having NDVI below 0.2 or having a 680 nm surface reflectance larger than 0.1 is flagged as being a bright surface and excluded from the retrievals.

Deleted: with

5 3.5 Look-up tables

Following Xu et al. (2017), the revised algorithm retrieves smoke ALH and AOD from EPIC measurements via a set of lookup tables (LUTs) constructed ~~using the~~ UNL-VRM model. The UNL-VRM ([https:// unl-vrtm.org](https://unl-vrtm.org)) is a radiative transfer testbed developed specifically for atmospheric remote sensing (Wang et al., 2014). By integrating the VLIDORT vector radiative transfer model (Spurr, 2006) and particulate scattering codes (Spurr et al., 2012), UNL-VRM can perform simulations for two or more sets of aerosol microphysical properties, typically with aerosols in one fine mode and one coarse mode. It also incorporates HITRAN spectroscopic gaseous absorption ~~with~~ up to 22 trace gases (Rothman et al., 2013), allowing for accurate hyperspectral simulations of remote sensing observations (Xu et al., 2018b).

Deleted: with

Deleted: the

Deleted: by

The LUTs ~~for~~ the current retrieval consist of simulated EPIC TOA reflectance at selected spectral bands for a set of AOD and ALH values under various atmospheric and observation ~~scenarios~~ (e.g., sun-earth-sensor geometry, surface reflectance, and surface pressure) as shown in Table 1. To build the LUTs, hyperspectral monochromatic radiances were simulated ~~using~~ UNL-VRM and convolved with EPIC instrumental filter response functions ~~for~~ the selected ~~six~~ bands. Table 2 provides the spectral range, interval, and resolution (in terms of full-width at half maximum; FWHM) of the UNL-VRM simulations.

Deleted: of

Deleted: circumstances

Deleted: by

Deleted: at

Deleted: 6

The UNL-VRM simulations assume a Gaussian-like aerosol profile characterized by a centroid altitude and a half-width parameter (Spurr and Christi, 2014; Xu et al., 2017). The centroid altitude is taken to represent ALH. A half-width of 1 km is assumed ~~following~~ Xu et al. (2017). This value was also used to derive AOD from UV observations by the Total Ozone Mapping Spectrometer (TOMS) and the Ozone Monitoring Instrument (OMI) (Torres et al., 1998). For this study, we implemented smoke optical properties calculated based on Lorenz-Mie scattering theory. Smoke refractive index was assumed to be $1.5 - 0.012i$, following Dubovik et al. (2002), and its spectral dependence was neglected. We assumed smoke particles followed a bi-lognormal size distribution, as adopted for the MODIS dark-target aerosol algorithm (Levy et al., 2013; Remer et al., 2005). Specifically, fine-mode volume median radius and standard deviation were prescribed as $0.14 + 0.01\tau_{680}$ μm and 0.44, respectively. The coarse-mode counterparts are $0.14 + 0.01\tau_{680}$ μm and 0.80, respectively. The volume ratio between fine and coarse modes is $(0.01 + 0.3\tau_{680}) / (0.01 + 0.09\tau_{680})$. Here, τ_{680} is ~~the~~ total AOD at 680 nm.

Deleted: by

Deleted: the wavelength of

3.6 Strategy of flexible spectral fitting

The retrieval procedure involves two steps over both ~~water~~ and land surfaces and, at the same time, incorporates flexible spectral fitting that accounts ~~for~~ the specifics of surface reflectivity. First, ~~the~~ TOA reflectance in ~~the~~ EPIC “atmospheric window” channels are matched with LUTs to determine AOD, ~~since the TOA reflectance does not depend on ALH in these~~

Deleted: s

Deleted: the

Deleted: s

Deleted: 's

Deleted: because at

channels. Second, based on this estimated AOD, the DOAS ratios around the O₂ bands are fitted to estimate ALH. In each step, least-squared fitting is applied, and fitting residuals are reported.

Deleted: the TOA reflectance is independent of ALH

In general, retrieving aerosol information from reflected solar radiation in vis-to-NIR wavelengths over land is more challenging than over ocean, since the satellite signal tends to be dominated by surface contributions over land. It is thus often the case that, for the same satellite instrument, separate over-land and over-ocean algorithms are developed with different strategies for characterizing surface reflectivity and band selection for fitting, e.g., MODIS aerosol algorithms (Remer et al., 2005). Similarly, our retrieval of smoke ALH from EPIC uses different band combinations over land versus over ocean, adjusted by surface type and spectral signature of smoke aerosol in the TOA reflectance.

Deleted: because over land

Deleted: d

Deleted: in

Deleted: the

In general, reflectance of water surfaces does not exceed 0.03 across the entire vis-to-NIR spectrum (Figure 4). Therefore, AOD can be determined by fitting TOA reflectance in all of the four “atmospheric window” channels at 443 nm, 551 nm, 680 nm, and 780 nm. In contrast, the 780 nm band is excluded for spectral fitting over vegetated land surface because of the high chlorophyll reflectance. Similarly, in the fitting of DOAS ratios, different weights are given for the ratios in the O₂ A and B bands for different surfaces, adjusted by the sensitivity of the DOAS ratio to ALH (e.g., Figure 3). The weighting coefficients are 0.4 for R688/R680 and 0.6 for R764/R780 over water surface, whereas these values are 0.9 and 0.1 over vegetated surface.

Deleted: band of

Deleted: the

Deleted: to

Deleted: are given differently

4. Retrieval demonstration

Deleted: Demonstration

We apply our algorithm to six EPIC scenes over the Hudson Bay – Great Lakes area obtained during 25 – 26 August 2017, with three consecutive scenes considered on each day. We chose these cases primarily because smoke plumes prevailed over both water and vegetated surface, and there were available aerosol vertical profile measurements from CALIOP that could be used to validate the ALH retrievals. As shown in the EPIC RGB images (Figures 7a and 8a), smoke aerosols appeared as diffuse plumes emitted from wildfires in western Canada and transported over the western and southern portions of Hudson Bay (Peterson et al., 2018). The retrieved smoke ALHs shown in Figures 7b and 8b, and retrieved 680-nm AOD in Figures 7c and 8c. It is noted that data gaps in the AOD maps represent screened-out bright pixels due to either cloud or bright land surface; ALH retrievals were only available when AOD was larger than 0.2.

Deleted: ,

Deleted: crossing

Deleted: are

Deleted: ;

Deleted: and

Obvious spatial variations are noted in retrieved smoke ALH and AOD. On August 25, smoke plumes had AOD values ranging from 0.1 to 0.45, with higher loading found at downwind regions in the south (Figure 7c). An ALH of 4 – 5 km was found over Hudson Bay, whereas the smoke altitudes decreased to 2 – 4 km over land off the bay’s western and southern shores (Figure 7b). Southward, the ALH increased rapidly to 4 – 6 km towards the Great Lakes. By August 26, the smoke plumes had traveled southeast (Figure 8c). The smoke altitudes remained at 3 – 5 km over the eastern part of Hudson Bay, and 2 – 4 km over the bay’s south side (Figure 8b). Altitudes of smoke plumes over the coast of northeastern U.S. were higher than 5 km. Aside from spatial variations, the retrievals also revealed diurnal changes of ALH and evolution of the

Deleted: towards

smoke plumes. For instance, the ALH of smoke plumes over the Hudson Bay and the north side of the Great Lakes rose by about half km within two hours from local morning to afternoon on August 25 (Figure 7b).

5. Retrieval validation

For the validation of the EPIC-retrieved ALH, we used CALIOP Level 2 version 4.10 aerosol extinction profiles at 5 km spatial resolution, retrieved from CALIOP observations of attenuated backscatter at 532 nm (Young and Vaughan, 2009). To facilitate quantitative comparison of aerosol altitude, we used a mean extinction height calculated from the CALIOP extinction profile, following Koffi et al. (2012):

$$ALH_{CALIOP} = \frac{\sum_{i=1}^n \beta_{ext,i} Z_i}{\sum_{i=1}^n \beta_{ext,i}} \quad (2)$$

Here, $\beta_{ext,i}$ is the aerosol extinction coefficient (km^{-1}) at 532 nm at level i and Z_i is the altitude (km) of level i . Thus, ALH_{CALIOP} represents an effective ALH weighted by aerosol extinction signal at each level and is consistent with ALH as defined in our EPIC algorithm.

In the CALIOP Level 2 products, aerosol extinction is only retrieved for the layers where aerosols are detected, as permitted by the instrument's signal-to-noise ratio (SNR). Atmospheric layers with no aerosols detected are classified as "clear air" and assigned an aerosol extinction coefficient of 0 km^{-1} . The detection limits are defined in terms of backscattering ratio, which depends on an aerosol lidar ratio (Vaughan et al., 2009). As such, aerosol layers that span a large altitude range frequently remain undetected (Toth et al., 2018), particularly for absorbing aerosols in the day time. As indicated by Winker et al. (2013), aerosol extinction threshold in a daytime CALIOP scan is $0.01 - 0.03 \text{ km}^{-1}$ for 80-km horizontal averaging resolution and increases to 0.07 km^{-1} for 5-km horizontal averaging resolution. In reality, aerosols are ubiquitous throughout the troposphere, though the concentration can be very low in the free troposphere. However, excluding the "clear-air" layers would lead to a biased estimate of ALH_{CALIOP} , particularly for cases of predominantly "clear-air" layers present below an elevated aerosol layer. To compensate for this bias, we use an exponentially-decayed background aerosol extinction profile to provide a proxy for aerosol extinction coefficients of these undetected aerosol layers within the troposphere. The background profile has a columnar AOD of about 0.07 at 532 nm, according to Tomasi and Petkov (2014), who use various lidar measurements to characterize the summertime Arctic atmosphere. Though aerosol extinction within the undetected aerosol layers by no means follows the background extinction profile identically, adding such a background aerosol extinction profile could, to the first order, correct the potential bias in ALH_{CALIOP} . Therefore, we consider ALH_{CALIOP} with and without the imposed background aerosol in the following validation analysis.

Figure 9 presents the evaluation of our ALH retrieval against CALIOP observations, showing that the ALH retrieval captures the overall spatial variability of ALH as seen by CALIOP. Quantitatively, 67% and 59% of the retrievals are within 0.5 km difference from the counterparts of ALH_{CALIOP} on August 25 and 26, respectively. Considering all EPIC-CALIOP ALH pairs, over 65% of ALH retrievals are within an uncertainty envelope of 0.5 km (Figure 10a-b) and EPIC ALH has a mean

Deleted: s

Deleted: in

Deleted: with

Deleted: that

Deleted: reaches up

Deleted: ate

Deleted: substitute of

Deleted: of the Arctic

Deleted: can

Deleted: of

Deleted:

Deleted: ;

bias of -0.13 km and a root-mean-squared-error (rmse) of 0.58 km, capturing 52% variation of the ALH_{CALIOP} (Figure 10a). If background aerosol is imposed in the CALIOP ALH calculation, our retrieved EPIC ALH is found to have a bias of 0.23 km and a rmse of 0.57 km (Figure 10b).

The EPIC 680 nm AOD retrievals during the two days were compared against 675 nm AOD observations at nine AERONET sites (Table 3). The collocation method follows Ichoku et al. (2002), but was modified to compare EPIC AOD retrievals over 3×3 pixels centered at the AERONET sites with 1-hour AERONET AOD observations around the EPIC scan time. Collocated AERONET AOD values at each site are shown as circles in Figures 7c and 8c. A comparison of EPIC and AERONET AODs is shown in Figure 10c. The collocated AOD pairs, though with limited data samplings, have over 77% falling in an uncertainty envelope of $\pm (0.05 + 0.1AOD)$ with a coefficient of determination (R^2) of 0.54. The EPIC AOD shows a positive bias of 0.03 and a rmse of 0.05, which was dominated for four subsets of EPIC AODs and was likely caused by cloud contamination.

6. Implication to the retrieval of UV absorbing aerosol properties

One important implication of this study is that the retrieved ALH can provide complementary height information for determining absorbing aerosol properties from EPIC's UV bands. EPIC also measures backscattered UV radiances at 340 nm and 388 nm, which was designed to detect and retrieve optical properties of UV-absorbing aerosols like mineral dust and smoke by using the UV aerosol index (UVAI) (Marshak et al., 2018). UVAI quantifies the difference in spectral dependence between measured and calculated near UV radiances assuming a purely molecular atmosphere (Torres et al., 1998). Physically, inferring aerosol properties from those UV bands requires the characterization of aerosol height, since UV radiance is sensitive to aerosol vertical distribution. For example, Jeong and Hsu (2008) retrieve aerosol single scattering albedo (SSA) from OMI radiance via synergic use of AOD from MODIS and ALH from CALIOP. Currently, EPIC's UV aerosol algorithm utilizes climatological aerosol height data set derived from CALIOP observations (Torres et al., 2013). However, these static climatological data may fail to capture the dynamic variation of aerosol height, and thus induce uncertainties in the UV-retrieved aerosol properties. With the aerosol height and loading available from the EPIC's O₂ A and B bands, these closures are now possible with measurements from a single instrument. Such synergy can be also applied to the TROPOspheric Monitoring Instrument (TROPOMI) and the Plankton, Aerosol, Cloud, ocean Ecosystem (PACE) satellite, both of which obtain hyperspectral measurements from the UV to the NIR covering the O₂ A and B bands (Omar et al., 2018; Veeffkind et al., 2012).

Here, we compare our retrieved ALHs with EPIC UVAI to illustrate the importance, as well the potential benefit, of the ALH retrievals to EPIC's UV aerosol products. The EPIC UVAI data are publicly available at <https://eosweb.larc.nasa.gov/project/dscovr/> (last accessed Nov 26, 2018). Figure 11 shows the UVAI of the same EPIC scenes on August 25. These smoke plumes are highlighted by large values of UVAI, which are in contrast to clouds that typically exhibit a UVAI close to zero (Torres et al., 1998). Since UVAI is a function of ALH, AOD, and SSA, its correlation with ALH varies with AOD. As shown in Figure 12, the sensitivity of UVAI to ALH, as well as the correlation

- Deleted: of 675 nm AOD
- Deleted: (
- Deleted: ,
- Deleted: associate
- Deleted: a subset of satellite retrievals within
- Deleted: a
- Deleted: AOD subset
- Deleted: each
- Deleted: to a subset of
- Deleted: C
- Deleted: AOD
- Deleted: are
- Deleted: b
- Deleted: error
- Deleted: 10%

- Deleted: because

- Deleted: perform
- Deleted: and cover
- Deleted: both
- Deleted: ;Omar et al., 2018

- Deleted: which
- Deleted: Because

between them, increases as AOD increases. In particular, high AOD values (e.g., over 0.4) may result in UVAI values from less than 1 to about 4, depending on the ALH (Figure 12, bottom-right panel). Therefore, the use of ALH derived from the EPIC O_2 bands is expected to improve EPIC UV aerosol retrievals.

Deleted: s

Deleted: a value

Deleted: 's

7. Summary and discussions

5 [We extend our retrieval algorithm for](#) retrieving over-water dust ALH from [EPIC measurements](#) (Xu et al., 2017), to biomass burning smoke aerosols over both water and vegetated land surfaces. The new algorithm uses earth-reflected radiances in EPIC's visible and NIR bands and incorporates flexible spectral fitting that accounts for the specifics of vegetation and water surface reflectivity. The fitting procedure first determines AOD using four EPIC atmospheric window bands (443 nm, 551 nm, 680 nm, and 780 nm), [and then uses the DOAS ratios](#) formulated in the O_2 A and B bands (688 nm and 764 nm) to derive the ALH that represents an optical centroid altitude. ALH retrieval over vegetated surface primarily takes advantage of the measurements in the O_2 B band, where surface reflectance is sufficiently low to yield aerosol height information. [Surface reflectance values are specified using](#) MODIS surface products over land and GOME-2 LER products over ocean.

Deleted: Based on our previous efforts in

Deleted: the

Deleted: , we extend the retrieval algorithm

Deleted: the

Deleted: s

15 We applied the algorithm to six EPIC images, with [three](#) images on each day, having biomass burning plumes over the United States and Canada acquired on 25 – 26 August 2017. [The algorithm is able to retrieve](#) AOD and ALH multiple times daily over both water and vegetated land surfaces. The retrieved ALHs were validated against CALIOP extinction weighted aerosol height (ALH_{CALIOP}), showing EPIC retrieved ALH has a rmse of 0.58 km and captures 52% variation of the ALH_{CALIOP} , and 65% of EPIC and CALIOP ALH pairs are within an uncertainty envelope of 0.5 km. EPIC retrieved AODs are validated against AERONET AOD observed at nine sites, indicating a [rmse](#) of 0.05 and over 77% of EPIC AOD retrievals fall within the error envelope of $\pm (0.05 + 0.1AOD)$. In addition, by comparing the retrieved ALH and EPIC UVAI, we show that the aerosol height information retrieved by the present algorithm can potentially benefit the retrieval of aerosol properties from the EPIC UV bands.

Deleted: And,

Deleted: s

Deleted: determined with

Deleted: 3

Deleted: We found the

Deleted: captures

Deleted: the

Deleted: rmse

Deleted: 10%

25 The over [three](#) years of data recorded thus far by EPIC (July 2015 – present) offers an opportunity to characterize aerosol height from diurnal to seasonal scales. Such datasets with well-characterized uncertainties are valuable for evaluating aerosol vertical distribution simulated by climate models. A follow-on study, based on the current work and Xu et al. (2017), will examine the long-term global retrieval of ALH for dust and smoke aerosols. Challenges for such a global retrieval include the heterogeneity of aerosol species, whereas our efforts have been mainly focused on single-species (dust or smoke) aerosol assumptions. In fact, mixtures of aerosol are more commonly found in nature. Furthermore, the currently available EPIC UVAI products can help identify elevated absorbing aerosols (mainly, dust and smoke), which can then be used to determine the retrieval targets on any EPIC image.

Deleted: 's

Deleted: 3

30 [Appendix A. Sensitivity and Error Analysis](#)

[In this appendix, we investigate the sensitivity of EPIC measurements in the \$O_2\$ A and B bands to aerosol vertical profile shape, aerosol optical properties, and surface reflectance assumed in the retrieval algorithm. Then, we estimate potential](#)

ALH retrieval errors due to the uncertainties in these parameters. For this case, ALH is derived from two DOAS ratios (ρ) in the O₂ A and B bands. Mathematically, the retrieval error (ϵ) of ALH can be estimated using the optimal estimation approach (Xu and Wang, 2015):

$$\epsilon^{-2} = \mathbf{K}^T \mathbf{S}_e^{-1} \mathbf{K}, \quad (\text{A.1})$$

5 where \mathbf{K} is the Jacobian matrix of ρ with respect to ALH, and \mathbf{S}_e is the observation error covariance matrix for ρ . In this study, we assume no error correlation between the two DOAS ratios; That is, \mathbf{S}_e is a diagonal matrix with its elements equal to the error variance for ρ . Observation error variance matrix \mathbf{S}_e can be characterized as a sum of two terms

$$\mathbf{S}_e = \mathbf{S}_\rho + \mathbf{S}_m, \quad (\text{A.2})$$

10 Here, \mathbf{S}_ρ is the error covariance matrix describing the uncertainty for EPIC measurements, \mathbf{S}_m is the covariance matrix for forward model errors caused by inaccurate model assumptions and uncertainties in model parameterizations, and \mathbf{S}_m can be calculated by

$$\mathbf{S}_m = \mathbf{K}_b^T \mathbf{S}_b \mathbf{K}_b, \quad (\text{A.3})$$

15 where \mathbf{S}_b is the error covariance matrix describing uncertainties of the vector of model parameters \mathbf{b} , and \mathbf{K}_b is Jacobian matrix of ρ with respect to \mathbf{b} . In the following analysis, we consider four parameters for the vector \mathbf{b} , i.e., the half-width parameter (σ_H) defining the quasi-Gaussian aerosol vertical profile, the 680-nm AOD (τ_{680}), aerosol single scattering albedo (SSA), and surface reflectance A_s .

20 Figure A1 presents the UNL-VRM simulated DOAS ratios ρ in EPIC O₂ A and B band (a), and their Jacobian gradients with respect to ALH (b) and with respect to the above-mentioned four model parameters (c-f) for the vector \mathbf{b} for various surface and aerosol loading conditions. Figure A2 shows the estimated ALH retrieval errors owing to EPIC observation error and uncertainties in the assumed model parameters for water (with $A_s = 0.05$ in both O₂ A and B bands) and vegetation (with $A_s = 0.30$ in A band and 0.05 in B band) surfaces. The findings from these Jacobian and error analysis results can be summarized as:

- The sensitivity of ρ with respect to ALH (Figure A1b) is weak for aerosols confined in the lower atmosphere (ALH below 1.5 km). The sensitivity increases rapidly with the increase of ALH, peaks for ALH about 4 km, and then decreases slightly for higher elevated aerosols. The sensitivity in the O₂ A band is stronger than that in the O₂ B band, and it is stronger for low surface reflectance and high aerosol loading. By considering 2% uncertainty for the EPIC DOAS ratios (Geogdzhayev and Marshak, 2018), the ALH retrieval error (ϵ_0) is less than 0.5 km for the water surface and is less than 0.75 km for the vegetated surface if ALH is over 1.5 km (black curves in Figure A2a-b). This error characterization is consistent with the retrieval error (0.57 – 0.58 km) as estimated through the validation with CALIOP data.
- The DOAS ratios ρ have a negative sensitivity to σ_H for elevated aerosol, and the sensitivity turns to positive for ALH below 1.5 km (Figure A1c). An error of 0.5 km σ_H may cause up to 0.3 km retrieval error for ALH (blue curves in Figure A2c-d).

- The DOAS ratios ρ have a positive sensitivity to τ_{680} , and the sensitivity increases as ALH increases (Figure A1d). An error of 0.05 in τ_{680} can lead to a retrieval error from 0 to 0.6 km for ALH, depending on the aerosol loading and surface type (orange curves in Figure B1c-d).
- DAOS ratios are sensitive to SSA, especially for large AOD values (Figure A1e). However, SSA only has marginal impact to the ALH retrieval error (green curves in Figure A2c-d).
- The DOAS ratio ρ has a negative sensitivity to A_s , and the sensitivity increases as ALH increases (Figure A1f). An error of 0.01 sensitivity to A_s can result in a retrieval error of 0.1 – 0.6 km for ALH over water surface and 0.1 – 0.4 km over vegetation surface, depending on the aerosol loading (purple curves in Figure B1c-d).
- If including possible uncertainties from these four parameters in vector \mathbf{b} and EPIC measurement error, the estimated ALH retrieval errors (ϵ_{AII}) are shown as red curves in Figure B2a-b. For elevated smoke aerosols with $\tau_{680} = 0.4$, ϵ_{AII} ranges from 0.4 km to 1.0 km for water surface and from 0.7 km to 1.25 km for vegetation surface; the ϵ_{AII} range for $\tau_{680} = 1.0$ is reduced to 0.3 – 0.5 km for water 0.4 – 0.6 km for vegetation surface.

15 Acknowledgements

This research is in part supported by the NASA DSCOVR Earth Science Algorithms Program (Grant No. NNX17AB05G managed by Richard S. Eckman) and in part supported by the Office of Naval Research (ONR) Multidisciplinary University Research Initiatives (MURI) Program under award No. N00014-16-1-2040. We acknowledge the computational support from the High-Performance Computing group at The University of Iowa and the UMBC High Performance Computing Facility (HPCF). The data presented in this paper can be obtained through email to the corresponding authors (X. Xu and J. Wang) of this paper. EPIC Level 1B (L1B) Version 02 imagery data is obtained from the NASA Atmospheric Science Data Center (ASDC). We also acknowledge the AERONET program for providing AOD data retrieved from sunphotometers.

Deleted: NASA

Deleted: 's

Deleted: the

Deleted: 's

Deleted: the

References:

- 25 Babu, S. S., Moorthy, K. K., Manchanda, R. K., Sinha, P. R., Satheesh, S. K., Vajja, D. P., Srinivasan, S., and Kumar, V. H. A.: Free tropospheric black carbon aerosol measurements using high altitude balloon: Do BC layers build “their own homes” up in the atmosphere?, Geophysical Research Letters, 38, L08803, 10.1029/2011gl046654, 2011.
- Baldrige, A. M., Hook, S. J., Grove, C. I., and Rivera, G.: The ASTER spectral library version 2.0, Remote Sensing of Environment, 113, 711-715, 10.1016/j.rse.2008.11.007, 2009.
- 30 Choi, J.-O., and Chung, C. E.: Sensitivity of aerosol direct radiative forcing to aerosol vertical profile, Tellus B, 66, 24376, 10.3402/tellusb.v66.24376, 2014.
- Chowdhary, J., Cairns, B., Mishchenko, M. I., Hobbs, P. V., Cota, G. F., Redemann, J., Rutledge, K., Holben, B. N., and Russell, E.: Retrieval of Aerosol Scattering and Absorption Properties from Photopolarimetric Observations over

- the Ocean during the CLAMS Experiment, *Journal of the Atmospheric Sciences*, 62, 1093-1117, 10.1175/JAS3389.1, 2005.
- Corradini, S., and Cervino, M.: Aerosol extinction coefficient profile retrieval in the oxygen A-band considering multiple scattering atmosphere. Test case: SCIAMACHY nadir simulated measurements, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 97, 354-380, 10.1016/j.jqsrt.2005.05.061, 2006.
- 5 Ding, S., Wang, J., and Xu, X.: Polarimetric remote sensing in oxygen A and B bands: sensitivity study and information content analysis for vertical profile of aerosols, *Atmos. Meas. Tech.*, 9, 2077-2092, 10.5194/amt-9-2077-2016, 2016.
- Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanr^v©, D., and Slutsker, I.: Variability of Absorption and Optical Properties of Key Aerosol Types Observed in Worldwide Locations, *Journal of the Atmospheric Sciences*, 59, 590-608, 10.1175/1520-0469(2002)059<0590:voaaop>2.0.co;2, 2002.
- Dubuisson, P., Frouin, R., Dessailly, D., Dufor^v™t, L., L^v©on, J.-F. B., Voss, K., and Antoine, D.: Estimating the altitude of aerosol plumes over the ocean from reflectance ratio measurements in the O2 A-band, *Remote Sensing of Environment*, 113, 1899-1911, 10.1016/j.rse.2009.04.018, 2009a.
- 15 Dubuisson, P., Frouin, R., Dessailly, D., Dufor^v™t, L., Léon, J.-F., Voss, K., and Antoine, D.: Estimating the altitude of aerosol plumes over the ocean from reflectance ratio measurements in the O2 A-band, *Remote Sensing of Environment*, 113, 1899-1911, 10.1016/j.rse.2009.04.018, 2009b.
- Dufor^v™t, L., Frouin, R., and Dubuisson, P.: Importance and estimation of aerosol vertical structure in satellite ocean-color remote sensing, *Appl. Opt.*, 46, 1107-1119, 10.1364/AO.46.001107, 2007.
- 20 Ge, C., Wang, J., and Reid, J. S.: Mesoscale modeling of smoke transport over the Southeast Asian Maritime Continent: coupling of smoke direct radiative effect below and above the low-level clouds, *Atmos. Chem. Phys.*, 14, 159-174, 10.5194/acp-14-159-2014, 2014.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Silva, A. M. d., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419-5454, 10.1175/jcli-d-16-0758.1, 2017.
- Geogdzhayev, I. V., and Marshak, A.: Calibration of the DSCOVR EPIC visible and NIR channels using MODIS Terra and Aqua data and EPIC lunar observations, *Atmos. Meas. Tech.*, 11, 359-368, 10.5194/amt-11-359-2018, 2018.
- 30 Hanel, R. A.: Determination of cloud altitude from a satellite, *Journal of Geophysical Research*, 66, 1300-1300, 10.1029/JZ066i004p01300, 1961.

- Ichoku, C., Chu, D. A., Mattoo, S., Kaufman, Y. J., Remer, L. A., Tanré, D., Slutsker, I., and Holben, B. N.: A spatio-temporal approach for global validation and analysis of MODIS aerosol products, *Geophysical Research Letters*, 29, 1616, 10.1029/2001GL013206, 2002.
- Jeong, M.-J., and Hsu, N. C.: Retrievals of aerosol single-scattering albedo and effective aerosol layer height for biomass-burning smoke: Synergy derived from “A-Train” sensors, *Geophysical Research Letters*, 35, L24801, 10.1029/2008gl036279, 2008.
- Kipling, Z., Stier, P., Johnson, C. E., Mann, G. W., Bellouin, N., Bauer, S. E., Bergman, T., Chin, M., Diehl, T., Ghan, S. J., Iversen, T., Kirkevåg, A., Kokkola, H., Liu, X., Luo, G., van Noije, T., Pringle, K. J., von Salzen, K., Schulz, M., Seland, Ø., Skeie, R. B., Takemura, T., Tsigaridis, K., and Zhang, K.: What controls the vertical distribution of aerosol? Relationships between process sensitivity in HadGEM3–UKCA and inter-model variation from AeroCom Phase II, *Atmospheric Chemistry and Physics*, 16, 2221–2241, 10.5194/acp-16-2221-2016, 2016.
- Koch, D., and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, 10.5194/acp-10-7685-2010, 2010.
- Koелеmeijer, R. B. A., de Haan, J., and Stammes, P.: A database of spectral surface reflectivity in the range 335–772 nm derived from 5.5 years of GOME observations, *Journal of Geophysical Research*, 108, 4070, 10.1029/2002jd002429, 2003.
- Koffi, B., Schulz, M., Bréon, F.-M., Griesfeller, J., Winker, D., Balkanski, Y., Bauer, S., Berntsen, T., Chin, M., Collins, W. D., Dentener, F., Diehl, T., Easter, R., Ghan, S., Ginoux, P., Gong, S., Horowitz, L. W., Iversen, T., Kirkevåg, A., Koch, D., Krol, M., Myhre, G., Stier, P., and Takemura, T.: Application of the CALIOP layer product to evaluate the vertical distribution of aerosols estimated by global models: AeroCom phase I results, *Journal of Geophysical Research: Atmospheres*, 117, D10201, 10.1029/2011jd016858, 2012.
- Kokaly, R. F., Clark, R. N., Swayze, G. A., Livo, K. E., Hoefen, T. M., Pearson, N. C., Wise, R. A., Benzel, W. M., Lowers, H. A., Driscoll, R. L., and Klein, A. J.: USGS Spectral Library Version 7, Reston, VA, Report 1035, 68, 2017.
- Kokhanovsky, A. A., Mayer, B., and Rozanov, V. V.: A parameterization of the diffuse transmittance and reflectance for aerosol remote sensing problems, *Atmospheric Research*, 73, 37–43, <https://doi.org/10.1016/j.atmosres.2004.07.004>, 2005.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmospheric Measurement Techniques*, 6, 2989–3034, 10.5194/amt-6-2989-2013, 2013.
- 30 Lewis, P., and ~~Barnsley, M. J.~~: Influence of the sky radiance distribution on various formulations of the earth surface albedo, in: *Proc. Conf. Phys. Meas. Sign. Remote Sens*, Val d'Isere, France, 707–715, 1994.
- Lucht, W., Schaaf, C. B., and Strahler, A. H.: An algorithm for the retrieval of albedo from space using semiempirical BRDF models, *Geoscience and Remote Sensing, IEEE Transactions on*, 38, 977–998, 10.1109/36.841980, 2000.

Deleted: J.,

Deleted: B. M.

- Maddy, E. S., DeSouza-Machado, S. G., Nalli, N. R., Barnet, C. D., L Strow, L., Wolf, W. W., Xie, H., Gambacorta, A., King, T. S., Joseph, E., Morris, V., Hannon, S. E., and Schou, P.: On the effect of dust aerosols on AIRS and IASI operational level 2 products, *Geophysical Research Letters*, 39, L10809, 10.1029/2012gl052070, 2012.
- 5 Marshak, A., Herman, J., Szabo, A., Blank, K., Carn, S., Cede, A., Geogdzhayev, I., Huang, D., Huang, L.-K., Knyazikhin, Y., Kowalewski, M., Krotkov, N., Lyapustin, A., McPeters, R., Meyer, K., Torres, O., and Yang, Y.: Earth Observations from DSCOVR/EPIC Instrument, *Bulletin of the American Meteorological Society*, 99, 1829-1850, 10.1175/bams-d-17-0223.1, 2018.
- Martins, J. V., Tanré, D., Remer, L., Kaufman, Y., Mattoo, S., and Levy, R.: MODIS Cloud screening for remote sensing of aerosols over oceans using spatial variability, *Geophysical Research Letters*, 29, 1619, 10.1029/2001GL013252, 10
2002.
- McClatchey, R. A., Fenn, R. W., Selby, J. E. A., Volz, F. E., and Garing, J. S.: *Optical Properties of the Atmosphere (Third Edition)*, Air Force Cambridge Research Labs, Hanscom AFB, MA, 110 pp., 1972.
- Ohtake, M., Matsunaga, T., Yokota, Y., Yamamoto, S., Ogawa, Y., Morota, T., Honda, C., Haruyama, J., Kitazato, K., Takeda, H., Iwasaki, A., Nakamura, R., Hiroi, T., Kodama, S., and Otake, H.: Deriving the Absolute Reflectance of
15 Lunar Surface Using SELENE (Kaguya) Multiband Imager Data, *Space Science Reviews*, 154, 57-77, 10.1007/s11214-010-9689-0, 2010.
- Omar, A. H., Tzortziou, M., Coddington, O., and Remer, L. A.: Plankton Aerosol, Cloud, ocean Ecosystem mission: atmosphere measurements for air quality applications, *J. Appl. Remote Sens.*, 12, 042608, 10.1117/1.JRS.12.042608, 2018.
- 20 Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick, G. P., Cossuth, J. H., and DeLand, M. T.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *npj Climate and Atmospheric Science*, 1, 30, 10.1038/s41612-018-0039-3, 2018.
- Remer, L. A., Kaufman, Y. J., Tanre, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R.-R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS Aerosol Algorithm, Products, and
25 Validation, *Journal of the Atmospheric Sciences*, 62, 947-973, doi:10.1175/JAS3385.1, 2005.
- Rothman, L. S., Gordon, I. E., Babikov, Y., Barbe, A., Chris Benner, D., Bernath, P. F., Birk, M., Bizzocchi, L., Boudon, V., Brown, L. R., Campargue, A., Chance, K., Cohen, E. A., Coudert, L. H., Devi, V. M., Drouin, B. J., Fayt, A., Flaud, J. M., Gamache, R. R., Harrison, J. J., Hartmann, J. M., Hill, C., Hodges, J. T., Jacquemart, D., Jolly, A., Lamouroux, J., Le Roy, R. J., Li, G., Long, D. A., Lyulin, O. M., Mackie, C. J., Massie, S. T., Mikhailenko, S.,
30 Müller, H. S. P., Naumenko, O. V., Nikitin, A. V., Orphal, J., Perevalov, V., Perrin, A., Polovtseva, E. R., Richard, C., Smith, M. A. H., Starikova, E., Sung, K., Tashkun, S., Tennyson, J., Toon, G. C., Tyuterev, V. G., and Wagner, G.: The HITRAN2012 molecular spectroscopic database, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 130, 4-50, 10.1016/j.jqsrt.2013.07.002, 2013.

- Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Penner, J. E., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, *Atmos. Chem. Phys.*, 13, 2423-2434, 10.5194/acp-13-2423-2013, 2013.
- 5 Satheesh, S. K., Moorthy, K. K., Babu, S. S., Vinoj, V., and Dutt, C. B. S.: Climate implications of large warming by elevated aerosol over India, *Geophysical Research Letters*, 35, L19809, 10.1029/2008GL034944, 2008.
- Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang, X., Jin, Y., Muller, J.-P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., d'Entremont, R. P., Hu, B., Liang, S., Privette, J. L., and Roy, D.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote Sensing of Environment*, 83, 135-148, [https://doi.org/10.1016/S0034-4257\(02\)00091-3](https://doi.org/10.1016/S0034-4257(02)00091-3), 2002.
- 10 Spurr, R.: VLIDORT: A linearized pseudo-spherical vector discrete ordinate radiative transfer code for forward model and retrieval studies in multilayer multiple scattering media, *Journal of Quantitative Spectroscopy & Radiative Transfer*, 102, 316-342, 2006.
- Spurr, R., Wang, J., Zeng, J., and Mishchenko, M. I.: Linearized T-matrix and Mie scattering computations, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 425-439, 10.1016/j.jqsrt.2011.11.014, 2012.
- 15 Spurr, R., and Christi, M.: On the generation of atmospheric property Jacobians from the (V)VLIDORT linearized radiative transfer models, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 142, 109-115, 10.1016/j.jqsrt.2014.03.011, 2014.
- Tilstra, L. G., Tuinder, O. N. E., Wang, P., and Stammes, P.: Surface reflectivity climatologies from UV to NIR determined from Earth observations by GOME-2 and SCIAMACHY, *Journal of Geophysical Research: Atmospheres*, 122, 4084-4111, doi:10.1002/2016JD025940, 2017.
- 20 Tomasi, C., and Petkov, B. H.: Calculations of relative optical air masses for various aerosol types and minor gases in Arctic and Antarctic atmospheres, *Journal of Geophysical Research: Atmospheres*, 119, 1363-1385, doi:10.1002/2013JD020600, 2014.
- 25 Torres, O., Bhartia, P. K., Herman, J. R., Ahmad, Z., and Gleason, J.: Derivation of aerosol properties from satellite measurements of backscattered ultraviolet radiation: Theoretical basis, *J. Geophys. Res.*, 103, 17099-17110, 10.1029/98jd00900, 1998.
- Torres, O., Ahn, C., and Chen, Z.: Improvements to the OMI near-UV aerosol algorithm using A-train CALIOP and AIRS observations, *Atmos. Meas. Tech.*, 6, 3257-3270, 10.5194/amt-6-3257-2013, 2013.
- 30 Toth, T. D., Campbell, J. R., Reid, J. S., Tackett, J. L., Vaughan, M. A., Zhang, J., and Marquis, J. W.: Minimum aerosol layer detection sensitivities and their subsequent impacts on aerosol optical thickness retrievals in CALIPSO level 2 data products, *Atmos. Meas. Tech.*, 11, 499-514, 10.5194/amt-11-499-2018, 2018.

- Vaughan, M. A., Powell, K. A., Winker, D. M., Hostetler, C. A., Kuehn, R. E., Hunt, W. H., Getzewich, B. J., Young, S. A., Liu, Z., and McGill, M. J.: Fully Automated Detection of Cloud and Aerosol Layers in the CALIPSO Lidar Measurements, *Journal of Atmospheric and Oceanic Technology*, 26, 2034-2050, 10.1175/2009jtecha1228.1, 2009.
- 5 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70-83, 10.1016/j.rse.2011.09.027, 2012.
- 10 Wang, J., Xu, X., Ding, S., Zeng, J., Spurr, R., Liu, X., Chance, K., and Mishchenko, M.: A numerical testbed for remote sensing of aerosols, and its demonstration for evaluating retrieval synergy from a geostationary satellite constellation of GEO-CAPE and GOES-R, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 146, 510-528, 10.1016/j.jqsrt.2014.03.020, 2014.
- Waquet, F., Cairns, B., Knobelspiesse, K., Chowdhary, J., Travis, L. D., Schmid, B., and Mishchenko, M. I.: Polarimetric remote sensing of aerosols over land, *J. Geophys. Res.*, 114, D01206, 10.1029/2008jd010619, 2009.
- 15 Wendisch, M., Hellmuth, O., Ansmann, A., Heintzenberg, J., Engelmann, R., Althausen, D., Eichler, H., Müller, D., Hu, M., Zhang, Y., and Mao, J.: Radiative and dynamic effects of absorbing aerosol particles over the Pearl River Delta, China, *Atmospheric Environment*, 42, 6405-6416, <http://dx.doi.org/10.1016/j.atmosenv.2008.02.033>, 2008.
- Wessel, P., and Smith, W. H. F.: A global, self-consistent, hierarchical, high-resolution shoreline database, *Journal of Geophysical Research: Solid Earth*, 101, 8741-8743, doi:10.1029/96JB00104, 1996.
- 20 Winker, D. M., Tackett, J. L., Getzewich, B. J., Liu, Z., Vaughan, M. A., and Rogers, R. R.: The global 3-D distribution of tropospheric aerosols as characterized by CALIOP, *Atmos. Chem. Phys.*, 13, 3345-3361, 10.5194/acp-13-3345-2013, 2013.
- [Xu, X., and Wang J.: Retrieval of aerosol microphysical properties from AERONET photopolarimetric measurements: 1. Information content analysis, *Journal of Geophysical Research: Atmospheres*, 120, 7059–7078, 2015.](#)
- 25 Xu, X., Wang, J., Wang, Y., Zeng, J., Torres, O., Yang, Y., Marshak, A., Reid, J., and Miller, S.: Passive remote sensing of altitude and optical depth of dust plumes using the oxygen A and B bands: First results from EPIC/DSCOVr at Lagrange-1 point, *Geophysical Research Letters*, 44, 7544-7554, 10.1002/2017GL073939, 2017.
- Xu, X., Wang, J., Wang, Y., and Kokhanovsky, A.: Chapter 1 - Passive Remote Sensing of Aerosol Height, in: *Remote Sensing of Aerosols, Clouds, and Precipitation*, Elsevier, 1-22, 2018a.
- 30 Xu, X., Wang, J., Zeng, J., Hou, W., Meyer, K. G., Platnick, S. E., and Wilcox, E. M.: A pilot study of shortwave spectral fingerprints of smoke aerosols above liquid clouds, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 221, 38-50, 10.1016/j.jqsrt.2018.09.024, 2018b.
- Yamamoto, G., and Wark, D. Q.: Discussion of the letter by R. A. Hanel, "Determination of cloud altitude from a satellite", *Journal of Geophysical Research*, 66, 3596-3596, 10.1029/JZ066i010p03596, 1961.

- Yang, Y., Marshak, A., Mao, J., Lyapustin, A., and Herman, J.: A method of retrieving cloud top height and cloud geometrical thickness with oxygen A and B bands for the Deep Space Climate Observatory (DSCOVR) mission: Radiative transfer simulations, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 122, 141-149, <http://dx.doi.org/10.1016/j.jqsrt.2012.09.017>, 2013.
- 5 Young, S. A., and Vaughan, M. A.: The Retrieval of Profiles of Particulate Extinction from Cloud-Aerosol Lidar Infrared Pathfinder Satellite Observations (CALIPSO) Data: Algorithm Description, *Journal of Atmospheric and Oceanic Technology*, 26, 1105-1119, 10.1175/2008jtecha1221.1, 2009.
- Zhang, J., Campbell, J. R., Reid, J. S., Westphal, D. L., Baker, N. L., Campbell, W. F., and Hyer, E. J.: Evaluating the impact of assimilating CALIOP-derived aerosol extinction profiles on a global mass transport model, *Geophys. Res. Lett.*, 38, L14801, 10.1029/2011gl047737, 2011.
- 10

Tables

Table 1. Adopted parameters for generating LUTs.

| Parameter* | Prescribed values for LUTs |
|-----------------------------|---|
| AOD at 680 nm | 0.0, 0.1, 0.2, 0.4, 0.7, 1.0, 1.5, 2.0, and 3.0 |
| ALH above surface (km) | 0 to 15 km at 1 km interval |
| Surface reflectance | 0, 0.05, 0.1, 0.2, 0.3, 0.4, and 0.6 |
| θ_0 and θ (°) | 0° to 72° at interval of 6°, and $ \theta_0 - \theta < 15^\circ$ |
| $\Delta\varphi$ (°) | 0° to 180° at interval of 12°, |
| Surface pressure (hPa) | 700, 800, 900, and 1050 |

* θ_0 indicates the solar zenith angle, θ the satellite viewing zenith angle, and $\Delta\varphi$ the relative azimuth angle

5

Table 2. Spectral settings of UNL-VRM for constructing the LUTs

| EPIC channel (nm) | Spectral range (nm) | Spectral interval (nm) | FWHM (nm) |
|----------------------|------------------------|---------------------------|--------------|
| 443 | 440 – 445 | 0.1 | 0.2 |
| 551 | 549 – 555 | 0.1 | 0.2 |
| 680 | 675 – 685 | 0.1 | 0.2 |
| 688 | 685 – 690 | 0.01 | 0.02 |
| 764 | 760 – 766 | 0.01 | 0.02 |
| 780 | 776 – 782 | 0.1 | 0.2 |

Deleted: in


10

Table 3. AERONET sites selected for AOD validation

| Site name | Latitude (°) | Longitude (°) | N_{AOD}^* |
|---------------|--------------|---------------|--------------------|
| Billerica | 42.53 | - 71.27 | 3 |
| Churchill | 58.74 | - 93.82 | 3 |
| Egbert | 44.23 | - 78.78 | 2 |
| Lake_Erie | 41.83 | - 83.19 | 2 |
| LISCO | 40.76 | - 73.34 | 4 |
| NEON_UNDE | 46.23 | - 89.54 | 2 |
| Pickle_Lake | 51.45 | - 90.22 | 1 |
| Thompson_Farm | 43.11 | - 70.95 | 3 |
| Toronto | 43.79 | - 79.47 | 2 |


* N_{AOD} is the number of collocated AERONET AOD values

Figures



5 **Figure 1.** EPIC instrument filter response function (blue) and atmospheric spectral transmission (orange). Panel (a) includes all ten EPIC bands, whereas panels (b) and (c) show zoom-ins for the 688-nm channel in the O₂ B band and the 764-nm channel in the O₂ A band, respectively. Here the atmospheric transmission is simulated by the UNL-VRTM model, with a spectral step size and a spectral full width at half maximum of 0.02 nm.

Deleted: F
 Deleted: s the EPIC instrument
 Deleted: the
 Deleted: band
 Deleted: -
 Deleted: -



5 **Figure 2.** Schematic diagram of the physical principle of remote sensing of aerosol layer height in the O₂ absorption bands. Shown is the same aerosol layer located at two different altitudes in the atmosphere. Due to the scattering by aerosol particles, photons scattered by a higher aerosol layer (right panel) would travel a shorter pathlength to reach the satellite sensor than those scattered by the lower-altitude aerosol (left panel), resulting in less absorption by O₂ and thus a higher radiance value detected by the satellite (i.e., $I_H > I_L$).

Deleted: al for

Deleted: of

Deleted: path way

Deleted: in

Deleted: a

Deleted: less

Deleted: chance of




Figure 3. Sensitivity of DOAS ratios of TOA reflectance around the O₂ A and B bands to smoke ALH as simulated by UNL-VRM for different AOD and surface reflectance (A_s) values. Simulations were performed for a typical biomass-burning aerosol model as observed by EPIC for the geometry of $[\theta_0, \theta, \Delta\phi] = [42^\circ, 37^\circ, 165^\circ]$, where θ_0 and θ are solar and viewing zenith angles, and $\Delta\phi$ the relative azimuth angle.

Deleted: at




Figure 4. Surface reflectance spectra (from the ASTER spectral library; Baldrige et al., 2009) for various surface types in the visible-to-NIR range, with selected EPIC spectral bands within this range shown in gray.





Figure 5. Flowchart of the AOD/ALH retrieval algorithm.






Figure 6. Determination of surface reflectance from MODIS surface albedo products. (a) USGS reflectance spectra for different vegetation samples, and (b) statistics for regression (i.e., slope, offset, and coefficient of determination, R^2) of reflectance between different EPIC bands, marked as red dotted lines in (a), and the respective closest MODIS bands, marked as blue lines in (a), according to the spectral library. (c-f) Scatterplots of reflectance in each EPIC band versus reflectance in the corresponding MODIS bands. Linear regression lines are shown in red.

5

Deleted: ir
 Deleted: at
 Deleted: at





Figure 7. Demonstration of ALH and AOD retrievals for three EPIC scenes acquired on 25 August 2017 around UTC times of 16:10, 17:26, and 18:32. (a) RGB composites of EPIC 443 nm, 551 nm, and 680 nm data. The gold line indicates the CALIOP sub-orbital track with an overpass time of 19:05 UTC. (b) Retrieved smoke ALH when 680-nm AOD is larger than 0.2. (c) Retrieved 680 nm AOD. Small circles on AOD maps represent AOD values observed at corresponding AERONET sites within two hours of the EPIC scan time.



5 **Figure 8.** Same as Figure 7 but for three EPIC scenes captured on 26 August 2017 at UTC times of 16:11, 17:17, and 18:22.

The CALIOP overpass was at 18:15 UTC.

Deleted: time




Figure 9. Comparison of ALH retrieved from EPIC and CALIOP level-2 aerosol extinction profile in both ‘curtain’ view (left column) and ‘scatterplot’ view (right column) for two-day overpasses. The CALIOP orbital tracks are marked on EPIC RGB images in Figures 7–8. Different EPIC scan times are marked with different line/symbol colors in the comparison. The error bar for EPIC ALH represents the standard deviation for an array of 3x3 24-km retrieval pixels, while that for CALIOP ALH represents the standard deviation of over 5 adjacent CALIOP 5-km footprints. Also shown in the left panels is the CALIOP* ALH (thin black curve) calculated with a background aerosol profile imposed for undetected aerosol layers (see text for detail), whereas bold black curve represents the CALIOP ALH without considering background aerosol.



Deleted:

Deleted: the ALH derived from

Deleted: E

Deleted: of

Deleted: the error bar of



Deleted:

Deleted: counterparts from

Deleted: C

Deleted: of

Deleted: b

Deleted: D

Deleted: by

Deleted: ting

Deleted:

Deleted: the

Deleted: s

Figure 10. Comparison of EPIC ALH and AOD with the corresponding CALIOP and AERONET measurements. (a) Scatterplot of EPIC ALH versus CALIOP ALH by including all EPIC-CALIOP pairs shown in Figure 9. The color of each scatter point represents the EPIC 680-nm AOD value for the same EPIC pixels. (b) Same as panel (a) but for CALIOP ALH calculated with a background aerosol profile imposed for undetected aerosol layers. (c) Scatterplot of EPIC 680-nm AOD versus AERONET 675-nm AOD, collocated at 9 AERONET sites listed in Table 3. The dotted lines in both the scatterplots represent error envelopes, i.e., ± 0.5 km for ALH and $\pm (0.05 + 10\%)$ for AOD. Also annotated are the one-to-one line (solid black line), linear regression fit (red line), number of scatter points (N), coefficient of determination (R^2), significance level (P), linear regression equation, bias, percentage of scatter points within error envelop (EE%), and root-mean-squared-error (rmse).




Figure 11. EPIC UV aerosol index (UVAI) for the three scenes on 25 August 2017 shown in Figure 7. Here the UVAI was
 5 obtained from EPIC Level-2 UV aerosol products.

Deleted: were




Figure 12. Scatterplots of EPIC UVAI (Figure 11) versus current retrievals of ALH (Figure 7) on 25 August 2017 (17:26 UTC) under different AOD values as indicated in the title of each panel. Also annotated are the linear regression fit, number of scatter points (N), coefficient of determination (R^2), and significance level (P).

Deleted: ting

Deleted: s




Figure A1. UNL-VRTM simulated DOAS ratios ρ in EPIC O₂ A and B band (a), and their Jacobian gradients with respect to ALH (b), half-width parameter σ_H for the quasi-Gaussian aerosol vertical profile (c), 680-nm AOD τ_{680} (d), SSA (e), and surface reflectance A_s (f). The y-axis represents aerosols being present at various ALH values. Three colors indicate ρ and its Jacobians for the O₂ A band with $A_s = 0.05$ (blue), O₂ A band with $A_s = 0.30$ (green), and O₂ B band with $A_s = 0.05$ (red). Two AOD loadings are indicated by the solid ($\tau_{680} = 0.4$) and dotted ($\tau_{680} = 1.0$) lines. Simulations are performed with the same aerosol model and observation geometry in Figure 3.

10




Figure A2. Estimated ALH retrieval errors caused by EPIC measurement errors and by uncertainties in aerosol and surface assumptions for two surface types. Left column is for a surface with $A_s = 0.05$ reflectance in both the O₂ A and B bands (close to a water surface), and right column is for $A_s = 0.30$ in the A band and $A_s = 0.05$ in the B band (close to a vegetation surface). Top panels show ALH retrieval errors due to 2% EPIC measurement error alone (black curves) and with model uncertainty added (red curves). Bottom panels show the error in retrieved ALH due to the uncertainties of 0.5 km in σ_H (blue), 0.05 in τ_{680} (orange), 0.03 in SSA (green), and 0.01 in A_s (purple). In all panels, the solid and dotted curves represent 0.4 and 1.0 τ_{680} , respectively.