

Reviewer #1

We thank the reviewer for the insightful comments (**bold font**) and we have replied to each of the comments and queries below each comment (regular font) and modifications to the manuscript are in *italic font*.

The manuscript attempts to answer the following questions: How many different cloud types co-exist within a particular area? What cloud type mixtures are more prevalent? How do answers to the above two questions depend on area size? (side question that emerges: at what spatial scale does one encounter the greatest diversity of distinct cloud type mixtures?). These all sound kind of philosophical questions, but the authors find practical relevance (at least for the first question) for AIRS (and AMSU) scales cloud retrievals. The link to AIRS allows the authors to make one the major compromises of the study: only cloud type identification of the topmost cloudy layer matters because that's where AIRS is most sensitive even though the data source identifying cloud type provides vertical profile information. The other major compromise is that when identifying cloud mixtures, the frequency of occurrence of each cloud type does not matter, in other words cloud mixtures consisting of the same cloud types are treated as equivalent even if the contributions of a cloud type are different. These two simplifications, along with an additional one where the spatial arrangement of the cloud types is ignored allow the authors to reduce the dimensionality of the problem and make the analysis tractable.

We appreciate the reviewer's synthesis of the paper and agree these are the major thrusts of the first aspect of the paper.

This is overall quite a difficult paper to read, but I find the results of the first part quite fascinating (I was less excited about the implications for AIRS retrievals—although I understand that these findings are important for understanding the quality of the AIRS retrievals), so I recommend acceptance of the article to AMT. As you can see below, I have some inquiries some of which are also of the philosophical kind I'd like the authors to consider.

We have made some edits and changes that follow from the reviewer suggestions that we hope make the manuscript more readable. We do appreciate the reviewer's point of view regarding how this paper may be viewed in two distinct pieces. However, our end goal of this work was to show practical relevance to the cloud scene variability and ultimately establish why it is important to describe cloud scenes on a pixel-by pixel basis. The reasons are hopefully clearer in the revised version as we have more carefully documented AIRS retrievals of cloud phase and ice cloud properties as a function of whether the scene contains one cloud type or multiple cloud types, or whether the scene is completely or partly cloudy, instead of reporting them in a slightly convoluted manner. These changes are closely coupled to those arising from reviewer #2's comments (please refer to replies to reviewer #2 for further detail).

– What does the cloud type from 2B-CLDCLASS mean? The names of cloud types are the same as the ones used by surface observers, but are they related? Some description of the physical meaning of the cloud types given their method of identification by the 2B-CLDCLASS algorithm is needed.

The 2B-CLDCLASS algorithm is described in Sassen and Wang (2005, 2008) that is referenced in the manuscript. As stated in Sassen and Wang (2008), the algorithm is based on earlier work by the same authors and combines the measurements of ground-based multiple remote sensors. They report having tested the results against surface observer cloud reports. We have included some additional text at the beginning of Section 2.2 to clarify:

“The CloudSat 2B-CLDCLASS product is used in this work and the algorithm is described in Sassen and Wang (2005, 2008). As summarized in Sassen and Wang (2008) and previous works, the algorithm uses methods developed from ground-based multiple remote sensors that have been tested against surface observer-based cloud typing reports. The cloud classification occurs in two steps. First, a clustering analysis is performed to group cloud profiles into cloud clusters. Secondly, classification methods are used to classify clouds into different cloud types. The decision trees guiding the classification are complex and are based on 23 variables derived from the clustering analysis of the first stage. Geometric quantities such as cloud base, top, and horizontal extents are present in decision trees (Sassen and Wang, 2005).”

I’m sure the authors are aware that another version of the product currently exists, 2B-CLDCLASS-LIDAR where the CALIPSO lidar assists in the identification of the cloud type. Why was this newer product not used? (I suspect the authors may have started the work before this product was released). If the authors were to use 2B-CLDCLASS-LIDAR and the results changed in a major way, how would that undermine the fundamentals and motivation for the first part of the study?

This is a good question and a fair one to ask. We are motivated by the results of Kahn et al. (2018) that suggests larger particle sizes in convective clouds compared to thin cirrus. The 2B-CLDCLASS product is better suited for differentiating cloud types other than small particle thin cirrus, in which 2B-CLDCLASS-LIDAR would excel. While AIRS is very sensitive to thin cirrus, the sensitivity of AIRS ice cloud particle size most strongly responds to clouds around an optical thickness on the order of 1.0. If we used 2B-CLDCLASS-LIDAR, the statistics would be weighted to the detection of vast areas of thin cirrus in layers above optically thicker clouds and elsewhere in absence of other cloud types. The Ci classification dominates in the 2B-CLDCLASS-LIDAR data set and will blur the signals of cumulus and deep convective cloud types capped by thin cirrus. As we describe in the paper, the simplifications required to make the approach tractable require us to quantify cloud type at cloud top, and thus we would lose the discriminatory ability for clouds that occur just below a thin layer of cirrus. We expect that the mixtures would be profoundly different in 2B-CLDCLASS-LIDAR, with much less ability to demonstrate AIRS’ skill at obtaining larger particle sizes at convective cloud tops (Kahn et al. 2018). Ultimately, we argue that 2B-CLDCLASS is a more appropriate tool for this work. We have included some additional text in Section 2.2 to clarify:

“Lastly, the results of Kahn et al. (2018) suggest larger ice cloud particle sizes occur at convective cloud tops compared to thin cirrus at the same cloud top temperature. Given the key assumption of cloud typing only at cloud top, the 2B-CLDCLASS product is better suited for identifying convective clouds in AIRS apart from stratiform clouds, the latter of which are dominant in 2B-CLDCLASS-LIDAR. If 2B-CLDCLASS-LIDAR was used in place of 2B-CLDCLASS, the statistics would be weighted towards the detection of vast areas of cirrus in thin layers above and in proximity to convective clouds. The Ci classification dominates in 2B-CLDCLASS-LIDAR at cloud top and will blur the signals of underlying cumulus and deep convective cloud types that are capped by thin cirrus.”

What if a completely different cloud type product was used, e.g., based on passive satellite observations where cloud type is identified by location in a cloud-top- pressure/cloud-optical-thickness joint histogram (the authors briefly touch on this in the last paragraph, but only with regard to the AIRS application – I’m more interested in the cloud scene climatology aspects)?

As the reviewer certainly knows, how one goes about “typing clouds” is not a settled research topic and is sensitive to the instrument sampling, radiance sensitivity, wavelength, underlying assumptions, and so forth. In the conclusions, we touch on the results of Wang et al. (2016) where comparisons of CloudSat cloud types as used here in this work are compared to ISCCP-like categories derived from the MODIS imager, which could be used in place of the CloudSat cloud typing. We expect that the CloudSat radar will have more skill in discriminating convective clouds from stratiform clouds than passive sensors, as these two types of clouds show strong differences in the AIRS microphysical retrievals. For many cloud types, the detection is similar between passive and active; please see Wang et al. (2016) for specifics.

– It seems to me that the results depend completely on how frequently 2B-CLDCLASS identifies certain cloud types based on its internal definitions. Yes, the authors do not often find mixtures containing stratus (St) simply because St is extremely rare in 2B-CLDCLASS, probably unrealistically so given other methods identifying St (I mean, cloud types will always be loosely defined).

We agree. Please refer above for our generalized perspective. It is well established that 2B-CLDCLASS contains very little stratus because of ground clutter in the bottom 3-4 bins.

I think one figure that the paper needs to include is the global frequency of the different cloud types according to 2B-CLDCLASS at its native resolution. This will give immediately clues on why certain cloud type mixtures (scenes) will be rare right off the bat (the authors kind of bring this this up already in some instances, e.g., p. 6, line 4). With DC, Cu, and St being rare according to 2B-CLDCLASS, one would expect that scenes containing those will also be rare.

With regard to plan view maps and zonal averaged plots of cloud type frequencies from 2B-CLDCLASS, these have been reported in the literature, in particular we are referring to Sassen

and Wang (2008), their Figure 1 for the plan view and their Figure 2 for the zonal average. We do not see a need to repeat these results in this paper. With regard to the relative histogram counts of cloud type as a function of length scale, these are depicted in Figure 6 in the manuscript. The percentages of cloud type and cloud scene frequencies are also reported in Tables 3-6 in the revised manuscript.

We have added the following statement in Section 2.2 to refer the reader for more detail on cloud type frequencies: *“Plan view and zonal average frequencies of 2B-CLDCLASS cloud types at its native resolution are reported in Sassen and Wang (2008).”*

It is unfortunate that the abbreviation for certain cloud types changes throughout the text, tables, and figures: As becomes AlSt, Ac becomes AlCu, Ci becomes ci, DC becomes Dc, and so forth. Please fix and make consistent throughout!

We have checked the notation throughout the revised manuscript and fixed any inconsistencies in the labeling. We intended to follow the notation cumulus (Cu), stratocumulus (Sc), stratus (St), altocumulus (Ac), altostratus (As), nimbostratus (Ns), cirrus (Ci), deep convective (Dc) clouds and a ninth classification of clear sky designated no cloud (nc). In addition to text changes, we corrected figure 1, 5, 6 and 7.

I don't understand panel d in Fig. 2. Whatever it depicts, it does not appear to have a very interesting pattern!

We agree that these are confusing and have removed panels b and d in figure 2 as they are not key pieces of information for the manuscript.

I recognize that the authors make a valiant effort in section 3.3, but that part of the paper remains a hard read. In this section, line 8 of p. 8 indicates that 200 possible mixed scenes were identified which seems to contradict the 194 figure quoted earlier (p. 5, line 23). Are these numbers for areas of different size (e.g., a third figure of 210 different scenes emerges for 105 km). Please clarify, 194, 200, and 210. Moreover, I found odd that the authors state (p. 7 line 2) that “The maximum number of observed cloud scenes (210) at a particular horizontal scale (105 km) remains unexplained” when the section that immediately follows tries to explain exactly that. Am I missing the subtle distinction? Section 3.3 tries to explain why the maximum number happens at 105 (not sure it succeeds), but why this maximum number is 210 remains as the unexplained mystery?

In Section 3.1, we describe the statistics of cloud scenes observed at the AMSU resolution at 45-km scales. There are 194 observed cloud scenes at this resolution. In section 3.2, we quantify the scale dependence of cloud scene statistics. The number of cloud scenes is shown to first increase with resolution then decrease as scales get large. The maximum number of cloud scenes is 210 and is observed at a resolution of 105 km. Both of the numbers 194 and 210 appear as 2 points on figure 3a. The former is indicated with the intersection of the red line and the curve, and the latter is just the value at the maximum.

The third number mentioned by the reviewer is 200 and it is a result of the procedure described in section 3.3. We shortened the name of this section 3.3 “Generalizing to all scales” as we understood from the reviewer’s comment that the title was potentially confusing or misleading. The number of 200 mixed scenes is identified *independently of a grid resolution*.

Therefore, there are 194 cloud scenes observed at a resolution of ~45 km, 210 scenes observed at a resolution of 105 km and 200 scenes identified with a procedure independent from a regular grid.

We have added the following text at the start of Section 3.2 to clarify: *“In Section 3.1, the relative frequencies of cloud scenes were derived for exact matches of AIRS and AMSU observations to the CloudSat ground track. As the CloudSat ground track can oscillate over several AIRS FOVs across a scan line within a given orbit, the numbers of coincident CloudSat profiles matching to AIRS and AMSU will vary. Below, cloud scenes are derived independently of the specific AIRS and AMSU collocation geometry.”*

As far as the quote mentioned by the reviewer on p. 7 line 2, that was intended to be a transition from Section 3.2 to 3.3 and as motivation for why we calculated the scale dependence of cloud scenes independently of a particular grid resolution. We have changed the text to the following to clarify: *“The reasons for the maximum number of observed cloud scenes (210) at a particular horizontal scale (105 km) are not immediately clear.”* Then we have changed the text on p.7 lines 4-5 to further clarify: *“A simple conceptual model is described below that is able to approximate the results of Fig. 3 and offers some insight for the observed maximum frequency of cloud scenes and the spatial scale at which it occurs.”*

We have rewritten some of the paragraph on p. 8 lines 11-20 as follows: *“A total of 200 out of 247 possible mixed scenes were identified. The minimum and maximum length occurrence frequencies of five cloud scenes (A_c, S_c), (A_s, S_c, C_u), (C_i, A_s, C_u, d_c), (A_s, A_c, N_s, d_c) and (C_i, A_s, A_c, S_t, S_c) selected randomly from the 200 present in the two-year record, are shown in Fig. 4a and 4c, respectively. Recall that the maximum length is defined from relation (2), while the minimum length is defined from relation (3), with an illustrative example shown for (A_c, S_c). From top to bottom, their respective ranks are 1, 26, 51, 76 and 101. It is striking that each frequency histogram in Fig. 4a and 4c is not monotonic and displays a frequency maximum between 100 and 1000 km. Consequently, the sum of all (200) observed mixed scenes across length scales will result in a curve with a maximum and these are shown in Fig. 4b and 4d. Both curves are very similar to Fig. 3a and have maxima for about 180 observed scenes at 77 km and 174 km, respectively. Using the methodology outlined in (1) to (3), the scale dependence of the number of observed scenes shows that the maximum will occur somewhere between 77 and 174 km.”*

– Can the same scale be used for the y-axis of Figs. 7 and 8? You say that that the common panels of these two figures (single cloud type scenes) should look very similar (inclusion of clear-sky notwithstanding), but the comparison is hampered by different y-axis range.

We regenerated figures 7 and 8 with the same ordinate scales. However, the reviewer should keep in mind that each bin in Figures 7-10 (revised) are normalized by the total number of cloud scenes (within an AIRS footprint) for single or multiple cloud types, or partly cloudy or fully cloudy scenes. We have added the following text to the caption of Figure 7: *“Each histogram sums to 1.0 and does not show the numbers of counts relative to another histogram. Relative counts could be inferred from the percentages listed in the 2nd to left column of Table 3.”*

– Somewhere in section 2 mention what the maximum optical thickness retrievable by AIRS is.

We added the following sentence on line 12 page 5: *“The AIRS sampling includes nearly all ice clouds with $\tau_i > 0.1$, while the maximum values of τ_i asymptote to values near 6–8 (e.g. Kahn et al., 2015).”*

Reviewer #2

We thank the reviewer for the insightful comments (**bold font**) and we have replied to each of the comments and queries below each comment (regular font) and modifications to the manuscript are in *italic font*.

In this paper AIRS measurements are combined with CloudSat observations to obtain statistics of the variation of cloud types within the AIRS field of view. These are then used to study the variation of cloud properties per cloud scene and the sensitivity of ice cloud properties retrievals to the variation of cloud types within the pixels. The study is interesting and the methods are generally sound. However, some issues with the presentation and the analysis need to be addressed before the paper can be accepted for publication. Below the major and minor comments are listed:

1) The CloudSat cloud types are used. However, these type names are not very quantitative. Please indicate how the different types are defined and summarize how they are derived from CloudSat observations. The reader should not have to dig through other papers to be able to understand the data presented here. For instance, what are the altitude boundaries distinguishing low, middle and high clouds? It is also not clear to me how cumulus and stratus can be distinguished using a single radar profile.

According to Sassen and Wang (2005), the cloud classification occurs in two steps. First, a clustering analysis is performed to group cloud profiles into cloud clusters. Secondly, classification methods are used to classify clouds into different cloud types. The decision trees guiding the classification are complex and based on 23 variables derived from the clustering analysis of the first stage. Conditions upon geometric quantities such as cloud base, top, and horizontal extents are present in decision trees (page 32 to 35 in Sassen and Wang, 2005).

One of our responses to reviewer #1 is relevant here and we added text to the manuscript (repeated from response to reviewer #1): *“The CloudSat 2B-CLDCLASS product is used in this work and the algorithm is described in Sassen and Wang (2005, 2008). As summarized in Sassen and Wang (2008) and previous works, the algorithm uses methods developed from ground-based multiple remote sensors that have been tested against surface observer-based cloud typing reports. The cloud classification occurs in two steps. First, a clustering analysis is performed to group cloud profiles into cloud clusters. Secondly, classification methods are used to classify clouds into different cloud types. The decision trees guiding the classification are complex and are based on 23 variables derived from the clustering analysis of the first stage. Conditions upon geometric quantities such as cloud base, top, and horizontal extents are present in decision trees (Sassen and Wang, 2005).”*


In the conclusions, we describe the results of Wang et al. (2016) where comparisons of CloudSat cloud types as used here in this work are compared to ISCCP-like categories derived from the MODIS imager, which could be used in place of the CloudSat cloud typing. For many cloud types, the detection is similar between passive and active; please see Wang et al. (2016) for specifics.

2) Please be consistent with cloud type names throughout the paper. For example, sometimes “As” is used and sometimes “AlSt”. Sometimes “clear sky” is used, sometimes “nc”. Also, mixtures are denoted with commas, although clear-sky is left out in this notation. I suggest to indicate mixtures of cloud types and clear sky consistently, e.g., “As, nc”. To give an example how this is confusing: Looking at Fig. 6, It was unclear how cumulus can have a length scale of 400 km, but I guess that’s possible because clear-sky is mixed in. This can be made clearer by naming this mixture “Cu, nc” (This then excludes pure “Cu” cases. If these are included too, then I suggest using something like “Cu + Cu, ns”).

We have checked the notation throughout the revised manuscript and fixed any inconsistencies in the labeling. We intended to follow the notation cumulus (Cu), stratocumulus (Sc), stratus (St), altocumulus (Ac), altostratus (As), nimbostratus (Ns), cirrus (Ci), deep convective (Dc) clouds and a ninth classification of clear sky designated no cloud (nc). In addition to text changes, we corrected figure 1, 5, 6 and 7.

Cumulus lengths reported in Figure 6 are for pure Cu and not a blend of clear and Cu (Cu,nc). Cumulus clouds of any length are possible according to the decision tree of low cloud classifier (page 34 of Sassen and Wang, 2005) when a cluster cloud fraction is smaller than 0.25. Sassen and Wang (2005) do not indicate that a maximum cluster length is prescribed, so long cumulus clouds might result from the unlikely situation where a long cumulus cloud is part of a cluster with cloud fraction less than 0.25.

The fact is that we observed long clouds in each cloud category of the 2B-CLDCLASS CloudSat product. One may quibble about this categorization but the algorithm most closely associates that cloud with Cu. For example, we found that the longest cumulus cloud is located in granule 03498 of the 2B-CLDCLASS product, 2006358131321_03498_CS_2B-CLDCLASS_GRANULE_P_R04_E02.hdf using the cloud type defined at the cloud top as described in the manuscript. The very long Cu cloud starts at profile # 25681 and ends at profile # 26118. It therefore has length of 438 CloudSat profiles or about 482 km. The location of this long cumulus cloud is shown in the figure below:



This fact emphasizes that no classification is perfect, that some misclassification is inevitable and that some cloud physical states are ambiguous for a given classification scheme. We used the 2B-CLDCLASS CloudSat product for our study but a comprehensive critical analysis of this product is beyond the scope of our paper.

3) Two different footprint sizes are considered, namely the AIRS/AMSU footprint of ~45 km and the AIRS footprint of ~15 km. It is a bit unclear throughout the paper which analysis is applied to which of the two footprints. In any case, the two different scales need to be addressed more consistently throughout the paper (in addition to the abstract). For example, Section 3.1 focuses on the AMSU footprint but not on the AIRS footprint. If I'm not mistaken, the AIRS cloud retrievals are performed on the AIRS ~15 km footprint, so the last section focuses only on that scale (If so, state this clearly in the paper.) Please expand the discussion in section 3.1 to the AIRS footprint as well and add a figure similar to figure 1 for the AIRS footprint. Furthermore, I assume Fig. 2 is showing statistics for the AMSU footprint. If this is true, please state that in the paper. I suggest adding figures similar to Fig 2. for the AIRS footprint, or at least discussing the (lack of) differences between AIRS and AMSU global distributions. In fig. 3, 4 and 6, indicate the size of AIRS and AMSU pixels.

We have added a second panel to Fig. 1 for the AIRS FOV scale and have labeled each panel clearly with the two scales depicted. Note that the number of scenes that explains >90% of the observed scenes is different between the two scales.

We describe in section 2 that the analysis is first conducted at the AMSU FOR spatial scale and that this scale is 45 km. We replaced the initial notation "AIRS/AMSU FOR" with "AMSU FOR" throughout the manuscript to avoid confusion between the two different resolutions and their corresponding scales. After discussing scene observations at this scale in section 3.1, we extend the analysis as a function of resolution in sections 3.2 and 3.3. After showing the cloud scene resolution dependence in Section 3, we now describe the dependence of AIRS cloud properties on the different types of cloud scene in section 4. We added a sentence at the beginning of section 4, to emphasize that the resolution used for cloud scenes determination, in this section is AIRS, i.e. about 15 km: *"In this section, the scenes are determined at the AIRS FOV resolution (approximately 15 km)."*

We agree with the referee and added a sentence in the caption of figure 2 to specify the resolution scale: *"These scenes were observed at the AMSU FOV resolution (~45 km)."* We note that the AIRS version of Figure 2 is nearly identical and are not included. We have added the following text to the Fig. 2 caption: *"Similar plots of the AIRS FOV resolution (~15 km) are nearly identical (not shown)."*

We think it is a good idea to indicate the AIRS and AMSU resolutions in figures 3 and 6. New versions of these figures with vertical lines indicating both resolutions are added to the revised manuscript.

4) If my interpretation is correct, one goal is to compare AIRS cloud retrievals for cases with a single cloud type in the footprint with clear sky mixed in versus without clear sky. However, a comparison is made between 1) cases with a single cloud type in the footprint and no clear sky and 2) cases with a single cloud type in the footprint either with and without clear sky. Thus, the case 2 set also contains the case 1 set in addition to the mixture of clear sky and a single cloud type. Therefore, the differences in properties listed in the paper are not representing the differences between cases without clear sky mixed in versus those with clear sky mixed in. From the information in the paper, we cannot deduce the relative number of cases per cloud type with versus without clear sky mixed in. If, for example, the number of cases without clear sky mixed in is much larger than those with clear sky, then the small differences shown between table 2 and 3 and figures 7 and 8 are surely expected. I suggest the following: 1) Include a table or figure showing the relative number of single type cases with and without clear sky mixed in. 2) For the single types listed in table 3 and figure 8 include only cases that also include clear sky. (Or add a table and figure showing this, leaving table 3 and fig. 8 as is.) To include a complete comparison between table 2 and 3 and figures 7 and 8, I suggest to also include the deep convection type (mixed with clear sky) in table 3 and figure 8. 3) Adjust the discussion of the differences accordingly.

These are very helpful comments by the reviewer and we have made some significant changes in the manuscript to make the categorization of cloud scenes, and the differences in the cloud phase and ice microphysical retrievals clearer.

First, we have added a new Table 2 and a new (brief) Section 4.1 that summarizes the new breakdown in cloud scenes: *“Table 2 summarizes five types of scenes at the 15-km AIRS FOV scale: (i) clear sky, (ii) cloudy sky with one cloud type, (iii) partly cloudy sky with one cloud type, (iv), cloudy sky with multiple cloud types, and (v) partly cloudy sky with multiple cloud types. The raw counts and the relative percentages for the two-year observing period are shown. The dominance of clear sky (30.7%) at 15-km is apparent and is consistent with an absence of thin cloud features in the 2B-CLDCLASS data set. Cloudy sky scenes with one cloud type (multiple cloud types) amount to 31.3% (10.2%) of all observed scenes, while partly cloudy sky scenes with one cloud type (multiple cloud types) amount to 23.5% (4.3%) of all observed scenes. A total of 41.5% of AIRS FOVs are completely cloudy while 27.8% are partly cloudy according to 2B-CLDCLASS. Below the differences in cloud thermodynamic phase detection and ice cloud property retrievals are quantified for the types of scenes summarized in Table 2.”*

Second, we have reworked the original Tables 2 and 3 into four different tables (3-6) that show the ice cloud property statistics for the scene classification (2), (3), (4) and (5) listed above. We have included clear sky in Table (3).

Third, we have similarly modified Figures 7 and 8 in the original manuscript to Figures 7-10 in the revised manuscript. They follow exactly the same scene classification divisions illustrated in Table 2.

We have added text in the manuscript that describe these new tables and figures and more clearly delineate the performance of cloud phase and ice cloud property retrievals by whether the cloud scene is completely versus partly cloudy, and whether the cloud scene contains a single cloud type or multiple cloud types. We refer the reviewer to the revised version of the manuscript to see every change made, but the most substantial for Section 4.2 and 4.3 follow:

The first paragraph of Section 4.2 that addresses cloudy sky with one cloud type now reads as: *“The occurrence frequency histogram of the sum of all thermodynamic phase tests is shown for cloudy sky with one cloud type in Fig. 7. Homogeneous cloud scenes serve as an ideal point of reference for establishing cloud phase sensitivity benchmarks. Overall, there is strong differentiation in the cloud thermodynamic phase among cloud scenes with single cloud types. Ice tests dominate Ci, Ns, Dc, and As, while liquid and undetermined tests dominate Ac, Sc, and Cu.”*

The fourth paragraph of Section 4.2 that addresses partly cloudy sky with one cloud type reads as: *“The occurrence frequencies of cloud phase for partly cloudy sky with one cloud type are shown in Fig. 8. The biggest change is the relative ordering of the ranks among cloud scene types between Figs. 7 and 8. Ac is now more common than As, Dc ranks higher than Ci as it is typically horizontally more extensive (Miller et al. 2014; Guillaume et al. 2018). There are more subtle changes in the cloud phase histograms that are consistent with partly cloudy sky. A weaker spectral signature for partly cloudy scenes results in slightly greater counts of unknown phase and also subtle shifts in liquid and ice phase tests in Fig. 8 compared to Fig. 7. In the Ac cloud scene histograms, there is a small but discernible increase in ice tests in Fig. 8 compared to Fig. 7. Horizontally heterogeneous Ac appears to have more frequent ice detection than horizontally homogeneous Ac.”*

There is a new fifth paragraph: *“The nine most frequent cloudy scenes with multiple cloud types are shown in Fig. 9. The (Ci,Sc) cloud scene ice phase histogram resembles a hybrid of histograms for Ci and Sc with undetermined phase the most frequent. (Ci,Sc) is a common cloud scene in the low latitudes as trade cumulus (Sc cloud type) and is frequently found under thin cirrus (Chang and Li, 2005). Furthermore, the spectral signatures of the two types of clouds frequently cancel, giving an undetermined phase result in the spectral tests used here (not shown). The (Ci,As) cloud scene shows a slight reduction in liquid detections and a slight increase in ice detections compared to As alone. While the As cloud scene in Fig. 7 is dominated by +2, the (Ci,As) cloud scene is dominated by +2 and +3. This suggests that a mixture of Ci and As together can trigger more ice tests in AIRS than As alone. Lastly, the mixed (As,Sc,nc) cloud scene in Fig. 9 closely resembles pure Ac and Sc cloud scenes in Fig. 7 since all three histograms are similar.”*

The sixth paragraph of Section 4.2 that addresses partly cloudy sky with multiple cloud types reads as: *“The nine most frequent partly cloudy scenes with multiple cloud types are shown in Fig. 10. As with the differences between Figs. 7 and 8, the biggest change is the relative ordering of the ranks among cloud scene types between Figs. 9 and 10. Furthermore, there are additional*

(yet subtle) changes in the phase test histograms for the cloud scenes that are common between Figs. 9 and 10.”

The first paragraph of Section 4.3 that addresses cloudy sky with single cloud type ice property retrievals reads: *“The mean ice cloud property retrievals are summarized in Table 3 for cloudy sky with one cloud type only for the ice only portions of the cloud phase histograms depicted in Fig. 7. Scenes identified as clear sky exhibit properties of a small population of thin cirrus detected by AIRS (Fig. 7) with mean values of $\tau_i=0.77$ and $r_{ei}=20.9 \mu\text{m}$ (Table 3). The AKs are notably lower and the relative error for τ_i is higher than other cloud scenes. The Sc cloud scene shows a small population of cirrus that go undetected in 2B-CLDCLASS (Fig. 7) and have mean values of $\tau_i=1.30$ and $r_{ei}=20.6 \mu\text{m}$ (Table 3). The AKs are also lowest in Table 3 for Sc relative to other cloud scenes with similarly high errors in τ_i and r_{ei} . Kahn et al. (2008; 2015) have shown that AIRS is very sensitive to thin cirrus, thus some ice clouds in CloudSat-identified Sc cloud scenes are expected. Because tenuous ice clouds have smaller values of τ_i and r_{ei} , the lower estimates of information content and larger error estimates are promising. These tenuous ice cloud retrievals are differentiated well from more robust retrievals within cloud scenes that are dominated by ice phase in the histograms (Fig. 7).”*

The sixth paragraph of Section 4.3 that addresses partly cloudy sky with single cloud type ice property retrievals reads: *“The mean ice cloud property retrievals are summarized in Table 4 for partly cloudy sky with one cloud type with cloud phase histograms depicted in Fig. 8. The biggest difference between Tables 3 and 4 is the relative frequency of occurrence with large differences between cloud scenes with or without clear sky. Another significant change is an overall reduction in AKs and magnitude of τ_i , with an increase in χ^2 in Table 4, consistent with partly cloudy scenes. The changes in r_{ei} AKs, magnitudes, and error estimates between Tables 3 and 4 are smaller than those for τ_i . Overall, the differences between Tables 3 and 4 are reassuring in that the AIRS retrieval is responding to partly cloudy scenes by reducing information content and the magnitude of τ_i , while χ^2 residuals are increasing somewhat.”*

The second to last paragraph summarizes the differences between Tables 5 and 6: *“The ice cloud property retrievals for cloud scenes that contain multiple cloud types are summarized in Table 5 for cloudy scenes and Table 6 for partly cloudy scenes. These tables list the nine most frequent cloud scene types as depicted in Figures 9 and 10. Seven of the nine cloud scenes are common between Tables 5 and 6. There is a general tendency for reductions of τ_i , increases in % relative error, and slight reductions in AKs in Table 6 for the seven common cloud scenes in Table 5. Changes in r_{ei} related variables are smaller than changes in τ_i related variables.”*

The last paragraph synthesizes Tables 3-6 and is: *“To summarize Tables 3-6, larger differences in ice cloud property retrievals are found between different cloud types than between cloudy and partly cloudy scenes. However, the differences between cloud scene types are the sharpest for the subset of cloudy scenes with one cloud type (Table 3). The AIRS cloud property retrievals are not greatly impacted by mixtures of cloud types within the AIRS footprint, and ice cloud property differences among cloud scenes are broadly consistent with the expected performance of infrared retrievals among these cloud types.”*

Minor comments:

At the start of Page 11 it is stated that “the AIRS footprint (~ 15 km) is commensurate with the dimension of a single cloud so that the most frequent observations involve the characteristics of one cloud.” What is meant by this statement? Many clouds, for example cumulus, have typical scales much smaller than that.

We agree that this statement could be confusing and we have removed the following from the revised manuscript: *“These differences between single and mixed cloud scenes can be explained by the fact that the AIRS footprint (~ 15 km) is commensurate with the dimension of a single cloud so that the most frequent observations involve the characteristics of one cloud. Cloud length statistics, such as median lengths and median absolute deviations calculated from the data shown in Fig. 6, corroborate this statement.”*

Table 1 and 2: Mean optical thicknesses are given here, but these are biased low because AIRS is not sensitive to any variation past an optical thickness of ~5. Please make this clear in the text. The true mean optical thickness of most types will be larger than the AIRS-retrieved means shown here.

We added the following sentence line 12 of page 5: *“The AIRS retrieval sample includes nearly all ice clouds with $\tau_i > 0.1$, while the maximum values of τ_i asymptote to values around 6-8 (e.g. Kahn et al., 2015).”*

We discovered an error in Table 1, which are now fixed, and now the values on the third row actually report the median absolute deviation.

I think the caption of figure 4 is missing the word “scenes” at end of first sentence. It is noted in the text that Fig. 4b and 4d are similar to Fig. 3, but it is not directly clear what the difference in the calculations are. Please explain.

We added the word “scene” at the end of the first sentence of the figure 4 caption. Figure 3 shows the number of scenes observed at different FOR scale whereas Figure 4b and 4d represents the number of scenes as a function of a cloud scale (not as a function of FOR scale).

I noted that the statistics listed in Table 3 are different than in Fig. 1, and this might be due to the different scales of AIRS vs AMSU. (See main comments.) However, for the ice properties to be retrieved, the phase needs to be identified as ice, so are these proportions for ice clouds only? Is that also explaining the difference in statistics compared to figure 1? Please explain in the text.

The referee is correct in that these are only for ice clouds (the blue bars in Figures 7-10). The statistics in Table 3 are obtained at the AIRS FOR whereas statistics in Figure 1 were obtained at the AMSU FOR. We note that the AIRS FOV is added as a second panel to Fig. 1.

Footprint-scale cloud type mixtures and their impacts on Atmospheric Infrared Sounder cloud property retrievals

Alexandre Guillaume¹, Brian H. Kahn¹, Eric J. Fetzer¹, Qing Yue¹, Gerald J. Manion¹, Brian D. Wilson¹, Hook Hua¹

5 ¹ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, 91109, USA

Correspondence to: Alexandre Guillaume (Alexandre.Guillaume@jpl.nasa.gov)

Abstract. A method is described to classify cloud mixtures of cloud top types, termed *cloud scenes*, using cloud type classification derived from the CloudSat radar (2B-CLDCLASS). The scale dependence of ~~the~~ cloud scenes is quantified. For spatial scales at 45 km (15 km), only 18 (10) out of 256 possible cloud scenes account for 90% of all observations and contain either one, two, or three cloud types. The number of possible cloud scenes is shown to depend on spatial scale with a maximum number of 210 out of 256 possible scenes at a scale of 105 km and fewer cloud scenes at smaller and larger scales. The cloud scenes are used to assess the characteristics of spatially collocated Atmospheric Infrared Sounder (AIRS) thermodynamic phase and ice cloud property retrievals within scenes of varying cloud type complexity. The likelihood of ice and liquid phase detection strongly depends on the CloudSat-identified cloud scene type collocated with the AIRS footprint. Cloud scenes primarily consisting of cirrus, nimbostratus, altostratus and deep convection are dominated by ice phase detection, while stratocumulus, cumulus, and altocumulus are dominated by liquid and undetermined phase detection. Ice cloud particle size and optical thickness are largest for cloud scenes containing deep convection and cumulus, and are smallest for cirrus. Cloud scenes with multiple cloud types have small reductions in information content and slightly higher residuals of observed and modelled radiance compared to cloud scenes with single cloud types. These results will help advance the development of temperature, specific humidity, and cloud property retrievals from hyperspectral infrared sounders that include cloud microphysics in forward radiative transfer models.

1 Introduction

There is increasing evidence of secular cloud trends at regional and global scales in both satellite observations (e.g., Norris et al. 2016) and climate general circulation model (GCM) simulations (e.g., Zelinka et al., 2013). The poleward migration of the extratropical storm tracks (Barnes and Polvani, 2013) is coupled with systematic changes in cloud thermodynamic phase partitioning in forced CO₂ experiments in climate GCMs (e.g., Mitchell et al. 1989; Ceppi et al. 2016). The spread in equilibrium climate sensitivity is also tightly coupled to the temporal evolution of phase partitioning in most climate GCMs (Tan et al. 2016). Obtaining reasonable observational estimates of the small-scale cloud phase partitioning at model sub-grid scales are critical for constraining the highly uncertain Wegener–Bergeron–Findeisen time scale parameter that is crucial for

Deleted: those

Deleted: studied

Deleted: near

Deleted: .

Deleted: 16

Deleted: 194

modelling mixed phase cloud and precipitation processes and (Tan and Storelvmo 2016). A new generation of probability distribution function-based parameterizations has shown promise for improving climate model simulations of cloud properties (e.g., Golaz et al. 2002) and would benefit from further exploitation of the information available in pixel-scale satellite observations. A rigorous assessment of the scale dependence of cloud types, and their mixtures, would also enhance climate GCM evaluation and parameterization development research efforts (Bony et al. 2006).

Kahn et al. (2018) showed that Atmospheric Infrared Sounder (AIRS) observations of ice cloud optical thickness (τ_i) and effective radius (r_{ei}) exhibit statistically significant temporal trends that are dependent on latitude and cloud type. Trends in Multi-angle Imaging SpectroRadiometer (MISR) observations of cloud texture have suggested that recent thinning of tropical cirrus has led to increased detection of trade cumulus (Zhao et al. 2016). Using high-spatial-resolution estimates of cloud thermodynamic phase obtained from the Hyperion instrument on Earth Observing 1 (EO-1), Thompson et al. (2018) showed that phase mixtures are highly variable at scales smaller than the AIRS footprint or typical GCM grid boxes. These studies (and many others) suggest that quantification of the scale dependence of cloud type mixtures could help explain satellite observations of cloud trends.

Statistical classification methods are commonly used to define weather states or cloud types (e.g., Rossow et al., 2005; Xu et al., 2005; Sassen and Wang, 2008; Wang et al. 2016). For instance, joint histograms of cloud top pressure and optical thickness from the International Satellite Cloud Climatology Project (ISCCP; Rossow and Schiffer 1999) are useful for relating cloud types to dynamical, radiation and precipitation variability, and in evaluating climate model simulations (e.g., Klein and Jakob, 1999; Jakob and Tselioudis 2003; Rossow et al. 2005; Tselioudis et al., 2013). Weather states are typically mixtures of conventional cloud types as shown by Rossow et al. (2005) and Oreopoulos et al. (2014). Partly inspired by this methodology, we introduce the concept of *cloud scenes* that are defined to be mixtures of CloudSat cloud types (2B-CLDCLASS; Sassen and Wang 2005) that vary with horizontal scale.

As cloud scenes will be matched to coincident A-train observations, we begin by defining cloud scenes with cloud types derived from CloudSat and observed within an AIRS/Advanced Microwave Sounding Unit (AMSU) (Chahine et al. 2006) field of regard (FOR) of roughly 45-km resolution. One AMSU FOR within an AMSU swath is spatially and temporally coincident with a ‘curtain’ of 94 GHz CloudSat radar profiles. The likelihood of observing clouds is resolution-dependent and is approximately 80–85% at the AIRS footprint scale of 15 km (Krijger et al. 2007; Kahn et al., 2008). The clouds in AMSU sounding FORs or AIRS footprints are more often broken or transparent, and less often uniform or opaque. Yue et al. (2013) showed that about 43% of the AMSU FORs are mixtures of CloudSat-identified cloud types, implying that roughly half of cloud soundings contain mixtures of cloud types.

Deleted: or reanalysis datasets

Deleted: AIRS/

Deleted: AIRS/

Deleted: AIRS/

Deleted: AIRS/

Our purpose in this work is to quantify the scale dependence of cloud type mixtures that are then used to understand the cloud complexity within AIRS cloud phase and ice cloud property data sets. The AIRS and CloudSat data and the collocation approach are described in Section 2. To quantify cloud type distributions and their dependence on horizontal scales, the cloud scenes are first characterized at the AMSU FOR resolution in Section 3.1, are extended to larger and smaller scales in Section 3.2, and key results of the scale dependence are placed into context in Section 3.3. The cloud scenes are used to partition AIRS cloud property retrievals into cloud types, specifically, cloud thermodynamic phase histograms in Section 4.2, and mean values of ice cloud microphysical parameters are described in Section 4.3. A discussion, summary, and suggestions for future investigation are found in Section 5.

Deleted: AIRS/

Deleted: 1

Deleted: 2

2 Data and Methodology

2.1 CloudSat and AIRS pixel-scale matching

The AIRS/AMSU/CloudSat matchup product described in Manipon et al. (2012) is used by Yue et al. (2013) and in this investigation. The matchup process uses a nearest neighbor approach to geolocate all CloudSat profiles within either an AMSU FOR at 45 km spatial resolution at nadir view, or a single AIRS footprint at 15 km spatial resolution at nadir view (Kahn et al. 2008). Approximately 45 to 50 (15–17) CloudSat profiles coincide with a single AMSU FOR (AIRS footprint), in a swath of width 30 FOR (90 footprints). The cloud scenes are first defined at the AMSU FOR scale and are then extended to other spatial scales. We use a two-year period of data extending from 01 July 2006 until 30 June 2008 which contains about 8 million AMSU FORs (or 24 million AIRS footprints).

Deleted: AIRS/

Deleted: AIRS/

Deleted: AIRS/

Deleted: AIRS/

2.2 CloudSat cloud types and their mixtures within the AIRS footprint

The CloudSat 2B-CLDCLASS product is used in this work and the algorithm is described in Sassen and Wang (2005, 2008). As summarized in Sassen and Wang (2008) and previous works, the algorithm uses methods developed from ground-based multiple remote sensors that have been tested against surface observer-based cloud typing reports. The cloud classification occurs in two steps. First, a clustering analysis is performed to group cloud profiles into cloud clusters. Secondly, classification methods are used to classify clouds into different cloud types. The decision trees guiding the classification are complex and are based on 23 variables derived from the clustering analysis of the first stage. Geometric quantities such as cloud base, top, and horizontal extents are present in decision trees (Sassen and Wang, 2005). Plan view and zonal average frequencies of 2B-CLDCLASS cloud types at its native resolution are reported in Sassen and Wang (2008).

There are eight CloudSat defined classes in the 2B-CLDCLASS files: cumulus (Cu), stratocumulus (Sc), stratus (St), altocumulus (Ac), altostratus (As), nimbostratus (Ns), cirrus (Ci), and deep convective (Dc) clouds, with a ninth classification of clear sky designated no cloud (nc). Since each AMSU FOR contains roughly 50 CloudSat profiles with 125 vertical levels each, there are $9^{50 \times 125}$ possible distinct cloud type combinations (although in practice there are fewer as many levels reside

Deleted: AIRS/

in the stratosphere) for each AMSU FOR. This number is too high to derive a classification that could be useful, i.e. where each cloud type combination could be populated with a significant number of samples for any climatological study. One particularly appealing way to reduce the dimensionality is to limit consideration of cloud type to cloud top only. This simplification is consistent with the capabilities of infrared sounders as the sampling of temperature and specific humidity is maximized in the atmosphere near and above cloud top, assuming the cloud is opaque and covers the entire sounder pixel area. There are 9^{50} possible cloud combinations defined in this manner including clear sky profiles. As a point of comparison, there are about 324,000 AIRS soundings per day or about 10^8 per year. Even when considering the 16 years of AIRS nominal operation, the number of cloud type combinations 9^{50} , or about 5×10^{47} , is many orders of magnitude greater than the number of AMSU FORs available, making impossible a statistically significant sampling of all combinations. This necessitates further assumptions to define a practical yet meaningful set of cloud scenes.

Deleted: AIRS/

Two additional simplifications are made here: variations in the count of each CloudSat cloud type is not considered, and the observation sequence of successive cloud types is disregarded. These two simplifications are applied to the AMSU FOV. We define a *cloud scene* as a list of the cloud types that are present within a given AMSU FOR. For example, the notation (Ci,Ac,Sc,Cu) is used to label a cloud scene that contains cirrus, altocumulus, stratocumulus, and cumulus clouds at cloud top in any frequency and in any sequence along the orbit segment. These simplifications greatly reduce the dimensionality of the classification problem and makes cloud scene identification tractable. We will show both partly cloudy and completely cloudy scenes in Section 4, so the clear sky (nc) type is both included and excluded in the analyses. Since each of the 8 cloud types is either present or absent, a cloud scene can also be represented by an 8 bit binary string. As a consequence, there are 2^8 possible cloud scenes that remain after taking into account the aforementioned simplifications. The number of possible cloud scenes is therefore reduced from $9^{50 \times 125}$ to a much more tractable 256. The limitations of this approach are: (i) a consideration of cloud tops only, (ii) the spatial sequence and frequency of individual cloud types are not considered, and (iii) equal weight is given to all cloud types within a cloud scene regardless of counts.

Deleted: AIRS/

Deleted: AIRS/

One advantage of using a classification to define cloud mixtures rather than an unsupervised learning technique, such as clustering, is that the size of the set of possible cloud mixtures is well defined and finite (here it is 256). A related and important advantage of classification is that one can use this set of classes (cloud scenes) with any parameter matched to any given scene. Here, the spatial scale dependence of those cloud scenes is described in section 3.2.

An alternative approach may consider the vertical layering of cloud types or cloud features, some form of weighting based on counts of each cloud type, or possibly the sequence of cloud types, which may result in different radiance measurements observed by the AIRS instrument (the radiance emitted within an AIRS footprint is non-uniform and channel-dependent, as described in Schreier et al., 2010). However, the simplified approach outlined above is broadly consistent with the sensitivity and sampling characteristics of nadir-viewing passive infrared sounders. Therefore, we consider the approach outlined above

Deleted: , yet simplifies and makes tractable the complexity of cloud scenes.

to be an appropriate compromise that retains the diversity of cloud scenes and makes the necessary data processing tractable by reducing the dimensionality for ease of interpretation.

5 Lastly, the results of Kahn et al. (2018) suggest larger ice cloud particle sizes occur at convective cloud tops compared to thin cirrus at the same cloud top temperature. Given the key assumption of cloud typing only at cloud top, the 2B-CLDCLASS product is better suited for identifying convective clouds in AIRS apart from stratiform clouds, the latter of which are dominant in 2B-CLDCLASS-LIDAR. If 2B-CLDCLASS-LIDAR was used in place of 2B-CLDCLASS, the statistics would be weighted towards the detection of vast areas of cirrus in thin layers above and in proximity to convective clouds. The Ci classification dominates in 2B-CLDCLASS-LIDAR at cloud top and will blur the signals of underlying cumulus and deep convective cloud types that are capped by thin cirrus.

2.3 AIRS thermodynamic phase and ice cloud properties

15 The AIRS version 6 cloud thermodynamic phase and ice cloud properties (Kahn et al., 2014) are geolocated to the CloudSat ground track and are binned by cloud scene. The cloud thermodynamic phase algorithm includes two liquid tests and four ice tests of brightness temperature (T_b) thresholds and T_b differences (ΔT_b) in the mid-infrared atmospheric windows. The T_b and ΔT_b thresholds are designed to exploit spectral differences in liquid and ice water indices of refraction. The two liquid and four ice tests are each assigned a value of -1 and +1, respectively, and a summed value that ranges from -2 to +4 is reported. Summed values -2 or -1 indicate liquid clouds, 0 is undetermined, and $\geq +1$ indicate ice, with the highest values indicating deeper, convective ice clouds (Naud and Kahn, 2015). Ice is detected in 26.5% of AIRS footprints by Kahn et al. (2014) and pixel-scale comparisons with estimates of ice from the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) lidar (Jin and Nasiri, 2014) are in agreement with AIRS more than 90% of the time. The success rate, however, is smaller for liquid cloud detection with AIRS using CALIPSO as a benchmark because of the small thermal contrast between low-lying liquid clouds and the surface. Despite this limitation in sensitivity, AIRS rarely misidentifies liquid clouds as ice (Jin and Nasiri, 2014). Furthermore, many liquid clouds are classified as undetermined phase. Low latitude shallow trade cumulus clouds generally fall within this category (Kahn et al., 2017).

25 Kahn et al. (2014) describe a retrieval algorithm that is based on optimal estimation (OE) theory and derives ice cloud optical thickness (τ_i) and effective radius ($r_{e,i}$) for AIRS footprints containing ice. The AIRS retrieval sample includes nearly all ice clouds with $\tau_i > 0.1$, while the maximum values of τ_i asymptote to values around 6-8 (e.g. Kahn et al., 2015). Scalar averaging kernels (AKs), χ^2 residuals from observed and simulated radiance fits, and values of relative error are also reported (Kahn et al., 2014). Values of AKs closer to 1.0 suggest higher information content while larger relative error estimates and values of χ^2 indicate increased uncertainty in retrieved parameters. Only the relative magnitude of error estimates should be considered since temperature, specific humidity, surface temperature, surface emissivity, and ice crystal habit and size distribution uncertainties are not included in the error covariance matrices of the AIRS version 6 algorithm (cf. Kahn et al., 2014). We

Deleted: In this study, the

Deleted: ice

Moved (insertion) [1]

Formatted: Font color: Text 1

focus on the differences in error estimates and χ^2 among cloud scenes and determine which cloud scenes contain higher or lower certainty in their ice cloud properties relative to other scenes.

3 Classification and characteristics of cloud scenes

3.1 Cloud scenes with ~45 km resolution

5 A cloud scene is assigned to every AMSU FOR along the CloudSat viewing path using the methodology outlined in Section
2. Using the two years of data, a total of 194 out of 256 possible cloud scenes are observed but only 18 of the cloud scenes
account for 90% of all observed scenes (Fig. 1a). The four most common scenes contain one cloud type with or without clear
sky, and the most common mixed cloud scene (Ac, Sc) is ranked as the fifth most common scene overall. Intuitively, the more
diverse a scene, the less frequently it should be observed. The scene that ranked last (18th) in Fig. 1a is (Ci, Ac, Sc). The least
10 frequently observed cloud scene with a ranking of 194 contains six cloud types (Ci, As, Ac, St, Cu, Ns) and was observed only
once in two years. Of the 256 possible types of cloud scenes, the number of unobserved cloud scenes is 62, of which 61 include
St. The unobserved cloud scenes include the only possible cloud scene with eight cloud types together, and the seven possible
cloud scenes with seven cloud types together.

15 The unobserved scenes in the two-year period contain a median of five different cloud types. This is consistent with the
improbability of particular cloud types occurring in rapid succession over a few tens of kilometers. The only unobserved cloud
scene that does not contain St is (Sc,Cu,Ns,Dc) and is consistent with the conclusion by Sassen and Wang (2008) that Dc
(1.8%) and Cu (1.7%) clouds are the least frequent of the cloud types. While Dc and Ns are typically associated with different
climatological regimes (tropical convection versus extratropical storm tracks), occasionally, Dc is embedded within
20 extratropical cyclones and Ns is classified in stratiform regions of mesoscale convective systems (MCSs). Given the prevalence
of Sc and Cu in Figure 1a, it is somewhat surprising that the combination (Sc,Cu,Ns,Dc) is not observed.

The relative ranking of cloud scenes within the AIRS FOV along the CloudSat track is shown in Fig. 1b for the same sets of
matched pixels. A total of 10 cloud scenes account for 90% of all observed cloud scenes (Fig. 1b). This shows that fewer cloud
25 scenes are found at the smaller AIRS FOV compared to the AMSU FOR.

Figure 2 depicts the geographic distribution of Sc at the AMSU FOR scale, the most observed scene after clear sky (nc), and
(Ac,Sc) is the most observed mixed cloud scene. The Sc classification is consistent with the prevalence of stratocumulus clouds
in subtropical subsidence regions and trade cumulus in the tropics and subtropics (e.g., Yue et al., 2011). The (Ac,Sc) cloud
30 scene is identified most frequently in the extratropical storm tracks and the transition from shallow cumulus to deep tropical
convection.

Deleted: AIRS/

Deleted: 1

Deleted: within the 90th percentile

Deleted:

Deleted: 1

Deleted: and clear,

Deleted: two

Deleted: scenes,

Deleted: The pattern of clear sky scenes is generally similar to the clear sky global distribution derived by Eguchi and Yokota (2008) from CALIPSO and Moderate Resolution Imaging Spectroradiometer (MODIS) data.

Deleted: low

Deleted: subtropical

3.2 Cloud scenes at 1 to 1000 km scales

In Section 3.1, the relative frequencies of cloud scenes were derived for exact collocated matches of AIRS and AMSU observations to the CloudSat ground track. As the CloudSat ground track can oscillate across several AIRS FOVs over a scan line within a given orbit, the numbers of coincident CloudSat profiles matching to AIRS and AMSU will vary. Below, cloud scenes are derived independently of the specific AIRS and AMSU collocation geometry.

To investigate the scale dependence of the number of cloud scenes, the approach described in Section 2 is modified for a range of horizontal extents between 1.1 and 1000 km. The number of observed cloud scenes calculated at each horizontal scale is shown in Fig. 3a for 10 to 1000 km. At the finest scale of 1.1 km, only eight possible observed cloud scenes or clear sky are expected. When the scale increases, as expected, the number of cloud scenes quickly increases with a total of 143 cloud scenes observed at a scale of 11 km. As horizontal scale is further increased, the probability of observing cloud scenes with only one or two cloud types is reduced. After a maximum number of cloud scenes is obtained at 105 km, the number of cloud scenes will decrease with increasing scale (e.g., 163 cloud scenes at 990 km) until a limiting case is reached at the largest scale with only one cloud scene with all observed cloud types. The number of cloud scenes observed at least once at the AMSU FOR horizontal scale (indicated by the red vertical line on Fig. 3a) is approximately 190.

The 90th percentile calculated at all horizontal scales is shown in Fig. 3b. The 90th percentile of the maximum number of cloud scenes is 33 between 303 to 440 km in horizontal scale. The number at the nominal 45-km AMSU footprint scale is 16 cloud scenes, while the average number at the AIRS footprint is 9 cloud scenes. (Note that these are slightly smaller than values of 18 and 10 using the exact AMSU and AIRS geometry, respectively, in Section 3.1.) While these results show that fewer cloud type mixtures are observed at a decreasing length of 45 km to 15 km, a variety of cloud type mixtures is still encountered. While infrared sounding at 15 km resolution does not eliminate the cloud scene complexity encountered for combined infrared and microwave sounding at 45 km, the vast majority of 15 km footprints contain a smaller subset of possible cloud mixtures. In Section 4, we will determine whether individual cloud types or cloud type mixtures have meaningful impacts on AIRS cloud property retrievals. (Impacts on temperature and specific humidity soundings are beyond the scope of this investigation.)

The reasons for the maximum number of observed cloud scenes (210) at a particular horizontal scale (105 km) are not immediately clear. The scale preference depends on the physical characteristics of cloud regimes and the degree to which cloud types are “mixed together” by region and furthermore depend on cloud length distributions (Guillaume et al., 2018). A simple model is described below that is able to approximate the results of Fig. 3 and offers some insight for the observed maximum frequency of cloud scenes and the spatial scale at which it occurs.

Formatted: Normal

Deleted: AIRS/

Deleted: at

Deleted: lengths

Deleted: large

Deleted: initial study

Formatted: Font: +Body (Times New Roman), English (UK)

Deleted: remains unexplained

Formatted: Font: +Body (Times New Roman), English (UK)

Formatted: Font: +Body (Times New Roman), English (UK)

Formatted: Font: +Body (Times New Roman), English (UK)

Deleted: 3

Formatted: Font: +Body (Times New Roman), English (UK)

3.3 Generalizing to all scales

Deleted : why a maximum number of cloud scenes at 105 km?

The goal of this section is to derive cloud scene scale statistics that are independent of any regular grid resolution, and explore whether these statistics can explain some features of the number of scenes as a function of scale observed in the previous section. In particular, we explore whether these statistics can explain the maximum observed around 105 km. There is however an inherent difficulty in defining the boundaries that delimit any given cloud scene in absence of a predefined horizontal extent. It is possible that within a given cloud scene there exists several scenes with the same cloud types but differing lengths making the scene identification ambiguous. To circumvent this problem, we define a cloud scene and its maximum length as follows:

- (1) We search for a cloud scene containing a pre-defined mixture of cloud types. The spatial extent of this scene is delimited by cloud types (or clear sky) on both ends that do not belong to the mixture.
- (2) The maximum length of a cloud scene is the sum of all the horizontal lengths of all the cloud types in the cloud scene.

For example, imagine that we will calculate the maximum length of the specific cloud scene (Ac,Sc). We then identify a location in the CloudSat data record with the following illustrative succession of cloud types: (Ci,Ac,Sc,Ac,Sc,Ac,Ns), with the number of CloudSat profiles associated with each cloud type of 10, 3, 6, 5, 7, 12, and 15, respectively. The Ci and Ns obviously do not belong to the (Ac,Sc) cloud scene and therefore delimit the scene as defined in (1) above. The maximum length of the cloud scene (Ac,Sc) will be the sum of the number of CloudSat profiles for (Ac,Sc,Ac,Sc,Ac), which is $3+6+5+7+12=33$ CloudSat profiles in total. Below, we define a minimum cloud length that is unequivocal:

- (3) If within a given cloud scene, there exist several cloud scenes with the same cloud types but smaller lengths than the maximum length, the minimum length of a cloud scene is defined as the smallest length of all those lengths.

In the example above, there are four possible sequences (in bold font) that could be the minimum length: (**Ac,Sc**,Ac,Sc,Ac), (**Ac,Sc,Ac**,Sc,Ac), (Ac,Sc,**Ac,Sc**,Ac) or (Ac,Sc,Ac,**Sc,Ac**). The corresponding lengths are $3+6=9$, $6+5=11$, $5+7=12$, and $7+12=19$, respectively. In this example, the minimum length would therefore be 9 CloudSat profiles. (The minimum and maximum may be equal for a particular mixed cloud scene.)

Before steps (1) and (2) are used to quantify the maximum and minimum lengths for each of the 247 mixed scenes (256 minus the 8 single cloud scenes and clear sky), the locations of each cloud scene must first be identified in the two-year data record. Starting at the first CloudSat profile, the presence of each of the 247 mixed cloud scenes is determined using (1). For each occurrence of each mixed cloud scene, (2) and (3) are then applied to determine the maximum and minimum lengths for each

individual cloud scene. After processing the maximum and minimum lengths for every mixed cloud scene, simple statistics are calculated.

A total of 200 out of 247 possible mixed scenes were identified. The minimum and maximum length occurrence frequencies of five cloud scenes (Ac,Sc), (As,Sc,Cu), (Ci,As,Cu,DC), (As,Ac,Ns,DC) and (Ci,As,Ac,St,Sc) selected randomly from the 200 present in the two-year record, are shown in Fig. 4a and 4c, respectively. Recall that the maximum length is defined from (2), while the minimum length is defined from (3), with an illustrative example shown for (Ac,Sc). From top to bottom, their respective ranks are 1, 26, 51, 76 and 101. It is striking that each frequency histogram in Fig. 4a and 4c is not monotonic and displays a frequency maximum between 100 and 1000 km. Consequently, the sum of all (200) observed mixed scenes across length scales will result in a curve with a maximum and these are shown in Fig. 4b and 4d. Both curves are very similar to Fig. 3a and have maxima for about 180 observed scenes at 77 km and 174 km, respectively. Using the methodology outlined in (1) to (3) to estimate numbers of cloud scenes, the scale dependence of the number of observed scenes shows that the maximum will occur somewhere between 77 and 174 km.

In order to shed additional light on why a maximum in the occurrence frequency of each cloud scene histogram is obtained, histograms of cloud length frequency of single cloud types (defined at cloud top) are calculated. An example CloudSat orbital segment is shown in Fig. 5. The distribution of lengths for each cloud type for the two-year period is then shown in Fig. 6 with corresponding median and median absolute deviation (or m.a.d.) values reported in Table 1. Note that these values are similar to but not exactly the same as those calculated in Guillaume et al. (2018), for which cloud length was derived from a 2-D curtain of cloud features. The main characteristic shared by all cloud types in Fig. 6 is that their distributions are heavily skewed towards small lengths.

The length of a mixed scene is the sum of the lengths of each cloud type within it. There are two aspects that will influence the number of scenes observed at a given length L. First, there are several combinations of different lengths that will sum to L and those lengths will be smaller than L (abscissa of Fig. 6). Second, the likelihood of observing a given scene depends on the frequency of occurrence of each cloud type (ordinate axis of Fig. 6). These two effects have opposite behaviors as a function of L: single cloud frequency decreases with L, whereas the number of cloud length combinations that sum up to L increases with length scale.

To illustrate the effects of these opposing behaviors, we consider the scene (As,Sc,Cu) length distribution. Since the minimum length of all cloud distributions in Fig. 6 is one CloudSat profile, there is only one possible cloud length combination (1+1+1) that will sum to the minimum possible length of the scene (As,Sc,Cu). This is indeed the value observed on the far left of each red-orange curve in Fig. 4a and 4c. Next, consider a measurement consisting of 4 CloudSat profiles with this particular scene, with three possible length combinations: (1+1+2), (1+2+1) or (2+1+1). The frequency of each individual cloud type, As, Sc

Deleted: AICu

Deleted: AISt

Deleted: ci,AISt

Deleted: DC), (AISt,AICu

Deleted: DC

Deleted: ci,AISt,AICu

Deleted: 4a and 4c, respectively.

Deleted: 3

Deleted: This demonstrates that

Deleted: is the result of the scale dependence of frequency distributions of each mixed cloud scene, and

Deleted: of the number of observed scenes

Deleted: , depending on the length of the cloud scene.

Deleted: mean

Deleted: AISt

Deleted: AISt

Deleted: AISt

or Cu, is smaller at the scale of 4 CloudSat profiles than it is at a length of 3 CloudSat profiles in Fig. 6. However, there are more (As,Sc,Cu) scenes at length 4 than at 3 in Fig. 4a and 4c. This indicates that the *increase of possible combinations is more important* than the individual cloud frequency decrease for larger scales. This reasoning applies for increasing lengths until the *decreasing frequency of individual cloud types between two consecutive lengths is more important*. There are very few single As, Sc or Cu clouds observed at large lengths (far right scale of Fig. 6) resulting in a very small number of observed (As,Sc,Cu) scenes in Fig. 4a and 4c, despite the large number of length combination possibilities that may contribute.

Deleted: AISt

Formatted: Font: Italic

Formatted: Font: Italic

Deleted: AISt

Deleted: AISt

4 Cloud scene dependence of AIRS cloud properties

We will now establish differences in the AIRS thermodynamic phase and ice cloud properties in the presence of complex and simple cloud types using coincident cloud scenes. In this section, the scenes are determined at the AIRS FOV resolution (approximately 15 km). We briefly summarize general categories of cloud scene statistics in Section 4.1. The AIRS cloud thermodynamic phase tests are discussed separately for single and mixed cloud scenes in Section 4.2. The AIRS ice cloud τ_i and r_{ei} , error estimates, averaging kernels (i.e., information content), and χ^2 residual fits between observed and simulated radiances are shown in Section 4.3.

Deleted: To summarize, we showed that the scale dependence of the number of mixed scenes is the result of two opposing contributions. First, the single cloud type cloud lengths distributions are heavily skewed toward small scales (Fig. 6), qualitatively explaining the scale dependence of a mixed scenes cloud lengths distributions (Fig 4a and 4c). Secondly, the scale dependence of all observed mixed scenes cloud lengths distributions (Fig 4a and 4c) explains the scale dependence of the number of mixed scenes (Fig 4b and 4d).¶

Deleted: 1

Deleted: 2

Formatted: Default Paragraph Font

4.1 Types of cloud scenes

Table 2 summarizes five types of scenes at the 15-km AIRS FOV scale: (i) clear sky, (ii) cloudy sky with one cloud type, (iii) partly cloudy sky with one cloud type, (iv), cloudy sky with multiple cloud types, and (v) partly cloudy sky with multiple cloud types. The raw counts and the relative percentages for the two-year observing period are shown. The dominance of clear sky (30.7%) at 15-km is apparent and is consistent with an absence of thin cloud features in the 2B-CLDCLASS data set. Cloudy sky scenes with one cloud type (multiple cloud types) amount to 31.3% (10.2%) of all observed scenes, while partly cloudy sky scenes with one cloud type (multiple cloud types) amount to 23.5% (4.3%) of all observed scenes. A total of 41.5% of AIRS FOVs are completely cloudy while 27.8% are partly cloudy according to 2B-CLDCLASS. Below the differences in cloud thermodynamic phase detection and ice cloud property retrievals are quantified for the types of scenes summarized in Table 2.

4.2 Cloud thermodynamic phase

The occurrence frequency histogram of the sum of all thermodynamic phase tests is shown for cloudy sky with one cloud type in Fig. 7. Homogeneous cloud scenes serve as an ideal point of reference for establishing cloud phase sensitivity benchmarks. Overall, there is strong differentiation in the cloud thermodynamic phase among cloud scenes with single cloud types. Ice tests dominate Ci, Ns, Dc, and As, while liquid and undetermined tests dominate Ac, Sc, and Cu.

Deleted: frequencies

Deleted: ranges between -2 and +4 for cloud scenes that contain only single cloud types (no clear sky and no mixtures of cloud types) are shown

Deleted: 7. (While these clouds account for a small percentage of the total number of AIRS FOVs, homogeneous

Deleted: .)

The ice tests dominate the Ci cloud scenes and reaffirm the sensitivity of AIRS to ice clouds. CloudSat-classified clear scenes contain occasional occurrences of AIRS-detected thin cirrus (+1 and +2), consistent with either thin cirrus that is undetected by the CloudSat radar, or thicker cirrus within the AIRS footprint but to the side of the CloudSat ground track (e.g., Kahn et al., 2008). A few occurrences of -1 and -2 may also arise from spatial mismatches between AIRS and CloudSat scenes, or from stratus below 1 km in altitude that is undetected by CloudSat. In the Sc cloud scenes, trade cumulus clouds dominate as previously shown by Yue et al. (2011) and Kahn et al. (2017). A larger proportion of liquid tests, and a smaller proportion of ice tests, is observed in the Sc cloud scenes compared to clear sky, but undetermined phase is dominant in both scene types. The Cu and Sc cloud scene histograms are generally similar with more undetermined cases for Cu, but with a slight reduction of liquid and slight increase in ice observed for Cu compared to Sc.

The As cloud scene histogram in Fig. 7 is overwhelmingly dominated by ice. The undetermined cases in part may result from supercooled liquid or mixed-phase clouds that potentially could be distinguished with an improved phase algorithm that factors in the spectral mid-infrared signature of supercooled liquid (e.g., Rowe et al., 2013). The Ac and As cloud scene histograms are very different from each other, with a majority of undetermined and liquid for Ac and a majority of ice for As, consistent with aircraft observations (Mazin, 2006). The preponderance of undetermined phase for Ac may indicate frequent supercooled liquid cloud tops (Zhang et al., 2010). Ham et al. (2013) showed that Ac are typically 2-3 km lower in altitude than As and this probably explains some of the difference in liquid and ice phase, as lower clouds are usually warmer. The Ns cloud scene histogram is dominated by ice detection with occasional liquid and undetermined cloud tops. The Ns cloud scene also has significant height overlap with Ac and As, with most tops for all three types typically located below 9 km. Ice tests dominate in the Dc cloud scene histogram although a very small proportion of -1, 0, and +1 occur. Inspection of AIRS granules (not shown) demonstrate that the spectral signatures used in thermal infrared phase tests break down in the presence of overshooting convection and other ice clouds within a few Kelvins of the tropopause (e.g., Kahn et al., 2018).

The occurrence frequencies of cloud phase for partly cloudy sky with one cloud type are shown in Fig. 8. The biggest change is the relative ordering of the ranks among cloud scene types between Figs. 7 and 8. Ac is now more common than As, and Dc ranks higher than Ci as it is typically horizontally more extensive (Miller et al. 2014; Guillaume et al. 2018). There are more subtle changes in the cloud phase histograms that are consistent with partly cloudy sky. A weaker spectral signature for partly cloudy scenes results in slightly greater counts of unknown phase and also subtle shifts in liquid and ice phase tests in Fig. 8 compared to Fig. 7. In the Ac cloud scene histograms, there is a small but discernible increase in ice tests in Fig. 8 compared to Fig. 7. Horizontally heterogeneous Ac appears to have more frequent ice detection than horizontally homogeneous Ac.

The nine most frequent cloudy scenes with multiple cloud types are shown in Fig. 9. The (Ci,Sc) cloud scene ice phase histogram resembles a hybrid of histograms for Ci and Sc with undetermined phase the most frequent. (Ci,Sc) is a common cloud scene in the low latitudes as trade cumulus (Sc cloud type) and is frequently found under thin cirrus (Chang and Li,

Deleted: nine most frequent

Deleted: scenes in Fig. 1

Deleted: While a few

Deleted: these

Deleted: scenes are marked as single cloud

Moved (insertion) [2]

Deleted: , we include clear sky here, unlike those shown in Fig. 7. In the case of Cu, Ns,

Deleted: Sc

Deleted: scenes, the

Deleted: are very similar when comparing Figs.

Moved up [2]: 7 and 8.

Deleted: In the Ci cloud scene histograms, there is a small but notable reduction in the frequency

Deleted: ice tests

Deleted: an increase

Deleted: undetermined

Deleted: This is consistent with the expectation that more heterogeneous Ci cloud scenes with mixtures of clear sky exert a weaker spectral infrared phase signature.

Deleted: therefore has a

Deleted: robust

Deleted: signature

2005). Furthermore, the spectral signatures of the two types of clouds frequently cancel, giving an undetermined phase result in the spectral tests used here (not shown). The (Ci,As) cloud scene shows a slight reduction in liquid detections and a slight increase in ice detections compared to As alone. While the As cloud scene in Fig. 7 is dominated by +2, the (Ci,As) cloud scene is dominated by +2 and +3. This suggests that a mixture of Ci and As together can trigger more ice tests in AIRS than

Deleted: putative

As alone.

Deleted: Lastly, the mixed (As,Sc,nc) cloud scene in Fig. 8 closely resembles pure Ac and Sc cloud scenes in Fig. 7 since all three histograms are similar.

The nine most frequent partly cloudy scenes with multiple cloud types are shown in Fig. 10. As with the differences between Figs. 7 and 8, the biggest change is the relative ordering of the ranks among cloud scene types between Figs. 9 and 10. Furthermore, there are additional (yet subtle) changes in the phase test histograms for the cloud scenes that are common between Figs. 9 and 10.

Deleted: There is strong differentiation in the cloud thermodynamic phase among cloud

Deleted: single

Deleted: . Ice tests dominate Ci, Ns, Dc,

Deleted: As, while liquid

Deleted: undetermined tests dominate Ac, Sc,

Deleted: Cu.

In most mixed cloud scenes in both Figs. 9 and 10, the characteristics of the histograms are similar either to single types or have combined characteristics of the multiple cloud types contained within the cloud scene. These results are encouraging and reaffirm the capabilities of thermal infrared cloud phase determination (Jin and Nasiri 2014) and exhibit consistency with cloud types from the CloudSat radar. We note, however, that the AIRS phase determination has some ambiguity in overlapping ice and liquid cloud layers as previously shown by Jin and Nasiri (2014).

Deleted: These differences between single and mixed cloud scenes can be explained by the fact that the AIRS footprint (~ 15 km) is commensurate with the dimension of a single cloud so that the most frequent observations involve the characteristics of one cloud. Cloud lengths statistics, such as median lengths and median absolute deviations calculated from the data shown in Fig. 6, corroborate this statement.

Deleted: Furthermore, the AIRS cloud phase determination is not adversely affected by complex cloud top morphology (but without considering vertical structure).

4.3 Ice cloud properties

The mean ice cloud property retrievals are summarized in Table 3 for cloudy sky with one cloud type only for the ice only portions of the cloud phase histograms depicted in Fig. 7. Scenes identified as clear sky exhibit properties of a small population of thin cirrus detected by AIRS (Fig. 7) with mean values of $\tau_i=0.77$ and $r_{ei}=20.9 \mu\text{m}$ (Table 3). The AKs are notably lower and the relative error for τ_i is higher than other cloud scenes. The Sc cloud scene shows a small population of cirrus that go undetected in 2B-CLDCLASS (Fig. 7) and have mean values of $\tau_i=1.30$ and $r_{ei}=20.6 \mu\text{m}$ (Table 3). The AKs are also lowest in Table 3 for Sc relative to other cloud scenes with similarly high errors in τ_i and r_{ei} . Kahn et al. (2008; 2015) have shown that AIRS is very sensitive to thin cirrus, thus some ice clouds in CloudSat-identified Sc cloud scenes are expected. Because tenuous ice clouds have smaller values of τ_i and r_{ei} , the lower estimates of information content and larger error estimates are promising. These tenuous ice cloud retrievals are differentiated well from more robust retrievals within cloud scenes that are dominated by ice phase in the histograms (Fig. 7).

Deleted: 2

Deleted: 2

Deleted: cloud scenes composed of single

Deleted: and are

Moved (insertion) [3]

Deleted: 36

Deleted: 3

Deleted: 2

Deleted: 2

Deleted: and Table 2

The Ci cloud scene has mean values of $\tau_i=1.91$ and $r_{ei}=25.4 \mu\text{m}$, an $AK=1.0$, the highest of any scene type, and lower errors compared to other types in Table 3. The As cloud scene has a larger mean of $\tau_i=2.42$ compared to the Ac cloud scene with a mean of $\tau_i=1.65$ (Table 3). Interestingly, the mean value and error estimate of r_{ei} is lower for Ac than As, exhibiting differentiation between these two mid-level cloud types. However, a much smaller proportion of Ac are ice compared to As (Fig. 7).

Deleted: 94

Deleted: 7

Deleted: .99

Deleted: 2

Deleted: 55, a slightly smaller error, and slightly larger AK than

Deleted: 2

The Ns cloud scene in Table 3 contains larger mean values of τ_i than Ci and Ac cloud scenes, but these values are similar to those for As; however, the mean values are lower than Cu and Dc cloud scenes (Table 3). A lower value of τ_i is characteristic of diffuse cloud tops where the infrared emission may originate several km deep within the cloud (e.g., see Kahn et al., 2008; Holz et al., 2008). The reduced AK=0.88 for the Ns cloud scenes illustrates that a diffuse cloud top is more problematic for ice cloud retrievals. Cu cloud scenes with ice cloud tops occur a small amount of the time (Fig. 7); furthermore, Cu is infrequent in CloudSat classification (1.7% of all clouds). The horizontal extent of Cu is also much smaller than Dc (see Table 1). Interestingly, τ_i is larger for Cu cloud scenes than for all categories except Dc cloud scenes (Table 2).

- Deleted: 2
- Deleted: and r_{ei}
- Deleted: 2
- Deleted: 89
- Deleted: shows

The mean Cu value of $r_{ei}=26.5 \mu\text{m}$ is larger than most cloud scenes. This is consistent with larger ice particles observed at the tops of convection instead of small ice particles in thin cirrus at the same cloud top temperature (e.g., Yuan and Li 2010; Protat et al. 2011; van Diedenhoven et al. 2014; Kahn et al., 2018). These Cu cases are likely transient cumulus congestus at altitudes cold enough for cloud top glaciation. The Dc cloud scene has the largest mean $\tau_i=5.47$ of all cloud scenes with a very dense cloud top that saturates the infrared emission signal in contrast to Ns. The values of r_{ei} for Dc are similar to Cu (Table 3) with a slight reduction in the r_{ei} AK=0.96 and τ_i AK=0.98 relative to Ci cloud scenes (Table 3). This is consistent with reduced sensitivity for high values of τ_i (e.g., Huang et al., 2004).

- Deleted: 27.9
- Deleted: the largest for all
- Deleted: 54
- Deleted: nearly as large as
- Deleted: 2
- Deleted: 2
- Deleted: single type
- Deleted: scenes

The relative variations between the ice cloud retrieval properties for cloudy sky with one cloud type in Table 3 are consistent with expectations of infrared sensitivity. CloudSat-observed Ci cloud scenes have smaller error estimates and higher information content in comparison to Sc, consistent with Sc scenes containing tenuous cirrus that goes undetected by 2B-CLDCLASS. Larger τ_i and r_{ei} are observed at the tops of convective ice clouds such as Dc and Cu compared to stratiform clouds such as As and Ci. Differences in ice cloud properties between Ac and As cloud scenes are consistent with observed differences in scene heterogeneity and cloud top height.

- Deleted: 2
- Deleted: CloudSat
- Deleted: clear
- Deleted: , mixed
- Deleted: , and cloud
- Deleted: with one cloud type mixed with clear sky, are summarized in Table 3.

The mean ice cloud property retrievals are summarized in Table 4 for partly cloudy sky with one cloud type with cloud phase histograms depicted in Fig. 8. The biggest difference between Tables 3 and 4 is the relative frequency of occurrence with large differences between cloud scenes with or without clear sky. Another significant change is an overall reduction in AKs and magnitude of τ_i , with an increase in γ^2 in Table 4, consistent with partly cloudy scenes. The changes in r_{ei} , AKs, magnitudes, and error estimates between Tables 3 and 4 are smaller than those for τ_i . Overall, the differences between Tables 3 and 4 are reassuring in that the AIRS retrieval is responding to partly cloudy scenes by reducing information content and the magnitude of τ_i , while γ^2 residuals are increasing somewhat.

- Moved up [3]: Scenes identified as clear sky exhibit properties of a small population of thin cirrus detected by AIRS (Fig. 7) with mean values of $\tau_i=0$.
- Deleted: 78 and
- Deleted: =20.9 μm (Table
- Deleted:). The AKs
- Deleted: notably lower
- Deleted: other cloud
- Deleted: and the relative error for τ_i is higher. The corresponding cloud phase histograms are shown in Fig. 8. The (Sc,Ne) cloud scene in Table 3 is similar to the Sc cloud scene in Table 2 except the (Sc,Ne) τ_i is slightly reduced in Table 3. This is consistent with some thin cirrus in the (Sc,Ne) cloud scene, expected since thin cirrus and shallow Cu frequently overlap in the subtropical and tropical trade cumulus regimes. The (As,Ne)
- Deleted: (

The As and Ac cloud scenes in Table 4 are very similar to As and Ac cloud scenes in Table 3, except for slight reductions in τ_i and r_{ice} . Scenes in Table 4 are partly cloudy, implying a weaker infrared cloud signal. The Ci cloud scene in Table 4 shows slight reductions in τ_i and r_{ice} from the Ci cloud scene in Table 3.

The differences between Tables 3 and 4 are more significant for the convective ice clouds, however. The Cu cloud scene τ_i and AK are smaller while errors are larger in Table 4 compared to the pure Cu cloud scene in Table 3. This is expected as Cu clouds are several kilometers in depth but often have small horizontal scales and are averaged with clear sky in an AIRS pixel. Mixed (Cu,Nc) cloud scenes are especially problematic for plane-parallel radiative transfer calculations. This results in more uncertain retrievals of ice cloud properties for partial cloud (Cu,Nc) in Table 3 than those for the pure Cu cases in Table 2, which are more likely to be completely fill an AIRS scene.

The ice cloud property retrievals for cloud scenes that contain multiple cloud types are summarized in Table 5 for cloudy scenes and Table 6 for partly cloudy scenes. These tables list the nine most frequent cloud scene types as depicted in Figures 9 and 10. Seven of the nine cloud scenes are common between Tables 5 and 6. There is a general tendency for reductions of τ_i , increases in % relative error, and slight reductions in AKs in Table 6 for the seven common cloud scenes in Table 5. Changes in r_{ice} related variables are smaller than changes in τ_i related variables.

To summarize Tables 3-6, larger differences in ice cloud property retrievals are found between different cloud types than between cloudy and partly cloudy scenes. However, the differences between cloud scene types are the sharpest for the subset of cloudy scenes with one cloud type (Table 3). The AIRS cloud property retrievals are not greatly impacted by mixtures of cloud types within the AIRS footprint, and ice cloud property differences among cloud scenes are broadly consistent with the expected performance of infrared retrievals among these cloud types.

5 Summary

A method is described to classify cloud mixtures of cloud top types, termed *cloud scenes*, using the 2B-CLDCLASS cloud type classification obtained from the 94 GHz CloudSat radar. The scale dependence of the cloud scenes is quantified. The method is initially applied to two years of CloudSat data collocated within the Atmospheric Infrared Sounder (AIRS)/Atmospheric Microwave Sounding Unit (AMSU) field of regard (FOR) at 45 km scale. Given the 45 km scale and approximately 50 coinciding CloudSat profiles, each with 125 levels, the total number of possible scenes within an AMSU FOR is $9^{50 \times 125}$. This very large number of possible scenes is reduced to 256 by making three assumptions in the classification.

First, only the cloud type at the cloud top is considered. Second, the occurrence frequency of each cloud type within the cloud scene is disregarded; thus, there is no consideration of the counts of each cloud type. Third, the sequence of cloud types along the orbit segment is not considered. These three assumptions make mixed cloud scene classification tractable and are broadly

Deleted: ,Nc)

Deleted: 3

Deleted: 2

Deleted: AK for

Deleted: and τ_i , and a slight increase in their estimated errors.

Deleted: 3

Deleted: The slight degradation in the fidelity of ice cloud for retrievals (As,Nc) and (Ac,Nc) is confirmed by small

Deleted: AKs and increases in χ^2 compared to the fully cloud scenes in Table 2. The (Ci,Nc) cloud scene in Table 3 shows only slight changes

Deleted: 2, consistent with some scenes containing thin cirrus undetectable by CloudSat and flagged as clear

Deleted: 2

Deleted: 3

Deleted: (

Deleted: ,Nc)

Deleted: 3

Deleted: 2

Deleted: Recall that any combination of Cu and Nc qualifies as (Cu,Nc), including mostly clear ones.

Deleted: 2 and

Deleted: pure and mixed cloud scenes.

Deleted: affected

Deleted: those

Deleted: studied

Deleted: that

Deleted: AIRS/

consistent with the sensitivity of infrared sounders to clouds. They are also independent of the spatial scale of a scene and therefore can be generalized to all horizontal scales. A total of 210 out of 256 possible cloud scenes are observed in a two-year period from 01 July 2006 to 30 June 2008. The maximum number of cloud scenes occurs at a horizontal scale of 105 km with fewer cloud scenes at larger and smaller scales, and the majority of observed cloud scenes contain single cloud types.

Deleted: 194

The cloud scenes are organized into five categories: i) clear sky, (ii) cloudy sky with one cloud type, (iii) partly cloudy sky with one cloud type, (iv), cloudy sky with multiple cloud types, and (v) partly cloudy sky with multiple cloud types.

Summarizing AIRS cloud top property retrievals for cloudy sky with one cloud type, there is strong differentiation in the cloud thermodynamic phase. Ice phase dominates Ci, Ns, Dc, and As, while liquid and undetermined phase dominate Ac, Sc, and

Deleted: scenes with no clear sky (Table 2 and Figure 7),

Deleted: for scenes with only one cloud type

Cu. The results are similar for partly cloudy sky with one cloud type with an increase in unknown cloud phase and χ^2 residuals, and a reduction in information content for some cloud types. A similar set of calculations were performed for both cloudy and partly cloudy skies with multiple cloud types. In most cloud scenes with multiple cloud types, the changes in the ice properties are generally either small or reflect the combined characteristics of the multiple cloud types contained within the cloud scene.

Deleted: While not a direct comparison, in most mixed cloud scenes...

The sensitivity of thermal infrared cloud phase determination is consistent with independently determined cloud typing from the CloudSat radar for clouds detected by CloudSat.

Deleted: Furthermore, the cloud phase determination from AIRS is not adversely affected by more complex cloud scene morphology as indicated by CloudSat cloud type determined at the cloud tops.

The relative magnitude of differences in r_{ci} and τ_i , and their averaging kernels (AKs) and error estimates, and the χ^2 residual between simulated and observed radiances, are consistent with expectations of infrared retrieval sensitivity to different cloud types. Smaller error estimates and higher information content (AKs) within Ci cloud scenes are observed in comparison to thin cirrus likely missed by CloudSat in clear sky and Sc scenes. Larger τ_i and r_{ci} are observed at the tops of convective ice clouds. Differences in retrieved cloud properties between Ac and As cloud scenes are consistent with differences in their scene heterogeneity and cloud temperature. Variations in ice cloud property retrievals are larger between types of cloud scenes than between cloudy and partly cloudy/mixed cloud scenes.

The fidelity of AIRS retrieved cloud phase and ice cloud microphysics was tested within scenes with both uniform and non-uniform cloud cover, and one or more cloud types within the scene. As with phase, retrieval differences are shown to be larger among cloud types rather than between uniform and mixed cloud scenes.

Deleted: uniform and mixed scenes.

Moved up [1]: Kahn et al., 2015

New methodologies for simultaneous retrievals of cloud microphysical properties and temperature and specific humidity profiles that include clouds in the forward radiative transfer (e.g., De Souza-Machado et al. 2018; Irion et al. 2018) necessitate careful investigation of the effects of cloud mixtures on retrieved cloud properties. The bias and root-mean square error of AIRS temperature and specific humidity soundings depend on cloud type (Yue et al., 2013; Wong et al., 2015). A more rigorous evaluation of scene complexity is necessary for optimizing the retrieval configuration of future sounding algorithms (Irion et al. 2018) and for validating their products.

Deleted: While not an emphasis of this particular investigation, previous studies have shown that phase identification (Jin and Nasiri 2014), ice microphysical retrievals (e.g.,

Formatted: Font color: Text 1

Deleted:), and thermodynamic profile soundings – especially within the planetary boundary layer (e.g., Irion et al. 2018) – will remain a significant challenge for multi-layer clouds.

This investigation shows that careful inspection of footprint-scale AIRS cloud property retrievals are consistent with expectations of infrared sensitivity to different cloud types defined with the 94 GHz CloudSat radar. Other cloud observations, such as MODIS, may be used in a similar analysis to the one described here. MODIS captures the off-nadir portion of the AIRS swath and the fine-scale variability within AIRS footprints. Wang et al. (2016) used the cloud typing in CloudSat to cross validate with cloud typing using MODIS-defined cloud types. This establishes a link between cloud types obtained from CloudSat and MODIS. A rigorous estimation of the pixel-scale relationships between cloud properties obtained from CloudSat, MODIS and AMSU will help to further advance multi-sensor and multi-variate geophysical retrievals (e.g., Irion et al. 2018).

Deleted: the

Deleted: AIRS/

Acknowledgments

Part of this research was carried out at the Jet Propulsion Laboratory (JPL), California Institute of Technology, under a contract with the National Aeronautics and Space Administration. The authors thank two reviewers for helpful and insightful comments that led to an improved manuscript. This project was supported by NASA's Making Earth Science Data Records for Use in Research Environments (MEaSUREs) program. CloudSat data were obtained through the CloudSat Data Processing Center (<http://www.cloudsat.cira.colostate.edu/>). The combined data files used in this work are available at the Goddard Earth Sciences Data and Information Services Center (ftp://measures.gsfc.nasa.gov/data/s4pa/AIRS_CloudSat/). © 2019. All rights reserved. Government sponsorship acknowledged.

Deleted: 2018

References

- Barnes, E. A., and Polvani, L.: Response of the Midlatitude Jets, and of Their Variability, to Increased Greenhouse Gases in the CMIP5 Models, *J. Climate*, 26, 7117–7135, 2013.
- Bony, S., Colman, R., Kattsov, V. M., and Coauthors: How Well Do We Understand and Evaluate Climate Change Feedback Processes?, *J. Climate*, 19, 3445–3482, 2006.
- Ceppi, P., Hartmann, D. L., and Webb, M. J.: Mechanisms of the Negative Shortwave Cloud Feedback in Middle to High Latitudes, *J. Climate*, 29, 139–157, 2016.
- Chahine, M. T., Pagano, T. S., Aumann, H. H., and Coauthors: The Atmospheric Infrared Sounder (AIRS): Improving weather forecasting and providing new insights into climate, *B. Am. Meteor. Soc.*, 87, 911–926, <https://doi.org/10.1175/BAMS-87-7-911>, 2006.
- Chang, F. L. and Li, Z.: A near global climatology of single-layer and overlapped clouds and their optical properties retrieved from TERRA/MODIS data using a new algorithm, *J. Clim.*, 18, 4752–4771, 2005.

DeSouza-Machado, S., Strow, L. L., Tangborn, A., Huang, X., Chen, X., Liu, X., Wu, W., and Yang, Q.: Single-footprint retrievals for AIRS using a fast TwoSlab cloud-representation model and the SARTA all-sky infrared radiative transfer algorithm, *Atmos. Meas. Tech.*, 11, 529–550, <https://doi.org/10.5194/amt-11-529-2018>, 2018.

Golaz, J.-C., Larson, V. E., and Cotton, W. R.: A PDF-Based Model for Boundary Layer Clouds. Part I: Method and Model Description, *J. Atmos. Sci.*, 59, 3540–3551, 2002.

Guillaume, A., Kahn, B. H., Yue, Q., Fetzer, E. J., Wong, S., Manipon, G. J., Hua, H., and Wilson, B. D.: Horizontal and vertical scaling of cloud geometry inferred from CloudSat data, *J. Atmos. Sci.*, 75, 2187–2197, doi:10.1175/JAS-D-17-0111.1, 2018.

Holz, R. E., Ackerman, S., Antonelli, P., Nagle, F., Knuteson, R. O., McGill, M., Hlavka, D. L., and Hart, W. D.: An improvement to the high spectral resolution CO₂ slicing cloud top altitude retrieval, *J. Atmos. Oceanic Technol.*, 23, 653–670, 2006.

Huang, H.-L., Yang, P., Wei, H., Baum, B. A., Hu, Y., Antonelli, P., and Ackerman, S. A.: Inference of ice cloud properties from high spectral resolution infrared observations, *IEEE Trans. Geosci. Remote Sens.*, 42, 842–853, 2004.

Irion, F. W., Kahn, B. H., Schreier, M. M., Fetzer, E. J., Fishbein, E., Fu, D., Kalmus, P., Wilson, R. C., Wong, S., and Q. Yue, Q.: Single-footprint retrievals of temperature, water vapor, and cloud properties from AIRS, *Atmos. Meas. Tech.*, 11, 971–995, <https://doi.org/10.5194/amt-11-971-2018>, 2018.

Jakob, C., Tselioudis, G.: Objective identification of cloud regimes in the Tropical Western Pacific, *Geophys. Res. Lett.*, 30, 21, 2082, doi:10.1029/2003GL018367, 2003.

Jin, H., and Nasiri, S. L.: Evaluation of AIRS cloud-thermodynamic-phase determination with CALIPSO, *J. Appl. Meteorol. Climatol.*, 53, 1012–1026, 2014.

Kahn, B. H., Chahine, M. T., Stephens, G. L., Mace, G. G., Marchand, R. T., Wang, Z., Barnet, C. D., Eldering, A., Holz, R. E., Kuehn, R. E., Vane, D. G.: Cloud type comparisons of AIRS, CloudSat, and CALIPSO cloud height and amount, *Atmos. Chem. Phys.*, 8, 1231–1248, doi:10.5194/acp-8-1231-2008, 2008.

Kahn, B. H., and Coauthors: The Atmospheric Infrared Sounder version 6 cloud products, *Atmos. Chem. Phys.*, 14, 399–426, 2014.

Kahn, B. H., Schreier, M. M., Yue, Q., Fetzer, E. J., Irion, F. W., Platnick, S., Wang, C., Nasiri, S. L., and L’Ecuyer, T. S.: Pixel-scale assessment and uncertainty analysis of AIRS and MODIS ice cloud optical thickness and effective radius, *J. Geophys. Res. Atmos.*, 120, doi:10.1002/2015JD023950, 2015.

Kahn, B. H., Matheou, G., Yue, Q., Fauchez, T., Fetzer, E. J., Lebsock, M., Martins, J., Schreier, M. M., Suzuki, K., and Teixeira, J.: An A-train and MERRA view of cloud, thermodynamic, and dynamic variability within the subtropical marine boundary layer, *Atmos. Chem. Phys.*, 17, 9451–9468, <https://doi.org/10.5194/acp-17-9451-2017>, 2017.

Kahn, B. H., Takahashi, H., Stephens, G. L., Yue, Q., Delanoë, J., Manipon G., and Manning, E. M.: Ice cloud microphysical trends observed by the Atmospheric Infrared Sounder, *Atmos. Chem. Phys. Discuss.*, 18, 10715–10739, doi:10.5194/acp-18-10715-2018, 2018.

Deleted: Eguchi, N., Yokota, T.: Investigation of clear-sky occurrence rate estimated from CALIOP and MODIS observations, *Geophys. Res. Lett.*, 35, L23816, doi:10.1029/2008GL035897, 2008.

- Klein, S. A., and Jakob, C.: Validation and Sensitivities of Frontal Clouds Simulated by the ECMWF Model, *Mon. Wea. Rev.*, 127, 2514–2531, 1999.
- Krijger, J. M., van Weele, M., Aben, I., Frey, R.: Technical Note: The effect of sensor resolution on the number of cloud-free observations from space, *Atmos. Chem. Phys.*, 7, 2881–2891, doi:10.5194/acp-7-2881-2007, 2007.
- 5 Manipon, G.: README document for A Multi-Sensor Water Vapor Climate Data Record Using Cloud Classification, 32 pp., https://disc.gsfc.nasa.gov/measures/documentation/README.AIRS_CloudSat.pdf, 2012.
- Mazin, I. P.: Cloud phase structure: Experimental data analysis and parameterization, *J. Atmos. Sci.*, 63, 667–681, 2006.
- Miller, S.D., and Coauthors, 2014: Estimating three-dimensional cloud structure via statistically blended satellite observations., *J. Appl. Meteor. Climatol.*, 53, 437–455. <https://doi.org/10.1175/JAMC-D-13-070.1>.
- 10 Mitchell, J. F. B., Senior, C. A., and Ingram, W. J.: CO₂ and climate: A missing feedback?, *Nature*, 341, 132–134, 1989.
- Norris, J. R., Allen, R. J., Evan, A. T., Zelinka, M. D., O'Dell, C. W., and Klein, S. A.: Evidence for climate change in the satellite cloud record, *Nature*, 536, 72–75, <https://doi.org/10.1038/nature18273>, 2016.
- Naud, C. M., and Kahn, B. H.: Thermodynamic phase and ice cloud properties in northern hemisphere winter extratropical cyclones observed by Aqua AIRS, *J. Appl. Meteor. Climatol.*, 54, 2283–2303, doi:10.1175/JAMC-D-15-0045.1, 2015.
- 15 Protat, A., Delanoë, J., May, P. T., Haynes, J., Jakob, C., O'Connor, E., Pope, M., and Wheeler, M. C.: The variability of tropical ice cloud properties as a function of the large-scale context from ground-based radar-lidar observations over Darwin, Australia, *Atmos. Chem. Phys.*, 11, 8363–8384, 2011.
- Oreopoulos, L., Cho, N., Lee, D., Kato, S., Huffman, G. J.: An examination of the nature of global MODIS cloud regimes, *J. Geophys. Res. Atmos.*, 119, 8362–8383, doi:10.1002/2013JD021409, 2014.
- 20 Rossow, W. B., Schiffer, R. A.: Advances in understanding clouds from ISCCP., *Bull. Amer. Meteor. Soc.*, 80, 2261–2287, doi:10.1175/1520-0477(1999)080<2261:AIUCFI>2.0.CO;2, 1999.
- Rossow, W. B., Tselioudis, G., Polak, A., Jakob, C.: Tropical climate described as a distribution of weather states indicated by distinct mesoscale cloud property mixtures, *Geophys. Res. Lett.*, 32, L21812, doi:10.1029/2005GL024584, 2005.
- Rowe, P., M., Neshyba, S., and Walden, V. P.: Radiative consequences of low-temperature infrared refractive indices for supercooled water clouds, *Atmos. Chem. Phys.*, 13, 11925–11933, 2013.
- 25 Sassen, K. and Z. Wang: Level 2 Cloud Scenario Classification Product Process Description and Interface Control Document, http://www.cloudsat.cira.colostate.edu/sites/default/files/products/files/2B537CLDCLASS_PDICD.P_R04.20070724.pdf, 2005.
- Sassen, K., Wang, Z.: Classifying clouds around the globe with the CloudSat radar: 1-year of results, *Geophys. Res. Lett.*, 35, L04805, doi:10.1029/2007GL032591, 2008.
- 30 Schreier, M. M., Kahn, B. H., Eldering, A., Elliott, D. A., Fishbein, E., Irion, F. W., and Pagano, T. S.: Radiance comparisons of MODIS and AIRS using spatial response information, *J. Atmos. Ocean. Tech.*, 27, 1331–1342, <https://doi.org/10.1175/2010JTECHA1424.1>, 2010.

- Tan, I. and Storelvmo, T.: Sensitivity Study on the Influence of Cloud Microphysical Parameters on Mixed-Phase Cloud Thermodynamic Phase Partitioning in CAM5, *J. Atmos. Sci.*, 73, 709–727, 2016.
- Tan, I., Storelvmo, T., and Zelinka, M. D.: Observational constraints on mixed-phase clouds imply higher climate sensitivity, *Science*, 352, 224–227, 2016.
- 5 Thompson, D. R., Kahn, B. H., Green, R. O., Chien, S. A., Middleton, E. M., and Tran, D. Q.: Global spectroscopic survey of cloud thermodynamic phase at high spatial resolution, 2005–2015, *Atmos. Meas. Tech.*, 11, 1019–1030, <https://doi.org/10.5194/amt-11-1019-2018>, 2018.
- Tselioudis, G., Rossow, W., Zhang, Y., Konsta, D.: Global Weather States and Their Properties from Passive and Active Satellite Cloud Retrievals, *J. Climate*, 26, 7734–7746, doi:10.1175/JCLI-D-13-00024.1, 2013.
- 10 van Diedenhoven, B., Fridlind, A. M., Cairns, B., and Ackerman, A. S.: Variation of ice crystal size, shape, and asymmetry parameter in tops of tropical deep convective clouds, *J. Geophys. Res. Atmos.*, 119, 11809–11825, 2014.
- Wang, T., Fetzer, E. J., Wong, S., Kahn, B. H., Yue, Q.: Validation of MODIS cloud mask and multilayer flag using CloudSat-CALIPSO cloud profiles and a cross-reference of their cloud classifications, *J. Geophys. Res. Atmos.*, 121, 11, 620–11, 635, doi:10.1002/2016JD025239, 2016.
- 15 Wong, S., Fetzer, E. J., Schreier, M., Manipon, G., Fishbein, E. F., Kahn, B. H., Yue, Q., and Irion, F. W.: Cloud-induced uncertainties in AIRS and ECMWF temperature and specific humidity, *J. Geophys. Res. Atmos.*, 120, doi:10.1002/2014JD022440, 2015.
- Xu, K., T. Wong, B.A. Wielicki, L. Parker, and Z.A. Eitzen: Statistical Analyses of Satellite Cloud Object Data from CERES. Part I: Methodology and Preliminary Results of the 1998 El Niño/2000 La Niña, *J. Climate*, 18, 2497–2514, 2005.
- 20 <https://doi.org/10.1175/JCLI3418.1>, 2005.
- Yuan, T., and Li, Z.: General macro- and microphysical properties of deep convective clouds as observed by MODIS, *J. Climate*, 23, 3457–3473, 2010.
- Yue, Q., Kahn, B. H., Fetzer, E. J., and Teixeira, J.: Relationship between marine boundary layer clouds and lower tropospheric stability observed by AIRS, CloudSat, and CALIOP, *J. Geophys. Res.*, 116, D18212, doi:10.1029/2011JD016136, 2011.
- 25 Yue, Q., Fetzer, E. J., Kahn, B. H., Wong, S., Manipon, G., Guillaume, A., Wilson, B.: Cloud-State-Dependent Sampling in AIRS Observations Based on CloudSat Cloud Classification, *J. Climate*, 26, 8357–8377, doi: 10.1175/JCLI-D-13-00065.1, 2013.
- Zhang, D., Wang, Z., and Liu, D.: A global view of midlevel liquid-layer topped stratiform cloud distribution and phase partition from CALIPSO and CloudSat measurements, *J. Geophys. Res.*, 115, D00H13, doi:10.1029/2009JD012143, 2010.
- 30 Zelinka, M. D., Klein, S. A., Taylor, K. E., and Coauthors: Contributions of different cloud types to feedbacks and rapid adjustments in CMIP5. *J. Climate*, 26, 5007–5027, doi:10.1175/JCLI-D-12-00555.1, 2013.
- Zhao, G., Di Girolamo, L., Diner, D. J., Bruegge, C. J., Mueller, K. J., and Wu, D. L.: Regional Changes in Earth's Color and Texture as Observed From Space Over a 15-Year Period, *IEEE Trans. Geosci. Remote Sens.*, 54, 4240–4249, 2016.

Cloud type	nc	Ci	As	Ac	St	Sc	Cu	Ns	Dc
median	6.6	8.8	12.1	1.1	3.3	1.1	1.1	15.4	14.3
Median absolute deviation	5.5	7.7	11.0	0.0	2.2	0.0	0.0	13.2	9.9

Table 1. Horizontal cloud chord length median and median absolute deviation for each cloud type (km).

5

- Deleted: cloud_
- Formatted: Centered
- Deleted: AISt
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: AICu
- Deleted: DC
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Centered
- Formatted Table
- Deleted: mad
- Formatted: Centered
- Deleted: 25.7
- Deleted: 28
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 29.6
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 39.2
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 11.1
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 6.1
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 9.8
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 6
- Formatted: Font: Times, 10 pt, Not Bold, English (UK)
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: 42.8
- Formatted: Font: Times, 10 pt, Not Bold, English (UK)
- Formatted: Font: Times, 10 pt, Not Bold
- Formatted: Font: Times, 10 pt, Not Bold
- Deleted: (m.a.d.)
- Deleted: , in kilometers.

Type of scene	Total count	percent	
clear	7175523	30.7	
cloudy sky with one cloud type	7332076	31.3	
partly cloudy sky with one cloud type	550607	4	23.5
cloudy sky with multiple cloud types	2377259	10	2
partly cloudy sky with multiple cloud types	1008158	4	3
all	23399090	100.00	

Table 2. Total counts and relative percentages of five cloud scene categories at the AIRS FOV scale: clear sky, cloudy sky with one cloud type, partly cloudy sky with one cloud type, cloudy sky with multiple cloud types, and partly cloudy sky with multiple cloud types.

5

- Deleted: ¶ ... [11]
- Deleted: Cloud¶ ... [2]
- Deleted: τ_i averaging kernel ... [10]
- Deleted Cells ... [10]
- Deleted: %¶ ... [11]
- Deleted Cells ... [3]
- Deleted Cells ... [12]
- Deleted: Mean r_{ei} ... [13]
- Deleted Cells ... [13]
- Deleted: r_{ei} relative error ... [14]
- Deleted Cells ... [14]
- Deleted: r_{ei} averaging kernel ... [15]
- Deleted Cells ... [15]
- Deleted: % passing QC for r_{ei} ... [16]
- Deleted Cells ... [16]
- Formatted Table ... [4]
- Deleted: χ^2 residual fit ... [17]
- Deleted Cells ... [17]
- Deleted: Single ... [6]
- Formatted ... [6]
- Deleted: proportion¶ ... [7]
- Formatted ... [7]
- Formatted ... [5]
- Deleted: Mean τ_i ... [8]
- Formatted ... [8]
- Deleted: τ_i relative error ... [9]
- Formatted ... [9]
- Deleted: C_i ... [18]
- Formatted ... [19]
- Moved down [4]: 57.0
- Deleted: 3.27
- Deleted: 5.29
- Deleted Cells ... [24]
- Deleted: 0.96
- Deleted Cells ... [25]
- Deleted: 27.9
- Deleted Cells ... [27]
- Deleted: 7.1
- Deleted Cells ... [28]
- Deleted: 0.96
- Deleted Cells ... [29]
- Deleted: 3.5
- Deleted Cells ... [23]
- Deleted Cells ... [31]
- Formatted Table ... [20]
- Deleted: 68.8
- Deleted Cells ... [30]
- Deleted: 1.0
- Formatted ... [21]
- Formatted ... [22]
- Deleted: 93
- Formatted ... [26]
- Deleted: N_s
- Formatted ... [32]

Cloud type	Cloudy single type proportion	Mean τ_i	τ_i relative error	τ_i averaging kernel	% passing QC for τ_i	Mean r_{ei}	r_{ei} relative error	r_{ei} averaging kernel	% passing QC for r_{ei}	χ^2 residual fit
nc	49.5	0.77	14.88	0.71	75.09	20.9	5.6	0.97	34.1	3.3
As	14.0	2.42	6.41	0.92	97.36	24.5	4.7	0.98	79.9	2.9
Sc	11.5	1.30	15.12	0.69	82.63	20.6	6.5	0.96	48.6	3.2
Ci	11.3	1.91	2.21	0.99	96.44	25.4	2.6	1.00	74.2	4.0
Ns	8.6	2.41	8.70	0.88	98.11	23.6	5.6	0.98	87.0	2.4
Ac	2.9	1.65	6.66	0.91	93.45	22.1	4.1	0.98	59.0	4.1
Dc	1.7	5.47	3.53	0.98	98.64	27.1	7.1	0.96	71.9	3.2
Cu	0.6	2.92	5.99	0.94	94.37	26.5	6.1	0.97	70.6	3.9
St	0.001	2.33	15.37	0.63	100.00	27.3	4.9	0.98	70.0	4.1

Table 3. Cloud ice properties for cloudy sky with one cloud type (i.e., all CloudSat profiles have the same cloud type). Proportions and relative errors are in percent. The effective radius is in μm .

Deleted: pixels within one AIRS footprint

Deleted: optical

<u>Cloud type</u>	<u>Cloudy single type proportion</u>	<u>Mean τ_i</u>	<u>τ_i relative error</u>	<u>τ_i averaging kernel</u>	<u>% passing QC for τ_i</u>	<u>Mean r_{ei}</u>	<u>r_{ei} relative error</u>	<u>r_{ei} averaging kernel</u>	<u>% passing QC for r_{ei}</u>	<u>χ^2 residual fit</u>
Sc	74.79	0.93	14.45	0.71	73.91	21.2	5.7	0.97	36.8	3.5
Ci	10.44	0.71	3.69	0.96	87.98	22.5	2.8	0.99	62.0	4.2
Ac	5.08	0.94	5.95	0.92	88.41	20.8	3.6	0.99	52.6	4.6
As	4.83	1.09	14.40	0.72	90.47	19.9	6.4	0.96	46.5	3.1
Cu	4.80	0.87	12.78	0.76	74.49	21.1	5.2	0.97	38.2	3.7
Ns	0.05	1.69	14.50	0.71	85.08	22.9	7.1	0.96	45.3	3.7
Dc	0.01	4.63	3.44	0.98	93.58	26.8	6.4	0.96	53.8	3.9
St	0.01	0.79	8.95	0.88	84.00	21.3	6.3	0.96	64.0	3.9

Table 4. Cloud ice properties for partly cloudy sky with one cloud type. All else the same as Table 3.

Formatted: English (US)


Cloud scene	Mixed scenes proportion	Mean τ_i	τ_i relative error	τ_i averaging kernel	% passing QC for τ_i	Mean r_{ei}	r_{ei} relative error	r_{ei} averaging kernel	% passing QC for r_{ei}	χ^2 residual fit
Ci As	16.7	2.52	4.05	0.96	97.56	24.7	3.8	0.99	74.6	3.5
Ac Sc	16.0	1.31	10.41	0.81	85.92	21.9	5.0	0.97	53.7	4.0
As Ns	11.0	2.07	10.66	0.83	97.01	22.4	5.7	0.97	81.6	2.6
Ci Ns	7.2	2.14	6.84	0.93	97.39	22.9	4.3	0.98	85.8	2.8
Ac Ns	6.1	1.75	12.83	0.77	91.58	21.2	6.0	0.97	60.0	3.1
Ci Sc	5.2	1.00	4.61	0.95	84.98	24.5	3.1	0.99	60.7	4.3
As Sc	4.8	1.34	16.31	0.66	91.55	19.7	6.6	0.96	57.0	2.9
As Dc	3.7	5.22	3.12	0.99	99.07	28.0	6.8	0.96	65.8	3.5
Ci Dc	3.4	4.30	2.82	0.99	98.61	27.4	4.9	0.98	62.0	4.4

Table 5. Cloud ice properties for cloudy sky with multiple cloud types for the first nine most observed cloud scenes at the AIRS FOV scale. All else the same as Table 4.

- Deleted: nc
- Formatted Table ... [59]
- Deleted: 30
- Deleted: 0.78
- Deleted: 14.8
- Deleted: 71
- Deleted: 20.9
- Deleted: 5.6
- Formatted ... [60]
- Deleted: 75.2
- Formatted ... [62]
- Deleted: 97
- Formatted ... [63]
- Deleted: 3
- Formatted ... [61]
- Deleted: 34.3
- Deleted: 24.7
- Formatted ... [64]
- Deleted: 0.98
- Deleted: 14.5
- Formatted ... [65]
- Deleted: 71
- Deleted: 75.0
- Formatted ... [66]
- Deleted: 1
- Formatted ... [67]
- Deleted: 4
- Formatted ... [68]
- Deleted: 3.
- Formatted ... [69]
- Deleted: 38.2
- Deleted: 9.8
- Formatted ... [71]
- Deleted: 35
- Deleted: 6.8
- Formatted ... [70]
- Formatted ... [72]
- Deleted: 91
- Formatted ... [73]
- Deleted: 0
- Deleted: 24.3
- Deleted: 4.8
- Formatted ... [74]
- Deleted: 98
- Formatted ... [75]
- Deleted: 9
- Deleted: 78.2
- Formatted ... [76]
- Deleted: 9.5
- Deleted: 1.71
- Deleted: 2.5
- Formatted ... [77]
- Deleted: 98
- Deleted: 94.9

Cloud scene	Mixed scenes proportion	Mean τ_i	τ_i relative error	τ_i averaging kernel	% passing QC for τ_i	Mean Γ_{ei}	Γ_{ei} relative error	Γ_{ei} averaging kernel	% passing QC for Γ_{ei}	χ^2 residual fit
Sc.Cu	18.0	1.03	14.00	0.73	76.20	21.8	5.9	0.97	39.1	3.7
Ac.Sc	20.1	0.87	6.34	0.92	84.99	21.4	4.0	0.98	54.9	4.4
Ci.Sc	17.8	0.70	4.07	0.95	86.31	22.6	2.9	0.99	61.1	4.2
As.Sc	5.2	1.16	14.36	0.72	88.08	20.4	6.3	0.96	47.9	3.4
Ac.Cu	3.4	0.94	4.84	0.95	86.14	23.0	4.1	0.98	56.3	4.7
Ci.Ac	3.3	1.18	2.93	0.98	93.16	22.3	2.5	0.99	56.4	5.0
Ac.Sc.Cu	3.2	0.83	5.36	0.94	85.47	22.7	4.3	0.98	59.5	4.4
Ci.As	2.7	1.22	7.10	0.89	92.41	21.3	4.2	0.98	52.5	3.9
As.Ac	2.1	1.14	8.07	0.87	90.54	21.3	4.5	0.98	48.6	4.3

Table 6. Cloud ice properties for partly cloudy sky with multiple cloud types for the first nine most observed cloud scenes at the AIRS FOV scale. All else the same as Table 5.






Figure 1. Histogram of cloud scenes containing relative counts of occurrence, observed at the AMSU FOR and AIRS FOV resolution (~45 km and ~15 km respectively). The cumulative sum of the relative counts of these 18 (10) cloud scenes amounts to more than 90% of all cloud scenes observed globally over a period of two years at AMSU (AIRS) resolution.





Figure 2. Geographic distribution of cloud scenes (Sc) and (Ac,Sc) in panels (a) and (b) respectively, in units of percentage with respect to all of the (194) observed cloud scenes. These scenes were observed at the AMSU FOR resolution (~45 km). Similar plots of the AIRS FOV resolution (~15 km) are nearly identical (not shown).

Deleted: .

Deleted: (at least)

Formatted: English (US)



Deleted:

Deleted: :

Deleted: on panel (a), (nc) on panel (b)

Deleted: on panel (c). Panel (d) show the geographic distribution...

Deleted: sum of counts of all the

Deleted: The counts of the three other panels are normalized in each grid cell with the count of the corresponding grid cell in

Deleted: lower right panel and are reported in percent.






Figure 3. (a) Number of observed scenes as a function of the horizontal length scale used to define the scene, (b) Number of scenes observed at the 90th percentile as a function of horizontal length scale used to define the scene. The vertical green (red) lines approximate the scale of the AIRS (AMSU) pixel size.

- Deleted:
- Deleted: scale
- Deleted: (a).
- Deleted: in
- Deleted: scale
- Deleted: (b).
- Deleted: /
- Deleted: FOR





Figure 4. Distribution of (a) minimum and (c) maximum length for 5 of the first 200 cloud scenes. The five scenes are, from top to bottom: (Ac,Sc) in blue, (As,Sc,Cu) in orange, (Cl,As,Cu,DC) in yellow, (As,Ac,Ns,DC) in purple and (Cl,As,Ac,St,Sc) in green and their respective ranks are 1, 26, 51, 76 and 101. In panels (b) and (d), the number of scenes were obtained by summing the number of scenes present a different lengths.

5



Deleted:

Deleted: AICu

Deleted: AIST

Deleted: ci,AIST

Deleted: DC

Deleted: AIST,AICu

Deleted: DC

Deleted: ci,AIST,AICu

Deleted: On





Figure 5. Cloud type vertical cross-section defined by the values of the cloud scenario variables of the 2B-CLDCLASS product. Each color corresponds to a different cloud type (legend on right). Color segments on top of the figure indicate the horizontal extent of a cloud measured at its top.



Deleted:

Deleted: profile

Deleted: 2b

Deleted: .







Figure 6. Horizontal cloud chord length frequency histograms for each of the eight CloudSat cloud types and clear sky. The cloud chord length was obtained at the cloud top (see Fig. 5) unlike that obtained in Guillaume et al. (2018).

5



Deleted:



Deleted:

Deleted: variable

Deleted: single

Formatted: English (US)

Deleted: types (



Deleted: pixels within one AIRS footprint

Deleted: but

Deleted: scenes

Figure 7. AIRS cloud_phase_3x3 histograms for cloudy sky with one cloud type (i.e., all CloudSat profiles have the same cloud type and no clear sky). The red, green, and blue bars indicate liquid, undetermined, and ice phase, respectively. Each histogram sums to 1.0 and does not show how many counts relative to another histogram. Relative counts could be inferred from the percentages listed in the 2nd to left column of Table 3.

5



Deleted:

5 Figure 8. AIRS cloud_phase_3x3 histograms for partly cloudy sky with one cloud type. All else equal to Figure 7.




Figure 9. AIRS cloud_phase_3x3 histograms for cloudy sky with multiple cloud types for the top 9 ranked cloud scenes in order of occurrence frequency.

Deleted: variable for the first mixed scenes of the 90th percentile. The first 'mixed' scene, with only clear fields, nc, was omitted

Deleted: clarity and because it is

Deleted: same as




Figure 10. AIRS cloud_phase_3x3 histograms for partly cloudy sky with multiple cloud types for the top 9 ranked cloud scenes in order of occurrence frequency.

- Deleted: nc
- Formatted: English (US)
- Deleted: _phase_3x3 barplot
- Deleted: pure nc (see Fig. 7).

Page 21: [1] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 21: [2] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 21: [3] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [4] Formatted Table Alexandre Guillaume 6/12/19 9:08:00 AM
Formatted Table

Page 21: [5] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM
Left

Page 21: [6] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM
Font: Times, English (US)

Page 21: [7] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM
Font: Times, English (US)

Page 21: [8] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM
Font: Times, English (US)

Page 21: [9] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM
Font: Times, English (US)

Page 21: [10] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [11] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 21: [12] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [13] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [14] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [15] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [16] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM
Deleted Cells

Page 21: [17] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Page 21: [18] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

partly cloudy sky with one cloud type	5506074	23.5
---------------------------------------	---------	------

Page 21: [19] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Left

Page 21: [20] Formatted Table Alexandre Guillaume 6/12/19 9:08:00 AM

Formatted Table

Page 21: [21] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times, English (US)

Page 21: [22] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times, English (US)

Page 21: [23] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [24] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [25] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [26] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 21: [26] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 21: [27] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [28] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [29] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [30] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [31] Deleted Cells Alexandre Guillaume 6/12/19 9:08:00 AM

Deleted Cells

Page 21: [32] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 21: [33] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times, English (US)

Page 21: [34] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times, English (US)

Page 21: [35] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 21: [36] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times, English (US)

Page 21: [37] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [38] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [39] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [40] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [41] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [42] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [43] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [44] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [45] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Left

Page 21: [46] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times, English (US)

Page 21: [47] Inserted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
-------------------------------------	----------------------------	---------------------------

Inserted Cells

Page 21: [48] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Page 21: [49] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times, English (US)

Page 21: [50] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [51] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [52] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [53] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [54] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [55] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [56] Deleted Cells	Alexandre Guillaume	6/12/19 9:08:00 AM
------------------------------------	----------------------------	---------------------------

Deleted Cells

Page 21: [57] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: 9 pt, Bold

Page 21: [58] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: 9 pt, Bold, English (US)

Page 24: [59] Formatted Table	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------------	----------------------------	---------------------------

Formatted Table

Page 24: [60] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [61] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [62] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [63] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [64] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Page 24: [65] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [66] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [67] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [68] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [69] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [70] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [71] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [72] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [73] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [74] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [75] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [76] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [77] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [78] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [79] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Font: Times Roman

Page 24: [80] Formatted	Alexandre Guillaume	6/12/19 9:08:00 AM
--------------------------------	----------------------------	---------------------------

Page 24: [81] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [82] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [83] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [84] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [85] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [86] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [87] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [88] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [89] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [90] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [91] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [92] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [93] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [94] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [95] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [96] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 24: [97] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [98] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [99] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [100] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

Page 24: [101] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [102] Formatted Table Alexandre Guillaume 6/12/19 9:08:00 AM

Formatted Table

Page 24: [103] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [104] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [105] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [106] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [107] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [108] Formatted Alexandre Guillaume 6/12/19 9:08:00 AM

Font: Times Roman

Page 24: [109] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

▼

Page 24: [109] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

▼

Page 24: [109] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

▼

Page 24: [109] Deleted Alexandre Guillaume 6/12/19 9:08:00 AM

▼

English (US)