



# Use of polarimetric radar measurements to constrain simulated convective cell evolution: A pilot study with Lagrangian tracking

Ann M. Fridlind<sup>1</sup>, Marcus van Lier-Walqui<sup>1,2</sup>, Scott Collis<sup>3</sup>, Scott E. Giangrande<sup>4</sup>, Robert C. Jackson<sup>3</sup>, Xiaowen Li<sup>5,6</sup>, Toshihisa Matsui<sup>6,7</sup>, Richard Orville<sup>8</sup>, Mark H. Picel<sup>4</sup>, Daniel Rosenfeld<sup>9</sup>, Alexander Ryzhkov<sup>10</sup>, Richard Weitz<sup>8</sup>, and Pengfei Zhang<sup>10</sup>

<sup>1</sup>NASA Goddard Institute for Space Studies, New York, NY, USA

<sup>6</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>7</sup>ESSIC/University of Maryland, College Park, MD, USA

<sup>8</sup>Texas A&M University, College Station, TX, USA

<sup>9</sup>The Hebrew University of Jerusalem, Jerusalem, Israel

<sup>10</sup>Cooperative Institute for Mesoscale meteorological Studies, University of Oklahoma, and NOAA/OAR National Severe Storms Laboratory, Norman, OK, USA

Correspondence to: ann.fridlind@nasa.gov

**Abstract.** To probe the potential value of a radar-driven field campaign to constrain simulation of isolated convection subject to a strong aerosol perturbation, convective cells observed by the operational KHGX weather radar in the vicinity of Houston, Texas, are examined individually and statistically. Cells observed in a single case study of onshore flow conditions during July 2013 are

- 5 first examined and compared with cells in a regional model simulation. Observed and simulated cells are objectively identified and tracked from observed or calculated positive specific differential phase  $(K_{\rm DP})$  above the melting level, which is related to the presence of supercooled liquid water. Several observed and simulated cells are subjectively selected for further examination. Below the melting level, we compare sequential cross-sections of retrieved and simulated raindrop size distribution pa-
- 10 rameters. Above the melting level, we examine time series of  $K_{\text{DP}}$  and radar differential reflectivity  $(Z_{\text{DR}})$  statistics from observations and calculated from simulated supercooled rain properties, alongside simulated vertical wind and supercooled rain mixing ratio statistics. Results indicate that the operational weather radar measurements offer multiple constraints on the properties of simulated convective cells, with substantial value added from derived  $K_{\text{DP}}$  and retrieved rain properties. The
- 15 value of collocated three-dimensional lightning mapping array measurements, which are relatively

<sup>&</sup>lt;sup>2</sup>Columbia University, New York, NY, USA

<sup>&</sup>lt;sup>3</sup>DOE Argonne National Laboratory, Argonne, IL, USA

<sup>&</sup>lt;sup>4</sup>DOE Brookhaven National Laboratory, Upton, NY, USA

<sup>&</sup>lt;sup>5</sup>GESTAR/Morgan State University, Baltimore, MD, USA





rare in the continental U.S., supports the choice of Houston as a suitable location for future field studies to improve the simulation and understanding of convective updraft physics. However, rapid evolution of cells between routine volume scans motivates consideration of adaptive scan strategies or radar imaging technologies to amend operational weather radar capabilities. A three-year clima-

- 20 tology of isolated cell tracks, prepared using a more efficient algorithm, yields additional relevant information. Isolated cells are found within the KHGX domain on roughly 40% of days year-round, with greatest concentration in the northwest quadrant, but roughly fivefold more cells occur during June through September. During this enhanced occurrence period, the cells initiate following a strong diurnal cycle that peaks in the early afternoon, typically follow a south-to-north flow, and
- 25 dissipate within an hour, consistent with the case study examples. Statistics indicate that ~150 isolated cells initiate and dissipate within 70 km of the KHGX radar during the enhanced occurrence period annually, and roughly ten times as many within 200 km, suitable for multi-instrument Lagrangian observation strategies. In addition to ancillary meteorological and aerosol measurements, robust vertical wind speed retrievals would add substantial value to a radar-driven field campaign.

#### 30 1 Introduction

Since the Intergovernmental Panel on Climate Change's first scientific assessment report (IPCC, 1990), the conclusion has been generally strengthened that aerosol pollution from anthropogenic activities is likely to have commonly offset regional and global radiative forcing of the Earth's climate by anthropogenic greenhouse gases to date, but uncertainty especially in aerosol effects on

- 35 cloud-related forcing still remains high (IPCC, 2013). Although such anthropogenic aerosol radiative forcing will be diminutive relative to that from build-up of anthropogenic greenhouse gases on century timescales under most scenarios, the variable degree to which anthropogenic aerosols offset greenhouse gas warming in simulations that reproduce the observational record of surface temperature change since pre-industrial times continues to be a leading factor limiting simulation constraints
- 40 on Earth's climate sensitivity (e.g., Kiehl, 2007). Fundamental understanding of the relationships between global cloud processes and atmospheric circulations and thermodynamics is another leading factor, as demonstrated by studies that find grossly differing predicted climate sensitivities associated with differing parameterization of fundamental processes such as convective mixing, convective aggregation or cloud glaciation (e.g., Sherwood et al., 2014; Mauritsen and Stevens, 2015; Tan et al., 2016).

45 2016).

Addressing aerosol-cloud-precipitation-climate interactions locally and regionally, Rosenfeld et al. (2014) describe how field campaigns designed to measure closed energy and moisture budgets over relatively large domains, referred to as box flux closure experiments, could help advance understanding of primary microphysical mechanisms and regional-scale dynamical feedbacks. Here we

50 further consider how observations designed for a box flux closure experiment could be amended to





aid attribution of primary cloud responses to aerosol variability, specifically in the case of convective cells responding to a boundary layer aerosol perturbation. The observational objective of establishing microphysical and dynamical differences across a population of evolving convective cells is considered an amendment because well observing the details of updraft cell evolution within a flux

- 55 closure campaign box poses a significant additional challenge beyond constraining fluxes at the box boundaries. However, there may be overlapping utility in the use of polarimetric radar systems to observe convective cell spatial evolution within the box and to provide state-of-the-art retrievals of surface precipitation rate at the lower box boundary, as discussed further below. Especially when coordinated with detailed high-resolution modeling, we argue that measurements optimized to observe
- 60 convective cell evolution would additionally be uniquely valuable for advancing understanding and accurate simulation of cloud processes such as entrainment and glaciation, thereby further addressing understanding of fundamental cloud processes relevant to climate sensitivity.

As understood for decades, cloud parcels rising in convective updrafts from a warm cloud base height pass through the melting level (0°C) carrying liquid water that does not instantaneously freeze owing to an energy barrier to ice crystal formation (Pruppacher and Klett, 1997). For individual drops, that barrier would not be spontaneously overcome in commonly occurring dilute solution drops until they are supercooled by at least 30°C (e.g., Herbert et al., 2015). However, relatively rare aerosol that commonly exhibit solid surfaces, such as dust, may serve as ice-nucleating particles

- (INPs) that lower the energy barrier to ice crystal formation (Vali et al., 2015). Such INPs may nucleate individual crystals with varying efficiencies that have been widely measured in the field over activation temperatures of roughly -10 to -35°C, for instance (DeMott et al., 2010); some biological agents may lead to primary ice formation at temperatures as warm as -3°C (e.g., Du et al., 2017). Once ice is present in an updraft parcel, whether via primary nucleation by INPs present or via some means of transport (sedimentation or entrainment), so-called secondary ice crystal formation
- 75 may potentially progress via ice-liquid or ice-ice collisions or some fracturing process related to ice expansion during drop freezing (e.g., Hallett and Mossop, 1974; Vardiman, 1978; Yano and Phillips, 2011; Lawson et al., 2015, and references therein). Based on several recent field campaigns, it has been argued that multiplication is likely a process that may commonly dominate ice size distribution evolution in warm-base convective updrafts and long-lived stratiform outflow (Lawson et al., 2015;
- 80 Ackerman et al., 2015; Fridlind et al., 2017; Ladino et al., 2017). In general, there is increasing evidence that the processes that determine updraft and outflow ice properties to first order, and by extension their relationship to environmental conditions, are still not yet well understood nor well represented in microphysics schemes to date (Lawson et al., 2015; Ackerman et al., 2015; Fridlind et al., 2017; Stanford et al., 2017).
- Although the latent heat of fusion is only roughly 15% of the latent heat of condensation, liquid water freezing does contribute to updraft buoyancy, and has been identified as a factor explaining updraft extent in tropical environments (Zipser, 2003). If rain formation leads to liquid water sed-





100

imentation from an updraft before it reaches the melting level or before it freezes at some colder temperature above, clearly no latent heat of fusion is contributed; to the extent that more liquid wa-

- 90 ter reaches freezing temperatures when rain formation is weaker, increasing aerosol loading will lead to stronger updrafts, all else being equal. This is borne out in simulations under some environmental conditions to an extent that may be dependent on the complexity of the microphysics scheme, and has been supported by large-domain statistical studies (e.g., Fan et al., 2013, and references therein). On the other hand, it has also been repeatedly demonstrated that differing microphysics schemes pre-
- 95 dict grossly differing updraft and cloud properties, at least in part owing to a lack of observational constraints on important factors such as liquid water content and ice properties, making it extremely challenging to establish whether any scheme is performing substantially better than any other for the correct reasons (e.g., Fridlind et al., 2012; Varble et al., 2014a, b; Wang et al., 2015).

Polarimetric radar systems such as those operated by the National Weather Service Next-Generation Radar (NEXRAD) program (NOAA, 2017) provide a rich source of information about the size dis-

- tribution, phase, and shape of hydrometeors (Zrnić and Ryzhkov, 1999), which is especially valuable for the study of convective updraft physics because of the paucity of such data available from aircraft (e.g., Loney et al., 2002; Fridlind et al., 2015; Snyder et al., 2015). Comparison of reflectivity and phase-shift differentials between horizontal and vertical radar polarizations yields differ-
- 105 ential reflectivity ( $Z_{DR}$ ) and specific differential phase ( $K_{DP}$ ), which are related to the presence of horizontally-aligned oblate or prolate hydrometeors when positive (Bringi and Chandrasekar, 2001). Vertically elongated columns of positive  $Z_{DR}$  and  $K_{DP}$  that extend above the environmental melting level ( $Z_{DR}$  and  $K_{DP}$  columns) have been generally attributed to the presence of supercooled liquid associated with a deep convective updraft that is not otherwise identifiable from reflectivity alone
- 110 (Bringi et al., 1996; Hubbert et al., 1998; Loney et al., 2002; Kumjian et al., 2014a). Recent studies suggest a strong connection between  $K_{\rm DP}$  and  $Z_{\rm DR}$  columns and other metrics of deep convective activity such as overshooting tops (Homeyer and Kumjian, 2015) and lighting flash rate and updraft mass flux (van Lier-Walqui et al., 2016). Observations also show differences in  $K_{\rm DP}$  versus  $Z_{\rm DR}$ column morphology (Zrnić et al., 2001; Loney et al., 2002; Kumjian and Ryzhkov, 2008), which
- 115 have been attributed to differing sensitivities to hydrometeor size distribution and phase characteristics (e.g., Kumjian et al., 2014b; Snyder et al., 2017b). However, precise attribution of specific morphological features at various wavelengths remains a challenge due to a paucity of colocated in situ measurements, the complexity of updraft microphysics, and uncertainties in calculating hydrometeor electromagnetic properties especially for mixed-phase particles (e.g., Loney et al., 2002;
- 120 Ryzhkov et al., 2011; Snyder et al., 2013, 2017b).

Many past studies have effectively examined deep convective cells using observations from individual radar scans (e.g., Hubbert et al., 1998; Loney et al., 2002). Tracking convective cells or other identified features in time increases the amount of information that can be gleaned from scanning radar because temporal aspects such as cell lifetime can be quantified (e.g., Stein et al., 2015). Fea-





- 125 ture identification and tracking have wide applications in the atmospheric sciences. Many studies have applied unsupervised feature identification to locating cloud regimes using satellite observations (e.g., Jakob and Tselioudis, 2003; Rossow et al., 2005; Pope et al., 2009). Automatic tracking is perhaps most widely applied in nowcasting using surface radar observations (Dixon and Wiener, 1993; Johnson et al., 1998; Scharenbroich et al., 2010; Limpert et el., 2015). Surface radar observa-
- 130 tions generally have a frequency on the order of 5–10 min, and the rate of successful tracking can be 60% to 90% according to Lakshmanan and Smith (2010). Here we investigate isolated convective cells, which have smaller sizes and shorter lifespans than the storm features in most radar weather tracking. The K<sub>DP</sub> column identification algorithm used in this pilot study was described by van Lier-Walqui et al. (2016). We also introduce a more efficient tracking algorithm for compilation of long-term statistics using parallel computing.

For the study of updraft microphysics we target conditions where a relatively strong aerosol perturbation exists and updrafts are not being strongly driven by synoptic flow, which are commonly satisfied in the vicinity of Houston when there is onshore flow. Such conditions increase the likelihood of observationally establishing the statistical relationship between aerosol and updraft proper-

140 ties, which can in turn be used as a constraint for evaluating and improving models. The Houston region currently offers the significant advantage of a Lightning Mapping Array (LMA; Orville et al., 2012), which can provide independent three-dimensional information on updraft location and phase (e.g., van Lier-Walqui et al., 2016). The objectives of this pilot study are to establish the lifetime and observable properties of typical isolated convective cells, and to demonstrate comparison of isolated 145 cell observations with an example regional model simulation.

Next in Sect. 2, we describe the data sources and methods of analyzing isolated cell features in a selected case study and in long-term statistics. Results are presented in Sect. 3. In Sect. 4, we discuss results in the context of pilot study objectives.

#### 2 Methodology

## 150 2.1 Data sources

Level II data from the NEXRAD KHGX radar on 8 June 2013 (NOAA, 1991) were mapped to Cartesian coordinates at 1-km resolution and approximately 5-min frequency using the Python ARM Radar Toolkit (Py-ART; Helmus and Collis, 2016). A linear programming (LP) phase processing algorithm based on Giangrande et al. (2013), and included in Py-ART, was used to unfold and process

155 raw differential phase into propagation differential phase. The LP phase processing algorithm imposes a monotonicity constraint on phase, which makes it inappropriate for estimating regions of decreasing propagation phase shift (i.e. regions where differential phase shift, or  $K_{\text{DP}}$ , is negative). Conversely, this fact makes the LP algorithm better suited for identifying regions of positive  $K_{\text{DP}}$ ,





such as those associated with oblate particles like rain and liquid-coated hail that are expected in 160 convective updrafts.

From a NASA Unified Weather Research and Forecasting (NU-WRF; Peters-Lidard et al., 2015) model simulation, 5-min frequency output were analyzed. The model is configured with a 600 x 600 km outer domain grid centered around Houston with 3-km horizontal grid spacing and, centered within the outer domain, a nested 300 x 300 km inner domain with 500-m horizontal grid spacing (Fig. 1).

- 165 Time steps of 12 and 2 s were used on the outer and inner grids, respectively. The same physics options were used on both grids. The planetary boundary layer parameterization used the Mellor-Yamada-Nakanishi-Niino level 2.5 turbulent kinetic energy scheme (Nakanishi and Niino, 2009). The Morrison et al. (2009) two-moment cloud microphysics scheme was used with a fixed droplet number concentration of 100 cm<sup>-3</sup>, and reflectivity at horizontal polarization ( $Z_{HH}$ ) was calculated
- 170 using the resulting hydrometeor size distributions with temperature-dependent refractive indices following Blahak et al. (2011). The Goddard broadband two-stream radiative transfer scheme was used to calculate radiative fluxes and atmospheric heating rates (Chou and Suarez, 1999, 2001; Matsui et al., 2018a). Model terrain was smoothed from the 30 s and 0.9 km U.S. Geological Survey topography data for both domains, and land cover was mapped from 30 s MODIS land use data.
- 175 Sea surface temperature and atmospheric initial and lateral boundary conditions are obtained from 6 hourly output of the Modern-Era Retrospective Analysis for Research and Applications Version 2 (MERRA-2; Bosilovich et al., 2015). Land surface initial conditions (soil moisture and temperature) were derived from a 6 month spin up (Kumar et al., 2007) of the Noah Land Surface Model (LSM) using the MERRA-Land meteorological forcing (Reichle et al., 2011). The LSM spin-up was con-

180 ducted at the identical grid configuration as that used in the simulation. The simulation was started at 0 UTC on 8 June 2013 and integrated for 24 h.

Additional data shown below are cloud condensation nucleus (CCN) number concentrations retrieved from satellite observations, raindrop size distribution parameters retrieved from NEXRAD measurements, and observed lightning flashes. CCN number concentration and associated supersat-

- uration at cloud base were retrieved from National Polar-orbiting Partnership (NPP) Visible Infrared Imaging Radiometer Suite (VIIRS) cloud top temperature and effective radius with an estimated uncertainty of 30% (Rosenfeld et al., 2016). Rain mass-weighted mean diameter ( $D_m$ ; the fourth moment of the drop number size distribution divided by the third moment) and generalized intercept parameter ( $N_w$ ) (cf. Testud et al., 2001) were retrieved from KHGX data at elevations below
- 190 the melting level following Ryzhkov et al. (2014), with an estimated uncertainty of roughly 5–10% in  $D_m$  and  $log(N_w)$  (Thurai et al., 2012). Collocated rain rate has been retrieved from the specific attenuation A using the R(A) methodology that is most efficient at S-band, with an estimated bias less than 6% (Ryzhkov et al., 2014; Zhang et al., 2017). Lightning flashes were estimated from raw very-high frequency (VHF) signals detected by the LMA. A simple set of heuristics was used to
- 195 cluster VHF sources into discrete flashes, similar to MacGorman et al. (2008): the first ten VHF





sources in a flash are required to be within 0.25 s and 3 km of one another, and each flash is not allowed to exceed 3 seconds in total duration and must be composed of at least 10 VHF sources.

For comparison with the additional observational data, we calculate several further quantities from standard NU-WRF outputs. From rain mixing ratio and number concentration, we use the 200 microphysics scheme assumptions and limitations on rain size distribution properties to calculate rain  $D_m$  and  $N_w$ . From  $D_m$  we further estimate rain  $Z_{DR}$  following Bringi and Chandrasekar (2001, their Eqn. 7.14a). From both  $D_m$  and  $N_w$ , rain  $K_{DP}$  is estimated as described in Appendix A.

#### 2.2 $K_{\rm DP}$ column tracking

205

Convective cells can be objectively identified and tracked using observed or forward-calculated radar variables ( $Z_{\text{HH}}$ ,  $Z_{\text{DR}}$ ,  $K_{\text{DP}}$ ) or model variables such as rain water mixing ratio ( $q_r$ ). From three-dimensional gridded KHGX data for this study,  $K_{\text{DP}}$ -based values were first calculated in each model column following

$$\xi = \int_{z_m}^{z_f} \phi(z)(z - z_m) \,\mathrm{d}z \tag{1}$$

where  $z_m$  and  $z_f$  are the melting and homogeneous freezing heights and  $\phi(z)$  is the value of  $K_{\text{DP}}$ 210 in each column grid cell, similar to the approach taken in van Lier-Walqui et al. (2016). Such a metric favors both the  $\phi(z)$  value and its height. Since hydrometeor size distribution assumptions made in bulk microphysics schemes such as that used in the NU-WRF simulation are not generally adequate to forward simulate fully realistic  $K_{\text{DP}}$  fields (e.g., Ryzhkov et al., 2011), analogous values were calculated from NU-WRF output first using rain water mixing ratio ( $q_r$ ), and then using rain

- 215  $K_{DP}$  estimated as described in Appendix A. In observations, fixed  $z_m$  and  $z_f$  grid cell bottom and top edge values of 4.5 and 9 km, respectively, were estimated from 0 and 12Z soundings at Lake Charles, Louisiana, such that the lowest gridded radar volume was above 0°C and the highest gridded radar volume below  $-40^{\circ}$ C. Similar heights were found from soundings at Corpus Christi, Texas. In NU-WRF output, fixed  $z_m$  and  $z_f$  values of 4.1 and 10 km were taken from the inner domain grid layer
- 220 mean temperature profile. The conclusions of this pilot study are not sensitive to the precise choice of  $z_m$  and  $z_f$  values. However, we note that obtaining accurate time- and space-dependent  $z_m$  and  $z_f$ values from observations could be challenging. It could conceivably be preferable to derive relevant integration limits from observed and forward simulated radar variables in future work.
- Using the two-dimensional fields of ξ values, features were identified and tracked using TrackPY, an open-source Python object tracking toolbox. Whether using observations or model data, regional relative maxima were identified and tracked using TrackPy's predictive tracker with a maximum tracked object velocity of 30 m s<sup>-1</sup> and a "memory" of three frames to allow for splitting or merging cells to be followed. Paths with five or fewer time frames were discarded.





When comparing observed and simulated reflectivity fields from tracked objects, the variables and
tracking parameters described above were subjectively deemed adequate for the purposes of this pilot
study. A more in-depth future study would motivate additional focus on optimizing such choices for
any specific conditions of interest. We also found that tracking performed on ξ values obtained from
simulated q<sub>r</sub> versus simulated K<sub>DP</sub> did not influence study results, likely attributable to the fact that
the K<sub>DP</sub> estimation approach used here is so closely linked to q<sub>r</sub> alone. In the following we focus
only on simulated objects tracked from ξ based on K<sub>DP</sub>.

### 2.3 Long-term cell tracking, introducing TINT

As the study of individual cell cases proceeded it became clear that a long-term study of suitable cell existence was needed. The aforementioned column tracking method did not lend itself well to implementation on large data sets and did not scale well on multi-processor computer clusters.

- 240 Therefore, motivated by this and several other projects, development commenced on a simple-touse and open-source tracking code-base designed specifically for atmospheric data. The TINT Is Not TITAN (TINT) package works directly with the Py-ART grid object in Python and is based on the Thunderstorm Identification, Tracking, Analysis and Nowcasting package (TITAN; Dixon and Wiener, 1993). While TITAN was designed to be used in operational settings and can be challenging
- 245 to configure, TINT is designed to simply take a temporal sequence of grids, a function that renders the 3D grids to a 2D binary mask (for example, a reflectivity threshold at a single level) denoting cell/no cell and returns a Pandas (McKinney (2010)) data-frame containing cell locations and characteristics as a function of time. TINT does not deal with splits or mergers but is thread-safe and pleasantly parallel when radar data is stratified by storm events. TINT uses an N step algorithm to
- 250 associate cells across time steps,  $t_0$  and  $t_1$ :
  - Cells are identified based on minimum thresholds for cell area and field value.
  - Phase correlation is performed in a neighborhood around each cell  $c_i$  to give a estimated translation vector,  $V_i$ , between the  $t_0$  and  $t_1$ . Example images (reflectivity) and their cross-correlation are given in Fig. 2.
- 255 Translation vector estimates are corrected based on prior cell movement.
  - For each identified cell in  $t_0$  the algorithm searches for cells in  $t_1$  at location  $V_i * (t_1 t_0)$ .
  - The Hungarian Algorithm is used to compare candidates and find optimal cell pairing. See Dixon and Wiener (1993) for details.
  - Cell positions are updated, and statistics are recorded.
- 260 New cells are assigned new unique identifiers.





The final product can then be analyzed and plotted either spatially (as tracks, as in Fig. 3) or time series. For the work presented in this study the binary mask was constructed by thresholding each level of the grid at 35 dBZ. If any level for a given latitude/longitude is above that threshold, then the 2D binary mask is marked as being part of a cell. Effectively this constitutes a 2D projection of a 3D binary field using a cascading AND operator.

Three years of KHGX data were processed (2015–2017) using aforementioned techniques, mapped to a Cartesian grid, and saved as CF-complaint NetCDF. Processing was parallelized by event, with events identified based on any radar grid exceeding a minimum reflectivity of 0 dBZ. Job scheduling and management was handled by the Dask library (Dask, 2016). Over the three years, 7 TB of

270 NEXRAD data resulted in 20 MB of cell track data in a CSV file, yielding an efficient data reduction. TINT tracks all convective cells. However, as we are interested in statistics of isolated cells, all TINT analyses presented here include only isolated cells. A cell is considered isolated if it is not connected to any other cell by a contiguous path of grid cells exceeding a field threshold. Isolated cells therefore contain at most one peak field value.

#### 275 3 Results

#### 3.1 Observed case study cell evolution

Starting with the 8 June 2013 case study, Fig. 1 illustrates the routes of three observed features tracked using radar  $K_{\rm DP}$  data, as well as the routes of three simulated features tracked using NU-WRF  $K_{\rm DP}$ . Most tracks are moving roughly northeastward, consistent with boundary layer onshore

- 280 flow conditions. Both observed and simulated tracks are roughly 10–20 km in distance. The only requirement for their selection was that they be isolated convective cells over land within 100 km of KHGX, such that cross-sections exhibited no near neighbors on a subjective basis (as demonstrated below). Relatively few such cells were found in the observations or the simulation owing to lack of isolated cells developing or lack of developing cells remaining isolated, respectively. Although
- the somewhat disorganized convection observed versus simulated differed, the tracking algorithm operating on  $K_{\text{DP}}$  fields yielded satisfactory results for the purposes of this pilot study.

From the three observed tracks numbered 9, 35 and 37 in Fig. 1, Fig. 4 shows the time series of several quantities from track start to track end time, with durations of roughly 30, 55 and 40 min, respectively. The top panels show lightning flash rate within 2.5 km of the track, as well as flash

290 occurrences within 2.5 km horizontally as a histogram in the two dimensions of height and time. Here we see that cells 9 and 35 become electrically active roughly halfway through the track, and flashes are most concentrated between 8 and 10 km, just below the homogeneous freezing level (below the  $-40^{\circ}$ C level). Flash activity in cell 37 remains very weak and is limited to elevations below 8 km. Cells 9 and 37 also show weak flash activity early in their tracks. We note that isolated





295 flashes below the estimated melting level may be artifacts of the processing algorithm used here, which could be refined in future work.

The second row of Fig. 4 shows two quantities calculated following Eqn. 1 where φ(z) is taken as the average value of K<sub>DP</sub> or Z<sub>DR</sub> within 2.5 km of the track location. The resulting ξ values, which we refer to as column strengths, allow a more robust measure of the feature K<sub>DP</sub> and Z<sub>DR</sub>
along the track than provided by the two-dimensional grids used by the tracking algorithm (in other words, a value applicable to the whole feature); we defer optimization of the averaging footprint to future work beyond this pilot study. Since we found that the selected tracks follow relatively

isolated and coherent reflectivity features in both the observations and the simulation (not shown), we also refer to the tracked features interchangeably as cells and columns, with the understanding that the features tracked contain continuous peaks of  $\phi(z)$  at the resolution observed or simulated (otherwise they would not have been tracked) but they do not necessarily correspond to individual isolated updrafts, however that may be defined. As shown in Fig. 4,  $K_{\text{DP}}$  column strength reaches sizable peaks roughly halfway through the track in cells 9 and 35, but cell 37 shows no such peak.

By contrast, all cells exhibit a  $Z_{DR}$  column peak during the first half of their track, and that peak is 310 greatest for cell 37. The robustness of the  $Z_{DR}$  column strength appears indicative that all cells loft sufficiently large raindrops sufficiently far above the melting level to generate a strong  $Z_{DR}$  signal

(e.g., Kumjian et al., 2012; Snyder et al., 2015).

The third row of Fig. 4 shows the median value and inner half of raindrop  $D_m$  values within 2.5 km of the track location at all elevations below the melting level, as well as rain rates retrieved

- at the lowest elevation angle  $(0.5^{\circ})$ . All cells show nearly continuous rain somewhere below  $z_m$  from track start to end, as evidenced by the continuity of  $D_m$  retrievals, but that rain does not reach the lowest scan elevation until at least 10 min after the start of each track. In cells 9 and 35, the near surface rain reaches peak rates during the second half of the track, beginning around the time that the  $K_{\text{DP}}$  column strength reaches its maximum. The considerable spread in surface rain rates
- 320 indicates localized heavy rain that exceeds 100 mm h<sup>-1</sup> in both cells. With the absence of a  $K_{DP}$  column, cell 37 by contrast exhibits weaker and shorter surface rain rate maxima before and after its  $Z_{DR}$  column maximum. The raindrop  $D_m$  is quite variable below all three tracked columns, but the behavior seen in cell 9 appears for many tracked cells (not shown), namely,  $D_m$  increasing with  $K_{DP}$  column strength and then plateauing at relatively high values along with decreased variance across

325 the analyzed volume.

Figures 5 and 6 show four sequential north-south cross sections across the tracks of cells 9 and 37, which remain within similar distances from the radar (roughly 60 to 75 km, see Fig. 1), although cell 37 occurs roughly two hours after cell 9. The cross-section times are indicated with vertical dotted lines in Fig. 4.

330 In cell 9 (Fig. 5), the first cross-section corresponds to the time of peak  $Z_{DR}$  and the last crosssection corresponds to the peak in electrical activity. The columns of Fig. 5 show  $Z_{HH}$ ,  $K_{DP}$ ,  $Z_{DR}$ ,





 $D_m$ , and  $\log(N_w)$ , respectively, with lightning flashes over-plotted in colors that indicate age from current to 10 min old. The first time shows the peak in  $Z_{DR}$  column strength, with elevated positive values (> 1dB) visible almost 3 km above the melting level. At this time the  $K_{DP}$  column is already

- visible, but lightning is absent. Within a rain shaft that is roughly 5–10 km in diameter,  $D_m$  is most commonly greater than 1.8 mm and  $N_w$  is most commonly less than 300 mm<sup>-1</sup> m<sup>-3</sup>. The second time slices capture the peak of the  $K_{DP}$  column strength, concurrent with the beginning of electrical activity. Already at this point the  $Z_{DR}$  column has decreased in height, with some evidence of negative  $Z_{DR}$  values associated with graupel or hail. By the third time, scarcely any  $Z_{DR}$  column
- remains, but the  $K_{\text{DP}}$  column remains visible and the lightning flash rate has intensified. By the fourth and last time, the  $K_{\text{DP}}$  has also largely vanished above the melting level. The lightning flash rate has not yet diminished in strength but flashes have lowered a bit in height (see also Fig. 4). The rain shaft has generally widened, with increasingly greater peak values of both  $D_m$  and  $\log(N_w)$ from track start to end.
- In cell 37 (Fig 6), a vigorous  $Z_{DR}$  column can be seen initially. The diameters of the cell as measured by radar reflectivity signal (roughly 10 km) and the  $Z_{DR}$  column as indicated by peak values (roughly 4 km) are similar to that of cell 9, but the  $Z_{DR}$  values are larger and the rain shaft appears weaker. However, in constrast to cell 9,  $K_{DP}$  enhancements are weak and almost isolated above the melting level at the first time. Retrieved rain parameters do not extend as high as seen in cell 9, con-
- sistent with a lower melting level that could be associated with its later time or more inshore location; owing to the absence of adequate meteorological observations, as discussed in Sect. 4, we assumed fixed  $z_m$  and  $z_f$  values here. Within the rain shaft initially, occurrences of  $D_m$  greater than 2 mm are less common than in cell 9 and  $N_w$  values are notably smaller. Enhanced  $K_{\text{DP}}$  subsequently descends over the course of the four radar volumes shown (roughly 15 min). At the third time, there
- is a sharp and localized peak of enhanced  $K_{\rm DP}$  roughly 1 km below the melting level, as in cell 9 at the third time shown, but the diameter of the region where  $K_{\rm DP}$  exceeds  $0.5^{\circ}$  km<sup>-1</sup> (often wider than 5 km) is roughly twice as great as that in cell 9 (roughly 3 km throughout). Perhaps related to the absence of a pronounced  $K_{\rm DP}$  column above the melting level, electrical activity remains weak at all times in cell 37. The evolving rain shaft remains generally weaker than in cell 9, with generally

360 smaller  $D_m$  and  $\log(N_w)$ .

A common pattern in observed tracked cells in this particular case study is that first  $Z_{DR}$  column strength peaks, followed by  $K_{DP}$  column strength, and then lightning activity. However, cell 37 and other tracked cells that initiate northeast (downwind) of Houston exhibit the following deviations from that pattern: a greater leading  $Z_{DR}$  peak followed by neglibigible  $K_{DP}$  peaks above

365 the melting level and much weaker lightning activity. According to satellite retrievals (see Fig. 1), the updraft cells in this latter class appear to exist within a region of generally more elevated aerosol concentration than is found upwind or adjacent to the Houston plume flow. All else being equal, enhanced aerosol could at least initially lead to larger and fewer raindrops (e.g., Storer and





van den Heever, 2013), potentially consistent with stronger  $Z_{DR}$  and weaker  $K_{DP}$  rain contributions in cell 37, whereas cleaner conditions could at least initially lead to a more active warm rain process with more numerous and therefore smaller raindrops. Rain shaft retrievals do show generally smaller  $N_w$  below cell 37, consistent with fewer raindrops. There is some hint that  $D_m$  at the top of the cell 37 rain shaft at intermediate times shown may be larger than at the top of the cell 9 rain shaft, but  $D_m$ values exceeding 2 mm are clearly more common in cell 9, which is producing consistently greater

- 375 rain rates than cell 37 (see Fig. 4). We note that although electrification mechanisms and lightning production are not well understood, increased aerosol concentrations have been more commonly associated with increased rather than decreased electrical activity (Murray, 2016, and references therein). Substantially more complex coupled microphysical and dynamical pathways could also be primary contributors to both  $Z_{DR}$  and  $K_{DP}$  column evolution (e.g., Ryzhkov et al., 2011; Kumjian
- et al., 2014a; Snyder et al., 2015, and references therein). Owing to the short cell life cycles here, even such basic and well observed factors as height trends below the melting level (e.g., Kumjian and Prat, 2014) could be more indicative of time trends than quasi-steady properties. We defer robust analysis to future work, which would also need to address the role of meteorology, topography, observability (distance from radar), and other factors. Here we make the limited conclusion that such patterns in observed convective cells around Houston can be effectively identified and analyzed in

#### 3.2 Simulated case study cell evolution

an integrated fashion.

Fig. 7 shows time series from three cells tracked from NU-WRF output in the same format as shown in Fig. 4 from observations. The times from track start to end for simulated cells 89, 116, and
188 are each roughly 25–30 min. The isolated cell tracks in the simulation are generally shorter than the isolated cells tracks in observations. Although many longer-lived cells were tracked in the simulation, they tended not to remain isolated. These general differences did not, however, hinder some basic comparisons of observed and simulated isolated cells as follows.

- Since we did not attempt forward simulation of lightning flashes here, the first row of panels 395 in Fig. 7 instead shows the 95th percentile of vertical wind speed (w) between the melting and freezing levels within 2.5 km of the track, which could potentially be retrieved from additional radar measurements in a future field campaign (e.g., Collis et al., 2013; North et al., 2017), and the column strength of supercooled  $q_r$ . The second row of panels in Fig. 7 shows the time series of column strengths of  $K_{\text{DP}}$  and  $Z_{\text{DR}}$  estimated from simulated  $D_m$  and  $N_w$  (see Sect. 2). The third
- 400 row of panels in Fig. 7 shows the median value and inner half of raindrop  $D_m$  values within 2.5 km of the track location at all elevations below the melting level, as well as rain rate at the surface.

All three simulated cells show surface rain beginning shortly after the track start and either declining or continuing after the track end time, as in observed cells, but the peak of median rain rates tends to be at least 5–10 times weaker than retrieved beneath observed cells. Despite weaker precipitation,





- 405 simulated raindrop  $D_m$  values are often near 2.5 mm, greater than observed values often near 2 mm; we note that a fixed droplet number concentration of 100 cm<sup>-3</sup> was used in the simulation (see Sect. 2) owing to the absence of aerosol size distribution measurements, which are discussed further in Sect. 4. Simulated cells show peaks of  $q_r$ ,  $K_{\text{DP}}$ , and  $Z_{\text{DR}}$  column strength roughly collocated in time, consistent with the simplified use of supercooled rain properties to estimate the polarimetric
- 410 quantities. Simulated  $Z_{DR}$  column strengths are a bit greater than those observed, with variable peak values of 1–2 dB km in each cell that are reached sometime during the inner half of the track duration. Simulated  $K_{DP}$  column strengths are by contrast roughly an order of magnitude weaker than observed, consistent with the rain-based estimate of  $K_{DP}$  resulting in underestimates aloft (Ryzhkov et al., 2011) that are amplified by height weighting (Eqn. 1). Whereas individual well-defined  $Z_{DR}$
- 415 peaks tend to consistently lead  $K_{\text{DP}}$  peaks in observations when the latter is present, all column strength peaks in the simulated cells tend to be coincident with one another, as well as with local peaks in the strongest colocated w values to a somewhat lesser degree. Simulated columns tend to show more peakedness than the w statistic, indicating that phase in this particular case study simulation is not as tightly controlled by local updraft strength as might be expected; future work could
- 420 examine whether a differing w statistic than shown here is more correlated. Simulated surface rain rate peaks with or shortly after the  $Z_{DR}$  column strength, similar to observed cells.

Figures 8 and 9 show north-south cross sections along the tracks of simulated cells 116 and 188 at the times indicated with vertical dotted lines in Fig. 7. In overall structure, both simulated cells are 5–10 km in diameter and their 45 dBZ echoes reach at least 6–8 km, similar to the observed cells

shown in Figs. 5 and 6. Simulated  $K_{\rm DP}$  structures appear generally similar to observed insofar as a single column is found at the center of each cell cross-section, but the simulated columns exhibit substantially higher peak values at column center (>1.75 ° km<sup>-1</sup>) and do not decrease as rapidly with height. In the case of cell 116, higher peak values abruptly decrease just below 6 km where rain appears to be rapidly frozen. The narrowness of simulated  $q_r$  columns is more similar to observed cell 9 than 37.

In contrast to observations, there is a strong increase in simulated rain  $Z_{DR}$  and  $D_m$  from the melting level to the surface, an expected signature of raindrop size sorting that can be severely overestimated in two-moment bulk microphysics schemes such as that used here (Kumjian and Ryzhkov, 2010, 2012). The presence of mixed-phase particles complicates interpretation above the melting

- 435 level. It can nonetheless be noted that simulated rain  $Z_{DR}$  greater than 2 dB reaches 5 km, as in both observed cells shown. Rain size distribution parameters shown from the mixed-phase region of the simulation, where they cannot be retrieved, indicate that weaker  $Z_{DR}$  approaching the homogeneous freezing level in cell 116 is associated with increasing rather than decreasing  $N_w$ . Simulated cells commonly exhibit isolated and narrow regions of high  $Z_{DR}$  at supercooled temperatures on their
- 440 north and south flanks, similar to a less prominent feature on the north flank of observed cell 9.





455

# 3.3 Long-term statistics of cell occurrence

Using TINT as described in Section 2 enables a long-term statistical analysis of isolated cell occurrence from KHGX observations. From a three-year climatology, Fig. 10 shows the total number of isolated cells that initiated as a function of month of the year. There is a pronounced period of

- enhanced occurrence between June and September (approximately the summer months). This raises the question: is the increase in cells over the enhanced period due to an increased density of cells on a given day or more days with convective initiation events? Figure 11 shows the percentage of days in a given month with an initiation event within range of the KHGX radar. There is only a weak seasonal cycle, ranging from a 35% chance of observing an isolated cell on a given day in December
  to just over 50% on a given day in July, indicating that the abrupt increase seen in June in Fig. 10
  - can be attributed to an increase in cell population density.

Focusing on the enhanced occurrence season, Fig. 12 shows the number of cells that initiate in that season as a function of time of day. The peak at a local time of 1 PM is consistent with a strong diurnal forcing. Furthermore, the lack of any apparent overnight maximum gives us confidence that we are effectively filtering out large-scale systems that have a nocturnal maximum (Nesbitt and Zipser, 2003).

The 2013 case study investigated above focused on observing the microphysical and dynamical evolution of convective cells in a Lagrangian frame of reference. When investigating the feasibility of deploying an agile radar system to Houston an important question arises: as a function of the radar's

- 460 unambiguous maximum range, how many cells will the radar see from initiation to dissipation? That is, how many full cell life cycles might the radar system collect? Figure 13 shows the total number of cells as a function of the cell lifetime that would occur within 70, 150 and 200 km of a radar placed at the KHGX site during the enhanced occurrence period. The totals are 441, 2442 and 4834 cells, respectively. If the assumption is made that the three years studied are typical, we
- 465 could therefore expect to see roughly 150, 800 and 1600 full life cycles in a single June through September deployment for a 70-km (e.g., X-Band), 150-km (C-Band) and 200-km (S-Band) radar range, scaling roughly with range area as would be expected if track density were geographically uniform.

We lastly investigate the initiation location and propagation direction of isolated cells, with relevance for both configuring multi-site deployments and contextualizing measurements. Cells preferentially initiate in the northwest sector of the KHGX range (Fig. 14). For each cell within and outside of the enhanced occurrence period a vector was calculated by linking the location of dissipation to the location of initiation, giving a mean propagation direction. Figure 15 shows the directional cumulative distribution including the enhanced occurrence period versus the remainder of the year.

475 During the enhanced period the cells are dominantly propagating slightly east of due southerly, in contrast to south-south westerly during the rest of the year. This indicates that most convective cells in the enhanced period might be expected to flow from a cleaner Gulf of Mexico air mass into a more





polluted Houston air mass. This quirk of climatology suggests that the enhanced period convection lends itself well to studying the impact of aerosols on isolated, precipitating convective cells.

## 480 4 Conclusions and discussion

The comparison of tracked cells from Houston NEXRAD observations and a NU-WRF simulation demonstrates the potential value of polarimetric weather radar observations for systematically observing and improving the understanding and simulation of convective cell physics. Factors related to the meteorological and aerosol environment, such as the structure of rain size distribution param-

- eters below the melting level, are particularly well suited to analysis using such data. Above the melting level, further investigation of the microphysical properties controlling  $K_{\rm DP}$  and  $Z_{\rm DR}$  signatures is likely to yield additional quantitative constraints on simulation physics; comparing observations with forward-simulated values from well-observed case studies is likely to yield substantial progress, especially using an integrative approach that also considers rain properties below the melt-
- 490 ing level and overall cell structural evolution. Future simulations could employ bin microphysics or other approaches to avoid errors associated with sedimentation or hydrometeor size distribution shape, as well as mixed-phase particle representation to improve forward simulation of polarimetric signatures (e.g., Ryzhkov et al., 2011; Kumjian et al., 2014a; Snyder et al., 2017a; Matsui et al., 2018b). Forward simulation of lightning flash rates (e.g., Barthe et al., 2010; Basarab et al., 2015)
- 495 may be simultaneously compared with collocated LMA observations to study the correlations of updraft physics and flash rate signatures such as those shown in Fig. 4.

However, cell tracking in both KHGX observations and a simulation also demonstrates the potential value of improved spatiotemporal resolution that could be achieved using mobile research radars. Isolated cell cores are relatively poorly resolved and their evolution is rapid compared with

- 500 the KHGX operational volume scan rate.  $K_{\text{DP}}$  and  $Z_{\text{DR}}$  columns associated with updraft cores rise and fall within 10–15 min time spans, as shown in Figs. 4 and 7. Similar conclusions have been reached in past studies (e.g., Loney et al., 2002). Future simulations can obtain arbitrarily higher spatial resolution and output timing whereas radar measurements are subject to cell distance from the radar and, in this study, the fixed scanning strategies of the operational weather radar. For the
- 505 purposes of studying isolated convection around Houston, we conclude that substantial value could be added by mobile research radars that could achieve higher resolution and faster scan rates (e.g., Isom et al., 2013; Pazmany et al., 2013; Snyder et al., 2013; Kumjian et al., 2014b). Stein et al. (2015) demonstrate the value of applying a statistical approach to convective cells that are tracked in simulations and in radar observations using an adaptive rapid scan strategy. Sufficient radar resources
- 510 to make wind vector retrievals (e.g., Collis et al., 2013; North et al., 2017) could supply observations adequate to statistically study the relationship of cell microphysics and updraft strength.





A statistical analysis of three years of Houston KHGX data indicates that there is a period of enhanced isolated convection from June through September, when the number of cells per day dramatically increases, indicating a most favorable season for studying such cells. During this period

- 515 approximately half of days can be expected to experience cell formation, and isolated cell number follows a strong diurnal cycle with a peak at 1 PM local time. During the June through September period, a hypothetical C-Band radar with a range of 150 km deployed to the site of the KHGX could be expected to observe the full life cycle of roughly 800 cells within range of the radar according to the statistics collected in our three-year sample. Finally, cells observed would have a dominant
- 520 propagation vector just west of southerly, indicating that cells forming along the shoreline would likely experience aerosol perturbations corresponding to their proximity to emission sources.

The demanding objectives of a box flux closure experiment (Rosenfeld et al., 2014) would require meteorological measurements at high spatiotemporal resolution at all domain boundaries, but even for the more limited study of updraft physics investigated here as an amendment to such a campaign,

- 525 we note that routine meteorological data are lacking in the Houston region. The nearest operational soundings are at Lake Charles and Corpus Cristi, roughly 200 km to the northeast and 300 km to the southwest, respectively. Obtaining soundings during convective activity, ideally at more than one site within the KHGX domain, would provide a foundation required to establish atmospheric structure, which is of first-order importance to convective cell development. To the extent that improving model
- 530 physics is an objective, it would be imperative to collect observations that are adequate to demonstrate that simulations are reproducing basic properties of atmospheric structure. The capability of state-of-the-art regional models to reproduce basic atmospheric structure should not be assumed a given even when using an assimilation-informed data set for inputs at domain boundaries. Owing to the relatively rapid evolution of the diurnal boundary layer properties with time and distance from
- 535 the coastline under the target conditions of onshore flow, ground-based in situ and remote-sensing measurements capable to establish boundary layer height and water vapor mixing ratio would also add great value.

Finally, in situ measurements remain the most robust means of observing cloud-active aerosol properties from surface to mid-free-troposphere. At a minimum, surface measurements of boundary

- 540 layer CCN spectra and total aerosol number concentration measurements from ideally more than one location in the KHGX domain would allow a means of constraining at least boundary-layer fields. To avoid the challenge and expense of a long aircraft campaign, past aircraft measurements from recent field campaigns in the Houston region could provide statistical guidance on expected discontinuities at the boundary layer top (e.g., Lance et al., 2009). Measurement of INPs from at least one surface
- site would add substantial value; we are aware of no past INP measurements in the Houston region.





# Appendix A: Calculation of K<sub>DP</sub>

Specific differential phase  $K_{\text{DP}}$  for liquid drops at S-band is calculated from the simulated massweighted diameter  $D_m$  and intercept  $N_w$  assuming an exponential drop size distribution

$$N(D) = N_w exp(-4D/D_m),\tag{A1}$$

550 consistent with model microphysics, and

$$K_{\rm DP} = N_w f(D_m),\tag{A2}$$

where

$$\log_{10} f(D_m) = -5.98 + 6.64 \log_{10} D_m - 1.28 (\log_{10} D_m)^2, \tag{A3}$$

with  $D_m$  in mm,  $N_w$  in m<sup>-3</sup> mm<sup>-1</sup>, and  $K_{DP}$  in ° km<sup>-1</sup>. Eqns. A2–A3 are derived for the radar 555 wavelength 11 cm.

## Appendix B: Code availability

Py-ART is available from http://arm-doe.github.io/pyart/. TrackPy is available from http://soft-matter.github.io/trackpy/v0.3.2/. TINT is available from https://github.com/openradar/TINT.

560 NU-WRF is available from http://nuwrf.gsfc.nasa.gov/.

## Appendix C: Data availability

KHGX NEXRAD data were downloaded from the National Climatic Data Center (NOAA, 1991). Lightning data, rain properties derived from NEXRAD data, and NU-WRF output are available upon request. TINT cell track data is available on request.

- 565 Author contributions. D. Rosenfeld provided CCN retrievals and selected suitable case study dates. M. van Lier-Walqui prepared KHGX data with assistance from S. Giangrande and S. Collis. M. Picel is the lead software engineer for TINT and was assisted by R. Jackson. T. Matsui and A. Fridlind prepared the NU-WRF simulation and outputs. A. Ryzhkov and P. Zhang provided rain retrievals. D. MacGorman and R. Weitz provided Lightning Mapping Array data. M. van Lier-Walqui, A. Fridlind, S. Collis, M. Picel and R. Jackson prepared
- 570 analyses and figures. A. Fridlind, S. Collis, M. van Lier-Walqui, and X. Li prepared the manuscript.

Acknowledgements. This work was partially supported by the Office of Science (BER), U.S. Department of Energy, under agreement DE-SC0006988, and by the NASA Radiation Sciences Program. NU-WRF is supported by the NASA Modeling, Analysis and Prediction (MAP) program. Resources supporting this work were





- provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing
  (NAS) Division at Ames Research Center and NASA's Center for Climate Simulation (NCCS) at Goddard
  Space Flight Center. The contribution of Scott Collis through Argonne National Laboratory was supported by
  the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under
  contract DE-AC02-06CH11357. We gratefully acknowledge the computing resources provided on Blues and
  Bebop, a high-performance computing cluster operated by the Laboratory Computing Resource Center at Ar-
- 580 gonne National Laboratory. Johannes Quaas co-organized the Aerosol-Cloud-Precipitation-Climate initiative workshop at the NASA Goddard Institute for Space Studies in April 2015, where this work was conceived with the additional participation of Jiwen Fan, Corinna Hoose, Philip Stier, Bastiaan van Diedenhoven, and Shuguang Wang. A companion regional model intercomparison study of another case study date is being led by Sue van den Heever (http://acpcinitiative.org).





# 585 References

- Ackerman, T. P. and Stokes, G. M.: The Atmospheric Radiation Measurement Program, Physics Today, 56, 38–44, 2003.
- Ackerman, A.S., Fridlind, A.M., Grandlin, A., Dezitter, F., Weber, M., Strapp, J.W., and Korolev, A.: High ice water content at low radar reflectivity near deep convection Part 2. Evaluation of microphysical path-
- ways in updraft parcel simulations, Atmos. Chem. Phys., 15, 11729–11751, doi:10.5194/acp-15-11729-2015, 2015.
  - Barthe, C., Deierling, W., and Barth, M. C.: Estimation of total lightning from various storm parameters: A cloud-resolving model study, J. Geophys. Res., 115(D24), D24202, doi:10.1029/2010JD014405, 2010.
- Basarab, B. M., Rutledge, S. A., and Fuchs, B. R.: An improved lightning flash rate parameterization developed
- 595 from Colorado DC3 thunderstorm data for use in cloud-resolving chemical transport models, J. Geophys. Res., 120: 9481–9499, doi:10.1002/2015JD023470, 2015.
  - Blahak, U., Zeng, Y., and Epperlein, D.: Radar forward operator for data assimilation and model verification for the COSMO-model: Preprint of 35th Conf. on Radar Meteorology, Amer. Meteor. Soc., Pittsburgh, PA, P10.119, 2011.
- 600 Bosilovich, M. G., Akella, S., Coy, L., Cullather, R., Draper, C., Gelaro, R., Kovach, R., Liu, Q., Molod, A., Norris, P., Wargan, K., Chao, W., Reichle, R., Takacs, L., Vikhliaev, Y., Bloom, S., Collow, A., Firth, S., Labow, G., Partyka, G., Pawson, S., Reale, O., Schubert, S., and Suarez, M.: MERRA-2: Initial Evaluation of the Climate, NASA/TM-2015-104606, in series: Technical Report Series on Global Modeling and Data Assimilation, vol. 43, 145 pg, 2015.
- 605 Bringi, V. N., Liu, L., Kennedy, P. C., Chandrasekar, V., and Rutledge, S. A.: Dual multiparameter radar observations of intense convective storms: The 24 June 1992 case study, Meteorol. Atmos. Phys., 59, 3–31, 1996.
  - Bringi, V. N., and Chandrasekar, V.: Polarimetric Doppler Weather Radar: Principles and Applications, Cambridge University Press, Cambridge, UK, 636 pp., 2001.
- 610 Chou, M.-D., and Suarez, M. J.: A solar radiation parameterization for atmospheric studies, NASA//TM-1999-10460, vol. 15, 38 pp.
  - Chou, M.-D., and Suarez, M. J.: A thermal infrared radiation parameterization for atmospheric studies, NASA/TM-2001-104606, in series: Technical Report Series on Global Modeling and Data Assimilation, vol. 19, 55 pp.
- 615 Collis, S., Protat, A., May, P. T., and Williams, C.: Statistics of storm updraft velocities from TWP-ICE including verification with profiling measurements, J. Appl. Meteorol. Climatol., 52(8), 1909–1922, doi:10.1175/JAMC-D-12-0230.1, 2013.

Dask Development Team (2016). Dask: Library for dynamic task scheduling URL https://dask.org

- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S.,
   Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts
  - on climate, Proc. Nat. Acad. Sci., 107(25), 11217–11222, doi:10.1073/pnas.0910818107, 2010.
    Dixon, M., and Wiener, G: TITAN: Thunderstorm identification, tracking, analysis and nowcasting—A radar-based methodology, J. Atmos. Ocean. Tech., 10, 785–797, 1993.





650

- Donner, L. J., O'Brien, T. A., Rieger, D., Vogel, B., and Cooke, W. F.: Are atmospheric updrafts a key to
   unlocking climate forcing and sensitivity?, Atmos. Chem. Phys., 16(20), 12983–12992, doi:acp-16-12983-2016, 2016.
  - Du, R., Du, P., Lu, Z., Ren, W., Liang, Z., Qin, S., Li, Z., Wang, Y., and Fu, P.: Evidence for a missing source of efficient ice nuclei, Sci. Rep., 7, 39673, doi:10.1038/srep39673, 2017.
- Fan, J., Leung, L. R., Rosenfeld, D., Chen, Q., Li, Z., Zhang, J., and Yan, H.: Microphysical effects determine macrophysical response for aerosol impacts on deep convective clouds, Proc. Nat. Acad. Sci., 110, E4581–E4590, doi:10.1073/pnas.1316830110, 2013.
  - Fridlind, A. M., Ackerman, A. S., Chaboureau, J.-P., Fan, J., Grabowski, W. W., Hill, A. A., Jones, T. R., Khaiyer, M. M., Liu, G., Minnis, P., Morrison, H., Nguyen, L., Park, S., Petch, J. C., Pinty, J. P., Schumacher, C., Shipway, B. J., Varble, A. C., Wu, X., Xie, S., and Zhang, M.: A comparison of
- 635 TWP-ICE observational data with cloud-resolving model results, J. Geophys. Res., 117, no. D5, D05204, doi:10.1029/2011JD016595, 2012.
- Fridlind, A. M., Ackerman, A. S., Grandin, A., Dezitter, F., Weber, M., Strapp, J. W., Korolev, A. V., and Williams, C. R.: High ice water content at low radar reflectivity near deep convection Part 1: Consistency of in situ and remote-sensing observations with stratiform rain column simulations, Atmos. Chem. Phys., 15, 11713–11728, doi:10.5194/acpd-15-11713-2015, 2015.
- Fridlind, A. M., Li, X., Wu, D., van Lier-Walqui, M., Ackerman, A. S., Tao, W.-K., McFarquhar, G. M., Wu, W., Dong, X., Wang, J., Ryzhkov, A., Zhang, P., Poellot, M. R., Neumann, A., and Tomlinson, J. M.: Derivation of aerosol profiles for MC3E convection studies and use in simulations of the 20 May squall line case, Atmos. Chem. Phys., 17, 5947–5972, doi:10.5194/acp-17-5947-2017, 2017.
- 645 Giangrande, S. E., McGraw, R., and Lei, L.: An application of linear programming to polarimetric radar differential phase processing, J. Atmos. Oceanic Tech., 30, 1716–1729, 2013.
  - Hallett, J., and Mossop, S. C.: Production of secondary ice particles during the riming process, Nature, 249, 26–28, 1974.
  - Helmus, J. J., and Collis, S. M.: The Python ARM Radar Toolkit (Py-ART), a library for working with weather radar data in the Python programming language, J. Open Res. Software, 4, doi:10.5334/jors.119, 2016.
  - Herbert, R. J., Murray, B. J., Dobbie, S. J., and Koop, T.: Sensitivity of liquid clouds to homogenous freezing parameterizations, Geophys. Res. Lett., 42(5), 1599–1605, doi:10.1002/(ISSN)1944-8007, 2015.
    - Homeyer, C. R., and Kumjian, M. R.: Microphysical characteristics of overshooting convection from polarimetric radar observations, J. Atmos. Sci., 72, 870–891, 2015.
- 655 Hubbert, J., Bringi, V. N., Carey, L. D., and Bolan, S.: CSU-CHILL polarimetric radar measurements from a severe hail storm in eastern Colorado, J. Appl. Meteorol., 37, 749–775, 1998.
  - Climate Change: The IPCC Scientific Assessment, in: Report Prepared for Intergovernmental Panel on Climate Change by Working Group I, edited by: Houghton, J. T., Jenkins, G. J., and Ephraums, J. J., Cambridge University Press, Cambridge, UK, 410 pp., 1990.
- 660 Climate Change 2013: The Physical Science Basis, in: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 1535 pp., 2013.





Isom, B., Palmer, R., Kelley, R., Meier, J., Bodine, D., Yeary, M., Cheong, B.-L., Zhang, Y., Yu, T.-Y., and
 Biggerstaff, M. I.: The atmospheric imaging radar: Simultaneous volumetric observations Using a phased
 array weather radar, J. Atmos. Oceanic Technol., 30(4), 655–675, doi:10.1175/JTECH-D-12-00063.1, 2013.
 Jakob, C., and Tselioudis, G.: Objective identification of cloud regimes in the tropical western Pacific, Geophys.
 Res. Lett., 30, 2082, doi:10.1029/2003GL018367, 2003.

- Johnson, J. T., Mackeen, P. L., Witt, A., Mitchell, D. W., Stmpf, G. J., Eilts, M. D., and Thomas, K. W.: The
  storm cell identification and tracking algorithm: An enhanced WSR-88D algorithm, Weath. Forecast., 13, 263–276, 1998.
  - Kiehl, J. T.: Twentieth century climate model response and climate sensitivity, Geophys. Res. Lett., 13, 263– 276, 1998.
  - Kumar, S. V., Peters-Lidard, C. D., Eastman, J. L., and Tao, W.-K.: An integrated high-resolution hydrometeorological modeling testbed using LIS and WRF, Environ. Modell. Softw., 23, 169–181, 2007.
- Kumjian, M. R., and Ryzhkov, A. V.: Polarimetric signatures in supercell thunderstorms, J. Appl. Meteorol. Climatol., 47(7), 1940–1961, doi:10.1175/2007JAMC1874.1, 2008.
  - Kumjian, M. R., and Ryzhkov, A. V.: The impact of evaporation on polarimetric characteristics of rain: Theoretical model and practical implications, J. Appl. Meteorol. Climatol., 49(6), 1247–1267, doi:10.1175/2010JAMC2243.1, 2010.
  - Kumjian, M. R., and Ryzhkov, A. V.: The impact of size sorting on the polarimetric radar variables, J. Atmos. Sci., 69(6), 2042–2060, doi:10.1175/JAS-D-11-0125.1, 2012.
  - Kumjian, M. R., Ganson, S. M., and Ryzhkov, A. V.: Freezing of raindrops in deep convective updrafts: A microphysical and polarimetric model, J. Atmos. Sci., 69(12), 3471–3490, doi:10.1175/JAS-D-12-067.1, 2012.
- 685 2

690

675

680

- Kumjian, M. R., and Prat, O. P.: The impact of raindrop collisional processes on the polarimetric radar variables,
  J. Atmos. Sci., 71(8), 3052–3067, doi:10.1175/JAS-D-13-0357.1, 2014.
- Kumjian, M. R., Khain, A. P., Benmoshe, N., Ilotoviz, E., Ryzhkov, A. V., and Phillips, V. T. J.: The anatomy and physics of z<sub>DR</sub> columns: Investigating a polarimetric radar signature with a spectral bin microphysics model, J. Appl. Meteorol. Climatol., 53(7), 1820–1843, 2014.
- Kumjian, M. R., Rutledge, S. A., Rasmussen, R. M., Kennedy, P. C., and Dixon, M.: High-resolution polarimetric radar observations of snow-generating cells, J. Appl. Meteorol. Climatol., 53(6), 1636–1658, doi:10.1175/JAMC-D-13-0312.1, 2014.
- Ladino, L. A., Korolev, A., Heckman, I., Wolde, M., Fridlind, A. M., and Ackerman, A. S.: On the role of
   ice-nucleating aerosol in the formation of ice particles in tropical mesoscale convective systems, Geophys.
   Res. Lett., 44, no. 3, 1574–1582, doi:10.1002/2016GL072455,2017.
  - Lakshmanan, V., and Smith, T.: An objective method of evaluating and devising storm-tracking algorithm, Weath. Forecast., 25, 701–709, 2010.
  - Lance, S., Nenes, A., Mazzoleni, C., Dubey, M. K., Gates, H., Varutbangkul, V., Rissman, T. A., Murphy, S. M.,
- 700 Sorooshian, A., Flagan, R. C., Seinfeld, J. H., Feingold, G., and Jonsson, H. H.: Cloud condensation nuclei activity, closure, and droplet growth kinetics of Houston aerosol during the Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS), J. Geophys. Res., 114, D00F15, doi:10.1029/2008JD011699, 2009.





Lawson, R. P., Woods, S., and Morrison, H.: The microphysics of ice and precipitation development in tropical
 cumulus clouds, J. Atmos. Sci., 72(6), 2429–2445, doi:10.1175/JAS-D-14-0274.1, 2015.

- Limpert, G., Houston, A., and Lock, N.: The advanced algorithm for tracking objects (AALTO), Meteor. Apps., 22, 694–704, doi:10.1002/met.1501, 2015.
- Loney, M. L., Zrnic, D. S., Straka, J. M., and Ryzhkov, A. V.: Enhanced polarimetric radar signatures above the melting level in a supercell storm. J. Appl. Meteorol., 41, 1179–1194, 2002.
- 710 MacGorman, D. R., Rust, W. D., Schuur, T. J., Biggerstaff, M. I., Straka, J. M., Ziegler, C. L., Mansell, E. R., Bruning, E. C., Kuhlman, K. M., Lund, N. R., Biermann, N. S., Payne, C., Carey, L. D., Krehbiel, P. R., Rison, W., Eack, K. B., and Beasley, W. H.: TELEX: The Thunderstorm Electrification and Lightning Experiment, Bull. Amer. Meteorol. Soc., 89, 997–1013, 2008.
- Mather, J. H., and Voyles, J. W.: The ARM climate research facility, Bull. Amer. Meteorol. Soc., 94(3), 377– 392, 2013.
- Matsui, T., Zhang, S. Q., Tao, W.-K., Lang, S., Ichoku, C., and Peters-Lidard, C.: Impact of radiation frequency, precipitation radiative forcing, and radiation column aggregation on convection-permitting west African monsoon simulations, Clim. Dyn., submitted, 2018.
- Matsui, T., Dolan, B., Rutledge, S., Tao, W.-K., Iguchi, T., Barnum, J., and Lang, S.: POLARRIS: Polarimetric
   Radar Retrievals and Instrumental Simulator, J. Geophys. Res., submitted, 2018.
  - Mauritsen, T., and Stevens, B.: Missing iris effect as a possible cause of muted hydrological change and high climate sensitivity in models, Nature Geosci., 8(5), 346–351, doi:10.1038/ngeo2414, 2015.

McKinney, W., 2010. Data Structures for Statistical Computing in Python, in: Walt, S. van der, Millman, J. (Eds.), Proceedings of the 9th Python in Science Conference. pp. 51–56.

725 Morrison, H., Thompson, G., Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: Comparison of one- and two-moment schemes, Mon. Wea. Rev., 137, 991–1007, 2009.

Murray, L. T.: Lightning NOx and impacts on air quality: Curr. Poll. Rep., 115-133, doi:10.1007/s40726-016-0031-7, 2016.

- 730 Nakanishi, M., and Niino, H.: Development of an improved turbulence closure model for the atmospheric boundary layer, J. Meteor. Soc. Japan, 87,895–87,912, 2009.
  - Nesbitt, S.W. and E.J. Zipser, 2003: The Diurnal Cycle of Rainfall and Convective Intensity according to Three Years of TRMM Measurements. J. Climate, 16, 1456–1475, doi:10.1175/1520-0442-16.10.1456

NOAA National Weather Service (NWS) Radar Operations Center: NOAA Next Generation Radar (NEXRAD)

735 Level 2 Base Data, NOAA National Centers for Environmental Information, doi:10.7289/V5W9574V accessed, 1991.

NOAA: WSR-88D meteorological observations, Part C, WSR-88D products and algorithms, Federal Meteorological Handbook 11, Report FCM-H11C-2017, Office of the Federal Coordinator for Meteorological Services and Supporting Research, 394 pp., 2017.

740 North, K. W., Oue, M., Kollias, P., Giangrande, S. E., Collis, S. M., and Potvin, C. K.: Vertical air motion retrievals in deep convective clouds using the ARM scanning radar network in Oklahoma during MC3E, Atmos. Meas. Tech., 10(8), 2785–2806, doi:10.5194/amt-10-2785-2017, 2017.





775

Orville, R. E., Cullen, M. R., Rodeheffer, D., Krehbiel, P. R., and Rison, W.: The Houston Lightning Mapping Array: Installation, operation, and preliminary results, AGU Fall Meeting Abstracts, 2012.

- 745 Pazmany, A. L., Mead, J. B., Bluestein, H. B., Snyder, J. C., and Houser, J. B.: A mobile rapid-scanning X-band polarimetric (RaXPol) Doppler radar system, J. Atmos. Ocean. Technol., 30(7), 1398–1413, doi:10.1175/JTECH-D-12-00166.1, 2013.
  - Peters-Lidard, C. D., Kemp, E. M., Matsui, T., Santanello, Jr., J. A., Kumar, S. V., Jacob, J. P., Clune, T., Tao, W.-K., Chin, M., Hou, A., Case, J. L., Kim, D., Kim, K.-M., Lau, W., Liu, Y., Shi, J., Starr, D., Tan, Q.,
- 750 Tao, Z., Zaitchik, B. F., Za- vodsky, B., Zhang, S. Q., and Zupanski, M.: Integrated modeling of aerosol, cloud, precipitation and land processes at satellite-resolved scales, Environ. Modell. Softw., 67, 149–159, doi:10.1016/j.envsoft.2015.01.007, 2015.
  - Pope, M., Jakob, C., and Reeder, M. J.: Objective classification of tropical mesoscale convective systems, J. Clim., 22, 578–5808, 2009.
- 755 Pruppacher, H. R., and Klett, J. D.: Microphysics of Clouds and Precipitation, 2nd Edition, Kluwer Academic Publishers, 954 pp., 1997.
  - Reichle, R. H., Koster, R. D., De Lannoy, G. J. M., Forman, B. A., Liu, Q., Mahanama, S. P. P., and Toure, A.: Assessment and enhancement of MERRA land surface hydrology estimates, J. Clim., 24, 6322–6338, doi:10.1175/JCLI-D-10-05033.1, 2011.
- 760 Rosenfeld, D., Andreae, M. O., Asmi, A., Chin, M., de Leeuw, G., Donovan, D. P., Kahn, R., Kinne, S., Kivekäs, N., Kulmala, M., Lau, W., Schmidt, K. S., Suni, T., Wagner, T., Wild, M., and Quaas, J.: Global observations of aerosol-cloud-precipitation-climate interactions, Rev. Geophys., 52(4), 750–808, doi:10.1002/2013RG000441, 2014.
- Rosenfeld, D., Zheng, Y., Hashimshoni, E., Pöhlker, M. L., Jefferson, A., Pöhlker, C., Yu, X., Zhu, Y., Liu, G.,
   Yue, Z. and Fischman, B.: Satellite retrieval of cloud condensation nuclei concentrations by using clouds as CCN chambers, Proc. Nat. Acad. Sci., 113(21), 5828–5834, doi:10.1073/pnas.1514044113, 2016.
  - Rossow, W. B., Tselioudis, G., Polak, A., and Jakob, C.: Tropical climate described as a distribution of weather states indicated by distinct mesoscale clour property mixtures, Geophys. Res. Lett., 32, L21812, doi:10.1029/2005GL024584, 2005.
- 770 Ryzhkov, A., Pinsky, M., Pokrovsky, A., and Khain, A.: Polarimetric radar observation operator for a cloud model with spectral microphysics, J. Appl. Meteorol. Climatol., 873–894, doi:10.1175/2010JAMC2363.1, 2011.

Ryzhkov, A., Diederich, M., Zhang, P., and Simmer, C.: Potential utilization of specific attenuation for rainfall estimation, mitigation of partial beam blockage, and radar networking, J. Atmos. Ocean. Tech., 31, 599–619, doi:10.1175/JTECH-D-13-00038.1, 2014.

- Scharenbroich, L., Nagnusdottir, G., Smyth, P., Stern, H., and Wang, C.-C.: A Bayesian framework for storm tracking using a hidden-state representation, Mon. Weath. Rev., 138, 2132–2148, 2010.
- Sherwood, S. C., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing, Nature, 505(7481), 37–42, doi:10.1038/nature12829, 2014.
- 780 Snyder, J. C., Bluestein, H. B., Venkatesh, V., and Frasier, S. J.: Observations of polarimetric signatures in supercells by an X-band mobile Doppler radar, Mon. Weath. Rev., 141(1), 3–29, doi:10.1175/MWR-D-12-00068.1, 2013.





Snyder, J. C., Ryzhkov, A. V., Kumjian, M. R., Khain, A. P., and Picca, J.: A ZDR column detection algorithm to examine convective storm updrafts, Weath. Forecast., 30(6), 1819–1844, doi:10.1175/WAF-D-15-0068.1, 2015.

- 785 2
  - Snyder, J. C., Bluestein, H. B., Dawson, D. T., II, and Jung, Y.: Simulations of polarimetric, X-band radar signatures in supercells. Part I: Description of experiment and simulated *ρ<sub>hv</sub>* rings, J. Appl. Meteorol. Climatol., 56(7), 1977–1999, doi:10.1175/JAMC-D-16-0138.1, 2017a.
- Snyder, J. C., Bluestein, H. B., Dawson, D. T., II, and Jung, Y.: Simulations of polarimetric, X-band radar
   signatures in supercells. Part II: Z<sub>DR</sub> columns and rings and K<sub>DP</sub> columns, J. Appl. Meteorol. Climatol., 56(7), 1977–1999, doi:10.1175/JAMC-D-16-0138.1, 2017b.
  - Stanford, M. W., Varble, A., Zipser, E., Strapp, J. W., Leroy, D., Schwarzenboeck, A., Potts, R., and Protat, A.: A ubiquitous ice size bias in simulations of tropical deep convection, Atmos. Chem. Phys., 17, 9599–9621, doi:10.5194/acp-17-9599-2017, 2017.
- 795 Stein, T. H. M., Hogan, R. J., Clark, P. A., Halliwell, C. E., Hanley, K. E., Lean, H. W., Nicol, J. C., and Plant, R. S.: The DYMECS project: A statistical approach for the evaluation of convective storms in high-resolution NWP models, Bull. Amer. Meteorol. Soc., 96(6), 939–951, doi:10.1175/BAMS-D-13-00279.1, 2015.
  - Storer, R. L., and van den Heever, S. C.: Microphysical processes evident in aerosol forcing of tropical deep convective clouds, J. Atmos. Sci., 70(2), 430–446, doi.org/10.1175/JAS-D-12-076.1, 2013.
- 800 Tan, I., Storelvmo, T., and Zelinka, M. D.: Observational constraints on mixed-phase clouds imply higher climate sensitivity, Science, 352(6282), 224—227, doi:10.1126/science.aad5300, 2016.
  - Testud, J., Oury, S., Black, R., Amayenc, P., and Dou, X.: The concept of "normalized" distribution to describe raindrop spectra: A tool for cloud physics and cloud remote sensing, J. Appl. Meteorol., 40(6), 1118–1140, 2001.
- 805 Thurai, M., Bringi, V. N., Carey, L. D., Gatlin, P., Schultz, E., and Petersen, W. A.: Estimating the accuracy of polarimetric radar–based retrievals of drop-size distribution parameters and rain rate: An application of error variance separation using radar-derived spatial correlations, J. Hydrometeorol., 13, 1066–1079, doi:10.1175/JHM-D-11-070.1, 2012.
- Vali, G., DeMott, P. J., Möhler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology,
  Atmos. Chem. Phys., 15(18), 10263–10270, doi:10.5194/acp-15-10263-2015-corrigendum.
- van Lier-Walqui, M., Fridlind, A. M., Ackerman, A. S., Collis, S., Helmus, J. J., North, K., Kollias, P., Mac-Gorman, D. R., and Posselt, D. J.: On polarimetric radar signatures of deep convection for model evaluation: columns of specific differential phase observed during MC3E, Mon. Weath. Rev., 144, 737–758, 2016.
- Varble, A., Zipser, E. J., Fridlind, A. M., Zhu, P., Ackerman, A. S., Chaboureau, J.-P., Collis, S., Fan, J., Hill,
  A., and Shipway, B.: Evaluation of cloud-resolving and limited area model intercomparison simulations
- using TWP-ICE observations: 1. Deep convective updraft properties, J. Geophys. Res. Atmos., 119, no. 24, 13891–13918, doi:10.1002/2013JD021371, 2014a.
  - Varble, A., Zipser, E. J., Fridlind, A. M., Zhu, P., Ackerman, A. S., Chaboureau, J.-P., Fan, J., Hill, A., Shipway,B., and Williams, C.: Evaluation of cloud-resolving and limited area model intercomparison simulations
- 820 using TWP-ICE observations: 2. Precipitation microphysics, J. Geophys. Res. Atmos., 119, no. 24, 13919– 13945, doi:10.1002/2013JD021372, 2014b.





Vardiman, L.: The generation of secondary ice particles in clouds by crystal-crystal collision, J. Atmos. Sci., 35, 2168–2180, 1978.

Wang, S., Sobel, A. H., Fridlind, A., Feng, Z., Comstock, J. M., Minnis, P., and Nordeen, M.L.: Simulations of

825 cloud-radiation interaction using large-scale forcing derived from the CINDY/DYNAMO northern sounding array, J. Adv. Model. Earth Syst., 7, no. 3, 1472–1498, doi:10.1002/2015MS000461, 2015.

Yano, J.-I., and Phillips, V. T. J.: Ice–ice collisions: An ice multiplication process in atmospheric clouds, J. Atmos. Sci., 68(2), 322–333, doi:10.1175/2010JAS3607.1, 2011.

Zhang, J., Qi, Y., Tang, L., Cocks, S., Wang, Y., Zhang, P., Ryzhkov, A., Langston, C., and Kaney, B.: MRMS

830 dual-polarization radar synthetic QPE, 38th Conference on Radar Meteorology, Chicago, IL, Amer. Meteor. Soc., 22B.3, 2017.

Zipser, E. J.: Some views on "hot towers" after 50 years of tropical field programs and two years of TRMM data, in: Cloud Systems, Hurricanes, and the Tropical Rainfall Measuring Mission (TRMM), edited by: W.-K. Tao and R. Adler, American Meteorological Society, 49–58, 2003.

835 Zrnić, D. S., and Ryzhkov, A. V.: Polarimetry for weather surveillance radars, Bull. Amer. Meteorol. Soc., 80(3), 389–406, doi:10.1175/1520-0477(1999)080<0389:PFWSR>2.0.CO;2, 1999.

Zrnić, D. S., Straka, J., Liu, Y., and Vivekanandan, J.: Testing a procedure for automatic classification of hydrometeor types, J. Atmos. Oceanic. Technol., 18, 892–913, doi:10.1175/1520-0426(2001)018,0892:TAPFAC.2.0.CO;2, 2001.







**Figure 1.** Map of Houston region (white symbol marks city center), NU-WRF inner domain boundaries (orange square), KHGX NEXRAD radar location (cyan symbol with 100 km and 200 km range rings), tracks of three example features from KHGX  $K_{\text{DP}}$  observations (cyan tracks numbered 9, 35, and 37) and from NU-WRF  $q_r$  output (orange tracks numbered 89, 116, and 188), and CCN number concentration and supersaturation retrieved from satellite data (within yellow boxes in cm<sup>-3</sup> and %, respectively).







**Figure 2.** Two reflectivity factor snapshots (gridded to a constant height of 1 km) from subsequent NEXRAD scans of KHGX and their cross-correlation. The peak in the cross-correlation gives a good indication of the image shift between the two time steps and is used as the position start of the search to identify cells in subsequent images.



**Figure 3.** An example of TINT-generated cell tracks from 7 July 2013. A constant-altitude plot of reflectivity at 1.5 km is shown for reference. Each line showing the path of each cell is given a randomly generated cell identification number. The data is loaded into memory as a Pandas data frame and saved to a comma-separated variable file for later analysis.







**Figure 4.** Time series from three  $K_{DP}$  column objects tracked from observations show (top-to-bottom) lightning flashes per KHGX volume time (grey line) and occurrence density as a function of height (see colorbar),  $K_{DP}$  and  $Z_{DR}$  column strength (calculated following Eqn. 1), and retrieved rain rate and drop  $D_m$ . Lightning and rain statistics collected within 2.5 km of the tracked  $K_{DP}$  column center. Vertical dashed lines indicate times of column 9 and 37 cross-sections shown in Figs. 5 and 6.







Figure 5. Sequential south-north cross-sections from observed K<sub>DP</sub> column 9 show (left to right) Z<sub>HH</sub>, K<sub>DP</sub>, Z<sub>DR</sub>, D<sub>m</sub>, and N<sub>w</sub>. In all panels, the approximate melting level is indicated with a dotted line and each identified lightning flash is overplotted with a dot color indicating time sequence (yellow to red). Full time series shown in Fig. 4.















**Figure 7.** Time series from three  $K_{\text{DP}}$  columns tracked from NU-WRF simulation output show (top-to-bottom) updraft strength,  $q_r$ ,  $K_{\text{DP}}$  and  $Z_{\text{DR}}$  column strengths (calculated following Eqn. 1), and rain rate and drop  $D_m$  and  $N_w$ . Updraft and rain statistics collected within 2.5 km of the tracked  $K_{\text{DP}}$  column center. Vertical dashed lines indicate times of column 116 and 188 cross-sections shown in Figs. 8 and 9.























**Figure 10.** Total number of isolated convective cells that initiated within a 200-km range of the KHGX radar during 2015–2017 by month.



**Figure 11.** Monthly percentage of days during 2015–2017 with at least one isolated cell initiation within a 200-km range of the KHGX radar.



**Figure 12.** Total number of isolated cells that initiated within a 200-km range of the KHGX radar as a function of time of day during June through September of 2015–2017. The peak at 18 UTC corresponds to 1 PM local time.







**Figure 13.** Number of isolated cells that both initiated and dissipated with 70, 150 and 200 km of the KHGX radar as a function of cell lifetime during June through September of 2015–2017. Integrated totals are 441, 2442 and 4843, respectively.



Figure 14. Distribution of isolated cell initiation location during 2015–2017.







Figure 15. Directional cumulative distribution of propagation direction during June through October versus November through May.