PLEASE SEE A REVISED VERSION OF THE PAPER AT THE END OF THESE COMMENTS THAT INCORPORATES THE CHANGES. YELLOW=REV 1 GREEN=REV2 GREY=AUTHOR CHANGES

Reviewer #2

This paper presents a comparison between OMI NASA v3.1 and PANDORA total NO2 VCD, showing a clear under-estimation of the OMI data at 7 long-term sites and 6 campaign-based sites. The results at most of the sites are presented and discussed and few arguments for the general underestimation result are mentioned. Although the paper is interesting and fulfill the scope of AMT, there is a lack of reference to literature (previous similar studies and scientific proof/reference of why such differences at the different sites). Sensitivity tests or further comparisons on OMI pixel sizes (edge and center of the swath, different position of the pixels, GB time-selection) could be done to help justifying the proposed conclusion. I recommend the publication after the suggested revisions.

General comments:

The paper is short and easy to read, but it lack some "proof" of the proposed explanation of the OMI under-estimation (argument

1= "Because of the local inhomogeneity of NO2 emissions, the large OMI FOV is the most likely factor when comparing OMI TCNO2 to retrievals from the small PANDORA effective FOV", line 20 and argument.

See page 2 Judd et al., 2018; Nowlan et al., 2016

2= "OMI estimated air mass factor, surface reflectivity, and the OMI 24x13 km2 FOV (field of view) are three factors that can cause OMI to underestimate TCNO2", line 18).

See the references Boersma et al., 2011; Lin et al., 2015; Nowlan et al., 2016; Lorente et al., 2018

Krotkov et al., 2017

Some sensitivity tests on the how much the choices made for the OMI pixel selection (FOV distance d<5km for an SZA<70, line 165) and PANDORA selection ("daily data matched to the OMI overpass times ±6miutes", line 87) affect the results would support argument.

It is the Pandora FOV is less than 5 km from the Pandora site for SZA <70 degrees. This is simple geometry. I have added a comment for this.

1 (or at least give an uncertainty range). Additional comparison (or at least further comments on other OMI retrievals, such as DOMINO (Boersma et al., 2011) or QA4ECV (Boersma et al., 2018)), would support argument

The DOMINO algorithm has some known problems (see reference) and the QA4ECV results are very similar to the NASA results. Because of this, I have put in a statement about the QA4ECV results and a reference.

2. Moreover, a lot of (redundant) figures are given (daily and monthly panels in Fig 3, 4, 5 and 9) could be simplified by plotting the mean and the variability – or a scatter plot of OMI vs PANDORA as often done in validation papers – while e.g., number of comparison points or impact of the Lowess(f) monthly running averages is not mentioned/discussed. How much this exercise results would change with a simple mean or median of the daily comparisons? This would allow putting an uncertainty number on the 1.8 and 1.7 PAN/OMI mentioned in page 11.

I disagree with referee about the redundancy. Figure 3A shows the daily data and Figure 3B shows the averages. Even though both show the difference, it is useful to see the daily data.

Consider adding a section or table with the different PANDORA site description, that would help the reader understanding the general differences among the stations (partially already described in the text, but bot for all sites – coordinated of the sites is also missing). This would be a good reference for future studies using these PANDORA data.

OK See Table 2

Please clarify how some justifying arguments are obtained (add references or explain not shown results). E.g. :

P14, line 278 "The relatively moderate TCNO2 values (0.4 to 0.8 DU) are probably a testament to the effectiveness of catalytic converters mandatory on all US automobiles in such a high traffic area"

Gary A. Bishop and Donald H. Stedman, Reactive Nitrogen Species Emission Trends in Three Light-/Medium-Duty United States Fleets, *Environmental Science & Technology* 2015 *49* (18), 11234-11240, DOI: 10.1021/acs.est.5b02392

add reference!;

 P17, line 290 "The highest amount of TCNO2 recorded during 2018 was about 5DU on 13 July 2018 from 11:20 and 12:30 EST (a time with very light winds (1 km/hr) and moderate temperature (25°C)"

☑ is the meteorology present at each site or only here? Could you shown some correlations? Or is this just a specificity of that time period?

Meteorology affects the amount of NO2 observed at all sites. I described the meteorology for this site on a 13 July 2018 because the amount, 5DU, was very unusual. In general, days with no winds show high values of NO2 near the sources for NO2.

Harkey, M., Holloway, T., Oberman, J., and Scotty, E., An evaluation of CMAQ NO2 using observed chemistry-meteorology correlations, J. Geophys. Res. Atmos., 120, 11,775–11,797, doi:10.1002/2015JD023316, 2015

Specific comments and Technical corrections

- Line 6: "14 sites" but only 13 are presented – 7 sites in table 1 and 6 in table 2. Same comment for line 13 "Eight additional sites..."

Corrected in Table 2

- Line 9 and 11: why mention sites in Northern Hemisphere and Southern hemisphere if this is never mentioned again in the manuscript? Same comment for line 16 "weekly or monthly average basis": weekly comparisons are never mentioned again. -

Now NH and SH mentioned in the text on Page 6. Even though true, I removed weekly from line 16.

Line 18 – 19 and 19-22: see general comment, these 2 arguments are not discussed a lot in the paper. –

Surface reflectivity was discussed on page 2. I added: "Accurately determining the AMF for TCNO₂ requires a-priori knowledge of the NO₂ profile shape, which is estimated from coarse resolution model calculations (Boersma et al., 2011)," The references give extensive discussions of these factors and their effects.

line 87 – 89: the explanation on how the comparison is done is mixed between this line and lines 165. Consider adding a paragraph grouping all the comparison selection choices (cloud free pixels? What is done with the row anomaly? Why is a 6 minutes time-selection selected for the PANDORA? What type of filtering is done for PANDORA ? (cf mention of impact of clouds in line 130), ...

The following paragraph has been added to page 3

OMI overpass data, <u>https://avdc.gsfc.nasa.gov/index.php?site=666843934&id=13</u>, are filtered for the row anomaly and cloudy pixels. The selection of a ± 6 minute window represents 720 seconds or 9 PANDORA measurements averaged together around the OMI overpass time to reduce the effect of any outlier points. PANDORA makes an NO₂ measurement every 80 seconds. The specific value of ± 6 minutes is arbitrary but increases the effective signal to noise ratio by a factor of 3. PANDORA data are filtered for significant cloud cover by examining the effective variance in sub-interval (20 seconds) measurements. Each PANDORA listed measurement is the average of up to 4000 (clear sky) individual measurement made over 20 seconds.

To my knowledge, the way the selection is done could have an impact on the results (size of OMI pixels, pixels covering the station or not, averaging the ground-based data (mean or median value?), ...), and this is only poorly/not discussed. What is the impact of "the Lowess(f) monthly running averages" choice? -

The overpass data set represents the closest filtered OMI pixel to the specified site on a given day. On any single OMI measurement, the OMI FOV may not be exactly over the site. This is an intrinsic characteristic of OMI data as used for practical purposes. An alternative would be to use gridded data on a fixed latitude x longitude grid. The result is an even wider area view of the specified site (an average of more OMI pixels). The point is that OMI data are used to represent the amount of NO₂ over a given location whether in comparison to PANDORA or a model study. Air quality decisions are made based on OMI data for urban and unpolluted regions that include intrinsic area averaging.

The impact of Lowess or adjacent averaging over a month's worth of data is to smooth out the daily variation and show an average difference. Daily data are presented as well. Weekly averages would show the same qualitative result. Lowess is preferable to adjacent averaging, since it is least-squares weighted and reduces the effect of possible outlier points

Line 117: change "." to ":". **Fixed** Same for line 137 giving the link to the data: introduce it in a sentence (e.g., Data can be found here: ...**It already says that**). Moreover, a table with coordinates and multiple names of the PANDORA stations would be helpful – "waterflow" overpass is e.g. found in the OMI link, but not on the PANDORA link. **Waterflow is labelled Four Corners on the website – I have added this name to the paper.**

Lines 142-147: give references and refer to this when discussing daily and monthly evolution of fig 7 and 8. –

(Lamsal et al., 2013; Bechle et al., 2013).

Line 172-174: why only give an illustration of O3 comparison for Busan?

I could give O3 plots for all sites at the expense of more figures. However, the appearance is very similar to that for Busan (except Mauna Loa because of altitude effects). The results for all sites are summarized in Table A1. The purpose is to show that all instruments were working properly.

Also in table A1, there are quite some differences in the percent difference from station to station (from 0 in Baltimore to 5.6 in Mauna Loa).

The Mauna Loa difference is caused by altitude for O3 with Pandora missing the lowest 3.4 km. NO2 differences are not related to ozone differences. This is stated in Table A1. The differences are not a function of the PANDORA instruments nor the retrieval algorithms.

For O3, the biggest error is the lack of effective O3 temperature in the algorithm. An average effective O3 temperature is used instead of a measured temperature. An example of this is give in Herman et al., 2015; 2017 for Boulder Colorado

How is the PAN/OMI here? Is the largest difference for in O3 also at the same stations than the largest differences for NO2? **No**

Is it in stations where we expect most of the NO2 in the stratosphere (Mauna Loa)? How is the NO2 tropo/strato ratio (seen by the satellite?)

For Mauna Loa, the Pandora saw more NO2 than is possible in the stratosphere. The NO2 is drifting upward from the coastal areas. This is mentioned in the paper.

Comment on table A1! (How to explain O3 differences of 2.5 to 2.8% at stations close to surface level?) if not here, at least in the Appendix. –

Without proof, I suspect that the incorrect average effective temperature is the cause of a part of the difference as it was at Boulder Colorado, since we use temperature dependent ozone cross sections for both Pandora and OMI. There is also the issue of field calibration to remove the reference amount of ozone (modified Langley calibration) for zero airmass. This is discussed in an earlier paper and not part of the scope of this paper. This procedure has not been done for City College nor for HUFS. If the instruemnts were not operating properly (e.g., pointing at the sun), the differences would be much larger.

Figures 3, 4 and 5: in the monthly averages, there is often peaks not seen in OMI (shortly discussed for some stations (lines 179-180 for Busan), but not for all of them. Regularly, there is also a divergent behavior of the monthly average at the edges of the time-series (e.g., end 2016 for Mauna Loa, in 2017 for NASA HQ, end of 2017 for Waterflow, end of 2017 to early 2018 for Boulder) or OMI columns at the end of the time-series as high as PANDORA (eg Buenos aires, NASA HQ). Is this real of is this related to the "Lowess(f) monthly running averages"? – '

I should exclude endpoints for running averages. I will change the figures. (NOT DONE YET)

Lines 195-196: "The calibration of the Mauna Loa PANDORA will be reviewed as part of a general data quality assurance program that is starting with the most recently deployed PANDORA instruments " - do you mean that the PANDORA data might be off? -

No, there is a new Pandora installed at Mauna Loa after the sun tracker broke down. At the time of this writing, data from the new Pandora are not used. The sentence has been changed.

Recently, the original Mauna Loa PANDORA has been replaced. The new instrument's calibration will be reviewed as part of a general data quality assurance program that is starting with the most recently deployed or upgraded PANDORA instruments at about 100 locations.

Lines 209-211: there is some repetition with previous paragraphs. -

This paragraph has been moved (Page 9)

Tables 1 and 2: add coordinates of the stations and measurement time-periods. How is the "average" among the stations performed? Mean? Median? Does it have a large effect? Consider giving the correlations. Comment on Seoul PAN = 1.2 (more than double of all the other sites!) New York value is missing. –

The average values are simply the arithmetic average of the daily points for each location. The overall average is the arithmetic average of the above averages. Seoul is the most polluted city considered, so

the average value is higher. However, the ratio with OMI is similar to most of the other sites. New York has been added to Table 2.

I have added correlation coefficients to Table 1 and the sentence on page 9

For example, the PANDORA at NASA Headquarters in Washington DC tracks the OMI measurement quite well on a monthly average basis with a correlation coefficient of $r^2(mn) = 0.7$ even though the daily correlation is low ($r^2(dy) = 0.17$). Other sites have only short periods of correlation and overall weak correlation (Table 1 showing daily, dy and monthly, mn, correlation coefficients for the graphs in Figures 4 and 5)

Line 220: give references of the Discover-AQ campaigns and discuss some of the outcomes (several PADORA on close locations; airborne flights; ...) Refer also to other studies dealing with PANDORA data for validation of NO2, eg., Judd et al., 2019 (https://www.atmos-meastech-discuss.net/amt-2019-161/) discussing heterogeneous NO2 situations. –

The Judd at al reference has been added on Page 2 and backs up the thesis that spatial resolution is a major cause of the underestimate by OMI compared to PANDORA.

Judd, L. M., Al-Saadi, J. A., Janz, S. J., Kowalewski, M. G., Pierce, R. B., Szykman, J. J., Valin, L. C., Swap, R., Cede, A., Mueller, M., Tiefengraber, M., Abuhassan, N., and Williams, D.: Evaluating the impact of spatial resolution on tropospheric NO₂ column comparisons within urban areas using high-resolution airborne data, Atmos. Meas. Tech. Discuss., https://doi.org/10.5194/amt-2019-161, in review, 2019.

Lines 245-267: consider re-organizing the paragraphs (order and repetition). Discuss first Fig 6 completely, and then comment on Fig 7. In the comments of figure 6, reference to literature trends is missing (e.g., Duncan et al., 2016; ...).

Duncan et al. (2016) estimated trends from OMI TCNO₂ time series and found that the Seoul metropolitan area had a decrease of -1.5 ± 1.3 %/Year (2005 – 2014) consistent with OMI estimated change of -1.4 ± 1 %/year (2012 -2018) in this paper. However, for the small area near Yonsei University, the decrease estimated from PANDORA is -5.8 ± 0.75 %/Year. Park (2019) estimates that metropolitan Seoul has decreased in population even as surrounding areas have increased population.

(see page 12)

It is a pity that only 6 of the 7 long term stations are shown in Fig 6.

While not showing an extra plot in Fig. 6 I have added the results for the 7th long-term site, Busan, in the text,

The results for Busan (from Fig. 3) show a least squares average for the percent difference of -48 \pm 0.8% for the 2012 – 2018 period with a slope of 6.8 \pm 1%/Year. There is a decrease in the percent difference after October 2015 (Fig. 3) that is mainly from PANDORA seeing less TCNO₂

than during the 2012 – 2014 period. There is a gap in the Busan time series from July 2014 until April 2015 when the original PANDORA was replaced with a new instrument. The calibrations of both PANDORAS appear to be correct. Because of the break in the time series it is not clear whether there was a change in local conditions around Pusan University compared to the wide area observed by OMI.

Move the discussion of the Boulder trend from the figure caption to the main text.

Done

Is there an explanation for the 3 classes of mean bias results (1) about -24 to -27% for Boulder, Mauna Loa and NASA HQ; 2) about -37% for Waterflow and 3) about -46% for Buenos Aires and Seoul)? –

I do not know the explanation for the differences between the narrow view trends (PANDORA and the wide area trends (OMI). The other long-term site considered, Busan, has gaps in the data record that are fairly large.

I added (page 14)

For some sites (see Fig. 6), PANDORA and OMI trends are the same (Waterflow, NM, Buenos Aires, and Mauna Loa) while the other 3 sites show significantly different trends (Boulder, NASA HQ, and Seoul).

Lines 258-259: consider giving all the correlation coefficients in the tables as suggested.

see Table 1

- Figure 7 and 8: pity that the figures are not presented for the same year (2018), so that we could compare NASA HQ Washington and New York NO2 levels.

I do not have a complete data record for NASA HQ in 2018 and only have 2018 for New York City

Moreover, the TCNO2 axis limit is changing from panel to panel, so it is not so easy to see the seasonal behavior. –

Making all of the scales the same will obscure the behavior relative to the OMI overpass time, which is the subject of this paper.

There is no easy way to represent the seasonal behavior vs time of day on a minute by minute basis or even an hourly basis for such complex highly variable behavior of TCNO2 shown in Figures 7 and 8. The seasonal variation at the OMI overpass time is given in Fig. 9 for NYC and in Fig. 4 for NASA HQ. The general seasonal variation is not the subject of this paper. However, while not part of this paper, I have added a graph at the end of this reply that shows the monthly average behavior of TCNO2 for CCNY for four different times of the day 10:00, 13:50, 14:00, 16:00. With variations in magnitude, the seasonal behavior is similar for the different times of the day.

Lines 278-279: "The relatively moderate TCNO2 values (0.4 to 0.8 DU) are probably a testament to the effectiveness of catalytic converters mandatory on all US automobiles in such a high traffic area". Is it purely speculative? Is there any correlation with when the regulation measures have been put in place? Give references! – **(Bishop and Steadman, 2015).**

Line 284: "the pollution levels are quite high, rivaling the pollution levels in Seoul, South Korea." It is not seen in Tables 1 and 2, and we don't have these kind of plots for Seoul, only Busan (fig 1). –

I added (see Fig. 5)

Line 293: "For both Washington DC (Fig. 7) and New York City (Fig. 8) there is strong day-today and month to month variability that depends on the local weather and the amount of automobile traffic in the area" – has the dependence on weather and traffic been tested or is this a guess or literature reference? –

Seo, J., Park, D.-S. R., Kim, J. Y., Youn, D., Lim, Y. B., and Kim, Y.: Effects of meteorology and emissions on urban air quality: a quantitative statistical approach to long-term records (1999–2016) in Seoul, South Korea, Atmos. Chem. Phys., 18, 16121-16137, https://doi.org/10.5194/acp-18-16121-2018, 2018.

Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang, D., Pöschl, U., Cheng, Y. F., and He, K. B.: Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous reactions, Atmos. Chem. Phys., 15, 2969–2983, <u>https://doi.org/10.5194/acp-15-2969-2015</u>, 2015.

Md. Shohel Reza Amin, Umma Tamima, and Luis Amador Jimenez, "Understanding Air Pollution from Induced Traffic during and after the Construction of a New Highway: Case Study of Highway 25 in Montreal," Journal of Advanced Transportation, vol. 2017, Article ID 5161308, 14 pages, <u>https://doi.org/10.1155/2017/5161308</u>, 2017

Andersen, M. Hvidberg, S.S. Jensen, M. Ketzel, S. Loft, M. Sørensen, A. Tjønneland, K. Overvad, O. Raa schou-Nielsen Chronic obstructive pulmonary disease and long-term exposure to traffic-related air pollution: A cohort study, Am. J. Respir. Crit. Care Med., 183, pp. 455-461, <u>10.1164/rccm.201006-09370C</u>, 2011.

Line 296: "Poor air quality affecting respiratory health would be improperly characterized by both the OMI average values being too low (Fig. 4) and by missing the extreme pollution events that occur frequently in the late afternoon". Also add a comment (with references) that here total columns are being analyzed, while tropospheric columns could be used, which anyway don't reflect systematically the surface concentrations important for air quality. –

Page 20 It should be noted that TCNO₂ does not accurately represent the NO₂ concentration at the surface, since it is mostly a measure of the amount in the lower 2 km. However, it is roughly proportional to the surface measurements close to the pollution sources (Bechle et al., 2013; Knepp et al., 2014) with the proportionality dependent on the profile shape near the ground.

Caption of figure 9: "Lowless(0.08)" it is the first time that the "f" is mentioned. Why is it different than in Herman et al., 2018 (e.g., caption of figure 9 "Lowless(0.1)")? –

I could have given the f-value for each graph. It is the fraction f of data points over which the Lowess(f) algorithm is applied to form an average local least squares fit. This is similar to the number of points included in an arithmetic running average. The exact fraction will depend on the number of points in a month's worth of data compared to the entire data set.

Line 308-309: "there is a period in March 2018 when OMI TCNO2 slightly exceeded that measured by PANDORA." Where are those pixels? Over the sea? What is their size? What is the wind condition? -

The OMI pixels for the March 2018 period are distributed over both land and water. I have replotted the data only using points less than 30 km from CCNY. The results are very similar, but not identical to when D < 80 km. The wind conditions were variable (I do not have the detailed meteorological data). The pixel size also is variable with the centers located less than 30 km from CCNY in the graph below.



Figure 9 has been replaced to exclude pixels further than 30 km. The results are almost identical. Most papers comparing OMI data with models related to air quality estimates use a gridded version of OMI data totally ignoring OMI pixel size in order to produce local area maps of TCNO₂

Line 2018-2019: "The OMI underestimate is much larger than error estimates for TCNO2 retrievals for either PANDORA or OMI". Consider adding the error on some of the graphs for illustration!

I added the error estimates for the least squares mean percent differences to the graphs in Fig. 6.

-

Add some discussion in the conclusion about new and upcoming satellites (eg TROPOMI with smaller pixels and geostationary that will be able to see the diurnal variation)

Done

and the uncertainties of this study (impact of the NASA product selection for OMI (wrt to DOMINO and QA4ECV) and related to the way the comparison is done (see general comment)).

See page 2

- Appendix: comment on table A1 O3 results (up to 2.8% also outside mountain conditions) -

The 2.8% offset is too large since the PANDORA calibration looks very good. Both data sets track each other quite well with high correlation on a monthly average basis. The most likely cause is an improper effective ozone temperature correction for PANDORA that was obtained from a model calculation

References: Boersma et al., 2011 is missing. Add suggested references. Mind the formatting!

Added

Suggested references:

Duncan, B. N., L. N. Lamsal, A. M. Thompson, Y. Yoshida, Z. Lu, D. G. Streets, M. M. Hurwitz, and K. E. Pickering (2016), A space-based, high-resolution view of notable changes in urban NOx pollution around the world (2005–2014), J. Geophys. Res. Atmos., 121, doi:10.1002/2015JD024121.

Judd, L. M., Al-Saadi, J. A., Janz, S. J., Kowalewski, M. G., Pierce, R. B., Szykman, J. J., Valin, L. C., Swap, R., Cede, A., Mueller, M., Tiefengraber, M., Abuhassan, N., and Williams, D.: Evaluating the impact of spatial resolution on tropospheric NO2 column comparisons within urban areas using high-resolution airborne data, Atmos. Meas. Tech. Discuss., https://doi.org/10.5194/amt-2019-161, in review, 2019.

Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen, J. H. G. M., Zara, M., Peters, E., Van Roozendael, M., Wagner, T., Maasakkers, J. D., van der A, R. J., Nightingale, J., De Rudder, A., Irie, H., Pinardi, G., Lambert, J.-C., and Compernolle, S. C.: Improving algorithms and uncertainty estimates for satellite NO2 retrievals: results from the quality assurance for the essential climate variables (QA4ECV) project, Atmos. Meas. Tech., 11, 6651-6678, https://doi.org/10.5194/amt-11-6651-2018, 2018.





1 Underestimation of Column NO₂ Amounts from the OMI Satellite Compared to Diurnally

2

Varying Ground-Based Retrievals from Multiple Pandora Spectrometer Instruments

Jay Herman¹, Nader Abuhassan¹, Jhoon Kim², Jae Kim³, Manvendra Dubey⁴, Marcelo Raponi⁵,
 Maria Tzortziou⁶

5 Abstract

6 Retrievals of Total Column NO_2 (TCNO₂) are compared for 14 sites from the Ozone Measuring 7 Instrument (OMI using OMNO2-NASA v3.1) on the AURA satellite and from multiple ground-8 based PANDORA spectrometer instruments making direct-sun measurements. The result is that 9 on a daily and monthly average basis, OMI almost always underestimates the amount TCNO₂ by 50 to 100%, while occasionally the daily OMI value exceeds that measured by PANDORA at very 10 clean sites. In addition to systematic underestimates, OMI always misses the frequently much 11 higher values of TCNO₂ that occur after the OMI overpass time. This suggests that OMI retrieved 12 13 TCNO₂ are not suitable for air quality assessments as related to human health, especially in 14 polluted urban areas. Six discussed Northern Hemisphere PANDORA sites have multi-year data 15 records (Busan, Seoul, Washington DC, Waterflow New Mexico, Boulder Colorado, and Mauna Loa) and one site in the Southern Hemisphere (Buenos Aires Argentina). The first four of these 16 17 sites and Buenos Aires frequently have high TCNO₂ (TCNO₂ > 0.5 DU). Eight additional sites have shorter term data records in the US and South Korea. One of these is a one-year data record from 18 19 a highly polluted site at City College in New York City with pollution levels comparable to Seoul, South Korea. OMI estimated air mass factor, surface reflectivity, and the OMI 24x13 km² FOV 20 21 (field of view) are three factors that can cause OMI to underestimate TCNO₂. Because of the local 22 inhomogeneity of NO₂ emissions, the large OMI FOV is the most likely factor for consistent 23 underestimates when comparing OMI TCNO₂ to retrievals from the small PANDORA effective FOV calculated from the solar diameter of 0.5[°]. 24

25

26

Key Words: Nitrogen dioxide, OMI, PAN, PANDORA, ground-based, satellite

Correspondence email: jay.r.herman@nasa.gov

¹University of Maryland Baltimore County JCET, Maryland

²Department of Atmospheric Sciences, Yonsei National University, South Korea

³Department of Atmospheric Science, Pusan National University, South Korea

⁴Earth Systems Observations, Los Alamos National Laboratory, Los Alamos, NM 87545

⁵Departamento de Investigaciones en Láseres y Aplicaciones (DEILAP), Instituto de Investigaciones

Científicas y Técnicas para la Defensa (CITEDEF), Ministerio de Defensa (MINDEF), Buenos Aires, Argentina ⁶City College of New York, New York City, NY

Underestimation of Column NO₂ Amounts from the OMI Satellite Compared to Ground-Based Retrievals from Multiple Pandora Spectrometer Instruments

29 1.0 Introduction

30 Retrieval of Total Column NO₂ (TCNO₂) from the Ozone Monitoring Instrument (OMI) has been a 31 scientific success story for the past 14 years. Near total global coverage from the well-calibrated OMI has 32 enabled observation of all the regions where NO₂ is produced and has permitted monitoring of the 33 changes during the 2004 to 2019 period, especially in regions where there is heavy and growing industrial 34 activity (e.g., China and India). TCNO₂ amounts (data used: OMNO2-NASA v3.1) retrieved from OMI over 35 various specified land locations show a strong underestimate compared to co-located Pandora Spectrometer Instruments (the abbreviation PAN is used for graph and table labels). The underestimate 36 37 of OMI TCNO₂ at the overpass time compared to ground-based measurements has previously been 38 reported at a few specific locations (Bechle, 2013; Lamsal et al., 2015; Ialongo et al., 2017; Kollonige, et 39 al., 2018; Goldberg et al., 2018; Herman et al., 2018). For any location, the OMI overpass local standard 40 time consists of the central overpass near the 13:30 hour equator crossing solar time and occasionally a 41 side viewing overpass from adjacent orbits within ±90 minutes of the central overpass time. 42 Independently from instrument calibration and retrieval errors, there are two specific aspects to the 43 underestimation of TCNO₂ pollution levels. First, the mid-day OMI observations do not see the large 44 diurnal variation of TCNO₂ that usually occur after the 13:30 overpass time, and second, because of spatial inhomogeneity the large OMI field of view (FOV) footprint 13 x 24 km² at OMI nadir view tends to average 45 46 regions of high NO₂ amounts (Nowlan et al., 2016; Judd et al., 2018) with those from lower pollution areas. An analysis by Judd et al., (2019, their Fig. 9) shows the effect of decreasing satellite spatial resolution on 47 48 improving agreement with PANDORA, with the best agreement occurring with an airborne instrument, GEO-TASO (resolution 3x3 km²) followed by TropOMI (5x5 km²) and then OMI (18x18 km²). Both OMI and 49

50 TropOMI show an underestimate of TCNO₂ compared to PANDORA.

51 There are other possible systematic retrieval errors with OMI TCNO₂. The largest of these is 52 determining the air mass factor (AMF) needed to convert slant column measurements into vertical column 53 amounts followed by the surface reflectivity Rs (Boersma et al., 2011; Lin et al., 2015; Nowlan et al., 2016; 54 Lorente et al., 2018). Accurately determining the AMF for TCNO₂ requires a-priori knowledge of the NO₂ 55 profile shape (Krotkov et al., 2017), which is estimated from coarse resolution model calculations 56 (Boersma et al., 2011), and using the correct Rs. Currently Rs is found using a statistical process of sorting 57 through years of data to find relatively clear-sky scenes for each location (Kleipool, et al., 2008; O'Byrne 58 et al., 2010). Boersma et al., 2004 gave a detailed error analysis for the various components contributing 59 OMI TCNO₂ retrievals resulting an estimated "retrieval precision of 35-60%" in heavily polluted areas 60 dominated by determining the air mass factor. An improved V2.0 DOMINO retrieval (Boersma et al., 2011) algorithm reduced the retrieval errors while increasing the estimated airmass factor, which reduces the 61 62 retrieved TCNO₂ up to 20% in winter and 10% in summer. The current version of OMNO2-NASA (Krotkov et al., 2017) and v2.0 DOMINO (Boersma et al., 2011) are generally in good agreement (Marchenko et al., 63 64 2015; Zara et al., 2018). However, the OMNO2-NASA TCNO₂ retrievals are 10 to 15% lower than the v2.0 65 DOMINO retrievals and with Quality Assurance for Essential Climate Variables (QA4ECV) retrievals. A subsequent detailed analysis of surface reflectivity (Vasilkov et al., 2017) shows that retrieval of TCNO₂ in

- 67 highly polluted areas (e.g., some areas in China) can increase by 50% with the use of geometry-dependent
- reflectivities, but only increase about 5% in less polluted areas. For PANDORA, calculation of the solar
- 69 viewing AMF is a simple geometric problem (AMF is approximately proportional to the cosecant of the
- solar zenith angle SZA) and is independent of R_s (Herman et al., 2009). For a polluted region with TCNO₂ = 5.34x10¹⁶ molecules/cm² or 2 DU, the PANDORA error is expected to be less than ±2.5% with the largest
- 72 uncertainty coming from an assumed amount of stratospheric TCNO₂ = 0.1 DU.

Accurate satellite TCNO₂ retrievals (and for other trace gases) are important in the estimate of the effect of polluted air containing NO₂ on human health (Kim and Song, 2017 and references therein), especially from the viewpoint of NO₂ as a respiratory irritant and precursor to cancer (Choudhari et al., 2013). Since NO₂ is largely produced by combustion, satellite observations of NO₂ serve as a proxy for changing industrial activity. Another important application requiring accurate measurements of the amount of TCNO₂ and its diurnal variation is atmospheric NO₂ contribution to nitrification of coastal waters (Tzortziou et al., 2018).

80 We show that the use of OMI TCNO₂ for estimating local air guality and coastal nitrification on a 81 global basis is misleading for most polluted locations, and especially on days when the morning or 82 afternoon amounts are higher than those occurring at the OMI overpass time near 13:30 hours standard 83 time. OMI TCNO₂ data are extremely useful for estimating regional pollution amounts and for assessing 84 long-term changes in these amounts. Modelling studies (Lamsal et al., 2017 Fig. 1) based on the Global 85 Modelling Initiative model (Strahan et al., 2007) simulating TCNO₂ diurnal variation over Maryland USA (37-40°N, 74-79°W) shows a late afternoon peak and shows that the stratospheric component does not 86 87 substantially contribute to this peak. Boersma et al. (2016) show that sampling strategy can cause 88 systematic errors between OMI TCNO₂ and model TCNO₂ with satellite results being up to 20% lower than 89 models. Duncan et al., (2014) reviews the applicability of satellite $TCNO_2$ data to represent air quality and 90 notes that TCNO₂ correlates well with surface levels of NO₂ in industrial regions and states that the portion 91 of TCNO₂ in the boundary layer could be over 75% of the total vertical column depending on NO₂ altitude 92 profile shape.

93 This paper presents 14 different site comparisons between retrieved OMI TCNO₂ overpass values 94 that are co-located with PANDORA TCNO₂ amounts from various locations in the world. Six of the 95 comparisons are where PANDORAs have long-term data (1-year or longer) records. The comparisons are 96 done using 80 second cadence data matched to the OMI overpass times ±6 minutes and with monthly 97 running averages calculated using Lowess(f) (Locally Weighted least squares fit to a fraction f of the data points, (Cleveland, 1981) of OMI-PANDORA time matched TCNO2. OMI overpass data, 98 99 https://avdc.gsfc.nasa.gov/index.php?site=666843934&id=13, are filtered for the row anomaly and 100 cloudy pixels. The selection of a ± 6 minute window represents 720 seconds or 9 PANDORA measurements averaged together around the OMI overpass time to reduce the effect of any outlier points. The specific 101 102 value of ±6 minutes is arbitrary but increases the effective signal to noise ratio by a factor of 3. PANDORA 103 data are filtered for significant cloud cover by examining the effective variance in sub-interval (20 seconds) 104 measurements. Each PANDORA listed measurement is the average of up to 4000 (clear sky) individual 105 measurement made over 20 seconds.

106 This paper gives a discussion and presentation of data on the effect of diurnal variation that are 107 always missed at the local OMI mid-day overpass times. We show that OMI TCNO₂ values are also 108 systematically lower than PANDORA values at sites with significant pollution (TCNO₂ > 0.3 DU). We present 109 a unique view of a year of fully time resolved diurnal variation of TCNO₂ at two sites, Washington DC and 110 New York City, which are similar to other polluted locations.

111

112 **2.0 Brief Instrument Descriptions**

113 For the purposes of TCNO₂ retrievals, both OMI and PANDORA are spectrometer-based 114 instruments using nearly the same spectral range and similar spectral resolution (about 0.5 nm). Both use 115 spectral fitting retrieval algorithms that differ (Boersma et al. 2011; Herman et al., 2009) because of the differences between direct-sun viewing retrievals (PANDORA) and above the atmosphere downward 116 117 viewing retrievals (OMI). The biggest difference is with the respective fields of view, 13 x 24 km² at OMI 118 nadir view and larger off-nadir FOV compared to the much smaller PANDORA FOV (1.2°) measured in m² 119 with the precise value depending on the NO₂ profile shape and the solar zenith angle. For example, if most 120 of the TCNO₂ is located below 2 km, then the PANDORA FOV is approximately given by $(1.2\pi/180)(2/\cos(SZA))$, which for SZA = 45° is about 59x59 m². If the solar disk (0.50) is used as the limiting 121 122 factor, then the FOV is smaller.

123 **2.1 OMI**

124 OMI is an east-west side (2600 km) and nadir viewing polar orbiting imaging spectrometer that 125 measures the earth's backscattered and reflected radiation in the range 270 to 500 nm with a spectral 126 resolution of 0.5 nm. The polar orbiting side viewing capabilities produce a pole to pole swath that is about 127 2600 km wide displaced in longitude every 90 minutes by the earth's rotation to provide coverage of 128 nearly the entire sunlit Earth once per day at a 13:30 solar hour equator crossing time with spatial gaps at 129 low latitudes. OMI provides full global coverage every 2 to 3 days. Additional gaps are caused by a problem 130 with the OMI CCD, "row anomaly" (Torres et al., 2018) that effectively reduces the number of near-nadir 131 overpass views. A detailed OMI instrument description is given in Levelt et al. (2006). TCNO₂ is determined 132 in the visible spectral range from 405 to 465 nm where the NO₂ absorption spectrum has the maximum spectral structure and where there is little interference from other trace gas species (there is a weak water 133 134 feature in this range). OMI TCNO₂ overpass data are available for many ground sites (currently 719) from 135 the following NASA website. <u>https://avdc.gsfc.nasa.gov/index.php?site=666843934&id=13</u> 136

137 **2.2 PANDORA**

138

PANDORA is a sun-viewing instrument for SZA < 80° that obtains about 4000 spectra for clear-sky views of the sun in 20 seconds for each of two ranges UV (290 – 380 nm using a UV340 bandpass filter) and visible plus UV (280 – 525 nm using no filter). The overall measurement time is about 80 seconds including a 20 second dark-current measurements between each spectral measurement throughout the day. About 4000 clear-sky spectra for the UV and visible portions are separately averaged together to achieve very high signal to noise ratios (SNR). The UV340 filter for UV portion of the spectra reduces stray light effects from the visible wavelength range. A detailed description of PANDORA and its SNR is given

in Herman et al., (2009; 2015). The effect of moderate cloud cover (reduction of observed signal by a 146 147 factor of 8) in the PANDORA FOV on TCNO₂ retrievals is small (Herman et al., 2018). Cloud cover also reduces the number of measurements possible in 20 seconds, which potentially increases the noise level. 148 149 PANDORA is driven by a highly accurate sun tracker that points an optical head at the sun and transmits the received light to an Avantes 2048 x 32 pixel CCD spectrometer (AvaSpec-ULS2048 from 280 - 525 nm 150 151 with 0.6 nm resolution) through a 50 micron diameter fiber optic cable. The estimated TCNO₂ error is approximately 0.05 DU (1 DU = 2.69 x 10^{16} molecules cm⁻²) out of a typical value of 0.3 DU in relatively 152 clean areas and over 3 DU in highly polluted areas. PANDORA data are available for 250 sites. Some sites 153 154 have multi-year data but these sites are short-term sets, many of 155 https://avdc.gsfc.nasa.gov/pub/DSCOVR/Pandora/DATA 01/. campaign sites. 156

130

157 **3.0 Overpass Comparisons and Diurnal Variation of TCNO**₂

158

159 The contribution of NO₂ to air quality at the Earth's surface is usually a proportional function of 160 TCNO₂ that varies with the time of day and with the altitude profile shape (Lamsal et al., 2013; Bechle et al., 2013). Most of the NO₂ amount is usually located between 0 and 3 km altitude with a small amount of 161 about 0.1±0.05 DU (Dirksen et al. 2011) in the upper troposphere and stratosphere. Because of the 162 163 relatively short chemical lifetime, 3-4 hours (Liu et al., 2016), in the lower atmosphere, most of the NO_2 is located near (0 to 20 km) its sources (industrial activity, power generation, and automobile traffic). At 164 higher altitudes or in the winter months, the life time of NO₂ is longer permitting transport over larger 165 166 distances from its sources.







Fig. 2. Monthly average values of TCNO₂ for OMI and PANDORA at OMI overpass times

168

During the South Korean campaign (KORUS-AQ) in the spring of 2016 the diurnal variations of TCNO₂ vs days of the year DOY were determined for 6 sites (Herman et al., 2018), one of which is reproduced here (Fig. 1) for the city of Busan showing relatively low values of TCNO₂ in the morning (0.5

172 DU), moderately high values during the middle of the day (1.3 DU), and very high values on some of the

afternoons (2 to 3 DU). Of these data, OMI only observes midday values near the 13:30 time marked on
the Local Time axis of Fig.1 thereby missing very high values (2 to 3 DU) that frequently occur later in the
afternoon coinciding with times when people are outdoors returning from work.

176

177 In addition to missing the TCNO₂ diurnal variation, the OMI values are about half those observed 178 by PANDORA (Fig. 2) at the OMI overpass time, so that using OMI values to estimate NO₂ pollution 179 seriously underestimates the air quality problem even at midday. The shaded area in Fig.2 corresponds to 180 the period covered in the KORUS-AQ campaign 7 April to 11 June 2016 shown in Fig. 1. An extended time 181 series for Busan location is shown in Fig. 2

181 series for Busan location is shown in Fig. 3.



Fig. 3 Extended time series for Busan. Left Panel: individual matching PANDORA and OMI data points for the overpass time \pm 6 minutes. Right Panel: monthly averages.

182

Because of the different effective NO₂ FOV of PANDORA (measured in meters²) while tracking the moving sun position located in the heart of Busan (FOV distance d < 5 km for an SZA < 70^o used for TCNO₂ retrievals), both the daily (Fig. 3, left panel) and PANDORA monthly average variation (Fig. 3, right panel), obtained at the OMI overpass time, differs from the variation in the OMI TCNO₂ because of the much larger OMI FOV (13 x 24 km² at OMI nadir view) retrieval. Because of this, the OMI time series has low correlation (r² = 0.1) with the PANDORA time series.

189

The extended OMI vs PANDORA time series from 2012 – 2017 for Busan (Fig. 3) shows the same magnitude of differences seen during the KORUS-AQ period. A similar OMI vs PANDORA plot for total column ozone TCO₃ (Appendix Fig A1) shows good agreement between PANDORA and OMI indicating that the PANDORA instrument was operating and tracking the sun properly. Because the spatial variability of TCO₃, which is mostly in the stratosphere, is much less than for TCNO₂, the effect of different FOV's is minimized for ozone.

196



Fig. 4. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Mauna Loa Observatory, Hawaii, NASA Headquarters, Washington DC, and Waterflow, New Mexico. Waterflow, a small town, is listed for PANDORA under Four Corners, NM, a nearby landmark.

- 199The same type of differences, TCNO₂(PAN) > TCNO₂(OMI), are seen at a wide variety of sites (e.g.,200see Fig.4) for Northern Hemisphere sites and one site in the Southern Hemisphere where PANDORA has201an extended time series. Comparing extended Busan multi-year time series, some broad-scale correlation202can be seen with peaks in February 2013, January 2014, and in 2016. The data from Busan are different203than from many sites, since Busan is located very near the ocean causing a portion of the OMI FOV to be204over the unpolluted ocean areas, whereas PANDORA is located inland (Pusan University) in an area of205dense automobile traffic and quite near mountains capable of trapping air.
- 207 Figures 4 and 5 show a variety of different sites, ranging from the Mauna Loa Observatory location 208 at 3.4 km (11,161 feet) on a relatively clean Hawaiian Island surrounded by ocean to a polluted landlocked 209 semi-arid site at Waterflow, New Mexico near a power plant. All the sites considered show a significant 210 underestimate of OMI TCNO₂. A summary of the monthly average underestimates is given in Tables 1 and 211 2. For some sites there is evident correlation between the two offset measurements. For example, the 212 PANDORA at NASA Headquarters in Washington DC tracks the OMI measurement quite well on a monthly average basis with a correlation coefficient of $r^{2}(mn) = 0.7$ even though the daily correlation is low ($r^{2}(dy)$ 213 214 = 0.17). Other sites have only short periods of correlation and overall weak correlation (Table 1 showing 215 daily, dy and monthly, mn, correlation coefficients for the graphs in Figures 4 and 5)
- 216

217 TCNO₂(PAN) comparisons with TCNO₂(OMI) from Mauna Loa Observatory (Fig. 4) are not those 218 that might be expected, since the PANDORA observations are in an area where there are almost no 219 automobile emissions and certainly no power plants, yet PAN > OMI and TCNO₂(PAN) values are large 220 enough so that the pollution values (0.18 DU) are well above the stratospheric values (approximately 0.1 221 DU). OMI, which mainly measures values over the clean ocean, has an average value of 0.1 DU. The 222 PANDORA values suggest upward airflow from the nearby circumferential ring road and resort areas. The 223 Mauna Loa TCNO₂ values do not show any correlation with the recent increased volcanic activity at Mt. 224 Kilauea after 2016. Recently, the original Mauna Loa PANDORA has been replaced. The new instrument's 225 calibration will be reviewed before being added to the time series as part of a general data quality 226 assurance program that is starting with the most recently deployed or upgraded PANDORA instruments 227 at about 100 locations.

228 An interesting inland site is near the very small town of Waterflow, New Mexico (Fig. 4), where 229 two power plants located near the PANDORA site ceased operation on December 30, 2013 (Lindenmaier 230 et al., 2014). According to a quote from AZCentral Newspaper (Tuesday 31 December 2013) "Three coal-231 fired generators that opened in the 1960s near Farmington, N.M., closed Monday as part of a \$182 million 232 plan for Arizona Public Service Co. to meet environmental regulations, the utility reported". The TCNO₂ 233 data suggests that the actual shutdown occurred near October 15, 2013. After the shutdown, air quality 234 improved in the area with TCNO₂ decreasing from 0.4 DU to 0.28 DU. The remaining more efficient 235 generators continued to produce smaller NO₂ emissions. These were shut down at the end of 2016 with 236 little additional observed change in TCNO₂, since these boilers used NO₂ scrubbers (Dubey at al., 2018 in 237 preparation). A nearby highway (Route 64) about 2 km from the PANDORA site has little automobile 238 traffic.



Fig. 5. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Seoul South Korea, Boulder, Colorado, and Buenos Aires, Argentina (Raponi et al. 2017).

Table 1 Values of TCNO ₂	for PANDORA and OMI from	monthly aver	ages in Figs.	<mark>4 and 5</mark>
Name	Location (Lat, Lon)	PAN (DU)	OMI (DU)	r² (dy, mn)
Mauna Loa Hawaii	19.536°, -155.5762°	0.16	0.11	0.01, 0.30
NASA HQ Washington DC	38.882°, -77.01°	0.34	0.25	0.17, 0.70
Waterflow New Mexico ¹	36.797°, -108.48°	0.32	0.18	0.13, 0.52
Seoul South Korea	37.5644° <i>,</i> 126.934°	1.2	0.58	0.11, 0.06
Busan South Korea	35.2353° <i>,</i> 129.0825°	0.68	0.32	0.09, 0.10
Boulder Colorado	39.9909° <i>,</i> -105.2607°	0.27	0.17	0.04, 0.09
Buenos Aires Argentina	-34.5554°, -58.5062°	0.50	0.26	0.16, 0.08
Average		0.49	0.27	

243

244

Table 2 Average values of TCNO ₂ f	or PANDORA and OMI for addit	ional sites	
Name	Location (Lat, Lon)	PAN (DU)	OMI (DU)
Essex Maryland	39.31083°, -76.47444°	0.30	0.28
Baltimore Maryland	39.29149°, -76.59646°	0.45	0.27
Fresno California	36.7854°, -119.7731°	0.42	0.17
Denver La Casa Colorado	39.778° <i>,</i> -105.006°	0.68	0.19
GIST ²	35.226°,126.843°	0.42	0.20
HUFS ³	37.338°,127.265°	0.61	0.51
City College New York City	40.8153°,-73.9505°	<mark>0.60</mark>	<mark>0.40</mark>
Average		0.50	0.29

¹Waterflow, NM is listed for OMI data as Four Corners, NM, a nearby landmark ²Gwangju Institute of Science and Technology S. Korea ³Hankuk University Foreign Studies South Korea

245

246 Table 2 contains a summary of some sites that were part of short-term Discover-AQ campaigns in 247 Maryland, Texas, California, and Colorado, two longer-term sites in South Korea, and one in New York 248 City. Essex, Maryland is located on the Chesapeake Bay 10 km east of the center of Baltimore. The site is relatively clean (PAN = 0.3 DU) compared to the center of Baltimore (PAN = 0.45 DU), while OMI measures 249 250 about the same amounts for both sites (0.28 and 0.27 DU) because the OMI FOV is larger than the distance 251 between the two sites. The Houston Texas site contains 7 months of data from January to July 2013 with widespread NO₂ pollution permitting PANDORA and OMI to measure the same average values even 252 253 though PANDORA observes episodes on many days when TCNO₂ exceeds 1.5 DU for short periods at times 254 not observed by OMI. Observations in the small city of Fresno, California were during January when 255 agricultural sources of NO₂ were at a minimum (Almaraz, 2018), but automobile traffic in the center of 256 Fresno was significant. In this situation, PANDORA recorded the effect of automobile traffic while OMI 257 averaged the city of Fresno and surrounding fallow agricultural areas. The Denver La Casa location is in 258 the center of the city in an area with high amounts of local automobile traffic and near the Cherokee

259 power generating plant. The result is a high level of average pollution (0.42 DU) while OMI measures both 260 the city center and the surrounding relatively clean plains areas. The HUFS South Korean site is southeast 261 of Seoul in a fairly isolated valley. However, Seoul and its surrounding areas are a widespread transported 262 source of pollution so that both PANDORA and OMI measure elevated TCNO₂ amounts. In contrast, the 263 PANDORA GIST site is on the outskirts of a small city in southwestern South Korea with significant traffic. 264 The result is significant amounts of localized $TCNO_2$ (PANDORA = 0.42) surrounded by areas that produce 265 little NO₂ leading to OMI observing a very clean 0.2 DU. The average of sites in the two tables are similar 266 leading to ratios of PAN/OMI of 1.8 and 1.7 respectively. The estimated 50% increase in OMI retrievals of 267 TCNO₂ from using the geometry-dependent reflectivity (Vasilkov, 2017) for the most polluted sites will 268 narrow the disagreement with PANDORA. For example, OMI Seoul TCNO2 may become 0.87 DU 269 (PANDORA = 1.2 DU) and Buenos Aires 0.39 DU (PANDORA = 0.5 DU) still underestimating the amount of 270 NO₂ pollution and missing the significant diurnal variation.

271 For the six sites shown, the average OMI underestimate of TCNO₂ is approximately a factor of 1.8 272 at the overpass time on a monthly average basis with occasional spikes that exceed this amount. The bias 273 values range from 1.1 to 3.6, with higher biases tending to be associated with higher $TCNO_2$ values. The 274 factor of 1.8 underestimate ignores the frequent large values of TCNO₂ at other times during the day (Fig. 275 7). In addition, averaging $TCNO_2(PAN)$ over each entire day yields average values for the whole period that 276 are 10 to 20% higher than just averaging over midday values that matched the OMI overpass time. Aside 277 from the absolute magnitude, the short-term variations (over several months) are similar for both OMI and PANDORA although mostly not correlated. If correlation coefficients r² are generated from linear fits 278 279 to scatter plots of TCNO₂ from OMI vs PANDORA, the correlation is mostly poor (Examples, $r^2 =:$ Seoul 280 0.06, Mauna Loa 0.3 NASA HQ 0.7, see Figs. 4 and 5). Additional sites with shorter PANDORA time series 281 of TCNO₂ show similar behavior.

Duncan et al. (2016) estimated trends from OMI TCNO₂ time series and found that the Seoul metropolitan area had a decrease of -1.5 ± 1.3 %/Year (2005 – 2014) consistent with OMI estimated change of -1.4 ± 1 %/year (2012 -2018) in this paper. However, for the small area near Yonsei University, the decrease estimated from PANDORA is -5.8 ± 0.75 %/Year. Park (2019) estimates that metropolitan Seoul has decreased in population even as surrounding areas have increased population.

287 The average percent differences between OMI and PANDORA shown in Fig. 6 are relatively 288 constant over time for each site with small changes over each multi-year observation period. The 289 differences between OMI and PANDORA are provided by forming the percent differences of the daily 290 TCNO₂ values (Fig. 6) in the form 100(OMI – PAN)/PAN. Also shown are the average percent differences 291 and the linear fit slopes in percent change per year of the percent differences over the multi-year period. 292 For example, the Boulder percent difference goes from -31% to -23% over 4 years. Of the six sites in shown 293 in Fig. 6, two have statistically significant slopes, Seoul South Korea 2.1±0.5 %/Year and NASA 294 Headquarters in Washington DC 3.4 \pm 0.9 %/Year at the 2 σ level suggesting a significant area average 295 increase in pollution compared to PANDORA's local values.



Fig. 6 Percent differences between OMI and PANDORA. The slopes are the absolute change in the percent difference. The LS Means are least squares means with the corresponding error estimates

297	For some sites (see Fig. 6), PANDORA and OMI trends are the same (Waterflow, NM,
298	Buenos Aires, and Mauna Loa) while the other 3 sites show significantly different trends (Boulder,
299	NASA HQ, and Seoul).
300	The results for Busan (from Fig. 3) show a least squares average for the percent difference
301	of -48 \pm 0.8% for the 2012 – 2018 period with a slope of 6.8 \pm 1%/Year. There is a decrease in the
302	percent difference after October 2015 (Fig. 3) that is mainly from PANDORA seeing less TCNO $_2$
303	than during the 2012 – 2014 period. There is a gap in the Busan time series from July 2014 until
304	April 2015 when the original PANDORA was replaced with a new instrument. The calibrations of
305	both PANDORAS appear to be correct. Because of the break in the time series it is not clear
306	whether there was a change in local conditions around Pusan University compared to the wide
307	area observed by OMI.
308	
309	
310	
311	
312	
313	
314	
315	
316	
317	
318	
319	3.1 Diurnal Variation at NASA HQ Washington DC
320	Figure 7 shows details of the daily diurnal variation of TCNO ₂ on the roof of NASA Headquarters

Washington, DC adjacent to a major cross-town highway (I695) for every day during each month of 2015
 for local time vs DOY. The midday observing local standard time for OMI is marked for each graph.



Fig. 7A. TCNO₂ diurnal variation (DU) from January to June, NASA Headquarters Washington, DC from January 2015 to June 2015. The approximate OMI overpass time near 13:30 hours is marked.





The amount of $TCNO_2$ is mostly from the adjacent highway and the surrounding urban area with heavy traffic. The relatively moderate $TCNO_2$ values (0.4 to 0.8 DU) are probably a testament to the effectiveness of catalytic converters mandatory on all US automobiles in such a high traffic area (Bishop and Steadman, 2015).



Fig. 8A TCNO₂ diurnal variation (DU) at CCNY in New York City January to June 2018. The approximate OMI overpass time near 13:30 hours is marked



Fig. 8B TCNO₂ diurnal variation (DU) at CCNY in New York City July to December 2018. The peak near 5 DU occurs on 13 July 2018 between 11:20 and 12:30 EST. The approximate OMI overpass time near 13:30 hours is marked.

Figure 8 contains the daily TCNO₂ diurnal variability vs DOY for each month measured by a PANDORA from the roof of a building on the CCNY (City College of New York) campus in the middle of Manhattan in New York City (NYC). From the values shown, the pollution levels are quite high, rivaling the

pollution levels in Seoul, South Korea (see Fig. 5). OMI at its mid-day overpass time would detect some of 335 336 the high-level pollution events, but miss many others occurring mostly in the afternoon. There are a 337 significant number of days in all the months where the $TCNO_2$ levels appear to be low (e.g., blue color in 338 July and October), but the blue color still represents significant pollution levels (TCNO₂(PAN) > 0.5 DU) 339 that are small only compared to the peak values during the month (TCNO₂(PAN) > 1 DU). The highest 340 amount of TCNO₂ recorded during 2018 was about 5DU on 13 July 2018 from 11:20 and 12:30 EST (a time with very light winds (1 km/hr) and moderate temperature (25° C). There were many smaller peaks 341 between 2 and 3 DU throughout the year. Extreme cases of high NO₂ amounts are frequently associated 342 343 with the local meteorology indications of stagnant air (Harkey et al., 2015),

344 For both Washington DC (Fig. 7) and New York City (Fig. 8) there is strong day-to-day and month 345 to month variability that depends on the local meteorological conditions (Seo et al., 2018; Zeng et al., 346 2015) and the amount of automobile traffic in the area (Andersen et al., 2011; Amin et al., 2017). High 347 TCNO₂ events occur most often in the afternoon such that the OMI overpass near 13:30 would miss most high TCNO₂ events. Poor air quality affecting respiratory health would be improperly characterized by 348 349 both the OMI average values being too low (Fig. 4) and by missing the extreme pollution events that occur 350 frequently in the late afternoon. The high value of TCNO₂ that occurred on 5 August (2.2 DU) at 07:45 EST 351 for Washington DC is not a retrieval error (SZA less than 70°), but is a one-time anomaly in 2015 compared 352 to more usual high values of 1.5 DU with an occasional spike to 2 DU. It should be noted that TCNO₂ does 353 not accurately represent the NO₂ concentration at the surface, since it is mostly a measure of the amount 354 in the lower 2 km. However, it is roughly proportional to the surface measurements close to the pollution 355 sources (Bechle et al., 2013; Knepp et al., 2014) with the exact proportionality dependent on the profile 356 shape near the ground.

Similar daily diurnal variation graphs of TCNO₂ (Figs. 7 and 8) could be shown for each site. However, the basic idea is the same for each site. OMI underestimates the amount of TCNO₂ because of its large FOV and misses most of the peak events at other times of the day. For some sites, such as Busan and Seoul, the peak values can reach 3 DU and above late in the afternoon, which are never seen by OMI (Herman et al., 2018).

362 Figure 9 for CCNY is similar to the graphs in Figs. 4 - 6 showing the relative behavior between PANDORA and OMI but including only OMI pixels that are at a distance D < 30 km from CCNY. The results 363 364 are almost identical to those when D < 80 km. There is a period in March 2018 when OMI TCNO₂ slightly exceeded that measured by PANDORA. OMI with its large FOV may be seeing part of the chemically driven 365 366 seasonal variation, while PANDORA is seeing a nearly constant source driven amount mostly from 367 automobile traffic. For most days during 2018, PAN(TCNO₂) > OMI(TCNO₂) with the average value for PAN 368 = 0.65 DU and for OMI = 0.45 DU (Fig. 9 Panel B). The percent difference plot shows that there is a 369 systematic increase between PANDORA and OMI TCNO₂ from a value 10% to a value of 50%.



Fig. 9 TCNO₂ overpass time series for CCNY in Manhattan, New York City. OMI pixels are at a distance D < 30 km from CCNY. Panel A: OMI overpass TCNO₂ (Black) compare with OMI (Red). Panel B: Monthly Lowess(f) fit to the daily overpass data. Panel C: Percent difference 100(OMI – PAN)/PAN calculated from the data in Panel A

371 4.0 Summary

372 Examination of long-term TCNO₂ monthly average time series from OMI satellite and PANDORA 373 ground-based observations show that OMI systematically underestimates the amount of NO₂ in the 374 atmosphere by an average factor of 1.5 to 2 at the local OMI overpass time near the equator crossing time of 13:30±1:30. As shown in Fig. 6 for TCNO₂, 100(OMI – PAN)/PAN least squares mean underestimates are 375 much larger than error estimates. These differences are reduced for the smaller pixel size TropOMI TCNO2 376 377 values (Judd et al.,2019). In addition, the PANDORA diurnal time series for every day during a year at each 378 site (only two typical sites are shown in this paper, NYC and NASA-HQ) shows peaks in TCNO₂ that are 379 completely missed by only observing at mid-day. The result is that estimates of air quality related to health effects from OMI observations are strongly underestimated almost everywhere as shown at all the 380 381 sites with a long PANDORA record. In comparisons to PANDORA, OMI data are mostly uncorrelated or weakly correlated (e.g., Seoul correlation coefficient $r^2 = 0.06$, Mauna Loa $r^2 = 0.3$), while NASA HQ in 382 383 Washington, DC shows a correlation on a seasonal basis (NASA HQ $r^2 = 0.7$) suggesting a wide area coordinated source of NO₂ (most likely automobile traffic). The data from CCNY shows some correlation 384

385 between the locations of the peaks and troughs. Seven short term TCNO₂ time series were examined 386 showing similar results (Table 1), except when the pollution region is widespread as in the Seoul South 387 Korea region. The conclusion is that while OMI satellite TCNO₂ data are uniquely able to assess regional 388 long-term trends in TCNO₂ and provide a measure of the regional distribution of pollutants, the OMI data 389 cannot properly assess local air quality or the effect on human health over extended periods in urban or 390 industrial areas. This will continue to be the case, but to a lesser degree, when the OMI TCNO₂ data are 391 improved by reprocessing with a new geometry-dependent reflectivity (Vasilkov, 2017) and by the smaller 392 FOV of TropOMI. The analysis shows that locating PANDORAs at polluted sites could provide quantitative 393 corrections for spatial and temporal biases that affect the determination of local air quality from satellite 394 data. Satellite detection of diurnal variation of TCNO2 will be improved with the upcoming launch of three 395 planned geostationary satellites over Korea, US, and Europe To verify the proper operation of the various 396 PANDORA instruments a similar analysis for Total Column Ozone TCO was performed (see Appendix) and 397 shows close agreement between OMI and PANDORA, with the largest difference occurring for Mauna Loa 398 Observatory at 3.4 km altitude, where PANDORA misses the ozone between the surface and 3.4 km.

399 Appendix

400 A1 Ozone This section shows the corresponding PANDORA total column ozone (TCO) values 401 compared to OMI TCO for Busan South Korea (Fig. A1) that shows close agreement for the entire 2012 – 402 2017 period. The different fields of view for OMI and PANDORA have a much smaller effect because of 403 the greater spatial uniformity of stratospheric ozone compared to tropospheric NO₂. Additional sites are 404 summarized in Table A1. The largest TCO difference (15 DU or 5.6%) occurs for Mauna Loa Observatory 405 (Altitude = 3.4 km) compared to OMI (Average altitude = Sea Level). The close results show that the 406 PANDORA was working properly and pointing accurately at the sun. The PANDORA TCO data shown here 407 use a mid-latitude effective ozone temperature correction from model calculations that may not be 408



Fig. A1 Monthly average values of TCO for OMI and PANDORA at OMI overpass times for Busan South Korea

Year

409

201

Year

	Table A1 Average values of TCO ₃ for PANDORA and OMI			
	Location	PAN	OMI	Percent
		(DU)	(DU)	Difference
	Mauna Loa Observatory Hawaii (3.394 km)*	254	269	5.6
	NASA HQ Washington DC (0.02 km)	308	314	1.9
	Waterflow New Mexico (1.64 km)	293	292	0.3
	Yonsei University Seoul South Korea (0.07 km)	317	325	2.5
	Busan University Busan South Korea(0.03 km)	313	315	0.6
	Boulder, Colorado (NOAA Bldg) (1.617 km)	299	302	1.0
	Buenos Aires, Argentina (0.025 km)	279	284	1.8
	Essex, Maryland (0.012 km)	299	301	0.7
	Baltimore, Maryland (0.01 km)	296	296	0.0
	Fresno, California (0.939 km)	306	309	1.0
	Denver La Casa Colorado (1.6 km)	292	294	0.7
	Gwangju Institute of Science and Technology (GIST) S. Korea (0.021 km)	302	307	1.6
	Hankuk University Foreign Studies (HUFS) South Korea (0.04 km)	318	326	2.5
	City College Manhattan New York City (0.04 km)	316	325	2.8
	Average	299	304	1.6
410				
411				
412	* OMI observes the sea level value of TCO_3			
413	The ozone retrievals shown here use an average effective ozone temperatu	re instead	of a local	Iv
414	measured ozone temperature (Herman et al., 2015:2017).			
415				
416				
417				
418				
44.0				
419				
420				
421	Acknowledgement: This project is supported by the Korea Ministry of Environment	t (MOE) as	Public Tec	hnology
422	Program based on Environmental Policy (2017000160001), by the Los Alamos Nati	onal Labo	ratory's L	aboratory

423 Directed Research and Development program and by the NASA Pandora project managed by Dr. Robert424 Swap.

425 **References**

Almaraz, Maya,Edith Bai, Chao Wang, Justin Trousdel, Stephen Conley, Ian Faloona and Benjamin Z.
Houlton, Agriculture is a major source of NOx pollution in California, SCIENCE ADVANCES,31,
DOI:10.1126/sciadv.aao3477, 2018.

429

430	Amin, Md. Shohel Reza, Umma Tamima, and Luis Amador Jimenez, "Understanding Air Pollution from
431	Induced Traffic during and after the Construction of a New Highway: Case Study of Highway 25 ir
432	Montreal," Journal of Advanced Transportation, vol. 2017, Article ID 5161308, 14 pages
433	2017. https://doi.org/10.1155/2017/5161308, 2017.

- 434
- 435 Andersen, M. Hvidberg, S.S. Jensen, M. Ketzel, S. Loft, M. Sørensen, A. Tjønneland, K. Overvad, O. Raasc
- 436 hou-Nielsen Chronic obstructive pulmonary disease and long-term exposure to traffic-related air
- 437 pollution: A cohort study, Am. J. Respir. Crit. Care Med., 183, pp. 455-461, <u>10.1164/rccm.201006</u>
- 438 **09370C, 2011**.

Bechle, M. J.; Millet, D. B.; Marshall, J. D. Remote sensing of exposure to NO2: Satellite versus groundbased measurement in a large urban area. Atmos. Environ., 69, 345-353, 2013

441

Bishop, Gary A. and Donald H. Stedman, Reactive Nitrogen Species Emission Trends in Three Light/Medium-Duty United States Fleets, *Environmental Science & Technology* 2015 *49* (18), 11234-11240,
DOI: 10.1021/acs.est.5b02392, 2015.

445

Boersma, K. F., H. J. Eskes, E. J. Brinksma, Error analysis for tropospheric NO₂ retrieval from space, J.

447 Geophys. Res. Atmos., <u>https://doi.org/10.1029/2003JD003962</u>, 2004.

448 Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V.,

449 Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved

450 tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, Atmos. Meas. Tech.,

- 451 4, 1905-1928, https://doi.org/10.5194/amt-4-1905-2011, 2011.
- 452 Boersma, K. F., Vinken, G. C. M., & Eskes, H. J., Representativeness errors in comparing chemistry
- transport and chemistry climate models with satellite UV–Vis tropospheric column retrievals.
- 454 Geoscientific Model Development, 9(2), 875-898, 2016.

455 Choudhari, Sheetal Korde, Minal Chaudhary, Sachin Bagde, Amol R Gadbai and Vaishali Joshi, Nitric oxide 456 а review, World Journal of Oncology 2013, 11:118, and cancer: Surgical http://www.wjso.com/content/11/1/118, 2013. 457

458

Cleveland, William S., LOWESS: A program for smoothing scatterplots by robust locally weighted
regression. The American Statistician. 35 (1): 54. JSTOR 2683591. doi:10.2307/2683591, 1981.

- 461
- 462

463 Dirksen, Ruud J., K. Folkert Boersma, Henk J. Eskes, Dmitry V. Ionov, Eric J. Bucsela, Pieternel F. Levelt, 464 Hennie M. Kelder, Evaluation of stratospheric NO2 retrieved from the Ozone Monitoring Instrument: 465 Intercomparison, diurnal cycle, and trending, J. Geophys. Res., 116, 466 https://doi.org/10.1029/2010JD014943, 2011.

467

Duncan, B. N., L. N. Lamsal, A. M. Thompson, Y. Yoshida, Z. Lu, D. G. Streets, M. M. Hurwitz, and K. E.
Pickering, A space-based, high-resolution view of notable changes in urban NOx pollution around the
world (2005–2014), J. Geophys. Res. Atmos., 121, 976–996, doi:10.1002/2015JD024121, 2016.

471

Goldberg, D. L., Saide, P. E., Lamsal, L. N., de Foy, B., Lu, Z., Woo, J.-H., Kim, Y., Kim, J., Gao, M., Carmichael,
G., and Streets, D. G.: A top-down assessment using OMI NO2 suggests an underestimate in the NOx
emissions inventory in Seoul, South Korea during KORUS-AQ, Atmos. Chem. Phys. Discuss.,
https://doi.org/10.5194/acp-2018-678, in review, 2018.

476

Harkey, M., Holloway, T., Oberman, J., and Scotty, E., An evaluation of CMAQ NO2 using observed
chemistry-meteorology correlations, J. Geophys. Res. Atmos., 120, 11,775– 11,797,
doi:10.1002/2015JD023316, 2015.

480

Herman, J., A. Cede, E. Spinei, G. Mount, M. Tzortziou, and N. Abuhassan, NO2 column amounts from
ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique:
Intercomparisons and application to OMI validation, J. Geophys. Res., 114, D13307,
doi:10.1029/2009JD011848, 2009.

485

489

Herman, J.R., R.D. Evans, A. Cede, N.K. Abuhassan, I. Petropavlovskikh, and G. McConville, Comparison
of Ozone Retrievals from the Pandora Spectrometer System and Dobson Spectrophotometer in Boulder
Colorado, Atmos. Meas. Tech., 8, 3407–3418, 2015 doi:10.5194/amt-8-3407-2015.

Herman, J., Evans, R., Cede, A., Abuhassan, N., Petropavlovskikh, I., McConville, G., Miyagawa, K., and
Noirot, B.: Ozone comparison between Pandora #34, Dobson #061, OMI, and OMPS in Boulder,
Colorado, for the period December 2013–December 2016, Atmos. Meas. Tech., 10, 3539-3545,
https://doi.org/10.5194/amt-10-3539-2017, 2017.

494

Herman, J., Spinei, E., Fried, A., Kim, J., Kim, J., Kim, W., Cede, A., Abuhassan, N., and Segal-Rozenhaimer,
M.: NO₂ and HCHO measurements in Korea from 2012 to 2016 from Pandora spectrometer instruments
compared with OMI retrievals and with aircraft measurements during the KORUS-AQ campaign, Atmos.
Meas. Tech., 11, 4583-4603, https://doi.org/10.5194/amt-11-4583-2018, 2018.

499

Ialongo, I., Herman, J., Krotkov, N., Lamsal, L., Boersma, K. F., Hovila, J., and Tamminen, J.: Comparison of
 OMI NO₂observations and their seasonal and weekly cycles with ground-based measurements in Helsinki,

502 Atmos. Meas. Tech., 9, 5203-5212, https://doi.org/10.5194/amt-9-5203-2016, 2016.

503

Judd, Laura M., Jassim A. Al-Saadi, Lukas C. Valin, R. Bradley Pierce, Kai Yang, Scott J. Janz, Matthew G.
Kowalewski, James J. Szykman, Martin Tiefengraber, and Moritz Mueller, The Dawn of Geostationary Air

506 Quality Monitoring: Case Studies From Seoul and Los Angeles, Environ. 507 Sci., https://doi.org/10.3389/fenvs.2018.00085, 2018. 508 509 Judd, L. M., Al-Saadi, J. A., Janz, S. J., Kowalewski, M. G., Pierce, R. B., Szykman, J. J., Valin, L. C., Swap, R., 510 Cede, A., Mueller, M., Tiefengraber, M., Abuhassan, N., and Williams, D.: Evaluating the impact of spatial 511 resolution on tropospheric NO₂ column comparisons within urban areas using high-resolution airborne 512 data, Atmos. Meas. Tech. Discuss., https://doi.org/10.5194/amt-2019-161, in review, 2019. 513 514 Kleipool, Q.L., M.R. Dobber, J.F. De Haan and P.F. Levelt, Earth Surface Reflectance Climatology from Three 515 Years of OMI Data, Journal of Geophysical Research, 113, doi:10.1029/2008JD010290, 2008. 516 517 Knepp, T., M. Pippin, J. Crawford, Jim Szykman, R. Long, L. Cowen, A. Cede, N. Abuhassan, J. Herman, R. 518 Delgado, J. Compton, T. Berkoff, J. Fishman, D. Martins, R. Stauffer, A. Thompson, A. Weinheimer, D. 519 Knapp, D. Montzka, D. Lenschow, AND D. Neil. Estimating Surface NO2 and SO2 Mixing Ratios from Fast-520 Response Total Column Observations and Potential Application to Geostationary Missions. JOURNAL OF 521 ATMOSPHERIC CHEMISTRY. Springer, New York, NY, 0(0):1-26, (2014). 522 523 Kollonige, D. E., Thompson, A. M., Josipovic, M., Tzortziou, M., Beukes, J. P., Burger, R., ... Laakso, L. (2018). 524 OMI satellite and ground-based Pandora observations and their application to surface NO2 estimations 525 at terrestrial and marine sites. Journal of Geophysical Research: Atmospheres, 123, 1441–1459. 526 https://doi.org/10.1002/2017JD026518, 2018. 527 528 Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., Chan, K. L., 529 Wenig, M., and Zara, M.: The version 3 OMI NO₂ standard product, Atmos. Meas. Tech., 10, 3133-3149, 530 https://doi.org/10.5194/amt-10-3133-2017, 2017. 531 532 Lamsal, L & Martin, Randall & Parrish, David & Krotkov, Nickolay., Scaling Relationship for NO₂ Pollution 533 and Urban Population Size: A Satellite Perspective. Environmental science & technology. 47. 534 10.1021/es400744g, 2013. 535 Lamsal, L. N., Krotkov, N. A., Celarier, E. A., Swartz, W. H., Pickering, K. E., Bucsela, E. J., Martin, R. 536 537 V., Philip, S., Irie, H., Cede, A., Herman, J., Weinheimer, A., Szykman, J. J., and Knepp, T. N.: Evaluation 538 of OMI operational standard NO₂ column retrievals using in situ and surface-based NO₂ observations, 539 Atmos. Chem. Phys. Discuss., 14, 14519-14573, doi:10.5194/acpd-14-14519-2014, 2014. 540 541 Lamsal, L. N., S. J. Janz, N. A. Krotkov, K. E. Pickering, R. J. D. Spurr, M. G. Kowalewski, C. P. Loughner, J. H. 542 Crawford, W. H. Swartz, and J. R. Herman, High-resolution NO₂ observations from the Airborne Compact 543 Atmospheric Mapper Retrieval and validation, J. Geophys. Res. Atmos., 122, 1953–1970, 544 doi:10.1002/2016JD025483, 2017 545

546 Levelt, P. F., Van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, 547 J. O. V., and Saari, H.: The Ozone Monitoring Instrument, IEEE T. Geosci. Remote, 44, 1093–1101, doi:10.1109/tgrs.2006.872333, 2006. 548

549

550 Lindenmaier, Rodica, Manvendra K. Dubey, Bradley G. Henderson, Zachary T. Butterfield, Jay R. Herman, 551 Thom Rahn · S.-H. Lee, Multiscale observations of CO2, (CO2)-C-13, and pollutants at Four Corners for emission verification and attribution, Proceedings of the National Academy of Sciences 06/2014; 552 553 111(23):8386-8391. DOI:10.1073/pnas.132188311, 2014

- 555 Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO_x lifetimes and emissions of cities and 556 power plants in polluted background estimated by satellite observations, Atmos. Chem. Phys., 16, 5283-557 5298, https://doi.org/10.5194/acp-16-5283-2016, 2016.
- 558

554

- 559 Lorente, A., Boersma, K. F., Stammes, P., Tilstra, L. G., Richter, A., Yu, H., Kharbouche, S., and Muller, J.-P.: 560 The importance of surface reflectance anisotropy for cloud and NO₂ retrievals from GOME-2 and OMI, 561 Atmos. Meas. Tech., 11, 4509-4529, https://doi.org/10.5194/amt-11-4509-2018, 2018.
- 562

563 Nowlan, C. R., Liu, X., Leitch, J. W., Chance, K., González Abad, G., Liu, C., Zoogman, P., Cole, J., Delker, T., 564 Good, W., Murcray, F., Ruppert, L., Soo, D., Follette-Cook, M. B., Janz, S. J., Kowalewski, M. G., Loughner, C. P., Pickering, K. E., Herman, J. R., Beaver, M. R., Long, R. W., Szykman, J. J., Judd, L. M., Kelley, P., Luke, 565 566 W. T., Ren, X., and Al-Saadi, J. A.: Nitrogen dioxide observations from the Geostationary Trace gas and 567 Aerosol Sensor Optimization (GeoTASO) airborne instrument: Retrieval algorithm and measurements 568 during DISCOVER-AQ Texas 2013, Atmos. Meas. Tech., 9, 2647-2668, https://doi.org/10.5194/amt-9-569 2647-2016, 2016.

570

571 O'Byrne, G., R. V. Martin, A. van Donkelaar, J. Joiner, and E. A. Celarier, Surface reflectivity from the Ozone 572 Monitoring Instrument using the Moderate Resolution Imaging Spectroradiometer to eliminate clouds: 573 Effects of snow on ultraviolet and visible trace gas retrievals, J. Geophys. Res., VOL. 115, D17305, 574 doi:10.1029/2009JD013079, 2010.

575

Marchenko, S., N. A. Krotkov, L. N. Lamsal, E. A. Celarier, W. H. Swartz, and E. J. Bucsela, Revising the 576 577 slant column densityretrieval of nitrogen dioxide observed by the Ozone Monitoring Instrument, J. 578 Geophys. Res. Atmos., 120, 5670–5692, doi:10.1002/2014JD022913, 2015

579 580 Park, Se Hoon, Seoul, The Wiley Blackwell Encyclopedia of Urban and Regional Studies. Edited by 581 Anthony Orum. John Wiley & Sons Ltd. Published 2019 by John Wiley & Sons Ltd. DOI: 10.1002/9781118568446.eurs0283, 2019.

582

583

Raponi, Marcelo, Cede, A, Santana Diaz, Daniel, Sanchez, R, A. Otero, L, O. Salvador, J, R. Ristori, P, Quel, 584 585 Eduardo. Total Column Ozone Measured In Buenos Aires Between March And November 2017, Using A 586 Pandora Spectrometer System. Anales AFA. 29(2), 46-50. 10.31527/analesafa.2018.29.2.46, 2018.

Seo, J., Park, D.-S. R., Kim, J. Y., Youn, D., Lim, Y. B., and Kim, Y.: Effects of meteorology and

emissions on urban air quality: a quantitative statistical approach to long-term records (1999–2016) in Seoul, South Korea, Atmos. Chem. Phys., 18, 16121-16137, https://doi.org/10.5194/acp-18-

- 587
- 588 589

16121-2018, 2018.



- 592 Torres, O., Bhartia, P. K., Jethva, H., and Ahn, C.: Impact of the ozone monitoring instrument row
- anomaly on the long-term record of aerosol products, Atmos. Meas. Tech., 11, 2701-2715,
- 594 https://doi.org/10.5194/amt-11-2701-2018, 2018.
- 595
- 596 Tzortziou, M.; Parker, O.; Lamb, B.; Herman, J.R.; Lamsal, L.; Stauffer, R.; Abuhassan, N. Atmospheric
- 597 Trace Gas (NO₂ and O₃) Variability in South Korean Coastal Waters, and Implications for Remote
- Sensing of Coastal Ocean Color Dynamics. *Remote Sens.*, *10*, 1587, ; doi:10.3390/rs10101587, 2018.
- 599

Vasilkov, A., Qin, W., Krotkov, N., Lamsal, L., Spurr, R., Haffner, D., Joiner, J., Yang, E.-S., and Marchenko,
S.: Accounting for the effects of surface BRDF on satellite cloud and trace-gas retrievals: a new approach
based on geometry-dependent Lambertian equivalent reflectivity applied to OMI algorithms, Atmos.
Meas. Tech., 10, 333-349, https://doi.org/10.5194/amt-10-333-2017, 2017.

604

Zara, M., Boersma, K. F., De Smedt, I., Richter, A., Peters, E., van Geffen, J. H. G. M., Beirle, S.,
Wagner, T., Van Roozendael, M., Marchenko, S., Lamsal, L. N., and Eskes, H. J.: Improved slant column
density retrieval of nitrogen dioxide and formaldehyde for OMI and GOME-2A from QA4ECV:
intercomparison, uncertainty characterization, and trends, Atmos. Meas. Tech., 11, 4033-4058,
https://doi.org/10.5194/amt-11-4033-2018, 2018.

- Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang,
 D., Pöschl, U., Cheng, Y. F., and He, K. B.: Exploring the severe winter haze in Beijing: the impact of
 synoptic weather, regional transport and heterogeneous reactions, Atmos. Chem. Phys., 15, 2969–
- 614 2983, <u>https://doi.org/10.5194/acp-15-2969-2015</u>, 2015
- 615
- 616
- 617

- 618 Figure Captions
- 619 Fig 1 Diurnal variation of TCNO₂ measured at Pusan University in Busan South Korea
- Fig. 2. Monthly average values of TCNO₂ for OMI and PANDORA at OMI overpass times
- Fig. 3 Extended time series for Busan. Left Panel: individual matching PANDORA and OMI data
- 622 points for the overpass time ± 6 minutes. Right Panel: monthly averages.
- 623 Fig. 4. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Mauna Loa
- 624 Observatory, Hawaii, NASA Headquarters, Washington DC, and Waterflow, New Mexico.
- Fig. 5. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Seoul South Korea,
 Boulder, Colorado, and Buenos Aires, Argentina (Raponi et al. 2018).
- Fig. 6 Percent differences between OMI and PANDORA. The slopes are the absolute change in the
 percent difference. For example, the Boulder percent difference goes from -31% to -23% over 4 years.
- 629 Fig. 7A TCNO₂ diurnal variation (DU) from January to June, NASA Headquarters Washington, DC
- 630 from January 2015 to June 2015. The approximate OMI overpass time near 13:30 hours is marked.
- Fig. 7B TCNO₂ diurnal variation (DU) from July to December, NASA Headquarters Washington, DC from
 July 2015 to December 2015. The approximate OMI overpass time near 13:30 hours is marked
- Fig. 8A TCNO₂ diurnal variation (DU) at CCNY in New York City January to June 2018. The approximate
 OMI overpass time near 13:30 hours is marked.
- Fig. 8B TCNO₂ diurnal variation at CCNY in New York City July to December 2018. The peak near 5 DU
- occurs on 13 July 2018 between 11:20 and 12:30 EST. The approximate OMI overpass time near 13:30hours is marked.
- Fig. 9 TCNO₂ overpass time series for CCNY in Manhattan, New York City. Panel A: OMI overpass
- TCNO₂ (Black) compare with OMI (Red). Panel B: Monthly Lowess(0.08) fit to the daily overpass
- 640 data. Panel C: Percent difference 100(OMI PAN)/PAN calculated from the data in Panel A
- Fig. A1 Monthly average values of TCO for OMI and PANDORA at OMI overpass times for Busan SouthKorea





Fig. 3 Extended time series for Busan. Left Panel: individual matching PANDORA and OMI data points for the overpass time ± 6 minutes. Right Panel: monthly averages.

651 FIGURE 3



Fig. 4. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Mauna Loa Observatory, Hawaii, NASA Headquarters, Washington DC, and Waterflow, New Mexico.

653 FIGURE 4



Fig. 5. PANDORA compared to OMI. Extended TCNO₂ overpass time series for Seoul South Korea, Boulder, Colorado, and Buenos Aires, Argentina (Raponi et al. 2017).

FIGURE 5

Fig. 6 Percent differences between OMI and PANDORA. The slopes are the absolute change in the percent difference. For example, the Boulder percent difference goes from -31% to -23% over 4 years. The LS Means are least squares means with the corresponding error estimates

FIGURE 6

Fig. 7A TCNO₂ diurnal variation (DU) from January to June, NASA Headquarters Washington, DC from January 2015 to June 2015. The approximate OMI overpass time near 13:30 hours is marked

FIGURE 7A

Fig. 7B TCNO₂ diurnal variation (DU) from July to December, NASA Headquarters Washington, DC from July 2015 to December 2015. The approximate OMI overpass time near 13:30 hours is marked

FIGURE 7B

Fig. 8A TCNO₂ diurnal variation (DU) at CCNY in New York City January to June 2018. The approximate OMI overpass time near 13:30 hours is marked

665 Figure 8A

Fig. 8B TCNO₂ diurnal variation (DU) at CCNY in New York City July to December 2018. The peak near 5 DU occurs on 13 July 2018 between 11:20 and 12:30 EST. The approximate OMI overpass time near 13:30 hours is marked.

667 Figure 8B

Fig. 9 TCNO₂ overpass time series for CCNY in Manhattan, New York City. OMI pixels are at a distance D < 30 km from CCNY. Panel A: OMI overpass TCNO₂ (Black) compare with OMI (Red). Panel B: Monthly Lowess(f) fit to the daily overpass data. Panel C: Percent difference 100(OMI – PAN)/PAN calculated from the data in Panel A

669 Figure 9

Fig. A1 Monthly average values of TCO_3 for OMI and PANDORA at OMI overpass times for Busan South Korea

671

672 FIGURE A1