Author's response to Anonymous Referee #2's comments (amt-2019-138)

Last updated: Tuesday $18^{\rm th}$ June, 2019 at 07:34 UT

The authors kindly thank the anonymous reviewer for their comprehensive review of the paper. We believe that the suggestions made that we have been able to address have significantly improved the paper. Below are our responses to the points raised, indicating the changes made to the manuscript where appropriate.

No.	Referee's comment	Author's response and changes
	1. As shown in "Spatial distribution of errors	
	associated with multistatic meteor radar", Earth,	
	Planets and Space, 70:93,	
	doi.org/10.1186/s40623-018-0860-2, 2018, some	
	data need to be excluded from calculations,	We acknowledge that horizontal wind/momentum
	notably those close to the midpoint between the	flux estimation errors arising from radial velocity
	radar transmitter and receivers. In Fig 1., lower	measurement errors are amplified for meteors
	right, (bistatic case) these correspond to the	close to the transmitter-receiver midpoint.
	region to the south-east of the transmitter, .	However, as the reviewer alludes to, there are so
	While modest in number in this region, the radial	few meteors detected here (less than around 10%
	velocities measured here relate mostly to the	per unit area than at the peaks of the distribution
	vertical velocities, and are best excluded for	in our case), that we did not consider filtering
	calculations of winds and momentum flux. Failing	them from the analysis or weighting them in any
	to exclude these points can adversely affect the	way. We believe that weighting the meteor
1.	errors. For a monostatic system, these are all	distribution to minimize the errors in the fitted
	data close to overhead of the transmitter-receiver	wind/momentum fluxes is a separate topic
	system (the blue-coloured cluster around the	warranting its own investigation, and so we have
	transmitter in the lower left figure). Typically	added a comment to this effect to the discussion
	data within 12-15 degrees of vertical are removed	section of the manuscript. Nevertheless, the
	for the monostatic case (e.g. see Radio Sci., 32,	simulations performed in this paper also do not
	833-865, 1997). A similar exclusion process (or at	incorporate any weighting of the meteors in the
	least a weighting which reduces the weight of	fits, so the momentum flux errors we have quoted
	these meteors) should be applied in this paper,	are still relevant for the analysis we performed on
	but no mention of it is made. While the numbers	the real data.
	of meteors in these regions are low (in the blue	
	colours), their effect can be disproportionate, so	
	some discussion about how these cases are treated	
	is warranted.	
	2. In the same spirit, Fig. 3 is a bad choice of	
	figure, since it seems to refer to a horizontal	
	meteor trail - which is the one case that should	
	always be avoided, as it produces measurements	
2.	of the vertical wind only, and consequently	Agreed, modified as suggested.
	division by zero when trying to deduce	
	"horizontal winds". I suggest using the figure,	
	but tilt it at some arbitrary (non-zero!) angle	
	from the horizontal.	

3.	3. The paper does not discuss how the Bragg angles are found. The paper by Stober et al. (2018) suggests they always point exactly to the centre-point of the transmitter-receiver line, but this is in error, and only approximately true (as some simple geometric calculations using high-school-level trigonometry will show). I believe the authors of the Stober et al. paper sent a correction to the journal in this regard. Incorrect calculation of the Bragg angle will bias the results.	By "Bragg angle", we assume the reviewer is referring to the direction of the vector that is perpendicular to the meteor trail. In this paper, we have assumed this is in the direction of the bisector of the vectors from the meteor to the transmitter and the meteor to the receiver (which have been referred to as t and b , respectively), and the way this has been estimated has already been discussed in Sect. 3.6. We agree with the reviewer's point that this vector does not necessarily point towards the midpoint between the transmitter and receiver, as suggested in <i>Stober</i> et al. (2018), Fig. 9. As part of this correction, we have made some modifications to how we discuss the use of the Bragg angle/vector in the section on wind (3.7) and covariance (3.9) estimation.
4.	4. The appendices (software) of Stober et al. contain couple of typos - the authors should confirm that they found these.	We found one typo in the appendices of this paper, and have now made note of this in the manuscript.
5.	5. The authors use a spherical-Earth calculation for the bistatic case, as they should. But it is unclear as to whether they did the same in the monostatic case. In some radars, a flat-Earth approximation is used for the monostatic case - the error in height is typically 0.5 km or so at zenith angles of 45 degrees. But it is unclear here whether the flat-Earth approximation was used in the monostatic case, or whether a full spherical Earth approximation was used in both cases. If a flat-earth approximation is used in one case, and a spherical-Earth system used in another, it could lead to biases in comparison. This may also impact the upper-right graph in Fig. 1	We have added comments in several places to clarify that we assumed an ellipsoidal Earth for both bistatic and monostatic cases.
6.	please add labels a,b,c,d - its getting annoying referring to "upper left", "lower right" etc.	Added labels to all figures where appropriate.
7.	Another point of note, though maybe less important, is the mean Bragg wavelength. The authors note that the bistatic procedure gives rise to a diversity of Bragg scales, but it must be work, so a shift to 47 MHz might be more dramatic (relatively) It would be of interest for the authors to look not just at diversity effects but also to consider the mean effective wavelength due to the bistatic arrangement.	This is a good point—we have added a histogram of the effective radar frequency to the paper.

	Page 6, line 10 - the term "said distributions"		
8.	seems clumsy - the word "said" is used in this		
	way several times - it does not add anything to	Removed.	
	the text, and I would recommend removing it.		
	Page 8, line 4 - the authors talk about		
	"well-known linear GW polarization relations".		
	There is no single set of "linear relations" -this is		
	just one of many, and it is the simplest. It all		
	depends on whether you include Coriolis terms,		
	Brunt-Vaisala terms, scale-height terms etc -these		
	cases still deal with linear waves, but the detailed	Thank you for raising this important point. We	
	formulas change So this misleading statement	have removed the "well-known" appellation. In	
	needs to be clarified, perhaps by saying "we use	addition, we only use wave periods up to 4 hours	
9.	the simplest polarization relations". Of course it	in the simulations, which is well below the inertial	
	needs to be recognized that the authors own work	and predominant tidal periods. Therefore, we	
	will be in error at periods close to the tidal and	believe it is unnecessary to qualify the potential	
	inertial frequencies if they use this simple	lack of realism associated with using the simpler	
	polarization relation - Coriolis terms should be	polarization relations.	
	included for a proper treatment, for example.		
	This is pertinent to discussions later in the text,		
	where the authors consider the optimum		
	bandwidth for studies and look at GW periods		
	close to the tides.		
	Page 9, line 9 — I would make the "dot product"		
	between vecef and e clearer - maybe use a bold		
	dot - it is important that readers do not miss this,		
10	and plays into the discussions introduced earlier	A much dama	
10.	regarding overhead meteors. The equation	Agreed, done.	
	emphasizes that when the meteors are close to		
	overhead, the radial velocity is dominated by the		
	vertical wind component		
	Incidentally, the authors used bold font to		
11.	represent both vectors and matrices. While not	We have tried doing this, but the Copernicus	
	wrong, since a vector is indeed a type of matrix,	LaTeX class does not permit arrows over vectors.	
	they may want to consider distinguishing these	All vectors have been changed to bold/italicized	
	cases e.g. by using an arrow over the vectors and	to distinguish them from matrices (bold).	
	keeping matrices bold.		
19	Page 10, line 18 - how "rare" is "rare"?	Thank you for picking up on this—we have	
12.		re-examined and clarified accordingly.	

	Page 10, lines 19 to 21, and page 11, lines 1 - 14.		
	These lines are very disturbing. First, the authors		
	seem to have the wrong Holdsworth et al. (2004)		
	paper in the References. The authors give		
	Holdsworth, D. A., Tsutsumi, M., Reid, I. M.,		
	Nakamura, T., and Tsuda, T.: Interferometric		
	meteor radar phase calibration using meteor		
	echoes, Radio Sci., 39, 2004. Yet line 19 on page	Thank you for picking up on the incorrect	
13.	10 discusses an "error code 3". I cannot see an	reference—this was a mistake on our part. This	
	"error code 3" in this paper. I CAN find one in	has been corrected.	
	the paper "Holdsworth, D. A., I. M. Reid, and M.		
	A. Cervera (2004), Buckland Park all-sky		
	interferometric meteor radar, Radio Sci., 39,		
	RS5009, doi:10.1029/2003RS003014". It appears		
	that the authors have referenced the wrong		
	paper, and I believe this is true for (almost?) all		
	references to Holdsworth et al., (2004).		
	But it gets worse. On page 11, line 3, the authors		
	give Holdsworth (2004) as a representative		
	example of "standard analysis", and then on line		
	11, make the statement " we follow the		
	iterative scheme PROPOSED by Holdsworth et		
	al., (2004).". Sorry , but this is very misleading.		
	This iterative scheme was proposed fully 7 years		
	earlier (Hocking and Thayaparan, Radio Sci., 32,		
	$833\text{-}865,1997,\mathrm{section}$ 4.1) and again discussed in	We enclosing for missing this reference, this has	
14.	Hocking et al., J. Atmos. Solar-Terr. Physics, 63,	we apologize for missing this reference—this has	
	155-169, 2001.). Holds worth may have made	now been added.	
	minor adjustments, but he did not "propose" the		
	idea - it was already proposed 7 years earlier.		
	Indeed much of Table 2 in Holdsworth et al (2004)		
	was also proposed 3 years before Holdsworth		
	$\left(2004\right)$ - in Hocking et al., $\left(2001\right).$ Repeated		
	reference to Holdsworth throughout the text in		
	regard to ideas which were proposed many years		
	earlier by others is unprofessional and misleading.		
		As discussed earlier in this reply, we did not	
	In regard to sections 3.7 and on - I again as k -	consider removing/weighting the meteors in this	
	were cases close to the zenith, where the radial	region given the small number of detections here.	
	velocity is dominated by vertical winds, removed,	Again, we believe that weighting the meteor	
15.	or given lower weighting? Or how were such cases	distribution to minimize the wind/momentum	
	dealt with? Nasty near-infinity terms can occur	flux estimation errors is a separate topic	
	when such terms are include, which could bias the	warranting its own investigation, and have	
	resultant mean winds and momentum fluxes.	mentioned this in the discussion section of the	
		manuscript.	

	Page 12, lines 1 to 9. This is another example of	
	poor citations, perhaps due to the first author's	
	inexperience. Line 1 says "The approach we apply	
	is identical to that presented by Thorsen et al.,	TT 1 1 . 1.1
	(1997) and subsequently Hocking (2005)." It gives	We have adjusted this section to avoid implying
	the impression that the ideas were developed by	that Hocking (2005) simply re-applied the
	Thorsen and simply re-applied by Hocking. This	estimator shown in Thorsen et al. (1997).
	is patently untrue. The current authors use the	However, it should be noted that the two
	matrix A' as discussed in lines 7 to 8. Thorsen et	approaches are fundamentally the same; the
	al. NEVER used this matrix. The matrix	Hocking (2005) paper is just more descriptive in
16.	Thorsen et al. used is given in their equation	that it outlines a way of inverting the A' matrix
	(18), section 3.3, page 714. It is a MUCH	(by directly finding the least squares solution to
	simplified version of the true A' The matrix that	the problem). We also note in this paper that we
	the current authors use comes straight out of	do not use the inversion method described by
	Hocking (2004) and was never used by Thorsen et	Hocking (2005), but rather SVD, due to the fact
	a) It makes a major improvement to the analysis	that SVD is known to provide more reliable
	So please replace "subsequently" in line 1 of page	solutions to poorly-conditioned problems.
	1 with "subsequently improved" and give a	
	specific reference to Hocking (2004) in regard to	
	lines 7 and 8	
	Page 12 line 26 "we have opted to compute the	
	age 12, line 20 - we have opted to compute the	
	"at the origin at the ground" or "at a point in the	
17.	at the origin at the ground of at a point in the	This has now been clarified.
	the ground". Lean see that either might work	
	but it poods to be clarified	
	Page 12 line 1. The statement that "This	
	rage 13, line 1. The statement that This	
	estimate represents what one would measure with	
	an anemometer at some fixed location in the	
	vicinity of the radar, is a little misleading. I	We have removed the statements about the
	but it ion't quite right and peads some sevents	anemometer and about the reference being "in
	added Einst "in the minimity of the reder" is	the vicinity of the radar". We have already briefly
	micloading. Do the authors mean "in the vicinity	discussed the point about the correlation of the
10	of the meteor perior immediately above the	wave field over the entire observation volume in
10.	of the meteor region immediately above the	Sect. 3.10; we found the covariances evaluated at
	radar ? As it reads, it could be at ground-level.	the coordinate system origin vs. at the
	in addition, in the distance case, which radar? the	positions/times of the meteors to be in very close
	than and me denote by the bet in this case	agreement. On the basis of this, we assumed use
	they are moderately close, but since the report is	of the "origin" reference point to be valid.
	supposed to somenow represent general	
	multistatic radars, where the transmitter and	
	receiver can be hundreds of km apart, the	
	question is no longer pedantic.	

		I do not understand the specific problem the	
19.	Furthermore, a suitable anenometer might truly	reviewer is discussing here. In any case, the whole	
	measure wind components, where with the radar	premise of the simulation has been to estimate	
	we have this problem that we deal with radial	the errors associated with propagating realistic	
	components. So while I get what the author is	radial velocity errors to the covariance estimates;	
	trying to say, I suggest it be mentioned with some	the "perfect anemometer" winds, assuming the	
	more caution than has been applied.	radar is attempting to measure these, are used as	
	T T	the reference for estimating the size of the error.	
		Thank you for picking up on the lack of clarity	
		have we have reworded this sentence to make it	
		clean that we are talling chaut the increase in	
		clear that we are taiking about the increase in	
		counts associated with combining the monostatic	
	Page 14, lines 9-10 - OK, but as discussed earlier,	and bistatic receiver data. There is indeed a	
	could some of the increase in count rates be due	slight increase in counts at the higher altitudes	
20.	to a new effective mean frequency? And I also	due to the effective lower operating frequency, but	
	ask again - did the monostatic radar use the same	this is not related to the point being made, and	
	"spherical Earth" geometry as the bistatic case?	has already been mentioned in Sect 2.1. As	
		discussed earlier in this reply, we also used the	
		same ellipsoidal Earth model to compute	
		meteor-related parameters in the monostatic and	
		bistatic cases.	
	Pages 15-21 seem quite good. However, the term		
	"gap" is used in Figs. 10, 11, in line 10, page 21.		
21.	"Gap" gives the impression of a "hole" - I would	Agreed, made suggested change.	
	say the term "temporal shift" might be more		
	appropriate.		
	Page 21, line 1 - the authors say " aside from		
	considering that the GWs may have propagated		
	from a region with weak eastward mesospheric		
	winds." Do they really mean this? Or do they	Thank you for picking up on this—changed to	
22.	mean " GWs may have propagated through a	"through" as suggested.	
	region with weak eastward "? It's the filtering		
	that matters - so either the statement needs some		
	expansion or the authors really mean "through"		
	Page 23 line 2 give a reference to the forcing		
23.	formula	Done.	
24.	Page 23, line 28 - suggest changing "Also like"	Done.	
	to "As for"		
25.	Page 24, lines 11-12 - seems a bit unclear - please	Thank you for picking up on this lack of	
	try to reword what you mean.	clarity—modified.	
	Page 28, conclusions. Some of these statements	None of the modifications affect the conclusion	
26.	may need revisiting following other comments	section, so we have left it unchanged.	
	discussed here-in.	seeden, so no have for it dichailged.	

Multistatic meteor radar observations of gravity wave-tidal interaction over Southern Australia

Andrew J. Spargo¹, Iain M. Reid^{1,2}, and Andrew D. MacKinnon¹

¹Department of Physics, School of Physical Sciences, The University of Adelaide, Adelaide, 5005, Australia ²ATRAD Pty. Ltd., 20 Phillips St., Thebarton, 5031, Australia

Correspondence: Andrew J. Spargo (andrew.spargo@adelaide.edu.au)

Abstract.

5

This paper assesses the ability of a recently-installed 55 MHz multistatic meteor radar to measure gravity wave-driven momentum fluxes around the mesopause, and applies it in a case study of measuring gravity wave forcing on the diurnal tide during a period following the autumnal equinox of 2018. The radar considered is in the vicinity of Adelaide, South Australia (34.9°S, 138.6°E) and consists of a monostatic radar and bistatic receiver separated by approximately 55 km.

The assessment shows that the inclusion of the bistatic receiver reduces the relative uncertainty of the momentum flux estimate from about 75% to 65% (for a flux magnitude of $\sim 20 \text{ m}^2\text{s}^{-2}$, one day's worth of integration, and for a gravity wave field synthesized from a realistic spectral model). This increase in precision appears to be entirely attributable to the increased number of meteor detections associated with the combined monostatic and bistatic receivers, rather than changes in the meteors' anatised distribution.

10 the meteors' spatial distribution.

The case study reveals large modulations in the diurnal tidal amplitudes, with a maximum tidal amplitude of $\sim 50 \text{ ms}^{-1}$ and an associated maximum zonal wind velocity of around 140 ms⁻¹. While the observed gravity wave forcing exhibits a complex relationship with the tidal winds during this period, the components of the forcing are seen to be approximately out of phase with the tidal winds above 88 km. No clear phase relationship has been observed below 88 km.

15 1 Introduction

It has been known for over three decades that the momentum deposition arising from the dissipation of atmospheric gravity waves (herein GW forcing) has a major influence on the background wind and thermal structure of the mesosphere-lower-thermosphere/ionosphere (MLT/I; \sim 80-100 km altitude) (Fritts, 1984). The small scales of the GWs relative to typical grid spacing in global climate models (GCMs) has led to a need to incorporate accurate parameterizations of the GW forcing within

20 the GCMs (Kim et al., 2003; Ern et al., 2011). To support this need, there have been dozens of ground-based, satellite and in-situ studies of the associated GW momentum fluxes in the MLT/I (see e.g., Fritts et al. (2012a) and Nicolls et al. (2012) and references therein). Even so, many of the effects of GWs in the MLT/I are still acknowledged to be poorly understood, which continues to motivate major observational campaigns (e.g., Fritts et al. (2016)).

In recent years, monostatic meteor radars have been the most widely deployed of those ground-based instruments (e.g., Hocking (2005); Antonita et al. (2008); Clemesha and Batista (2008); Beldon and Mitchell (2009, 2010); Clemesha et al. (2009); Fritts et al. (2010a, b, 2012a, b); Vincent et al. (2010); Placke et al. (2011a, b, 2014, 2015); Andrioli et al. (2013a, b, 2015); Liu et al. (2013); de Wit et al. (2014b, a, 2016); Matsumoto et al. (2016); Riggin et al. (2016); Jia et al. (2018)). This is

5 largely due to the low cost and ease of installing and continuously running meteor radars relative to other instruments capable of making the same measurements, such as partial reflection radars (e.g., Vincent and Reid (1983)), coherent radars (e.g., Reid et al. (2018b)), incoherent scatter radars (e.g., Nicolls et al. (2012)), and Doppler lidars (e.g., Agner and Liu (2015)).

Like all other ground-based radar observations of momentum fluxes (see e.g., the discussions in Fritts et al. (2012a), Spargo et al. (2017) and Reid et al. (2018b)), there are concerns around the accuracy and precision of the estimates derived from meteor

- 10 radar. As shown by Vincent et al. (2010), the measurement uncertainties are dependent on both the meteor detection rates and the complexity of the GW spectrum. Their results showed that even at the altitude of the peak of the meteor distribution, integration times of the order of a month or longer may be needed to definitively estimate the sign of the flux, for typical flux magnitudes. Fritts et al. (2012a) and Andrioli et al. (2013a), who also incorporated real time and spatial meteor distributions and a wider variety of GW fields in their simulation, reach similar qualitative conclusions, although Fritts et al. (2012a) in
- 15 particular argue that their measurement uncertainties for a composite day of data comprising measurements spanning one month may be much smaller than those reported in Vincent et al. (2010), due to the use of a larger total number of meteors and an assumption that the wave field in the MLT/I is often dominated by large amplitude monochromatic waves.

Given the demonstrated sensitivities of momentum flux estimation uncertainties, it is important that all users of meteor radars appreciate the uncertainties specific to their radar configuration (the count rates and count distribution, the radar location, and

20 the time of year, and the likely GW field) prior to interpretation of their measurements. This study considers such a simulation of momentum flux measurement uncertainties from a 55 MHz meteor radar in a mid-latitude Southern Hemisphere (SH) site in Australia, and bears those uncertainties in mind in the interpretation of a case study of GW forcing on the diurnal tide. The aspects of this study that are unique can be summarized as follows:

- we consider a multistatic meteor radar configuration consisting of a monostatic radar and a bistatic receiver separated by \sim 55 km;

- we propagate realistic levels of receiver noise and mean phase bias to the angle-of-arrival (AOA) and radial velocity estimates, that are used in the subsequent momentum flux estimation;

- a realistic GW spectral model is used to synthesize the wind field from which the momentum fluxes arise.

25

Section 2 briefly overviews the radar configuration, the count rates obtained and the phase calibration offsets applied. Section
30 3 gives a detailed description of the simulation that estimates the momentum flux measurement uncertainties, and its results. Section 4 presents a case study of momentum fluxes estimated using the radar during the Austral winter, and attempts to validate them by looking at the interaction between the measured fluxes and the tidal winds. Discussion and conclusions follow.

Table 1. Experiment parameters used for the BP meteor radar transmitter, for all data presented in this paper.

Parameter	Value	
Frequency	55 MHz	
Pulse width	7.2 km	
Pulse code	4-bit complementary	
Pulse shape	Gaussian	
PRF	440 Hz	
Range sampling	68.4-309.6 km	
Range sampling interval	1.8 km	
Peak power	40 kW	
Polarisation	Circular	

2 Instrumentation

The multistatic meteor radar considered in this study consists of a Stratosphere-Troposphere (ST)/Meteor radar located at the Buckland Park (BP) field site (34.6° S, 138.5° E) (briefly described by Reid et al. (2018a)), and a remote receiving system located near the township of Mylor, South Australia (35.1° S, 138.8° E) (about 55 km to the South-East of BP).

5

In meteor mode on the BP system, a single crossed folded dipole is used for transmission and a five-element interferometer arranged in a configuration identical to that of Jones et al. (1998) is used for reception. Three element Yagi antennas are used for the interferometer's receive antennas. A peak power of 40 kW is used on transmission. Other experimental parameters used are summarized in Table 1, and a detailed description of the radar hardware is given in Dolman et al. (2018).

The remote receiver system consists of a six receive channel digital transceiver identical to the transceiver system of the 10 BP ST/Meteor radar. In the current configuration, only five of those receive channels are used. The same five receiver antenna arrangement is used at the remote site. To permit accurate range and Doppler estimates at the remote site, the system timing, frequency and clocks at both sites are synchronized with GPS-disciplined oscillators (GPSDOs).

The techniques used to estimate various data products from the received meteor echoes, including radial velocity, meteor position, signal-to-noise ratio (SNR), and decay time, follow those outlined in Holdsworth et al. (2004b)Holdsworth et al. (2004a).

15

20

The dataset considered spans 17th March 2018 to 9th September 2018, with few interruptions (the number of meteors detected per day on both receivers for this interval are shown in Fig. 1).

2.1 Receiver channel phase calibration

Compensating for any systematic receiver channel phase offsets plays an important role in ensuring the accuracy of the position and height estimates of the detected meteors. To calibrate the phases of the receive channels for both of the meteor receiver interferometers used in this study, we have followed the approach suggested in Pholdsworth et al. (2004b).



Figure 1. Upper rowa): the meteor detection rates for the BP and Mylor receivers over the 2018 campaign(left) and, b): the associated distribution of meteors in altitude (right), c): histogram of the effective radar frequency at the Mylor bistatic receiver (i.Lower rowe., $f \cos(\beta/2)$, where f is the operating frequency and β is the forward scatter angle), and d): the horizontal distribution of meteors for the BP (left) and e) Mylor (right) receivers . The (the transmitter location in each case is denoted by a white cross).

The ?-Holdsworth et al. (2004b) approach determines the offsets to apply to the phase differences between the centre and each of the other receive antenna channels that maximise the number of meteors within a range of heights that the meteors are expected to occur (see Chau and Clahsen (2019) for a generalized approach to this). For the BP system, we have used minimum and maximum permissible heights of 70 and 110 km, respectively, and 70 and 120 km for the Mylor system. A slightly larger height interval has been used for the Mylor system to allow for the effect of the distribution of Bragg wavelengths (see Fig. 1, panel c)) on the meteor height distribution width (see, e.g., Thomas et al. (1986) and Stober and Chau (2015), Sect. 3 for a description of this effect).

The phase offsets applied to the Mylor system, and the variability variability of the offsets for the BP system (for which a fixed calibration was used) are shown in Fig. 2. We note a stable calibration for BP, but a few sudden "shifts" in the Mylor case; this has subsequently been determined to be due to a slight rotation of the antenna elements by local wildlife. We do

- not expect isolated shifts like this to have an adverse impact on the analysis performed in this paper, although to somewhat compensate for it, we have performed a daily re-calibration of the receiver channels (using the calibration results for each day) before subsequent processing of the data.
- 10

5



Figure 2. Phase offsets applied to Mylor meteor radar antennas as a function of time (top), and the phase offsets indicated by the ?-Holdsworth et al. (2004b) calibration procedure on BP meteor radar data (bottom).

3 Simulation of wind covariance estimation

3.1 Simulation overview

5

The aim in developing this simulation has been to quantify the uncertainties in the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariance components derived from meteor echoes received in an arbitrary network of meteor radar transmitters and receivers, and to be able to characterize the dependence of those uncertainties on the network shape and the spectrum of the GWs constituting the input wind field. The basic workflow of the simulation (all components of which are elaborated upon in subsequent subsections) may be summarized as follows:

- 1. Produce a sample of meteors in space and time for each site under consideration, by sampling from realistic spatiotemporal meteor detections corresponding to each site.
- 10 2. Specify a wind field based on the superposition of monochromatic gravity waves derived from a realistic GW spectrum, and compute the wind velocities at each of the simulated meteors.
 - 3. Compute the "radial" wind velocity measured at the receiver associated with each meteor detection.
 - 4. For each meteor-site combination, synthesize in-phase and quadrature (IP and Q)-time series for each receiver at the site, based on the "radial" velocity and AOA of the meteor.

- 5. Add a realistically-sized phase bias and noise floor to each receiver channel.
- 6. Estimate the "radial" velocity and AOA of the meteor from the simulated time series.
- 7. Estimate the wave field covariances using the meteors retrieved from different combinations of sites.
- 8. Return to 1., and repeat for the number of realisations required to produce covariance error distributions (in the next step)

```
5
```

- of the desired statistical significance and resolution.
- 9. Compare the estimated covariances with those computed directly from the 3D wind velocities at the meteors and those calculated at 2-minute resolution at the origin of the coordinate system.

3.2 Meteor position specification

To incorporate the dependence of the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ uncertainties on the temporal and spatial characteristics of the meteor

10 distribution, we have based said the distributions used in the model on real measurements. For both the BP and Mylor sites, we constructed a composite day of 2D histograms of the meteor position distributions at 5 km spatial and hourly time resolution, using measurements from April-July 2018. These 2D histograms were taken to represent probability distributions for the meteor positions.

The sampling from these probability distributions at the beginning of each realisation was done according the following 15 process:

- 1. Prescribe a number of meteor detections for the day of measurements and altitude in question (e.g., 1340 per day at 90 km for the BP radar case).
- 2. Use rejection sampling to distribute those meteors across the day, according to the relative number of meteors in each hour in the input probability distribution.
- Distribute the number of meteors prescribed in each hour of measurements according to the spatial probability distribution for that hour, again using rejection sampling.
 - 4. Return to 1., and repeat for the number of days prescribed in this realisation (for results presented in this paper, 1 or 10).

The horizontal position coordinates assigned to each meteor in the probability distribution (and subsequently the model) are based on the distances from the receiver site in Transverse Mercator coordinates, calculated using the method of Bowring

25 (1989). The altitudes assigned to the meteors are derived from a uniform probability distribution, with a centre value of 90 km and a full-width of 2 km (such that the simulation emulates the idea of analysing meteors from a single height bin).

3.3 Meteor detection rate specification

To clarify the effect of a variable number of meteor radial velocity/AOA pairs on the covariance error distribution, a variety of meteor detection rates have been simulated. We have endeavoured to make the detection rates used resemble the number

Table 2. Meteor detection rates used for the simulations in this paper. The rates shown are per day, in 2 km-wide bins centred at the altitude specified.

Altitude (km)	BP	Mylor
76	140	20
80	510	130
82	780	180
84	1080	380
86	1360	540
88	1480	640
90	1340	690
92	1010	640
96	300	350

of meteors detected across a range of heights by the combined BP-Mylor radar link (we note again though that the simulation itself is performed around a single altitude). The detection rates we have used for different heights, listed in Table 2, correspond to those averaged over April 2018 for the two receive sites, in 2 km-wide bins.

3.4 Wind field specification

15

5 The wind field in the simulation is comprised of tidal components and a superposition of monochromatic GWs whose amplitudes have a vertical wavenumber and frequency dependence. Diurnal and semidiurnal tidal components are assumed, with amplitudes of 25 and 10 ms⁻¹ respectively. Random phases from a uniform distribution spanning the interval [0, 2π) are added to the phase of the zonal component of the tides at the beginning of each realisation, and the meridional component is set to be in quadrature with the zonal component. The 3D wind velocity associated with the GWs at a given time t and Cartesian position vector **r**-r can be written as:

$$\underline{\mathbf{v}}\boldsymbol{v} = \sum_{i=1}^{n_m} \sum_{j=1}^{n_\omega} A(m_i, \omega_j) \underline{\mathbf{v}'} \boldsymbol{v'}_{ij} \sin\left(\boldsymbol{\kappa}_i \underline{\mathbf{r}} \cdot \boldsymbol{r} - \omega_j t + \phi_{ij}\right), \tag{1}$$

where *m* is the vertical wavenumber, ω is the wave's angular frequency, n_m and n_ω are the number of vertical wavenumbers and angular frequencies respectively in the spectral grid, *A* is the joint vertical wavenumber-angular frequency spectral amplitude, $\mathbf{v}' = [u', v', w'] \cdot \mathbf{v}' = [u', v', w']$ is the vector of wind component fluctuation sizes, $\boldsymbol{\kappa} = [k, l, m]$ is the 3D wave vector, and ϕ represents a (random for each unique $[m_i, \omega_j]$ pair) phase offset.

As per Sect. 3.2, the coordinate system used to specify horizontal position with respect to a reference location (i.e., that embodied by the r_{τ} vector) is based on the Transverse Mercator distances evaluated using the Bowring (1989) method (which follow the Earth's surface and take into account its ellipsoidal shape). This is used in preference to line-of-sight distances, the use of which would result in "stretching" of the horizontal scales of the waves at large distances from the coordinate system

origin. Furthermore, the calculated wind velocities are assumed to be in the local East-North-Up (ENU) coordinates at the associated meteor positions.

To ensure that the correlations between the horizontal and vertical winds take on physically reasonable values, we have allowed the component fluctuation amplitudes to be related by the well-known-linear GW polarization relations, i.e. relation 5 $w' = \frac{v_h k_h}{m}$, where $v_h = \sqrt{u'^2 + v'^2}$ and $k_h = \sqrt{k^2 + l^2}$. The horizontal components are determined by the wave propagation azimuth φ , through the relations $[k, l] = k_h [\sin \varphi, \cos \varphi]$ and $[u', v'] = v_h [\sin \varphi, \cos \varphi]$.

In order to give the wind field a level of "spatially-correlated randomness" akin to what is seen in mesospheric wind fields when no predominant wave scales are present, we have opted to let $A(m, \omega)$ take on values from a gravity wave spectral model. The vertical wavenumber spectrum we have used (Gardner et al. (1993), eqn. (7), and following their nomenclature) is given by:

10

$$F_u(m) = 2\pi\alpha N^2 \begin{cases} m_*^{-3} (\frac{m}{m_*})^s & m \le m_* \\ m^{-3} & m_* \le m \le m_b \\ m_b^{-3} (\frac{m_b}{m})^{5/3} & m_b \le m \end{cases}$$
(2)

where *m* is the vertical wavenumber of the wave, and following Gardner et al. (1993), Fig. 1, we let $\alpha = 0.62$, $N = \frac{2\pi}{3 \times 10^2} \text{ s}^{-1}$, $m_* = \frac{2\pi}{1.5 \times 10^4} \text{ m}^{-1}$, $m_b = \frac{2\pi}{5 \times 10^2} \text{ m}^{-1}$, and s = 2. The frequency spectrum we have used (Gardner et al. (1993), eqn. (24)) is given by:

15
$$B(\omega) = \frac{p-1}{f} \left(\frac{f}{\omega}\right)^p$$
, (3)

where ω is the angular frequency of the wave, and following Gardner et al. (1993), Fig. 2, we let $f = \frac{2\pi}{7.2 \times 10^4} \text{ s}^{-1}$ and p = 2. We then simply assume that the joint vertical wavenumber-angular frequency spectrum is given by the product of these two spectra, i.e.,

$$A(m,\omega) = F_u(m)B(\omega).$$
⁽⁴⁾

20 The 2D spectrum we used for results presented in this paper consisted of 80 different vertical wavelengths and wave periods, spanning the ranges 0.5–20 km and 5–240 minutes (uniformly sampled in vertical wavenumber and frequency), respectively. These limits largely encompass the waves responsible for the majority of the momentum deposition in the MLT-region (see e.g., Fritts and Alexander (2003)), whose momentum fluxes are of principal interest in this study.

The wave propagation azimuths were sampled from a uniform random distribution spanning $[0, 180^\circ]$ in bearing, with the intention being to emulate a wave field whose westward-propagating waves have been removed from the spectrum through selective filtering. This led to true values for the estimates of $\langle u'w' \rangle$ that were on average positive, and values of $\langle v'w' \rangle$ that were on average zero. Testing a wider variety of wave field configurations was considered beyond the scope of the paper.



Figure 3. Bistatic meteor reception geometry. Following the nomenclature Using similar terminology to that in Protat and Zawadzki (1999), t t is a unit-vector in the direction of from the meteor to the transmitter, **b** b is a unit vector in the direction of from the meteor to the bistatic receiver, and $e \cdot e$ is a unit vector that is perpendicular to the meteor trail axis (and therefore, assuming specular reflection from the trail, is a bisector of t t and b). β is the so-called "forward scatter" angle.

The absolute values taken by $A(m, \omega)$ were normalized in a way that resulted in mean values of $\langle u'w' \rangle$ in the vicinity of 20 m²s⁻², which is a typical value for this parameter in the MLT-region (see e.g., the discussion in Fritts et al. (2012a)). An example distribution of "true" covariances evaluated in the simulation is shown in the <u>lower panels panel b</u>) of Fig. 5.

3.5 Projection of the wind velocity onto the Bragg vector

5 A diagram summarising the bistatic reception geometry is shown in Fig. 3. Following the development of Protat and Zawadzki (1999), the so-called "radial" velocity measured by a bistatic receiver corresponds to the projection of the 3D wind velocity onto **e** (also referred to *e* (which is in the same direction as the Bragg vector in e.g., Stober and Chau (2015)), in turn projected onto **b***b*. Mathematically, this velocity is expressed as:

$$v_{\rm rrm} = \cos(\beta/2) \cdot \underline{v} v_{\rm ecef} \cdot \underline{e} \,, \tag{5}$$

- 10 which is the velocity that is used to produce a phase progression in the simulated receiver time series, discussed in Sect. 3.6. It should be noted that \mathbf{t}, \mathbf{b} and $\mathbf{e}, \mathbf{t}, \mathbf{b}$ and \mathbf{e} are expressed in Earth-centred, Earth-fixed (ECEF) coordinates, and that the wind velocities \mathbf{v} or computed in the simulation are in the local ENU coordinates of each meteor. The "ecef" subscript on \mathbf{v} is to denote $\mathbf{v}v$'s rotation to the ECEF coordinate system; we have followed a slightly modified version of the approach discussed in detail by Stober et al. (2018) to do this (see App. A). We note that we apply the same procedure for both bistatic and monostatic
- 15 receivers (though of course, in the monostatic case t = b).



Figure 4. Probability distributions of SNR and decay time used in producing the receiver time series discussed in Sect. 3.6.

3.6 Receiver time series generation and parameter re-estimation

To ensure that realistic radial velocity and position estimation errors are propagated to the covariance estimation, we have opted to generate synthetic receiver time series based on the observables discussed in the previous sections, and to then attempt to re-estimate the observables from the time series. The complex time series for the jth receiver is written as:

5
$$V_i(t) = e^{i(2\pi\mathbf{A}\cdot\mathbf{d} - 4\pi v_{\rm r}t/\lambda + \Phi_j)i(2\pi\mathbf{A}\cdot\mathbf{d} - 4\pi v_{\rm rm}t/\lambda + \Phi_j)} e^{-t/\tau} + n_i(t), \qquad (6)$$

where $\mathbf{A} = [\sin\theta\sin\phi, \sin\theta\cos\phi, \cos\theta]$ (where θ and ϕ are the zenith and azimuth angles of the meteor, respectively, as measured from the receiver), $\mathbf{d} \cdot \mathbf{d}$ is a three-element vector of Cartesian displacements to the receiver antenna in question, λ is the radar wavelength, Φ_j is a phase calibration offset for the *j*th receiver, τ is the e^{-1} decay time of the meteor, and $n_j(t)$ is a background noise function.

The background noise function consists of values derived from a Gaussian distribution, with a root-mean-square (RMS) value derived from a probability distribution of meteor echo SNRs from the monostatic 55 MHz meteor radar at BP. The values used for τ are also derived from a probability distribution from this radar's data. In both cases, the data used to generate the probability distributions spanned 1-30 April 2018, and altitudes 70-110 km. Plots of these distributions are shown in Fig. 4.

The phase calibration offsets Φ_j , which are set for each receiver at the beginning of each simulation realisation, are intended to embody the consequences of incorrectly estimating the true phase calibration offsets between the receiver channels. Based on the phase calibration offset time series shown in Fig. 2, we have chosen to apply to each receiver Gaussian-distributed phase offsets with an RMS value of 2° .

Radial velocities and meteor positions are estimated from the noise and phase-offset time series following the procedures outlined in Holdsworth et al. (2004b) Holdsworth et al. (2004a) (Sects. 3.11 and 3.12, respectively), with the exception that the

20 radial velocity is corrected for the forward scatter angle in the case of bistatic reception. Using the definitions in Fig. 3 and

following the approaches outlined in Stober et al. (2018) to compute the $\frac{t}{t}$ and $\frac{b}{and e} = \frac{t+b}{|t+b|} t$ and $\frac{b}{and e} = \frac{t+b}{|t+b|}$) vectors, the forward scatter angle may be estimated using:

$$\beta = \cos^{-1} \left(\frac{-\mathbf{t} \cdot \mathbf{b}}{|\mathbf{t}||\mathbf{b}|} \frac{-\mathbf{t} \cdot \mathbf{b}}{|\mathbf{t}||\mathbf{b}|} \right), \tag{7}$$

and then eqn. (5) may be rearranged for $\frac{\mathbf{v}_{ecef} \cdot \mathbf{e}}{\mathbf{v}_{ecef} \cdot \mathbf{e}}$, to get said the radial velocity.

5 It should be noted that in rare cases, it becomes (~ 1 detection in every 3000) cases, we found it became impossible to estimate the AOA of the meteor unambiguously when the phase biases and noise are-were incorporated into the receiver time series (i.e., the error code 3 discussed in Holdsworth et al. (2004b) is Holdsworth et al. (2004a) was encountered). In these particular cases, the echo in guestion is was simply discarded from the subsequent calculation of mean winds and covariances.

3.7 Mean horizontal wind and tidal component estimation

The way we have estimated mean horizontal winds in this simulation is similar to that typically applied to meteor radars in the 10 literature (e.g., Holdsworth et al. (2004b)Hocking and Thayaparan (1997); Holdsworth et al. (2004a)). Our approach has been to use singular value decomposition (SVD) to solve the following inverse equation in the least squares sense for $\mathbf{v}v$:

$$\underline{\mathbf{v}_{\mathbf{r}}} \boldsymbol{v}_{\boldsymbol{r}} = \underline{\mathbf{A}} \underline{\mathbf{v}}_{\underline{\mathbf{A}}} \boldsymbol{v}, \tag{8}$$

where $\frac{\mathbf{v}_{\mathbf{r}} \text{ is a } n_{met} \times 1}{\mathbf{v}_{\mathbf{r}} \text{ is a } n_{met} \times 1}$ vector of radial velocities ($n_{met} n_{met}$ being the number of meteors in the time bin under consideration), $\mathbf{v} \cdot \mathbf{v}$ is a 2 × 1 vector of wind velocities, and A is a $n_{met} \times 2 \cdot n_{met} \times 2$ matrix whose rows take the 15 same form as that described in eqn. (6) (without the vertical component). However, it is important to note in this case that the θ and ϕ defined in A need to be modified to account for the small rotation of the ENU coordinate systems between the receiver and meteor positions (and be valid for the meteor's position). The rotation that is to be applied to θ and ϕ is outlined by Stober et al. (2018), App.A4represent the orientation of the e (or "Bragg") vector (see Fig. 3) in the ENU coordinate system

20 at the location of the meteor. The velocities in v_r of course also represent the projection of the wind vector on the e vector; i.e., $v_{\rm rm}/\cos{(\beta/2)}$, where $v_{\rm rm}$ is the radial velocity measured at the receiver.

In order to remove outliers from the input radial velocity distribution, we follow the iterative scheme proposed by Holdsworth et al. (2004 subsequently used by e.g., Holdsworth et al. (2004a)). This involves performing an initial fit for the wind velocities, removing the radial velocities whose value differs from the horizontally-projected radial wind by more than 25 ms⁻¹, and repeating the procedure until no outliers are found or until less than 6 meteors remain.

25

Removal of background wind and tides 3.8

To remove the previously estimated mean winds and tides from the time series, we have calculated a low-pass filtered version of the hourly-averaged horizontal wind time series using an inverse wavelet transform with a Morlet wavelet basis, and have subtracted the projection of this on linearly interpolated a wind estimate at the time of each meteor, and subtracted the radial projection of the wind from the radial velocity time series. This is in principle similar to the approach of Fritts et al. (2010a), who applied an S-transform (in preference to a least squares sinusoidal fit) in order to more completely remove transient spectral features around the tidal periods from the time series. The application of the inverse wavelet transform is described in

5 App. B.

To ensure that the filtered time series pertain to tidal (or longer)-like wind oscillations (and not short-period GWs), we select a minimum scale size in the reconstruction of 6 hours and a total number of scales of 250. The reconstructed time series is then interpolated to the times of each of the meteors in question, and the radial component of this wind at each of the meteor positions is subtracted from the measured radial velocity.

10 3.9 Covariance estimation

Following the removal of the mean and tidal components of the horizontal wind from the radial velocities, covariances that pertain predominantly to gravity wave-driven wind perturbations are estimated. The approach we apply is identical to that based on those presented by Thorsen et al. (1997) and subsequently-Hocking (2005); much like in the wind estimation, it involves using SVD to least-squares solve the following inverse equation:

15
$$\underline{\mathbf{v}} v_{\mathrm{r}}^{\prime 2} = \underline{\mathbf{A}' \mathbf{v}'} \underline{\mathbf{A}}' v',$$
 (9)

where $\frac{v_r'^2}{r}$ is a $n_{met} \times 1$ $v_r'^2$ is a $n_{met} \times 1$ vector containing the squares of the perturbation component of the radial velocities,

$$\underline{\mathbf{v}'} \mathbf{v'} = \left[\langle u'^2 \rangle, \langle v'^2 \rangle, \langle w'^2 \rangle, \langle u'v' \rangle, \langle u'w' \rangle, \langle v'w' \rangle \right]^{\mathrm{T}}$$

is the vector of covariance components, and A' is a $n_{met} \times 6 n_{met} \times 6$ matrix whose rows read:

$$\begin{split} & [\sin^2\theta\sin^2\phi, \sin^2\theta\cos^2\phi, \cos^2\theta, \sin^2\theta\sin 2\phi, \\ & \sin 2\theta\sin\phi, \sin 2\theta\cos\phi]. \end{split}$$

It is noted that, as per the wind estimation case, the θ and ϕ terms must be defined according to represent the orientation of the ENU coordinate system at the meteor in question, not the receive site vector in ENU coordinates at the location of each meteor, and that the velocities in $v_{L}^{\prime 2}$ are based on the wind velocities' projection onto *e*.

A two-step radial velocity outlier rejection procedure is utilized to remove <u>meteors with</u> dubious square radial velocity/AOA pairs from the input distribution in an attempt to reduce the bias in the resulting covariance estimates. The first step is to

25 discard all radial velocity/AOA pairs that have a projected horizontal velocity of $\geq 200 \text{ ms}^{-1}$ (by virtue of which we argue that measured horizontal velocities above this threshold are "nonphysical"). The second step iteratively discards the pairs that satisfy the criterion:

$$|v_{ri}'^{2} - v_{rpi}'^{2}| \ge \left[\text{median} \left(\sqrt{|v_{r}'^{2} - v_{rp}'^{2}|} \right) + 5 \times 1.4826 \times \text{MAD} \left(\sqrt{|v_{r}'^{2} - v_{rp}'^{2}|} \right) \right]^{2}$$
(10)

where $v_{rpi}^{\prime 2} = A_i^{\prime} i * \mathbf{v}^{\prime} v_{rpi}^{\prime 2} = A_{i*}^{\prime} v_{i}^{\prime}$ is the *i*th "projected" square radial velocity, "MAD" indicates the median absolute deviation operator, and 1.4826 is the factor to convert a MAD to a standard deviation, assuming the input has a Gaussian distribution. 5 In practice, we have found that the 5 "standard deviations" criterion removes outliers that are large enough to substantially bias the resulting covariance estimates, without iteratively removing an excessive number of samples that are "good". The intention of using the median and MAD statistics (as opposed to mean and standard deviation) has been to reduce the bias outlying points inflict on the "measured standard deviation" of the distribution of $|v_{ri}^{\prime 2} - v_{rpi}^{\prime 2}|$.

The performance of the second outlier rejection criterion on simulated data is briefly summarised in Sect. 3.11.3.

10 3.10 "Truth value" of the simulated covariances

To evaluate the "truth value" of the simulated covariances—i.e., that used to estimate the accuracy and precision of the covariances derived through inversion of eqn. 9(9)—we have opted to compute the covariances at the origin of the coordinate system (in the meteor region directly above the receiver at BP) at 2 min time resolution. This estimate represents what one would measure with an "anemometer" at some fixed location in the vicinity of the radar. We found this estimate to agree extremely

15 closely with that computed at the positions and times of the meteors incorporated in the simulation, which in turn represents the most accurate and precise estimate one could hope to obtain when inverting eqn. (9).

In the case of using wave fields generated from the previously discussed gravity wave spectral model, we found that the covariances estimated by inverting eqn. (9) are more correlated with those calculated using the above two methods than those computed by summing the covariances associated with each wave in the spectrum. Therefore, while the latter method gives

20 the covariances that would be measured over an infinitely large sampling area/time (in a sense the "expectation value" of the covariances), we have refrained from using it as a "truth" value with a view to not overestimating the size of the simulated technique's measurement errors.

3.11 Simulation results

3.11.1 Spectrum of gravity waves

25 This section considers the covariance bias distributions associated with a wind field generated using the GW spectral model discussed in Sect. 3.4. Three different time integration cases (that are later employed in this paper on real data) are tested: 1 day (which could be considered fairly "high time resolution" sampling of day-to-day variations), 10 days (which sacrifices time resolution for measurement precision), and a 20-day composite (which intends to gather enough meteors in each time-of-day bin for a precise covariance estimate, but in doing so ignores day-to-day variations entirely).



Figure 5. Simulated wind covariance bias distributions for 1 day of integration (upper row a)) and the simulated covariance distributions (lower row b)). As discussed in Sect. 3.10, biases are calculated with respect to a "reference" value computed at 2 min resolution at the coordinate system origin. The lower row shows the distribution for the "reference" covariance in a dotted black line, and the "true" covariances in coloured lines. The different line colours in each plot represent different simulated heights, which are a subset of those shown Table 2 (red represents 76 km, yellow 80 km, green 84 km, black 88 km, blue 92 km, and violet 96 km). Thick lines show the distribution for the multistatic case (i.e., by combining data from BP and Mylor), and thinner lines show the monostatic case (i.e., just BP data). The mean and standard deviation evaluated from the samples' MAD are shown in the left and right columns respectively of the arrays of numbers in each plot figure.

1-day integration

and spatial distribution of the meteors.

The biases for 15,000 realisations of 1-day integrated covariance estimations are shown in Fig. 5. It is clear that the $\langle u'w' \rangle$ term is systematically underestimated, with larger biases present at lower count rates. The width of the bias distribution is also larger at lower count rates. For a simulated mean $\langle u'w' \rangle$ value of $\sim 21 \text{ m}^2 \text{s}^{-2}$, the distribution widths imply a 1-sigma measurement uncertainty of $\sim 65\%$ at the peak of the height distribution, and $\sim 145\%$ at the edges of the distribution, for a multistatic configuration. The same uncertainties are $\sim 72\%$ and 168%, respectively, for a monostatic configuration.

The width of the bias distributions for $\langle v'w' \rangle$ are also essentially identical to those for $\langle u'w' \rangle$. The relative uncertainties in the measurements of this term are meaningless, as the wave propagation directions have been chosen in a way that the mean truth value of $\langle v'w' \rangle$ is zero. What the results do illustrate, however, is that there is no bias in the case of estimating a covariance

with a zero mean, and that there is no change in the measurement uncertainty of the two components arising from the temporal

5

It should be noted that $\langle u'w' \rangle$ is systematically underestimated for both configurations and for all count rate sets investigated, especially at lower count rates (the absolute error ranges from about 20% to 50%). Subsequent investigation has confirmed that



Figure 6. As per Fig. 5, but for 10 days of integration.



Figure 7. As per Fig. 5, but for single monochromatic GWs.

5

10

this occurs when an attempt is made to remove the tidal effects incorporated in the simulated wind field (i.e., the tides are largely removed, but so is some of the variance due to the GWs). The larger biases at low count rates arise from the inability to define the tidal amplitudes and phases correctly in the presence of wind estimates with larger uncertainties and/or missing wind estimates for particular time bins. Overall, we consider the bias an unavoidable consequence of ensuring that tidal effects are not included in the measured covariances. Further discussion of this point is taken up in Sect. 5.2.

It also appears that there is no clear dependence of covariance uncertainty on the use of a monostatic or multistatic configuration, for a fixed detection rate. This is evidenced by the uncertainties at 84 km for the multistatic configuration (1460 detections) being 14.4 m²s⁻² and 14.5 m²s⁻² for $\langle u'w' \rangle$ and $\langle v'w' \rangle$ respectively, and the corresponding uncertainties at 88 km for the monostatic configuration (1480 detections) being 15.2 m²s⁻² and 14.6 m²s⁻². In other words, since these uncertainties are essentially the same, we surmise that the multistatic configuration combining the detections from the monostatic



Figure 8. Means and standard deviations of the simulated $\langle u'w' \rangle$ (top row **a**)) and standard deviations $\langle v'w' \rangle$ (lower row **b**)) of the simulated covariance bias distributions for a 20-day composite, as a function of height, for the BP-Mylor link.



Figure 9. Covariance bias distributions for different combinations of outlier contamination and outlier rejection. Black is no rejection or outliers, red is rejection with no outliers, blue is outliers without rejection, and green is outliers with rejection.

and bistatic receivers only offers a lower measurement uncertainty at a given height because of the higher number of meteor detections, not because of the altered Bragg vector distribution associated with having two receiver sites.

10-day integration

Figure 6 shows the bias distribution for 1,500 realisations of 10-day integrated covariance estimates. It is clear that the relative uncertainties in both $\langle u'w' \rangle$ and $\langle v'w' \rangle$ are considerably smaller than for 1 day's integration, ranging from ~50% at the peak of the distribution, to ~60% at the edges. Interestingly, it appears as though the uncertainty is asymptoting to a minimum value, implying that the use of integration times longer than 10 days will lead to diminishing gains in measurement precision. For this reason, we have not opted to use integration times longer than this in the analysis of the BP-Mylor data in this paper.

As per the 1-day integration case, $\langle u'w' \rangle$ has been systematically underestimated, increasingly so at low meteor detection rates. There is also no clear advantage or disadvantage associated with using the bistatic receiver, meteor detection rates aside.

5 20-day composite

Figure 8 shows expected values of the covariance bias' mean and standard deviation for 300 realisations of a composite day spanning an interval of 20 days, with three hour time bins, as a function of height from 82-92 km. The highest standard deviations for both $\langle u'w' \rangle$ and $\langle v'w' \rangle$ occur in the 6-9 and 9-12 UT bins, and the lowest in the 18-21 UT bin. The mean value for $\langle u'w' \rangle$, which is again $\sim 21 \text{ m}^2\text{s}^{-2}$, implies a relative uncertainty at the peak of the height distribution of about 70% in the 18-21 UT bin, and about 85% in the 6-9 UT bin. It should be noted that the uncertainty is as high as $\sim 100\%$ in the 6-9 UT bin.

at 82 km.

10

25

Once again, a systematic underestimation of $\langle u'w' \rangle$ is present, which as discussed in is Sect. 3.11.1 is an artefact of attempting to remove tidal effects.

3.11.2 Monochromatic gravity wave

15 The previous section considered a wind field containing a multitude of waves whose spatial/temporal scales spanned a large part of the spectrum atmospheric gravity waves are expected to occupy. This section briefly addresses the other limiting case, which is that of a wind field consisting of a single monochromatic wave.

In all simulation realisations for this case, we have set the single monochromatic wave's propagation direction to 45°T, so as to make the true $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariances equal. A horizontal wavelength and phase speed has been randomly selected

for each realisation, from a uniform distribution with bounds [10, 60] km and [10, 40] ms⁻¹, respectively. A 1-day integration is used for the covariance estimate.

The bias distributions for 15,000 realisations are shown in Fig. 7. As per the spectral wave field case, the distribution widths are largest at the edges of the height distribution, and narrowest at the peak. However, the widths are far smaller than in the spectral wave field case. Across all wavelengths and phase speeds, the simulated mean true covariance was $\sim 38 \text{ m}^2 \text{s}^{-2}$, which translates to uncertainties of about 8% and 44% at the peak and lower edge of the height distribution respectively for the multistatic configuration. For the monostatic configuration, the same uncertainties are about 10% and 52%, respectively.

Similarly to the spectral wave field case, both covariance terms are systematically underestimated (ranging from about 2% to 26% for $\langle u'w' \rangle$ in the multistatic configuration, at the peak and lower edge of the height distribution, respectively). Interestingly, $\langle v'w' \rangle$ is underestimated to a slightly lesser degree than $\langle u'w' \rangle$. Once again, there is also no clear advantage or V advantage of the height distribution of the height distribution of the height distribution.

30 disadvantage of using the bistatic receiver (meteor detection rates aside).

3.11.3 Outlier rejection criteria performance

This section shows the effect of the application of the outlier rejection criterion of eqn. (10), in the absence of tidal effects and attempted removal of them.

To emulate a radial velocity time series "partially corrupted" with outliers in this section, Gaussian-distributed noise with a 5 standard deviation of 50 ms⁻¹ has been added to a randomly selected 5% of the radial velocity estimates in a given realisation. We note that radial velocity errors of this size are rare in practice; they have been used to test the rejection criterion's robustness, and to allow us to highlight potential downsides of not having the criterion in place.

Figure 9 shows the covariance bias distributions for the same spectral gravity field as applied in Sect. 3.11.1 and for 1 day of integration, for four cases: rejection not applied with no outliers present, rejection applied with no outliers present, rejection

10 not applied with outliers present, and rejection applied with outliers present. The mean true values for $\langle u'w' \rangle$ and $\langle v'w' \rangle$ are the same as in Sect. 3.11.1, i.e., ~ 21 and 0 m²s⁻², respectively.

The application of the criterion is clearly beneficial in the presence of outliers, resulting in a reduction in relative uncertainty of the $\langle u'w' \rangle$ estimate from about 214% to 74%. Interestingly, the application of the criterion in the presence of no outliers also results in a slight reduction in relative uncertainty (from about 86% to 73%), although it does result in $\langle u'w' \rangle$ being underestimated (by about 20%). This point is revisited in Sect. 5.3.

Despite the fact that it appears to introduce a small measurement bias, we still apply the criterion in the subsequent analysis of BP-Mylor data, so that we can be assured that anomalous radial velocities do not contribute to the covariance measurement errors.

4 Momentum flux retrievals

15

20 This section uses the methodology described in the previous section to estimate covariances from the BP-Mylor meteor radar link from 17th March 2018 through to 9th September 2018. The aim of this analysis was originally to verify that the estimated covariances and flow acceleration derived from them were physically reasonable; however, in observing an apparent tidal modulation of the covariances, we realised that the results themselves may be of more general interest.

4.1 Covariances during the Austral winter

- Plots of the mean horizontal winds and the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariance terms from 17th March through to 9th September 2018 are shown in Fig. 10. Both quantities have been sampled using 2 km, non-oversampled altitude bins. We chose to evaluate the covariance terms using 10-day-long windows, with a <u>gap time shift</u> of 2 days between the centres of adjacent windows, in attempt to resolve the planetary-wave induced modulation of the covariances. A low-pass wavelet filter with a cut-off of 2 days and a 10-day moving average has been applied to the hourly horizontal winds to evaluate the winds shown; the filtering was
- 30 performed to avoid the aliasing of GW activity and tides into the wind's variability, and the moving average in order to more



Figure 10. Mean horizontal winds (top-row a)) and the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariance components (row b)) measured using the BP-Mylor link between 17th March and 9th September 2018. As discussed in Sect. 4.1, the winds shown correspond to a 10-day moving average of the hourly-averaged winds with tidal components removed, and the covariances have been evaluated over 10-day windows, with a gap time shift of 2 days between the centres of adjacent windows.

closely match the temporal sampling of the two parameters. Therefore, the winds shown should provide a good measure of the "background mean winds" responsible for selective filtering of the gravity wave spectrum.

As is expected for this time of year at a mid-latitude SH site (see e.g., Vincent and Ball (1981)), the eastward winds around 80 km generally increase with time from the autumnal equinox to the winter solstice (\sim days 80 and 170 respectively) and decrease toward the vernal equinox (\sim day 265). A wavelet analysis (not shown here) reveals that much of the shorter term zonal wind variability evident in the figure is transient, and encompasses a spectrum of periods between about 10 and 60 days. The meridional wind, conversely, has a mean much closer to zero. Much of its variability is confined to periods around 10, 20, 25 and 40-50 days below 90 km, with variability in the 50-100 day period becoming increasingly dominant above 90 km.

5

The level of (anti)correlation between the covariance terms and the winds is highly variable. The $\langle u'w' \rangle$ term appears to 10 be anticorrelated with the zonal wind between 80 and 84 km around the winter solstice, as does $\langle v'w' \rangle$ with the meridional wind above 88 km across a similar time interval. While pronounced levels of anticorrelation between these quantities in the mesospheric region arising from the selective filtering mechanism are typical (see e.g., the recent summary provided by Jia et al. (2018))—particularly in the zonal component—departures from these predictions are also not uncommon. As Jia et al. (2018) explains, it is difficult to conceive a mechanism for departures from this theory in the zonal component (given the dominance



Figure 11. As per Fig. 10, but for April 2018. Also, in this case no moving average has been applied on the winds post-tide-removal, and the covariances have been evaluated over windows of length 1 day, with a **gap-time shift** of 6 hours between the centres of adjacent windows.

of eastward winds in the lower mesosphere during winter), aside from considering that the GWs may have propagated from through a region with weak eastward mesospheric winds.

The feature we focus the remainder of this discussion on concerns the coincident enhancement in the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ terms in the interval spanning days 100 to 120, around 90-94 km. Peak values of $\sim 50 \text{ m}^2\text{s}^{-2}$ and 100 m²s⁻² for $\langle u'w' \rangle$ and

5 $\langle v'w' \rangle$ respectively are obtained during this interval. Interestingly, they coincide with a brief enhancement in the zonal winds at the same height, and the peak of the northward phase of an oscillation in the meridional winds with periods spanning 50-100 days.

Figure 11 shows an inset of Fig. 10, spanning April 2018 (which the aforementioned covariance enhancement is centred on). In an attempt to increase the temporal resolution, the covariances in this figure have been evaluated with 1-day windows, with

10 a gap time shift of 6 hours between adjacent windows. Tidal components have also been removed from the winds as per Fig. 10 (i.e. in order to not alias tidal/GW activity into the winds), and for a closer match to the time sampling of the covariances, no moving average has been applied.

This figure shows evidence of a pronounced periodicity around 10 days in the zonal wind, which attains its highest amplitude at approximately day 110 around 85 km. At this time and in the same altitude region, the mean meridional winds abruptly (over

15 a period of a few days) switch from northward to southward. All of this variability is likely attributable to a superposition of planetary waves. Albeit noisy (owing to the relatively short integration time), the $\langle u'w' \rangle$ covariance term shows an enhancement between days 105 and 110, and attains especially high positive values (exceeding 100 m²s⁻²) at around 90 km altitude.



Figure 12. Amplitude of the diurnal (top-row a)) and semidiurnal (second-row b)) tides, and and phase of the diurnal (third-row c)) and semidiurnal (fourth-row d)) tides as measured by the BP-Mylor meteor radar during April 2018.

Interestingly, the $\langle v'w' \rangle$ enhancement lags that of $\langle u'w' \rangle$ by several days, with a peak again in excess of 100 m²s⁻² around day 110.

We have also noted that this interval is associated with an abrupt enhancement of the amplitudes of the diurnal and semidiurnal tides. Figure 12 shows the amplitude of the horizontal wind time series reconstructed from a inverse wavelet transform (see eqn. B1), for scales between 0.4 and 0.6 days for the semidiurnal tide, and 0.8 and 1.2 days for the diurnal tide. The diurnal tide in the zonal wind is seen to reach an amplitude of $\sim 50 \text{ ms}^{-1}$ during day 107 at a height of around 92 km, and 35-40 ms⁻¹ in the meridional component around 88 km during day 109. It should be noted that the hourly averaged zonal wind velocity (not shown here) reached a maximum of about 140 ms⁻¹ at 92 km during this period. The semidiurnal tide, whose

- 5 amplitude is known to rarely exceed 10 ms⁻¹ at Adelaide's location (e.g., Vincent et al. (1998)), also reached an amplitude of 35-40 ms⁻¹ during day 104 in both the zonal and meridional components, at a height of around 94 km. The figure additionally shows that the phase of the diurnal tide is modulated, with the time scale of those modulations appearing to follow the phases of the planetary wave activity in Fig. 11—although there are no noteworthy phase changes at the times of the sudden amplitude enhancements. The semidiurnal tidal phase is persistent, and also with a well-defined vertical progression, during the few days
- 10 in which its amplitude is large, but clearly has little meaningful structure at other times.

The large tidal amplitudes during this period lead us to expect the propagation directions of the GWs removed from the wave spectrum by the winds to exhibit a diurnal variation. A complicating factor is that these waves may also amplify, dampen or shift the phase of the tide, depending on the waves retained in the spectrum at the wave breaking height; the large variability in the tidal amplitudes during this period indicates that this may have indeed occurred. To provide some clarity on the extent

15 to which the GWs have been modulated by the tide and vice versa, in the next section we examine a composite day of the tidal winds, covariances and the implied flow accelerations over a 20-day interval spanning the interval in which the diurnal tide has a reasonably consistent phase and an enhanced amplitude.

4.2 Observed GW-tidal interaction

30

Figure 13 shows a composite day of the horizontal winds, covariances, and flow accelerations implied by the covariances, over
 5-25 April 2018 (i.e., days 95-115). The composite day consists of time windows of width 3 hours, with a gap time shift of 30 minutes between the centres of adjacent windows. The height binning again consists of 2 km-width bins with centres separated by 1 km.

The flow accelerations (e.g., in the case of the zonal direction) have been evaluated using the expression (e.g., Fritts (1984)):

25
$$\langle F_x \rangle = -\frac{1}{\rho(z)} \frac{\partial}{\partial z} \left(\rho(z) \langle u'w' \rangle \right),$$
 (11)

where $\rho(z)$ represents the neutral density as a function of height z. The density climatology we have used has been derived from the Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) satellite instrument (see App. C for details). Similarly to Liu et al. (2013), we also apply a low-pass filter with a cut-off wavelength of 10 km to the vertical profile of the covariance prior to evaluating its density-weighted derivative, in order to remove small-scale fluctuations from it that are clearly not associated with tidal modulation.



As expected from the amplitudes in Fig. 12, both horizontal wind components show a predominantly diurnal variation, with the meridional component lagging the zonal's by approximately 6 hours across the observed height region. The time of the zonal wind maximum occurs around 0 UT at 92 km, and 8-9 UT at 82 km.

In contrast, the $\langle u'w' \rangle$ covariance term shows a predominantly semidiurnal variation with little vertical phase progression, 5 maximising at around 0 and 12 UT, and minimising around 5 and 20 UT. The $\langle v'w' \rangle$ term is more variable with altitude, exhibiting a semidiurnal variation between 82 and 84 km, and a largely diurnal variation above this. The semidiurnal variation between 82 and 84 km is associated with positive covariances for the entire day except between about 18 and 24 UT, and the diurnal variation above is associated with negative values covariances between about 8 and 15 UT and positive otherwise.

Between about 88 and 92 km, the zonal flow acceleration shows a pronounced minimum between 4 and 6 UT, a maximum

10 around 13 UT at about 88 km, and a weaker minimum around 19 UT. The maximum occurs at a similar time to the corresponding zonal wind minimum, whereas the first minimum lags the zonal wind maximum by about 5 hours, and the second minimum precedes it by about 5 hours. Conversely, there is little flow acceleration structure below 87 km, other than a broad maximum at about 85 km around 1 UT. These observations are difficult to reconcile for three reasons: (1) the wave forcing is consistent with a rapid deceleration of the zonal wind from 4-6 UT at around 90 km, but there appears to be no positive forcing around 20 UT to accelerate the wind, (2) the strong positive forcing which does occur around 13 UT appears to result in little wind variability, and (3) the positive forcing around 85 km between 23 and 4 UT is associated with an acceleration of the zonal wind be table as a doe loop loop.

5 wind, but this acceleration is much smaller than that around 90 km.

From 88-92 km, the meridional flow acceleration shows a small maximum around 4 UT, a minimum at about 10 UT, and a large maximum around 20 UT. As per the zonal case, this leads to a peculiar relationship with the meridional wind; the forcing's large maximum occurs at a similar time to the wind minimum, the minimum corresponds roughly with a rapid wind deceleration, and the smaller maximum corresponds with a rapid wind acceleration. Also like in As for the zonal component,

10 there is little meridional flow acceleration structure below around 86 km.

5 Discussion

5.1 Uncertainties in $\langle u'w' \rangle$ and $\langle v'w' \rangle$ estimates

In the simulations section of this paper, we have tried to conclusively define estimates for the absolute and relative uncertainties of the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariance terms as measured by the multistatic BP-Mylor meteor radar, for typical time and height sampling cases. We subsequently replicated these sampling schemes on the case study data. Even with this replication, we have noticed that there are three main caveats in applying the uncertainties directly to the observations:

- 1. As shown by Kudeki and Franke (1998), the covariance estimation uncertainty is proportional to the geometric mean of the horizontal and vertical variances, in the case of sampling the wind field using a perfect "anemometer". Assuming this holds for a meteor radar-like detection distribution, this means that the absolute uncertainties of \langle u'w' \rangle and \langle v'w' \rangle reported in this paper should be similar for a given wave field, regardless of the value of |\langle u'w' \rangle| | |\langle v'w' \rangle. Therefore, the likelihood of correctly estimating the sign of one component at any given time/height may be very different to the other of the components in the presence of an anisotropic wave field may not be the same as for the other component.
- 2. As evidenced by the differences in the distribution widths of Figs. 5 and 7 for given detection rates, the relative uncertainties of a non-zero covariance term appear to be dependent on the total frequency/scale span of all the associated waves. In our example, the relative uncertainty in the covariance for a spectral GW field is around 8 times that for a single monochromatic GW. This finding, which is qualitatively consistent with the conclusion reached by Vincent et al. (2010) (for high meteor detection rates) and Fritts et al. (2012a), makes it impossible to accurately define the covariance measurement uncertainty for this radar without a-priori knowledge of the GW field and its variation with time.
- 3. The spectral components of the wave field may vary during the integration period. This is particularly problematic for the 10-day window; for example, during a period of intense but short-lived monochromatic wave events followed by more "complex" wave activity, increasing the integration time may actually increase the uncertainty in the covariance

20

25

estimate of the monochromatic wave activity—not only because of the likely change in the mean covariance, but also because of the "noise" added to the radial velocity time series by the more "complex" activity.

Despite these caveats, we can broadly conclude that the 10-day integrated covariances (Fig. 10), except where the absolute values are smaller than about 10-15 $m^2 s^{-2}$, are likely to be of the correct sign. The correlation length of the features in both the

5 tim

time and height domains also indicates that the noise component in the signal is considerably smaller than the sum of all the modes of geophysical variability. Additionally, at this time integration there is likely to be little difference in the uncertainty at the peak and edges of the height region analysed.

The 1-day integrated covariances (Fig. 11), in contrast, are far more affected by measurement noise. There is still some degree of temporal-height correlation, especially in the region of consistently high values of $\langle u'w' \rangle$ between days 105 and 110

10 above about 86 km, but very little below 84 km. The excursions below 84 km are of the same order as the simulations predict for 1 day of integration in a spectral wave field, so it may be that the noise component at these heights is considerably larger than the signal.

The 20-day composite covariances (Fig. 8), while clearly affected by measurement noise, do not show fluctuations from bin-to-bin of the same size as the uncertainties predicted in the corresponding simulation. This gives weight to the covariance

15 structures observed, and also suggests that the wave field being observed over the 20-day period was not as complex as the simulation's, nor particularly variable.

Unfortunately, it is impossible to know (using the meteor observations alone) if the discrepancies between the 1-day and the 10-day integration (for example, the absolute values of the covariances during the enhancement between days 105 and 110) are a result of statistical noise in the 1-day estimate or a precise estimate of a strong, transient monochromatic wave event using

20 the 1-day integration. The observation of waves in the MLT airglow may aid in the interpretation of how "monochromatic" the background wave field is; in future, we intend to complement these meteor radar case studies with images of the sodium and hydroxyl airglow taken nearby the BP site. This, in conjunction with the random resampling method employed by Liu et al. (2013), may lead to more refined uncertainty estimates.

In the 10-day integrated results, the small difference in measurement error at the peak and lower edge of the height distribution (around 20%, for an order of magnitude increase in detections) places an important question on the usefulness of further increasing the integration times/detection rates. On this point, Fritts et al. (2012a) argued that the covariance measurement error should decrease with the square root of the number of detections, and by extrapolating from the 250% error for a 1-hour integration presented in Vincent et al. (2010), concluded that their relative error for a one-month composite should have been

30 detection rate and a 3-hour bin in our 20 day composite of a spectral model-derived wave field shown in Fig. 8, we obtain a minimum relative error of about 70%. In saying this, we note of course that a relative error of 10% is possible for a considerably less complex wave field.

as low as 10%. Our simulations suggest that an increase in precision of this magnitude cannot occur. Moreover, using a similar

5.2 Effects of tides on covariance estimates

All of our simulations have shown that a systematic underestimation of non-zero covariances arises when an attempt is made to remove tidal effects. This clearly becomes more of a problem in the presence of large amplitude GWs with ground-based periods close to those of the tides. A number of questions about the process of tidal removal could be raised:

- 5 1. What is the importance of incorporating the momentum fluxes of gravity waves with ground-based periods close to the tides in climate models?
 - 2. If those longer-period waves are unimportant, what is an appropriate frequency cut-off for covariance measurements?
 - 3. If those waves are important, what is the optimal way to remove the tides?

With regard to 3., it may be that a wavelet/S-transform has insufficient frequency resolution to define solely tidal features; a
long-windowed harmonic fitting (as used by e.g., Andrioli et al. (2013a)) may be more appropriate if there is a specific interest in GW features close to or between the tidal periods. Of course, this method assumes no variability in the tidal amplitudes, tidal periods, or in the GW spectrum. The best way forward may be to simply apply both of the methods independently and contrast their effects.

5.3 Radial velocity outlier removal

- 15 In Sect. 3.11.3 we showed that the radial velocity outlier rejection scheme of eqn. (10) substantially increases the covariance measurement precision in the presence of outliers. However, we note that the criterion used (especially the "5 standard deviations" aspect) has not been rigorously tested; we merely selected it on the basis of it removing points in the distribution of $|v_{ri}'^2 - v_{rpi}'^2|$ (real and simulated) that we had noticed were spuriously affecting the covariance estimates. A more rigorous scheme would adaptively modify the thresholding based on observed characteristics of the wind field, rather than simply the
- 20 residual of the fit.

A complication arises from the fact that the criterion results in a more precise (albeit less accurate) covariance estimate in the absence of outliers. This also illustrates an important point about the sensitivity of the eqn. (9) inversion to the input: it is as though the data that contribute to the accuracy of the measurement actually increase the measurement's uncertainty, if they are associated with large radial velocity perturbations.

25 5.4 Weighting of meteors in the wind/covariance estimation fits

A subject we have not addressed in this paper is the application of weights to the meteors in the inversion of eqns. (8)/(9) to minimize the errors in the resulting winds/covariance estimates. In particular (as discussed by Hocking (2018)), at the midpoint between the transmitter and receiver sites the e vector (see Fig. 3) is vertical, meaning that the measured radial velocity corresponds to the true wind velocity projected onto the vertical. Large errors in the inverted horizontal winds/covariances may

30 result in the presence of radial velocity errors here, and at nearby locations where e is close to vertical. We decided to ignore



Figure 14. Simulated errors in flow acceleration estimates, using the bias mean and standard deviations in the Fig. 8 covariances.

the issue on the basis of there being a small number of meteors with sufficiently oblique entrance angles to be detected in this region; at Mylor, we found about 0.3% of all detected meteors to have effective zenith angles (that is, the zenithal orientation of the e vector) of less than 20°. Nevertheless, there is still a need to quantify the usefulness a weighting scheme may have in minimizing errors arising from these meteors.

5 5.5 Observed GW-tidal interaction

Our aim in analysing the GW-induced flow accelerations in Sect. 4.2 has been to verify that the estimated momentum fluxes were physically reasonable and devoid of tide-induced biases, and to contribute to the well-known gap in knowledge of GW effects on tides. Our analysis, which was centred on a 20-day interval containing an abrupt enhancement in tidal amplitudes, has yielded inconclusive results on whether the GW momentum deposition has on the whole enhanced, dampened or changed

10 the phase of the tidal motions. Nevertheless, the expected uncertainties in the flow accelerations based on the bias mean and standard deviations in the Fig. 8 covariances, shown in Fig. 14, indicate that the signal components between 84 and 90 km shown in Fig. 13 will have well exceeded the noise levels.

The results are complex, illustrating tidal enhancement at some times of day, dampening at others, and that there are also times in which a forcing is present but no apparent effect on the tide is clear. A broad observation is that the forcing components

15 have a more pronounced diurnal variability between about 86 and 92 km, with the result that the forcing dampens the tide at the tide's minimum (i.e. westward and southward phase), and shifts its phase at its maximum. Of course, our interpretation is complicated by the fact that we have no knowledge of what the tidal features may have looked like without any GW forcing.

It is widely accepted in modelling studies that GW forcing plays a role in the observed seasonal variation of the migrating diurnal tide (DW1) amplitudes (i.e. equinoctial maxima and solsticial minima), and that whether amplification or dampening of

20 the amplitude occurs depends on the GW source spectrum (e.g., Ortland and Alexander (2006); Yiğit and Medvedev (2017)). However, there is still ongoing debate about whether or not the forcing is responsible for all of DW1's observed amplitude and phase variability. For example, both Mayr et al. (1998) and Watanabe and Miyahara (2009) have concluded that the forcing is in phase with DW1 during the equinoxes and out of phase during the solstices, leading to DW1's amplification at the equinoxes and dampening at the solstices. Yiğit and Medvedev (2017) reached the same conclusion for the September equinox, but stated that Watanabe and Miyahara (2009) may have significantly underestimated the magnitude of the forcing. In contrast, for the March equinox Lu et al. (2012) has argued that the tidal variability is caused by a superposition of GW forcing and advection terms that varies with altitude and latitude, and that GW forcing exclusively dampens tidal amplitudes in the MLT/I. Moreover,

5 Lu et al. (2012) has reported considerably larger GW forcing magnitudes than in a related modelling study by McLandress (2002).

The small number of recent observational studies that have sought to quantify the effect of GW forcing on the DW1 amplitude and phase have also yielded contradictory results. For example, using TIMED satellite data Lieberman et al. (2010) showed that while the zonal and meridional GW forcing maximises at the equinoxes and minimises at the solstices, the zonal forcing

- 10 is in quadrature with the zonal tidal wind, and the meridional forcing is out of phase with the meridional tidal wind, leading to a zonal tide with advanced phase and a dampened meridional tide. They noted that the zonal advection due to variability in the meridional DW1 amplitude also, like the GW forcing, maximised at the equinoxes and minimised at the solstices, but were not able to reconcile if this variability was a cause or an effect of the seasonal DW1 variation. Also using TIMED data, Xu et al. (2009) concluded that the GW-induced dampening of tidal amplitudes is largest during equinoxes, and therefore that
- 15 dampening cannot cause the observed seasonal variation in tidal amplitudes. In contrast, using measurements from a groundbased meteor radar in Hawaii (20.7°N, 156.3°W), Liu et al. (2013) noted that GW forcing tends to slightly dampen the DW1 amplitude below 90 km, but enhance it above 90 km. Using a similar approach on LIDAR data from Starfire Optical Range (35.0°N, 106.5°W), Agner and Liu (2015) also noted that GW forcing can amplify or dampen the DW1 amplitudes, depending on the altitude.
- Tides may also interact with GWs through the diurnal variations in atmospheric stability they induce (i.e., making conditions more favourable for GW breaking and hence GW forcing at particular times of day). For example, Fritts et al. (1988) showed from observations at Scott Base, Antarctica that the highest levels of turbulence due to convective instability occurred at the times that the vertical component of the tidal wind induced the most negative value of dT/dz (the vertical temperature gradient). Using temperature perturbations from the GSWM-98 model for the BP site, Holdsworth et al. (2001) also showed
- that maximum negative values of dT/dz were in phase with the maximum values of the turbulent velocity measured by the BP MF around the autumnal equinox. Using GSWM-00 output, we have noted that the maximum negative dT/dz (of ~ -1 K km⁻¹) should occur between 1-3 UT across the 85-92 km region at the BP site during the period of our composite day analysis; curiously, we observe large positive values of $\langle F_x \rangle$ at this time just below this region, and an abrupt shift in the sign of $\langle F_x \rangle$ above it. As Holdsworth et al. (2001) notes, while a dT/dz of this size is too small to result in static instability, it still
- 30 corresponds with a large level of GW forcing and the maximum eastward phase of the diurnal tide, which we have observed to be particularly large during this interval.

6 Conclusions

This study has defined limits on the expected uncertainties in estimates of the $\langle u'w' \rangle$ and $\langle v'w' \rangle$ covariance terms made using a multistatic meteor radar, and has presented an example case study of using the radar to measure the GW forcing on the diurnal tide that arises from the height variation of the measured covariances. We have concluded that the extra detections offered by

- 5 the bistatic receiver appreciably improve the precision of the covariance measurements, although little of that improvement can be attributed to the increased Bragg vector diversity associated with having two viewing perspectives. The winds observed in the case study revealed substantial variations in the amplitude of the diurnal tide, but we were unable to conclusively show if GW forcing caused this variation. Nevertheless, our simulations have indicated that the bulk of the variability in the covariance and GW forcing we have seen far exceeds the expected measurement uncertainties, and therefore that GW forcing has not been
- 10 the only contributor to the tidal variability. We note that studies concerning GW forcing on tides are few, and that there is a clear need for further studies at other locations. Furthermore, there is a need for a definition of the part of the GW spectrum that is most likely to contribute to forcing on the tides; this will inform what periodicities in the time series should be filtered out prior to making a covariance estimate.
- Our simulations showed that 10-day-integrated covariance estimates could broadly be considered as reliable for our 55 MHz 15 multistatic radar configuration; shorter integration times may of course be possible for lower frequency radars with higher 16 meteor detection rates. However, we did note that the uncertainty appears to asymptote towards a minimum value after about 10 days of integration; this value is clearly governed by the wave field characteristics. We also suggest that the accuracy and precision of the covariance estimates may be able to be improved slightly by using a more rigorous radial velocity outlier rejection scheme than applied here.
- 20 *Code and data availability.* The simulation code developed in this study is available on request from Andrew J. Spargo, as is the data from the BP and Mylor meteor radars.

Appendix A: Procedure used for converting between coordinate systems

To embody the ellipticity of the Earth's surface in the estimation of meteor altitudes, Bragg vector orientations, and wind field components (for both bistatic and monostatic receiver cases), we followed the coordinate system conversion algorithms outlined by Stober et al. (2018). However, we note that we applied a correction to their reported expression of the radius of

$$N(\phi) = \frac{a}{\sqrt{1 - e^2 \sin^2 \phi}},$$

curvature of the Earth $N(\phi)$, viz.:

25

where a is the semi-major axis of the Earth, and e^2 is the first numerical eccentricity of the Earth ellipsoid.

Furthermore, in the interests of reducing computational overhead we applied the Olson (1996) method for converting ECEF coordinates to geodetic coordinates, rather than the Heikkinnen (1982) method used by Stober et al. (2018).

Appendix B: Extraction of tidal features through the use of a wavelet transform

The time series reconstructed from the wavelet transform can be expressed as (Torrence and Compo (1998), eqn. (11)):

5
$$x_n = \frac{\delta j \, \delta t^{1/2}}{C_\delta \, \psi_0(0)} \sum_{j=0}^J \frac{\Re(W_n(s_j))}{s_j^{1/2}},$$
 (B1)

where δj describes the wavelet scale separation, δt represents the time separation between adjacent points, J is the number of wavelet scales, C_{δ} is a reconstruction factor (0.776 for the Morlet wavelet), $\psi_0(0)$ is an energy scaling factor ($\pi^{-1/4}$ for the Morlet wavelet), s_j are the wavelet scales, and $W_n(s_j)$ contains the complex wavelet transform coefficients at scale s_j . In reconstructing the hourly-averaged wind time series (regardless of the time series length), we have taken $\delta j = 0.02$, in contrast

to Torrence and Compo (1998) (Sect. 2f), who choose $\delta j = 0.125$ in their example with the Morlet wavelet; we have done this to reduce the spacing between adjacent wavelet scales and hence improve the accuracy of the reconstruction. Also in contrast to Torrence and Compo (1998) (Sect. 2g), we have not applied any zero padding in the application of the wavelet transform. This was done given our finding that the magnitude of artefacts at the ends of the wind time series appeared to be larger with zero padding applied.

15 Appendix C: SABER-derived density climatology creation

To create a climatology of the diurnal variability in density from SABER instrument data that was representative of conditions around Adelaide during the autumnal equinox, we acquired densities from individual limb scans with tangent point latitudes spanning 28°S-42°S, longitudes 108°E-168°E, days 01-March to 31-May inclusive, and years 2008-2018 inclusive. Measurements falling into given time-of-day (hourly) and height (0.5 km) bins were averaged.

A spatial sampling region and measurement time-of-year span of this size was necessary to fill all time-of-day bins with measurements. An average over 11 years of data was performed to reduce the level of aliasing arising from GW-induced perturbations occurring in individual scans.

The climatology produced using this method had features that were qualitatively consistent with the same time averaging on NRLMSISE-00 model output from Adelaide's location. However, we did note that given density surfaces from SABER

25 were, on average, 2 km lower than NRLMSISE-00's predictions between about 80 and 95 km. Nevertheless, the use of the SABER-derived density climatology in the production of Fig. 13 yielded almost identical flow accelerations as the use of said the NRLMSISE-00 output.

Author contributions. Andrew Spargo carried out the model development and data analysis, and wrote the paper. Iain Reid contributed to the Instrumentation section, and made other minor revisions to initial drafts of the paper. Iain Reid is the principal supervisor of Andrew Spargo's postgraduate candidature, and Andrew MacKinnon the co-supervisor.

Competing interests. The meteor radars used in this study were designed and manufactured by ATRAD Pty. Ltd., and Iain Reid is the 5 executive director of this group of companies.

Acknowledgements. Andrew Spargo would like to thank Jorge Chau, Chris Adami, Bob Vincent, David Holdsworth, <u>Gunter Stober</u>, Joel Younger, Richard Mayo, Andrew Heitmann, Yi Wen, Tom Chambers and Baden Gilbert for useful discussions regarding this work. Andrew Spargo is supported by an Australian Government Research Training Program Scholarship.

The BP ST/Meteor radar is supported by ATRAD Pty. Ltd. and the University of Adelaide. The Mylor receiving site and equipment is supported solely by ATRAD Pty. Ltd.

10

References

- Agner, R. and Liu, A. Z.: Local time variation of gravity wave momentum fluxes and their relationship with the tides derived from LIDAR measurements, J. Atmos. and Sol.-Terr. Phys., 135, 136–142, https://doi.org/https://doi.org/10.1016/j.jastp.2015.10.018, 2015.
- Andrioli, V. F., Fritts, D. C., Batista, P. P., and Clemesha, B. R.: Improved analysis of all-sky meteor radar measurements of gravity wave variances and momentum fluxes, Ann. Geophys., 31, 889–908, https://doi.org/doi:10.5194/angeo-31-889-2013, 2013a.
- Andrioli, V. F., Fritts, D. C., Batista, P. P., Clemesha, B. R., and Janches, D.: Diurnal variation in gravity wave activity at low and middle latitudes, Ann. Geophys., 31, 2123–2135, https://doi.org/doi:10.5194/angeo-31-2123-2013, 2013b.
 - Andrioli, V. F., Batista, P. P., Clemesha, B. R., Schuch, N. J., and Buriti, R. A.: Multi-year observations of gravity wave momentum fluxes at low and middle latitudes inferred by all-sky meteor radar, Ann. Geophys., 33, 1183–1193, https://doi.org/doi:10.5194/angeo-33-1183-
- 10 2015, 2015.

5

- Antonita, T. M., Ramkumar, G., Kumar, K. K., and Deepa, V.: Meteor wind radar observations of gravity wave momentum fluxes and their forcing toward the Mesospheric Semiannual Oscillation, J. Geophys. Res. Atmos., 113, 2008.
- Beldon, C. L. and Mitchell, N. J.: Gravity waves in the mesopause region observed by meteor radar, 2: Climatologies of gravity waves in the Antarctic and Arctic, J. Atmos. Sol. Terr. Phys., 71, 875–884, 2009.
- 15 Beldon, C. L. and Mitchell, N. J.: Gravity wave–tidal interactions in the mesosphere and lower thermosphere over Rothera, Antarctica (68°S, 68°W), J. Geophys. Res. Atmos., 115, 2010.
 - Bowring, B. R.: Transverse Mercator equations obtained from a spherical basis, Survey Review, 30, 125–133, https://doi.org/10.1179/sre.1989.30.233.125, 1989.

Chau, J. L. and Clahsen, M.: Empirical phase calibration for multistatic specular meteor radars using a beamforming approach, Radio Sci.,

20 54, https://doi.org/https://doi.org/10.1029/2018RS006741, 2019.

Clemesha, B. R. and Batista, P. P.: Gravity waves and wind-shear in the MLT at 23°S, Adv. Space Res., 41, 1472–1477, 2008.

- Clemesha, B. R., Batista, P. P., da Costa, R. A. B., and Schuch, N.: Seasonal variations in gravity wave activity at three locations in Brazil, Ann. Geophys., 27, 1059–1065, 2009.
- de Wit, R. J., Hibbins, R. E., and Espy, P. J.: The seasonal cycle of gravity wave momentum flux and forcing in the high latitude northern
- 25 hemisphere mesopause region, J. Atmos. Sol. Terr. Phys., 127, 21–29, 2014a.
 - de Wit, R. J., Hibbins, R. E., Espy, P. J., Orsolini, Y. J., Limpasuvan, V., and Kinnison, D. E.: Observations of gravity wave forcing of the mesopause region during the January 2013 major Sudden Stratospheric Warming, Geophys. Res. Lett., 41, 4745–4752, 2014b.
 - de Wit, R. J., Janches, D., Fritts, D. C., and Hibbins, R. E.: QBO modulation of the mesopause gravity wave momentum flux over Tierra del Fuego, Geophys. Res. Lett., 43, 4049–4055, https://doi.org/10.1002/2016GL068599, 2016GL068599, 2016.
- 30 Dolman, B. K., Reid, I. M., and Tingwell, C.: Stratospheric tropospheric wind profiling radars in the Australian network, Earth, Planets and Space, 70, 170, 2018.

Ern, M., Preusse, P., Gille, J. C., Hepplewhite, C. L., Mlynczak, M. G., Russell, J. M., and Riese, M.: Implications for atmospheric dynamics derived from global observations of gravity wave momentum flux in stratosphere and mesosphere, J. Geophys. Res. Atmos., 116, n/a–n/a, https://doi.org/10.1029/2011JD015821, d19107, 2011.

35 Fritts, D. C.: Gravity wave saturation in the middle atmosphere: A review of theory and observations, Reviews of Geophysics, 22, 275–308, 1984.

Fritts, D. C. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, Rev. Geophys., 41, 2003.

- Fritts, D. C., Smith, S. A., Balsley, B. B., and Philbrick, C. R.: Example of gravity wave saturation and local turbulence production in the summer mesosphere and lower thermosphere during the STATE experiment, J. Geophys. Res., 93, 7015–7025, 1988.
- Fritts, D. C., Janches, D., and Hocking, W. K.: Southern Argentina Agile Meteor Radar: Initial assessment of gravity wave momentum fluxes, J. Geophys. Res. Atmos., 115, 2010a.
- 5 Fritts, D. C., Janches, D., Iimura, H., Hocking, W. K., Mitchell, N. J., Stockwell, R. G., Fuller, B., Vandepeer, B., Hormaechea, J., Brunini, C., et al.: Southern Argentina Agile Meteor Radar: System design and initial measurements of large-scale winds and tides, J. Geophys. Res. Atmos., 115, 2010b.
 - Fritts, D. C., Janches, D., Hocking, W. K., Mitchell, N. J., and Taylor, M. J.: Assessment of gravity wave momentum flux measurement capabilities by meteor radars having different transmitter power and antenna configurations, J. Geophys. Res. Atmos., 117, 2012a.
- 10 Fritts, D. C., Janches, D., Iimura, H., Hocking, W. K., Bageston, J. V., and Leme, N. M. P.: Drake Antarctic Agile Meteor Radar first results: Configuration and comparison of mean and tidal wind and gravity wave momentum flux measurements with Southern Argentina Agile Meteor Radar, J. Geophys. Res. Atmos., 117, 2012b.
 - Fritts, D. C., Smith, R. B., Taylor, M., Doyle, J. D., Eckermann, S. D., Dörnbrack, A., Rapp, M., Williams, B. P., Pautet, D., Bossert, K., et al.: The Deep Propagating Gravity Wave Experiment (DEEPWAVE): An Airborne and Ground-Based Exploration of Gravity Wave
- 15 Propagation and Effects from their Sources throughout the Lower and Middle Atmosphere, Bull. Amer. Meteor. Soc., 97, 425–453, 2016. Gardner, C. S., Hostetler, C. A., and Franke, S. J.: Gravity wave models for the horizontal wave number spectra of atmospheric velocity and density fluctuations, J. Geophys. Res. Atmos., 98, 1035–1049, https://doi.org/10.1029/92JD02051, 1993.
 - Heikkinnen, M.: Geschlossene Formeln zur Berechnung räumlicher geodätischer Koordinaten aus rechtwinkligen Koordinaten, Z. Ermess., 107, 207–211, 1982.
- 20 Hocking, W. K.: A new approach to momentum flux determinations using SKiYMET meteor radars, Ann. Geophys., 23, 2433–2439, 2005. Hocking, W. K.: Spatial distribution of errors associated with multistatic meteor radar, Earth, Planets and Space, 70, 93, https://doi.org/10.1186/s40623-018-0860-2, https://doi.org/10.1186/s40623-018-0860-2, 2018.
 - Hocking, W. K. and Thayaparan, T.: Simultaneous and co-located observation of winds and tides by MF and meteor radars over London, Canada (43°N, 81°W), during 1994-1996, Radio Sci., 32, 833–865, https://doi.org/https://doi.org/10.1029/96RS03467,, 1997.
- 25 Holdsworth, D. A., Vincent, R. A., and Reid, I. M.: Mesospheric turbulent velocity estimation using the Buckland Park MF radar, Ann. Geophys., 19, 1007–1017, 2001.
 - Holdsworth, D. A., Reid, I. M., and Cervera, M. A.: Buckland Park all-sky interferometric meteor radar, Radio Sci., 39, https://doi.org/doi:10.1029/2003RS003014, 2004a.
 - Holdsworth, D. A., Tsutsumi, M., Reid, I. M., Nakamura, T., and Tsuda, T.: Interferometric meteor radar phase calibration using meteor echoes, Radio Sci., 39, https://doi.org/doi:10.1029/2003RS003026, 2004b.
 - Jia, M., Xue, X., Gu, S., Chen, T., Ning, B., Wu, J., Zeng, X., and Dou, X.: Multiyear Observations of Gravity Wave Momentum Fluxes in the Midlatitude Mesosphere and Lower Thermosphere Region by Meteor Radar, J. Geophys. Res. Space Phys., 123, 5684–5703, https://doi.org/10.1029/2018JA025285, 2018.

Jones, J., Webster, A., and Hocking, W.: An improved interferometer design for use with meteor radars, Radio Sci., 33, 55–65, 1998.

35 Kim, Y.-J., Eckermann, S. D., and Chun, H.-Y.: An overview of the past, present and future of gravity-wave drag parametrization for numerical climate and weather prediction models, Atmos. Ocean, 41, 65–98, 2003.

Kudeki, E. and Franke, S. J.: Statistics of momentum flux estimation, J. Atmos. Sol. Terr. Phys., 60, 1549–1553, 1998.

30

- Lieberman, R. S., Ortland, D. A., Riggin, D. M., Wu, Q., and Jacobi, C.: Momentum budget of the migrating diurnal tide in the mesosphere and lower thermosphere, J. Geophys. Res. Atmos., 115, https://doi.org/10.1029/2009JD013684, 2010.
- Liu, A. Z., Lu, X., and Franke, S. J.: Diurnal variation of gravity wave momentum flux and its forcing on the diurnal tide, J. Geophys. Res. Atmos., 118, 1668–1678, 2013.
- 5 Lu, X., Liu, H.-L., Liu, A. Z., Yue, J., McInerney, J. M., and Li, Z.: Momentum budget of the migrating diurnal tide in the Whole Atmosphere Community Climate Model at vernal equinox, J. Geophys. Res. Atmos., 117, https://doi.org/10.1029/2011JD017089, 2012.
 - Matsumoto, M., Shinbori, A., Riggin, D. M., and Tsuda, T.: Measurement of momentum flux using two meteor radars in Indonesia, Ann. Geophys., 34, 369–377, https://doi.org/doi:10.5194/angeo-34-369-2016, 2016.
 - Mayr, H. G., Mengel, J. G., Chan, K. L., and Porter, H. S.: Seasonal variations of the diurnal tide induced by gravity wave filtering, Geophys.
- 10 Res. Lett., 25, 943–946, 1998.
 - McLandress, C. L.: The seasonal variation of the propagating diurnal tide in the mesosphere and lower thermosphere. Part I: The role of gravity waves and planetary waves, J. Atmos. Sci., 59, 893–906, 2002.
 - Nicolls, M. J., Fritts, D. C., Janches, D., and Heinselman, C. J.: Momentum flux determination using the multi-beam Poker Flat Incoherent Scatter Radar, Ann. Geophys., 30, 945–962, 2012.
- 15 Olson, D. K.: Converting Earth-centered, Earth-fixed coordinates to geodetic coordinates, IEEE Transactions on Aerospace and Electronic Systems, 32, 473–476, https://doi.org/10.1109/7.481290, 1996.
 - Ortland, D. A. and Alexander, M. J.: Gravity wave influence on the global structure of the diurnal tide in the mesosphere and lower thermosphere, J. Geophys. Res., 11, https://doi.org/10.1029/2005JA011,467, 2006.
- Placke, M., Hoffmann, P., Becker, E., Jacobi, C., Singer, W., and Rapp, M.: Gravity wave momentum fluxes in the MLT—Part II: Meteor
 radar investigations at high and midlatitudes in comparison with modeling studies, J. Atmos. Sol. Terr. Phys., 73, 911–920, 2011a.
- Placke, M., Stober, G., and Jacobi, C.: Gravity wave momentum fluxes in the MLT—Part I: seasonal variation at Collm (51.3°N, 13.0°E), J. Atmos. Sol. Terr. Phys., 73, 904–910, 2011b.
 - Placke, M., Hoffmann, P., Latteck, R., and Rapp, M.: Gravity wave momentum fluxes from MF and meteor radar measurements in the polar MLT region, J. Geophys. Res. Sp. Phys., 120, 736–750, https://doi.org/10.1002/2014JA020460, 2014.
- 25 Placke, M., Hoffmann, P., and Rapp, M.: First experimental verification of summertime mesospheric momentum balance based on radar wind measurements at 69°N, Ann. Geophys., 33, 1091–1096, 2015.
 - Protat, A. and Zawadzki, I.: A Variational Method for Real-Time Retrieval of Three-Dimensional Wind Field from Multiple-Doppler Bistatic Radar Network Data, J. Atmos. Oc. Tech., 16, 432–449, https://doi.org/10.1175/1520-0426(1999)016<0432:AVMFRT>2.0.CO;2, 1999.
 - Reid, I. M., McIntosh, D. L., Murphy, D. J., and Vincent, R. A.: Mesospheric radar wind comparisons at high and middle southern latitudes,
- 30 Earth, Planets and Space, 70, 84, 2018a.
 - Reid, I. M., Rüster, R., Czechowsky, P., and Spargo, A. J.: VHF radar measurements of momentum flux using summer polar mesopause echoes, Earth, Planets and Space, 70, 129, 2018b.
 - Riggin, D. M., Tsuda, T., and Shinbori, A.: Evaluation of momentum flux with radar, J. Atmos. Sol. Terr. Phys., 2016.
- Spargo, A. J., Reid, I. M., MacKinnon, A. D., and Holdsworth, D. A.: Mesospheric gravity wave momentum flux estimation using hybrid
 Doppler interferometry, Ann. Geophys., 35, 733–750, https://doi.org/10.5194/angeo-35-733-2017, 2017.
- Stober, G. and Chau, J. L.: A multi-static and multi-frequency novel approach for specular meteor radars to improve wind measurements in the MLT region, Radio Sci., 50, 2015.

Stober, G., Chau, J. L., Vierinen, J., Jacobi, C., and Wilhelm, S.: Retrieving horizontally resolved wind fields using multi-static meteor radar observations, Atmos. Meas. Tech., 11, 4891–4907, https://doi.org/10.5194/amt-11-4891-2018, 2018.

Thomas, R. M., Whitham, P. S., and Elford, W. G.: Frequency Dependence of Radar Meteor Echo Rates, Proc. ASA, 6, 303–306, 1986.

- Thorsen, D., Franke, S. J., and Kudeki, E.: A new approach to MF radar interferometry for estimating mean winds and momentum flux, Radio Sci., 32, 707–726, 1997.
- Torrence, C. and Compo, G. P.: A Practical Guide to Wavelet Analysis, Bulletin of the American Meteorological Society, 1998.
- Vincent, R. A. and Ball, S. M.: Meospheric winds at low- and mid-latitudes in the southern hemisphere, J. Geophys. Res. Space Phys., 86, 9159–9169, 1981.

Vincent, R. A. and Reid, I. M.: HF Doppler measurements of mesospheric gravity wave momentum fluxes, J. Atmos. Sci., 40, 1321–1333,

10 1983.

5

- Vincent, R. A., Kovalam, S., Fritts, D. C., and Isler, J. R.: Long-term MF radar observations of solar tides in the low-latitude mesosphere: Interannual variability and comparisons with the GSWM, J. Geophys. Res. Atmos., 103, 8667–8683, https://doi.org/10.1029/98JD00482, 1998.
- Vincent, R. A., Kovalam, S., Reid, I. M., and Younger, J. P.: Gravity wave flux retrievals using meteor radars, Geophys. Res. Lett., 37, 2010.
- 15 Watanabe, S. and Miyahara, S.: Quantification of the gravity wave forcing of the migrating diurnal tide in a gravity wave-resolving general circulation model, J. Geophys. Res., 114, https://doi.org/10.1029/2008JD011218, 2009.
 - Xu, J., Smith, A. K., Liu, H.-L., Yuan, W., Wu, Q., Jiang, G., Mlynczak, M. G., and Russell, J. M.: Estimation of the equivalent Rayleigh friction in mesosphere/lower thermosphere region from the migrating diurnal tides observed by TIMED, J. Geophys. Res., 114, https://doi.org/10.1029/2009JD012209, 2009.
- 20 Yiğit, E. and Medvedev, A. S.: Influence of parameterized small-scale gravity waves on the migrating diurnal tide in Earth's thermosphere, J. Geophys. Res. Space Phys., 122, 4846–4864, https://doi.org/10.1002/2017JA024089, 2017.