Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2019-144-AC2, 2019 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.





Interactive comment

Interactive comment on "Quantifying organic matter and functional groups in particulate matter filter samples from the southeastern United States, part I: Methods" by A. J. Boris et al.

A. J. Boris et al.

aboris08@gmail.com

Received and published: 7 August 2019

Referee comment:

Based on a series of previous studies (such as Takahama), the author improved the current FTIR measurement method for quantifying organic aerosols. The research topic is of great significance. It can be used not only to quantify the mass concentration of organic compounds, but also quantify the chemical functional group information of organic aerosols.

Author response:

Printer-friendly version

Discussion paper



Thank you for your kind summary of, and comments regarding, our work.

Referee comment:

Here is a suggestion to improve this paper. It is hoped that the author can analyze and determine the functional groups in at least a few different samples by other instruments, such as NMR or HR-AMS. The results were used to confirm that the author's FTIR method for the determination of functional groups can be matched with the nonconventional analysis methods.

Author response:

We appreciate this insightful suggestion. Although a comparison to other direct measurements of OM in the same sampling time periods and locations such as NMR or AMS was not possible for this work, we have compared to various southeastern US aerosol measurements: AMS measurements of O/C and H/C ratios (using the van Krevelen space; Section 3.4.4), various measurements of OM/OC, as well as residual OM and TOR OC measurements (Sections 3.4.1 and 3.4.3). In addition to these comparisons, our introduction mentions the work of previous studies directly comparing AMS fragment and FT-IR spectrometry functional group abundances. We have responded to a similar comment made by Reviewer #1, and respectfully request that Reviewer #2 also view our response to Reviewer #1.

References

El-Zanan, H. S., Zielinska, B., Mazzoleni, L. R. and Hansen, D. A.: Analytical determination of the aerosol organic mass-to-organic carbon ratio, J. Air Waste Manag. Assoc., 59(1), 58–69, doi:10.3155/1047-3289.59.1.58, 2009.

Hand, J. L., Prenni, A. J., Schichtel, B. A., Malm, W. C. and Chow, J. C.: Trends in remote PM 2.5 residual mass across the United States: Implications for aerosol mass reconstruction in the IMPROVE network, Atmos. Environ., 203(January), 141–152, doi:10.1016/j.atmosenv.2019.01.049, 2019.



Interactive comment

Printer-friendly version

Discussion paper



Kamruzzaman, M., Takahama, S. and Dillner, A. M.: Quantification of amine functional groups and their influence on OM/OC in the IMPROVE network, Atmos. Environ., 172, 124–132, doi:10.1016/j.atmosenv.2017.10.053, 2018.

Liu, J., Russell, L. M., Ruggeri, G., Takahama, S., Claflin, M. S., Ziemann, P. J., Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., McKinney, K. A., Budisulistiorini, S. H., Bertram, T. H., Nenes, A. and Surratt, J. D.: Regional Similarities and NOx-Related Increases in Biogenic Secondary Organic Aerosol in Summertime Southeastern United States, J. Geophys. Res. Atmos., 123(18), 10,620-10,636, doi:10.1029/2018JD028491, 2018.

Pitchford, M., Malm, W., Schichtel, B., Kumar, N., Lowenthal, D. and Hand, J.: Revised algorithm for estimating light extinction from IMPROVE particle speciation data, J. Air Waste Manag. Assoc., 57(11), 1326–1336, doi:10.3155/1047-3289.57.11.1326, 2007.

Ruthenburg, T. C., Perlin, P. C., Liu, V., McDade, C. E. and Dillner, A. M.: Determination of organic matter and organic matter to organic carbon ratios by infrared spectroscopy with application to selected sites in the IMPROVE network, Atmos. Environ., 86, 47–57, doi:10.1016/j.atmosenv.2013.12.034, 2014.

Simon, H., Bhave, P. V, Swall, J. L., Frank, N. H. and Malm, W. C.: Determining the spatial and seasonal variability in OM/OC ratios across the US using multiple regression, Atmos. Chem. Phys., 11, 2933–2949, doi:10.5194/acp-11-2933-2011, 2011.

Takahama, S. and Ruggeri, G.: Technical note: Relating functional group measurements to carbon types for improved model-measurement comparisons of organic aerosol composition, Atmos. Chem. Phys., 17(7), 4433–4450, doi:10.5194/acp-17-4433-2017, 2017.

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2019-144, 2019.

AMTD

Interactive comment

Printer-friendly version

Discussion paper

