

First comment 1:

“However, I have, besides some minor remarks, a major concern: you do not take into account any secondary radical-radical reaction with the argument, that your radical concentrations are low enough. I do not agree with this point, even though it is not always easy to get enough details from the manuscript to judge. So my comment is based on your statement page 6, that the typical initial OH concentration is between 2×10^{11} and $5 \times 10^{13} \text{ cm}^{-3}$. In the below graph are shown two simulations with $[\text{OH}]_0 = 1 \times 10^{12}$ and $\text{H}_2\text{O}_2 = 5 \times 10^{14}$ (left) and $[\text{OH}]_0 = 1 \times 10^{13}$ and $[\text{H}_2\text{O}_2] = 1 \times 10^{15}$ (right graph). The blue symbols show the simple model $\text{OH} + \text{H}_2\text{O}_2 \rightarrow \text{HO}_2 + \text{H}_2\text{O}$, while the green symbols include on top the reaction of $\text{OH} + \text{HO}_2 \rightarrow \text{H}_2\text{O} + \text{O}_2$ with $1 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$. t / s [OH, HO₂] 0.000 0.002 0.004 0.006 0.008 0.010 0 5.0×10¹ 1 1.0×10¹ 2 HO₂ OH with secondary reactions OH w/o secondary reactions Y₀ Plateau K oh 1.001e+012 = 0.0 887.4 x 10.000e+011 = 0.0 848.7 HO₂_ini k_slow HO₂_sec k_fast ho2 = 0.0 3.015 8.923e+011 949.9 xy = 0.0 -0.03680 9.998e+011 849.2 HO₂ t / s [OH, HO₂] 0.000 0.001 0.002 0.003 0.004 0.005 0 5.0×10¹ 2 1.0×10¹ 3 Y₀ Plateau K oh 1.002e+013 = 0.0 2003 x 9.998e+012 = 0.0 1687 HO₂_ini k_slow HO₂_sec k_fast ho2 = 0.0 20.16 6.379e+012 2596 xy = 0.0 -0.09674 9.993e+012 1691 ho2 oh x It can very clearly be seen that even under the relatively low initial radical concentration of 1×10^{12} (which is at your lower end) already the HO₂ yield is not 100% anymore, situation gets much worse with 1×10^{13} OH: only 60% of the initial OH is converted to HO₂. This has also an influence on the OH decay rate, as well as on the retrieved HO₂ rise time (both get faster). This “problem” has been discussed in detail by Assaf et al, JPCA 2016, when using this system to retrieve the OH absorption cross section. In your case not taking into account secondary chemistry will lead to an overestimation of the HO₂ yield.

Of course taking into account this chemistry is possibly only if you know the absolute initial OH concentration. Maybe you did some experiments where you varied the photolysis energy? Because this would give you an idea if secondary reactions are important or not under your conditions. In the case of the OH + CH₃OH experiments, secondary chemistry might play a role as well. Very recently, Assaf et al (PCCP, 20, 10660, 2018) have measured the rate constant of CH₃O + HO₂ and CH₃O + CH₃O, both have found to be very fast (1.1×10^{-10} and $7 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$). The result is that even under moderate high initial radical concentrations, some CH₃O will react away before it is converted into HO₂. You find a yield in good agreement with literature, either your initial radical concentration are at the lower end of the indicated range, or maybe the internal calibration, tending to overestimate the yield, makes up for this underestimation. Please give more information on the estimated initial radical concentration for the different experiments and check, if your systems are really free from secondary chemistry. In any case, before I can agree to the sentence that your instrument can accurately measure HO₂ yields, I would like to see a more detailed discussion on possible secondary chemistry.”

Response

Thank you for this very pertinent question. Prior to submission many checks for radical-radical effects were made by varying the repetition rate and photolysis laser power, and no observed differences were seen in the HO₂ yields. We had based our statement on the empirical observations rather than a review of the possible secondary chemistry, however, your careful review and analysis, does suggest that we ought to see a significant difference. The literature does imply we should see a change in HO₂ yield between OH + H₂O₂ and OH + CH₃OH/O₂ (n.b. note that the same high [O₂] is used in both experiments). As the [OH] is increased we should get an ~50 % yield when $[\text{OH}] \sim 3 \times 10^{13} \text{ cm}^{-3}$ and OH + H₂O₂ is used as the OH → HO₂ conversion reaction. Also the observed OH removal kinetics of the OH + H₂O₂ reaction should increase with [OH]; >20% faster when $[\text{OH}] \sim 3 \times 10^{13} \text{ cm}^{-3}$. Because of this inconsistency of our result with the literature, we have carried a number of new experiments, where the [OH] is

varied over a greater range, varying pump laser power from 0.5-60 mJ cm⁻². The take home message is that we cannot reproduce the literature, and our HO₂ yields / kinetics for the reaction OH + H₂O₂ are close to unchanged over all [OH] from 2E11 up to 5E13 molecule cm⁻³.

We are in agreement with the reviewer as to the implications of the literature. From a model (as detailed by the reviewer), when [OH]₀ = 1E13 cm⁻³, the HO₂ yield from OH + H₂O₂ is about 50% compared to when a large excess of methanol is added, see Fig 1.




Figure 1 A simulation of the expected HO₂ yields for reaction of 9E12 OH with 1E15 H₂O₂ with 6E18 O₂, in the presence and absence of 1E16 methanol. Where the removal of OH by reaction with the HO₂, OH were included, and accounting for the loss of HO₂ via reaction with HO₂, OH, CH₃O and diffusion.

The crucial reaction in attenuating the HO₂ yield in the OH + H₂O₂ reaction is OH + HO₂. According to the literature the HO₂ yield will only be close to unity when [OH] < 1E12 cm⁻³. Also, it is noted that when OH + HO₂ is significantly occurring the OH + H₂O₂ kinetics will be significantly faster than the literature. Based on these predictions we have done further experiments.

By comparing HO₂ yield when CH₃OH(O₂) is present we can assign yields without knowing the absolute radical concentration; it is wholly reasonable to assign the HO₂ yield in the presence of sufficient CH₃OH(O₂) as 100%. Comparing the HO₂ yield from OH and methanol in the presence of high oxygen, to the yield in the absence of methanol allowed for the assignment of HO₂ yields from the reaction of OH with hydrogen peroxide.

The literature predicts a large decrease in the HO₂ yield from hydrogen peroxide as the [OH]₀ is increased, see Fig 1. Below is our yield for HO₂ from OH + H₂O₂ and is compared to when a large amount of CH₃OH(O₂) is added. It is clear that no attenuation of the HO₂ yield is observed in our system. Many other experiments were carried out as [OH]₀ was varied over a factor of ~300, and the HO₂ yield from all the experiments was the same, within error, for OH + H₂O₂ compared to when CH₃OH(O₂) is added. The [H₂O] and [CH₃OH] in the system is too small for significant complexation to HO₂. Our experiments assign yields as 100% (101 ± 7 at 1E11 [OH], 101 ± 2 at 3E12 [OH] and 102 ± 3 at 3E13 [OH]).




Figure 2 HO₂ growth profiles collected with 2.5E13 cm⁻³ [OH], 6E18 cm⁻³ O₂, 2.7E15 H₂O₂ in the presence and absence of 2.5E16 CH₃OH.

Also, the impact of the HO₂ + OH on the observed OH + H₂O₂ rate constant is to make it significantly faster as the initial [OH] is increased. From our literature model, measurable changes in the rate constant should be observed as [OH] is increased >20 % for 1 Hz experiments; in experiments carried out at 10 Hz where there is HO₂ present from the previous laser flash at time zero, this should lead to the observation of an increase in the OH removal rate by up to 50 %. The precision of the system means that we can readily see changes in the rate constant to ~ 1% when initial OH radical concentrations are >1E12 cm⁻³. The results are summarised in the graph below.




Figure 3 Expected and observed OH removal rates with $2.7E15 \text{ cm}^{-3} \text{ H}_2\text{O}_2$ and $1\text{-}60 \text{ mJ cm}^{-2}$ photolysis energy at 248 nm and 10Hz.

Again, this kinetics test versus [OH] demonstrates that under our conditions $\text{HO}_2 + \text{OH}$ is having little impact on the $\text{OH} + \text{H}_2\text{O}_2$ reaction. The measurable increase in the figure 3 (6.8%) can be assigned to $\text{OH} + \text{OH}$ ($\approx 1E\text{-}11 \text{ cm}^3 \text{ s}^{-1}$, at 1600 Torr).

We recognise that our results are in contradiction with the literature rate coefficient for $\text{HO}_2 + \text{OH}$. The IUPAC literature value is $1.1E\text{-}10 \text{ cm}^3 \text{ s}^{-1}$. In order to reconcile our experiments we require this rate coefficient to be $< 1E\text{-}11 \text{ cm}^3 \text{ s}^{-1}$. However, our result is wholly consistent with the previous paper on the reaction of OH with H_2O_2 (Wine et al. 1981 J. Chem. Phys). In Wine et al. the removal kinetics were not perturbed by additional HO_2 added to the system. In this work, with additional $[\text{HO}_2] \sim 1E13$ added, no measurable change in the $\text{OH} + \text{H}_2\text{O}_2$ was observed. This result is in agreement with our present study. We note that Wine et al study used flash photolysis study, as used in our present study. Most literature assignments on $\text{HO} + \text{HO}_2$ were carried out in low pressure, flow tubes; very different conditions. The flash photolysis is less prone to interference.

Comment 2:

“Figure 3 : the black squares are difficult to distinguish from the blue triangle. Better chose other symbols or other colours.”

Response:

In part, that these are hard to distinguish is due to these traces showing no evidence of back diffusion of NO into the region where the OH is probed in the first detection axis. However, we will adjust this figure to highlight the different traces better.

Comment 3:

“Figure 7: Who is who? I guess red is HO₂ and black is OH? What was the reaction system in Figure 7 and what was the estimated initial radical concentration? Because from the above model, one would expect a faster HO₂ decay compared to OH decay if secondary reactions are taken into account (2003 s⁻¹ for OH against 2596 s⁻¹ for HO₂ in the example of the right graph above).”

A legend will provide for clarity in this figure, and the OH concentrations were 1-3 E12 cm⁻³. The experimental detail has now been included in the description.

Please see the response above for comment on observations of secondary reactions.

Comment 4:

Figure 10: what are the different colored symbols? Different experiments? Or is the blue line a fit to different data points?

Response:

The red fit is an exponential fit to the data, the blue fit is a multi-exponential fit that allows for assignment of the returned OH. The three colours of symbols are merely three different time scans of the same experimental conditions to allow for correct assignment of both the fast and slow loss processes. A legend will add clarity to this figure.