

Authors' Reply

The authors would like to thank the referees for their constructive feedback, that helped in improving the manuscript. In the following, all revision points are addressed and the resulting text edits are included in the following way:

The comments are repeated and the responses are given below. Changes made in the manuscript are indicated in blue. Figure numbers with “R” correspond to figures in this reply not included in the manuscript. In the manuscript red indicates removal and blue addition.

Reply to Anonymous Referee #1

Comment: A slightly clearer distinction between total liquid water path, cloud liquid water path and rain liquid water path could be made throughout the paper. I would also like to see some more details of the distinction between cloud and rain liquid water path. This is hinted at on P7 line 17, which implies that it is taken from the ICON microphysics scheme, but it would be helpful to specify the difference in terms of the different size distributions etc.

Response: To clarify the discussion we added the following to the the second paragraph of Sec. 1: “**Thus, we define LWP as the sum of CLWP and RWP.**” We checked throughout the manuscript the consistency and corrected especially Sec. 1 accordingly. Furthermore, it is made clear that the size distributions from ICON microphysics. They are specified in Sec. 2.3, paragraph 2, now. “**Cloud and rain particles are simulated with a 20 μm diameter mono-disperse and exponential distribution of water spheres, respectively. The exponential distribution has its intersect N_0 classically fixed to 0.08 cm^{-4} (Marshall and Palmer, 1948).**”

Comment: Why are all the available microwave channels not used in the retrieval? I would expect that particularly the 183 GHz channels would contain additional information on the IWV, including its vertical distribution, and the quasi-window channels on the far wings of the 118 and 50-60GHz O2 bands will also respond strongly to liquid water. Since the data are already screened for cloud ice then scattering at 183GHz should not be a concern here.

Response: We thank the reviewer for proposing this retrieval extension. It certainly is an interesting objective for a follow-up study, but we chose the frequencies for two main reasons. First, we wanted to use only frequency bands that are currently also used by spaceborne microwave sensors to obtain a comparable product. Second, including additional channels to the LWP retrieval inhibits the calibration crosscheck with the IWV retrieval. This is important to avoid additional bias errors, as the calibration and the absorption characteristics in the O2 band are still uncertain (Maschwitz et al., 2013). Furthermore, the 183 GHz channels unfortunately suffered from hardware instabilities during both campaigns as pointed out by Konow et al. (2018a) and their data is not available all the time.

Comment: I would like to see some further discussion on the impact of surface wind speed (and the minor impact of surface temperature) on the retrieval of LWP and IWV. How does the frequency-variation of the brightness temperatures differ for surface wind speeds compared to that for IWV/LWP/RWP shown in figure 2? Is there any independent information content on the wind speed contained in the radiometer measurements, or does it effectively just add noise to the LWP/IWV retrievals?

Response: The error of the LWP retrieval shows no dependence on the 10 m wind speed as it can be seen in Fig. R1. A slight mean underestimation of the retrieval between 5 and 10 g m^{-2} for calm wind decreases toward wind speed of about 6 m s^{-1} . There seems to be a retrieval overestimation for stronger wind speed above 12 m s^{-1} , but this effect is of minor significance, as wind speed above 12 m s^{-1} occurs only rarely (as determined by ICON).

The IWV retrieval error as a function of wind speed is shown in Fig. R2. A slight systematic dependence of the error on wind speed is notable. However, the largest errors are again related to rare wind speeds.

To address this issue included a variation of the surface wind speed in Fig. 2. It shows a brightness temperature change of up to 2 to 3 K per 5 m s^{-1} wind speed change. The potential to retrieve surface wind speed from this rather weak signal, however, is low. The surface temperature influence of less than 1 K difference for sea surface temperature (sst) variation of 4 K is too weak for any reasonable sst retrieval with the HAMP channels. We added to the 3rd paragraph of Sec. in the manuscript: “**The near surface wind speed slightly alters the BTs through modification of surface reflectivity and emissivity as also shown in Fig. 2.**”

This influence will act as a random source of error to the LWP and IWV retrievals as no independent information to correct for wind influence is available.”




Figure R1. Neural network LWP retrieval error as function of 10 m wind speed.




Figure R2. Left: Neural network IWV retrieval error as function of 10 m wind speed. Right: 10 m wind speed frequency distribution of the test database.

Comment: I find it slightly surprising that there is more liquid water during the dry season than the wet. I would like to see some more discussion about how the results in sec. 6 may be influenced by the choice of flight paths during the two campaigns. If specific conditions were either targeted or avoided then this could significantly bias the results.

Response: Indeed we find the higher LWP during the dry season to be one of our most interesting results. While you are right that the flight patterns could be responsible for part of the difference, also changes in the organization of clouds could cause the differences in cloud fraction and LWP. The fact that the medium LWP range from 100 to 400 gm-2 is less frequent in the wet season could be due to the higher degree of organization causing more clear sky areas and more intense clusters with higher amounts of precipitation. In that sense the latter would be missed by our flight patters as we avoided strongly convective scenes with formation of large ice particles. The aspect of organization is currently investigated by different LES modeling groups and will be a major objective of the EUREC⁴A campaign.

The discussion was added to the paper as follows: “The higher LWP in the dry season might partly be explained by the choice of flight patterns. However, an analysis of ground-based LWP measurements at the Barbados Cloud Observatory (Stevens et al., 2016) over the years 2013-2018 confirms the generally higher LWP values during December than August (not shown). Thus, also changes in the organization of clouds could cause the differences in cloud fraction and LWP. The fact that the medium

- 5 LWP range from 100 g m^{-2} to 400 g m^{-2} is less frequent in the wet season could be due to the higher degree of organization causing more clear sky areas and more intense clusters with higher amounts of precipitation. In that sense the latter would be missed by our flight patterns as we avoided strongly convective scenes with formation of large ice particles.”
- The abstract was extended by “We hypothesize that higher degree of cloud organization on larger scales in the wet season reduces the overall cloud cover and observed LWP.”
- 10 Within the conclusions we added: “An extension of the NARVAL observations is planned by the EUREC⁴A field study in early 2020 (‘elucidate the couplings between clouds, convection and circulation’; Bony et al. (2017)) which among other objectives will investigate convective aggregation. The algorithms presented here will be applied and together with additional measurements a better understanding of the governing processes that cause differences between dry and wet season will be analyzed. For that the campaign will provide additional observations of large scale dynamics, horizontally resolved remote sensing ob-
- 15 servations by a second and in situ observations by additional aircraft in the cumulus layer. Also, more locally targeted flights, distributed over the daytime are planned to study the diurnal cycle. Together with ship, drone and buoy measurements a unique dataset for a better understanding of precipitation onset will be generated.”

- Comment:** The authors note that the WALES IWV measurements are only available in clear sky conditions so they only
- 20 provide validation for the MWR retrievals when there are no confounding effects from liquid water. Is it possible to split the dropsondes into clear and cloudy scenes to demonstrate if there is any impact of liquid water on the quality of the IWV retrievals?

- Response:** The separation between clear sky and cloudy sondes is the reason, why we separate Tab. 2 in “observed pairwise” (all sky) and “observed by all” (WALES available, clear sky). We added the following paragraph to make this more explicit.
- 25 “A small confounding effect from liquid water in cloudy scenes can be derived from the separation of the HAMP – dropsonde comparison into all (‘observed pairwise’) and clear sky (‘observed by all’, i.e., when also WALES is available) in Tab. 2. In the NARVAL1 dataset, the bias for cloudy sky sondes (0.24 kg m^{-2}) is somewhat smaller than that for clear sky (0.32 kg m^{-2}). However, RMSD and STD in cloudy scenes are about 0.3 kg m^{-2} larger than in clear sky. NARVAL2 also shows a larger bias in cloudy sky of about 0.53 kg m^{-2} in comparison to clear sky (0.28 kg m^{-2}). The cloudy sky RMSD and STD of 1.32 kg m^{-2} and
- 30 1.21 kg m^{-2} , respectively, are only slightly larger than their clear sky counterparts. An increase of the random error for cloudy scenes is expected as also higher water vapor variations are expected in heterogeneous cloud fields.”

- Comment:** P6 line 18 Why is there a need to convert from water vapour number density to volume mixing ratio? It is the former that is required to calculate integrated water vapour mass.
- 35 **Response:** Indeed, you are right. We note this conversion from water vapor density to water vapor mixing ratio because we want to show mixing ratio in Fig. 4.

- Comment:** P8 figure 2 I suggest using a logarithmic colour scale to show the relative frequency to highlight any detail away from the strong “clear sky” line
- 40 **Response:** We followed the suggestion and changed the color scale of Fig. 3, that was on page 8 of the previous manuscript version and shows BT frequency distribution.

- Comment:** P9 line 31 Biases with respect to what?
- Response:** The biases are related to the deviation of LWP from 0 during clear sky scenes. We adjusted the sentence accordingly. “When retrieval algorithms are applied to HALO measurements, slight biases of LWP from 0 with slow changes over
- 45 time are observed during clear sky scenes.”

- Comment:** It would be useful to have an indication of along-track distance on figure 4 rather than just time
- Response:** We followed the suggestion and updated Fig. 4 as well as Fig. 10 accordingly.
- Comment:** P13 final paragraph – it would be nice to refer to fig 6 early in this discussion.
- 5 **Response:** We added an earlier reference.
- Comment:** P14 discusses the impact of negative LWP retrieval values on the bias. These could be avoided by performing the retrieval in logarithmic space (I.e. retrieving $\log(\text{LWP})$). Would this have a significant impact on the results?
- Response:** Retrieving $\log(\text{LWP})$ would avoid negative values, but it would inevitably result in a overestimation of the retrieved mean LWP as noise is always positive. Another reason for not retrieving the logarithm is, that the natural first order approximation relation between brightness temperatures and LWP is linear.
- 10 **Comment:** In figure 10 it might be clearer to plot the LWP and RWP on a logarithmic scale – in the current plot it is hard to see the cloud LWP retrieved by the MWR between 17:38:30 and 17:39:10 that is discussed in the text at the end of page 17
- 15 **Response:** We changed Fig. 10 by using a piecewise linear scale now, that magnifies the range from -20 to 20 g m^{-2} . We decided against a logarithmic scale as this can not display negative retrieval artifacts.
- Comment:** The paper is clearly written and generally easy to follow, although I find a number of sentences do not read well and should be redrafted. There are also a few typos: [...]
- 20 **Response:** All remaining minor corrections by the referee were agreed on and are incorporated in the revised manuscript.
- Comment:** P1 line 1 "...identified especially marine low level clouds to play a critical role for the climate."
- Response:** We removed the first sentence on recommendation from referee #2.
- 25 **Comment:** P1 line 5 "...to better understand *the* LWP of warm clouds..."
- Response:** "the" is added.
- Comment:** P2 line 5 "Especially, shallow marine clouds are attributed to contribute largely to intermodel spread of climate models"
- 30 **Response:** We redrafted this sentence to: "[Sherwood et al. \(2014\)](#) attribute especially shallow marine clouds to contribute largely to intermodel spread of climate models."
- Comment:** P2 line 28 "Visible/near infrared techniques *such* as those applied to MODIS..."
- Response:** "such" is added.
- 35 **Comment:** P3 line 9 "...allow to study clouds with similar, however, more sensitive and higher spatially resolving instrumentation than available on satellites." Perhaps "...allow the study of clouds with similar, but more sensitive and higher spatially resolving, instruments to these available on satellites."
- Response:** We thankfully incorporated this comment.
- 40 **Comment:** P3 line 13 "Their study shows *the* sub-footprint variability of spaceborne Special Sensor Microwave Imager/Sounder..."
- Response:** We added "the".
- 45 **Comment:** P3 line 24 "The assessment of LWP (Sec. 4) reveals the importance *of using* ancillary measurements, e.g. lidar measurements for low LWP values and cloud radar measurements for lightly precipitating cases."
- Response:** We thankfully incorporated this comment.

- Comment:** P3 line 26 ". . . between dry and wet *seasons*"
Response: Changed "season" to "[seasons](#)".
- Comment:** P4 figure 1 I think the caption mis-labels the thin and thick lines based on the dates in the legend (I.e. NARVAL
5 1 looks like it should be the thick lines)
Response: Corrected.
- Comment:** P5 line 12 "more dominant in the higher *frequency* window channels"
Response: Corrected "window" to "[frequency](#)".
- 10 **Comment:** P5 line 16 remove the comma after "both"
Response: Removed comma.
- Comment:** P6 line 9 "cloudy conditions as *well* as possible"
15 **Response:** Corrected "good" to "[well](#)".
- Comment:** P6 line 25 "...prevent having data during some flights..."
Response: We redrafted that sentence to: "[While all research flights during NARVAL1 can be used, no data is available for some NARVAL2 flight days due to hardware issues as summarized in Tab. 1.](#)"
- 20 **Comment:** P7 line 10 is ambiguous. Are all profiles with ice excluded, or only ones with ice water path above 1000 g/m² ?
Response: We reordered that sentence for better understanding. "[In general, the training and test data excludes cases with LWP greater than 1000 g m⁻² and cases with ice, i.e., 86 % of all profiles over the ocean are used.](#)"
- 25 **Comment:** P7 line 11 "...the ocean *are* used"
Response: Corrected "is" to "[are](#)".
- Comment:** P8 line 3 "...are *visible* as a line..."
Response: Corrected "visibly" to "[visible](#)".
- 30 **Comment:** P9 line 11 "...never deviates *more* than..."
Response: Added the missing "[more](#)".
- Comment:** P14 line 7 "...decided *to use* a retrieval"
35 **Response:** Replaced "we decided for" by "[we chose](#)". Further we slightly changed the beginning of the following sentence to: "[Instead to include clear sky directly in the retrieval, we make use of lidar measurements, which are better suited than MWR for cloud masking.](#)"
- Comment:** P21 line 11 "...consistent *with*"
40 **Response:** Replaced "to" by "[with](#)".

Reply to Anonymous Referee #2

Comment: I think a bit more discussion of the science possible with the integration of the radar and Lidar datasets, and of the radar data itself would enhance the scientific impact of the manuscript, but these constitute minor comments, incorporated into those provided below.

- 5 **Response:** We added an outlook to combine the results of the present study with radar and lidar datasets to study the presence and condensate loads of different shallow trade wind cumulus types. We added to Sec. 7: “With respect to trade wind cumuli, the products of the present study in combination with cloud boundary estimations from the radar and backscatter lidar will be used to evaluate the condensate loads of different shallow trade wind cumulus types in large eddy simulations. For example, radar and lidar both detect shallow convection or shallow outflow anvils as depicted in Fig. 10. In addition, the lidar also allows
10 detecting boundary layer driven clouds, which have tops around 1 km and are below the radar sensitivity.”

Comment: Also, while I am not sure of the Copernicus standards, I would recommend that DOIs be generated for the datasets and included within the manuscript.

- 15 **Response:** The DOI assignment was in preparation and is completed now. DOIs are added to the references.

Comment: p.2 line 14: also mention the clear-sky contribution to the field of view (it is mentioned later but the sentence suggest precip is the major error source).

- 20 **Response:** We added two sentences to that paragraph to address this point. “Furthermore, the observed LWP per se is an average over the sensors field of view, which is affected by cloud and rain inhomogeneity, and clear sky contribution. Therefore, the spatial resolution is a key information to interpret LWP statistics.”

Comment: p. 2 Line 27: what do Greenwald et al and other conclude for the tropical Atlantic region you are interested in?

- 25 **Response:** It is difficult to give any quantitative estimate from the coarse figures provided by Greenwald et al. (2018) for the North Atlantic tropical region. However, our study region in the tropics behaves close to the global average conditions. Nevertheless, from Elsaesser et al. (2017) we added two values. “Elsaesser et al. (2017) additionally estimate the contribution RWP to the total LWP by a simple parametrization and recommend to only use those values with a ratio RWP:LWP of less than 0.2. The average MAC RWP:LWP ratio in our area of interest is 0.23 and 0.30 in December 2013 and August 2016, respectively. Therefore, a more detailed assessment of the rain cloud partitioning is important to better interpret satellite measurements in our study area.”

30 **Comment:** P. 3. some where I think the adiabatic constraint on LWP is worth mentioning. Is it possible to construct an adiabatic estimate from the Lidar cloud top height and dropsonde RH-derived cloud base do you think? This is an earnest question - I am not sure how well this would work. But it would provide an additional constraint on the retrieval that might be more physical than the imposed 1000 g/m^2 (and its relaxation), and could also provide some additional physical insights. For example, in clean marine stratocumulus regions, the adiabatic constraint on LWP seems to hold well until about 200 g/m^2 , at which point precipitation begins to deplete LWP (Zuidema et al., 2005, fig. 8 and 9). I think during RICO the adiabaticity deviated more quickly from the theoretical maximum because of mixing with environment air (Rauber et al., 2007). Related to this I do not see any discussion on the radar or cloud top height at which precipitation becomes discernible later on in the manuscript - perhaps I missed it.

- 35 **Response:** Thanks for these interesting thoughts. However, two aspects limit the applicability of the adiabatic theory in our case. First, the adiabatic assumption requires, that the cloud develops through vertical transport, i.e. is buoyancy driven, and without horizontal exchange. This is more realistic in stratocumulus situations as addressed by Zuidema et al. (2005) than in the trade wind cumuli cases as in our study. Quite often we see clouds in the radar, that are not buoyancy driven which is also the case in the example of Fig. 10. The radar echo between 17:40:00 and 17:41:40 looks more like a shallow outflow of the nearby precipitating core. This shallow outflow anvil is not buoyancy driven, as its radar echo shows no link to the lifting condensation level (lcl), which is roughly at 700 m. Second, the vertical cloud extend Δz is very important as the adiabatic water content is proportional to Δz^2 , but the exact estimation of Δz is difficult. The estimation of the cloud base height using the lcl derived from near surface dropsonde data has an accuracy of 50 to 100 m. The cloud top height estimation of shallow
45

cumulus adds additional uncertainty as the lidar sees the cloud top typically 250 m higher in altitude than the radar, in cases in which the radar sees a cloud at all. Thus we think, that a comparison to a somehow estimated adiabatic LWP raises more methodological questions than it would help to constrain the LWP.

5 **Comment:** P. 3 lines 11-19: what is approximately the spatial footprint of the HAMP instrumentation? It would be nice to see this number in relation to the satellite spatial footprints. On p. 6 you mention that the different footprints and sensitivities of the instruments are covered in Stevens et al 2019, but a brief summary here would be useful.

Response: The reference to Schnitt et al. (2017) reads now: “[Their study uses the 1 km resolution HAMP data to show the sub-footprint variability of spaceborne CLWP estimation of about 30 km resolution. Further they illustrate how MODIS products at 1 km resolution likely underestimate CLWP of thick clouds due to MODIS’ sensitivity towards the upper part of the cloud.](#)”

Comment: P. 7 line 23: how is scattering off of the ocean surface dealt with?

Response: We understand the “scattering off of the ocean” as surface reflection. Surface reflectivity is also calculated using FASTEM5. The related sentence in Sec. 2.3 reads now: “[The emissivity and reflectivity of the sea ocean surface is calculated by the FAST microwave Emissivity Model version 5 \(FASTEM5; Liu et al. \(2011\)\), which is a modification of the Fresnel coefficients including corrections for ocean surface roughness and foam building as a function of wind speed.](#)”

Comment: p.9 line 24: where is the ocean emissivity represented? It would be nice to see a bit more discussion of the ocean surface microwave radiation characteristics in general. A figure of the emission/scattering as a function of SST and wind speed, for the 2 frequencies would be nice, for example. How much does error in the surface characterization contribute to the overall error?

Response: Emission by the surface is only implicitly included in the retrieval. When generating the database, ocean emissivity is calculated by FASTEM5 using sea surface temperature (sst) and 10 m wind speed as input. Based on a comment by referee #1 we investigated the uncertainty due to sst and 10 m wind speed. (See Fig. R1 and R2 and discussion in our answer to referee #1.)

Comment: p.13 line 4: Is there any cloud fraction within a model column? At a grid spacing of 0.5 degree, clouds will not necessarily fill the full grid box.

Response: This is probably a misunderstanding. The ICON simulations were run at 1.25 km resolution. Afterwards, the data used for the retrieval database was coarse grained to 0.5° as explained in the first paragraph of Sec. 2.3.

Comment: p. 17 : it would be nice to see the retrieved LWP/RWP as a function of the vertically integrated reflectivity from both campaigns as part of fig. 11. They should look the same, if not, that may tell you something about the cloud droplet number concentration variation between the two seasons.

Response: The RWP retrieval already includes the vertically integrated reflectivity Z_{int} . Therefore, the empirical relation between RWP and Z_{int} would be established from two dependent variables. Each of our retrievals was trained for each campaign individually by using simulations for the respective period. Thus, differences in the RWP- Z_{int} relation would also represent the different training datasets.

Technically, we have to exclude scenes with no radar echo above noise level because $Z_{\text{int}} = 0$ can not be represented on a decibel scale from such comparison. A logarithmic scale is required for displaying in analogy to dBZ. This means that such a figure excludes clouds, that were too thin to be detectable by the radar but were detected by the lidar. Nevertheless, we prepared Fig. R3 showing the relation between LWP and Z_{int} for scenes, where $10\log_{10}(Z_{\text{int}}) > -30$. The scatter plot shows, that there are less scenes with $\text{LWP} > 300 \text{ g m}^{-2}$ during NARVAL1 than NARVAL2 as it can also be seen on Fig. 11. For NARVAL1, there is a pronounced maximum of combinations for Z_{int} from 15 dB at 100 g m^{-2} increasing to 40 dB at $\text{LWP} > 400 \text{ g m}^{-2}$. A similar relation can be seen also for NARVAL2. In addition there is a second mode for $\text{LWP} < 200 \text{ g m}^{-2}$ with Z_{int} being about 0 dB. Scenes with lower Z_{int} most likely consist of smaller droplets for the same LWP as $Z_{\text{int}} \propto D^6 \Delta z$ and $\text{LWP} \propto D^3 \Delta z$. So, probably clouds with smaller droplets were slightly more prominent during NARVAL1 than NARVAL2.

Figure R3 is an interesting starting point for a microphysical study. However, to present this topic in an appropriate manner more work has to be done, which will be included on a follow-up study.




Figure R3. Decibel of vertically integrated reflectivity ($10\log_{10}(Z_{\text{int}})$) versus LWP during (left) NARVAL1 and (right) NARVAL2.

Comment: How does RWP spatial heterogeneity affect the retrieval do you think?

Response: The spatial heterogeneity of rain affects the airborne HAMP measurements less than microwave satellites as the HAMPs spatial resolution is at least an order of magnitude better than satellite resolution. To illustrate the scale on that HAMP resolves precipitation, we added a km scale to Fig. 10. Also we added to the fourth paragraph of Sec. 5: “The figure shows how HAMP is able to resolve spatial features of showering cells, which were observed with a cross section of several HAMP footprints.”

Comment: P. 18: how does WVP vary in this example?

Response: See Fig. R4. We added the summary “the IWV varies around $31.5 \pm 1.5 \text{ kg m}^{-2}$ in this scene” to the Fig. 10 caption in the manuscript, as the main aspect of the figure is the liquid phase and IWV variation is only of secondary interest in that example.




Figure R4. As Fig. 10 but with additional time series of IWV.

Comment: p. 19, lines 17-18: Some discussion of the sampling of the diurnal cycle (I presume HALO only flew during the day, were cumuli more prevalent in the afternoon?), and how that might alias into the results from the 2 seasons would be nice. I presume the BCO LWP measurements mentioned are diurnal averages

Response: Yes you are right, we added that “flights were scheduled during local daytime” to the second paragraph of Sec. 2.1. In the conclusions we added that “sound conclusions on the diurnal cycle can not be drawn from the data presented here, as the spatial variability of the clouds on the observed mesoscale was higher than an expected effect of the diurnal cycle.” Radar time-height-plots of the NARVAL1 flights that were flown from Barbados to the East and back are rather symmetrical to the return point due to large-scale patterns. This means, a potentially diurnal cycle during the flights is overlaid by the changes

in the larger scale cloud field. Additionally, we added in the outlook the following with respect to EUREC4A: “Also, more locally targeted flights, distributed over the daytime are planed to study the diurnal cycle.”

Comment: P.11: an easy additional plot would be how LWP and RWP vary with lidar-derived cloud top height. This would be of scientific interest. How would that compare to, e.g., Byers and Hall, 1955?

Response: Indeed, liquid condensate load versus cloud extend is an interesting comparison. Figure R5 depicts the relations during the two campaigns. The overall impression of increasing rain amount with increasing cloud top height during NARVAL1 agrees with the findings by Byers and Hall (1955). The dry winter season during the NARVAL1 campaign compares best to their pioneer study according to their description. Differences exist in details and but are also partially due to the analysis approach. Byers and Hall subjectively identified cloud objects while we analyze the data profile-wise. For example, Byers and Hall (1955) found the lowest cloud top of a precipitating cloud near 1.8 km, whereas we already observed $RWP > 10 \text{ g m}^{-2}$ for cloud top heights near 1.0 km. Stevens et al. (2019, in press) present in Fig. 7 a cloud object oriented analysis of NARVAL, that directly uses radar reflectivity instead of the RWP retrieval presented in our manuscript. A more detailed analysis of cloud dimensions in relation to their LWP and RWP will follow in a subsequent study, which is in preparation at the moment.




Figure R5. Lidar (WALES) derived cloud top height in relation to LWP (left) and RWP (right) for both campaigns.

Comment: P. 23, data availability: do the datasets have dois? They should.

Response: The DOI assignment was in preparation and is completed now. DOIs are added in the references.

Comment: The writing overall is fine, but there are small awkward uses of the English language sprinkled throughout that reflect English as a second language. If it is possible to find a native English speaker to read it that would polish the manuscript.

Response: Referee #1 pointed out problematic sentences and typos that are remedied now.

Comment: In particular the abstract and its first sentence needs a revisit (you could consider just removing the first sentence).

Other comments on the abstract: mention the frequencies you use. You don't mention the linear regression approach, is that intentional? Mention clear-sky frequency and LWP statistics, as opposed to focusing on IWV - the title only mentions LWP after all. Overall the abstract seems to have been written in a hurry.

Response: We removed the first sentence from the abstract as suggested and rearranged most parts of the abstract for improved comprehensibility. We incorporated the radiometer frequencies into the abstract. We don't mention the linear regression approach on purpose, as it is mainly used as reference to classical retrievals. Averages for LWP, RWP and cloudiness were added and the sentence order was rearranged such that it has the dry and wet season in a consistent order. Further, we reformulated the closing two sentences of the abstract after the discussion of the flight patterns question by referee #1 and reconsidering that issue. We conclude that our former formulation was too negative. The revised abstract reads now:

“Liquid water path (LWP) is an important quantity to characterize clouds. Passive microwave satellite sensors provide the most direct estimate on global scale, but suffer from high uncertainties due to large footprints and the superposition of cloud

and precipitation signals. Here, we use high spatial resolution airborne microwave radiometer (MWR) measurements together with cloud radar and lidar observations to better understand the LWP of warm clouds over the tropical North Atlantic. The nadir measurements were taken by the German High Altitude and Long range research aircraft (HALO) in December 2013 (dry season) and August 2016 (wet season) during two Next generation Advanced Remote sensing for VALidation (NARVAL) campaigns.

Microwave retrievals of integrated water vapor (IWV), LWP and rain water path (RWP) are developed using artificial neural network techniques. A retrieval database is created using unique cloud-resolving simulations with 1.25 km grid spacing. The IWV and LWP retrievals share the same eight MWR frequency channels in the range from 22 GHz to 31 GHz and at 90 GHz as their sole input. The RWP retrieval combines active and passive microwave observations and is able to detect drizzle and light precipitation. The comparison of retrieved IWV with coincident dropsondes and water vapor lidar measurements shows root-mean-square deviations below 1.4 kg m^{-2} over the range from 20 kg m^{-2} to 60 kg m^{-2} . This comparison raises the confidence in LWP retrievals which can only be assessed theoretically. The theoretical analysis shows that the LWP error is constant with 20 g m^{-2} for LWP below 100 g m^{-2} . While the absolute LWP error increases with increasing LWP, the relative one decreases from 20 % at 100 g m^{-2} to 10 % at 500 g m^{-2} . The identification of clear sky scenes by ancillary measurements, here backscatter lidar, is crucial for thin clouds ($\text{LWP} < 12 \text{ g m}^{-2}$) as the microwave retrieved LWP uncertainty is higher than 100 %.

The analysis of both campaigns reveals that clouds were more frequent (47 % vs. 30 % of the time) in the dry than in the wet season. Their average LWP (63 g m^{-2} vs. 40 g m^{-2}) and RWP (6.7 g m^{-2} vs. 2.7 g m^{-2}) were higher as well. Microwave scattering of ice, however, was observed less frequently in the dry season (0.5 % vs. 1.6 % of the time). We hypothesize that higher degree of cloud organization on larger scales in the wet season reduces the overall cloud cover and observed LWP. As to be expected, the observed IWV clearly shows that the dry season is on average less humid than the wet season (28 kg m^{-2} vs. 41 kg m^{-2}). The results reveal that the observed frequency distributions of IWV are substantially affected by the choice of the flight pattern. This should be kept in mind when using the airborne observations to carefully mediate between long-term ground-based and spaceborne measurements to draw statistically sound conclusions. ”

Other minor corrections

We corrected two transfer errors of one value in Sec. 6 and one value in Tab. 3.

Investigating the liquid water path over the tropical Atlantic with synergistic airborne measurements

Marek Jacob¹, Felix Ament², Manuel Gutleben³, Heike Konow², Mario Mech¹, Martin Wirth³, and Susanne Crewell¹

¹Institute for Geophysics and Meteorology, University of Cologne, Albertus-Magnus-Platz, 50923 Cologne, Germany

²Meteorological Institute, Universität Hamburg, Bundesstrasse 55, 20146 Hamburg, Germany

³German Aerospace Center, Münchener Str. 20, 82234 Oberpfaffenhofen-Wessling, Germany

Correspondence: Marek Jacob (marek.jacob@uni-koeln.de)

Abstract. ~~Clouds are a strongly variable component of the climate system and several studies have identified especially marine low-level clouds to play a critical role for the climate.~~ Liquid water path (LWP) is an important quantity to characterize clouds. Passive microwave satellite sensors provide the most direct estimate on global scale, but suffer from high uncertainties due to large footprints and the superposition of cloud and precipitation signals. Here, we use high spatial resolution airborne microwave radiometer (MWR) measurements together with cloud radar and lidar observations to better understand the LWP of warm clouds over the tropical North Atlantic. The nadir measurements were taken by the German High Altitude and Long range research aircraft (HALO) in December 2013 (dry season) and August 2016 (wet season) during two Next generation Advanced Remote sensing for VALidation ~~campaigns~~ (NARVAL) campaigns.

Microwave retrievals of integrated water vapor (IWV), LWP and rain water path (RWP) are developed using artificial neural network techniques ~~and a unique database based on.~~ A retrieval database is created using unique cloud-resolving ~~model~~ simulations with 1.25 km grid spacing. The IWV and LWP retrievals share the same eight MWR frequency channels in the range from 22 GHz to 31 GHz and at 90 GHz as their sole input. The RWP retrieval combines active and passive microwave observations and is able to detect drizzle and light precipitation. The comparison of retrieved IWV with coincident dropsondes and water vapor lidar measurements shows root-mean-square deviations below 1.4 kg m^{-2} over the range from 20 kg m^{-2} to 60 kg m^{-2} .

This comparison raises the confidence in LWP retrievals which can only be assessed theoretically. The theoretical analysis shows ~~the dependency of the uncertainty on LWP itself as the error is below that the LWP error is constant with~~ 20 g m⁻² for LWP below 100 g m⁻² and below 20 % above. While the absolute LWP error increases with increasing LWP, the relative one decreases from 20 % at 100 g m⁻² to 10 % at 500 g m⁻². The identification of clear sky scenes by ancillary measurements, here backscatter lidar, is crucial for thin clouds ($\text{LWP} < 12 \text{ g m}^{-2}$) as the microwave retrieved LWP uncertainty is higher than 100 %.

~~The RWP retrieval combines active and passive microwave observations and is able to detect drizzle and light precipitation.~~

The analysis of both campaigns reveals that clouds were more frequent (47 % vs. 30 % of the time) in the dry than in the wet season ~~and their LWP and RWP were higher, but microwave.~~ Their average LWP (63 g m⁻² vs. 40 g m⁻²) and RWP (6.7 g m⁻² vs. 2.7 g m⁻²) were higher as well. Microwave scattering of ice ~~was observed more,~~ however, was observed less frequently in the ~~wet season (1.6 % vs. 0.5 %~~ dry season (0.5 % vs. 1.6 % of the time). We hypothesize that higher degree of cloud organization on larger scales in the wet season reduces the overall cloud cover and observed LWP. As to be expected,

the observed IWV clearly shows that the ~~wet season ($\overline{IWV} = 41 \text{ kg m}^{-2}$) is more~~ dry season is on average less humid than the ~~dry season ($\overline{IWV} = 28 \text{ kg m}^{-2}$)~~ wet season (28 kg m^{-2} vs. 41 kg m^{-2}). The results reveal that the observed frequency distributions of IWV are ~~strongly~~ substantially affected by the choice of the flight pattern. ~~Therefore, This should be kept in mind when~~ using the airborne observations ~~need to be used carefully to~~ carefully mediate between long-term ground-based and spaceborne measurements to draw statistically sound conclusions.

1 Introduction

Clouds and precipitation are a fundamental part of the Earth's climate system and significantly contribute to the water and energy cycle. However, the great variability of clouds, the complex interaction of small-scale processes involved, and their coupling to atmospheric circulation make them a major source of uncertainty in numerical climate and weather models (e.g., Bony et al. (2015), Boucher et al. (2013)). ~~Especially, Sherwood et al. (2014) attribute especially~~ shallow marine clouds ~~are attributed~~ to contribute largely to intermodel spread of climate models ~~(Sherwood et al., 2014). These~~. Such clouds are particularly difficult to assess from spaceborne sensors due to their small size, with about 70 % appearing in sizes of less than 2 km over the tropical North Atlantic (Schnitt et al., 2017). The accurate observation of thin liquid clouds is an ongoing and important challenge as they cover more than a quarter of the globe and are an important contribution to Earth's energy balance (Turner et al., 2007).

Liquid water content (LWC) is the key parameter to describe clouds in atmospheric models. Due to the even higher difficulty in observing LWC profiles (Crewell et al., 2009), we focus on the liquid water path (LWP). It describes the total mass of all liquid water droplets in an atmospheric column above a unit surface area. However, care has to be taken whether LWP only denotes the contribution by cloud droplets, later on called CLWP, or whether it also includes the contribution by liquid precipitation, i.e., drizzle and rain drops (RWP). Thus, we define LWP as the sum of CLWP and RWP. Furthermore, the observed LWP per se is an average over the sensors field of view, which is affected by cloud and rain inhomogeneity, and clear sky contribution. Therefore, the spatial resolution is a key information to interpret LWP statistics.

Few global (C)LWP datasets exist and differences in global mean (C)LWP of a factor of two are reported by Lohmann and Neubauer (2018). These findings reflect the different sensing principles, i.e., microwave radiometry and visible/near-infrared techniques. Satellite microwave imagers such as the Special Sensor Microwave Imager (SSM/I) provide ~~LWP-CLWP~~ estimates for several decades but are limited to the ice free oceans where the background signal is low. (C)LWP is mainly derived from the thermal emission signal in window regions with low water vapor contribution. Microwave receivers also sense rain water within the satellite footprint which can be as large as several tens of kilometer. Recently, the Multisensor Advanced Climatology of Liquid Water Path (MAC-LWP; Elsaesser et al. (2017)) covering the period 1988 to 2016 has been generated. Elsaesser et al. (2017) additionally estimate the contribution of ~~rain-water-path (RWP)~~ RWP to the total LWP by a simple parametrization and recommend to only use those values with a ratio RWP:LWP of less than 0.2. The average MAC RWP:LWP ratio in our area of interest is 0.23 and 0.30 in December 2013 and August 2016, respectively. Therefore, a more detailed assessment of the rain cloud partitioning is important to better interpret satellite measurements in our study area. Greenwald et al. (2018) evaluate

MAC-LWP using measurements by the Moderate Imaging Spectroradiometer (MODIS), the CloudSat Profiling Radar (CPR; Stephens et al. (2002)) and the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP; Winker et al. (2007)). They found in some cases a net LWP bias of more than 50 percent of the mean CLWP due to the combined effects of the in-cloud and adjacent precipitation biases as well as the cloud-rain partition.

5 Visible/near infrared techniques such as those applied to MODIS exploit the spectral response of reflected sun light to derive ~~LWP~~CLWP from optical depth and effective radius retrievals and are therefore limited to daytime. As the signal mainly relates to the upper part of the cloud, assumptions of the cloud vertical structure introduce uncertainties (Zhou et al., 2016). The horizontal MODIS resolution of about one kilometer is much better than those of microwave satellites. Therefore, MODIS data also have been used to assess the clear sky bias of microwave retrievals (Greenwald et al., 2018), to combine them with
10 microwaves for a better assessment of low clouds (Masunaga et al., 2002), and to detect the ratio of rain and cloud water in low-latitude shallow marine clouds via combination with CPR (Lebsock et al., 2011). In summary, quantifying the accuracy of ~~LWP~~CLWP and RWP observations is a major challenge as no absolute reference exists. While shipborne microwave observations have potential for satellite ~~LWP~~CLWP evaluation (Painemal et al., 2016), they fail during precipitation events, due to a wet radome.

15 In this study, we use the Next generation Advanced Remote sensing for VALidation studies (NARVAL; Stevens et al. (2019, in press)) expeditions for investigating LWP and its uncertainty ~~in-over~~ the tropical North Atlantic. The NARVAL missions aim at improving the understanding of clouds, their role for the distribution of water in the atmosphere, and their interaction with the environment (Bony et al., 2015). Within NARVAL, the German High Altitude and LOng range research aircraft (HALO; Krautstrunk and Giez (2012)) was configured as an airborne cloud observatory combining active and passive
20 microwave instruments with water vapor lidar, solar reflectance measurements and dropsondes. Measurements taken during two campaigns in December 2013 (dry season) and in August 2016 (wet seasons) allow ~~to study the study of~~ clouds with similar, ~~however,~~ but more sensitive and higher spatially resolving instrumentation than available on satellites.

Schnitt et al. (2017) demonstrate the ability of the HALO NARVAL 2013 instrumentation to characterize shallow clouds ~~in-over~~ the tropical North Atlantic in terms of size, integrated water vapor (IWV), ~~LWP~~CLWP, and surface reaching precipitation using classical regression algorithms. Their study ~~shows-uses the 1 km resolution HAMP data to show the~~ sub-footprint variability of spaceborne ~~Special Sensor Microwave Imager/Sounder (SSMIS) and illustrates~~ CLWP estimation of about 30 km resolution. Further they illustrate how MODIS products ~~likely underestimate LWP at 1 km resolution likely underestimate CLWP~~ of thick clouds due to ~~its~~ MODIS' sensitivity towards the upper part of the cloud. In this study, we refine the (C)LWP retrieval by making use of high resolution simulations that start to resolve cloud scale circulations and were
30 performed over the full tropical North Atlantic with the ICON (ICOsahedral Non-hydrostatic) weather model to support the analysis (Klocke et al., 2017). We further assess the ~~LWP-total-LWP~~ retrieval accuracy over a wide range of cases, extend the retrieval towards a separation of rain and clouds, and reanalyze the dry season measurements in relation to the wet season campaign.

First, we aim to provide an accurate LWP dataset including uncertainty estimates to support the NARVAL overall goals. For
35 this purpose, we develop retrieval algorithms using multi-channel microwave radiometer measurements as input for LWP and

- based on the similar principle - IWV. The novel cloud resolving ICON simulations serve as a training dataset (Sec. 2). In contrast to LWP, IWV can be evaluated using simultaneous measurements by dropsondes and water vapor lidar. The evaluation is presented in Sec. 3. The assessment of LWP (Sec. 4) reveals the importance ~~to use of using~~ ancillary measurements, ~~i.e., lidar measurements~~ e.g. lidar measurements, for low LWP values and cloud radar measurements for lightly precipitating cases. For the latter an RWP retrieval is developed and assessed (Sec. 5). Finally, the campaign data are analyzed to investigate differences between dry and wet ~~season~~ seasons (Sec. 6).

2 Material and methods

This section presents the data and methods used in this study. That includes an introduction to the two NARVAL campaigns and the relevant measurements that were conducted during both campaigns. Furthermore, the generation of the retrieval database and the subsequent retrieval development are presented.

2.1 Campaign overview

During the NARVAL expeditions HALO was operated out of Grantley Adams International Airport on Barbados to observe trade wind cumuli and their environment over the tropical North Atlantic (Fig. 1). Different flight patterns were chosen to perform satellite underflights, survey the area, probe the environment of a tropical cyclone, and to determine the large scale vertical motion by launching several dropsondes within circles of approx. 170 km diameter (Bony and Stevens, 2019). In total eight research flights were performed during NARVAL1-South in December 2013 and ten flights during NARVAL2 in August 2016. NARVAL1 also included research flights in the northern sector of the Atlantic which are not considered here. For simplicity we refer to the southern part as NARVAL1 in the following.

Flights were scheduled during local daytime. Flight altitudes varied between 6.4 km and 15.0 km with an average speed above ground of 237 m s^{-1} and 207 m s^{-1} during NARVAL1 and NARVAL2, respectively. All further analyses refer to the area from 37°W to 60°W and 7°N to 20°N . A detailed description of the different research flights can be found in Klepp et al. (2014) for NARVAL1 and Stevens et al. (2019, in press) for NARVAL2.

2.2 Measurements

The microwave radiometer (MWR) being part of the HALO Microwave Package (HAMP; Mech et al. (2014)) provides the key measurements for this study. HAMP was installed in a belly pod below the HALO fuselage in nadir looking configuration. While the 26-channel MWR includes channels from 22 GHz to 195 GHz, we only use the seven K-band channels (22.24 GHz to 31.40 GHz) and the 90 GHz channel to retrieve LWP or IWV in the present work. At these frequencies ice particles do not influence the microwave signal substantially with the exception of precipitation sized particles.

As we focus on warm clouds only, cases of ice precipitation are filtered using the differential response of two frequencies along the 60 GHz and 118 GHz oxygen lines. The channels at 53.75 GHz and $118.75 \pm 1.4 \text{ GHz}$ have similar temperature weighting functions but the higher frequency is more affected by ice scattering. The difference between a moving median of




Figure 1. NARVAL1 (thin-thick lines) and NARVAL2 (thick-thin lines) flight patterns. The study area of interest is depicted by sub-sampled ICON LWP from August 19, 2016 14:00 UTC. A grid point is shown every 0.5° as present in the retrieval database.

differential brightness temperature (BT) to the instantaneous differential BT is used to define the "ice flag". This procedure flags 1.2 % of the measurements of both campaigns.

Both liquid water and water vapor emit microwave radiation across the full microwave spectrum albeit with different spectral sensitivity (Fig. 2). BTs around the 22.235 GHz water vapor rotational line increase with increasing water vapor. The effect is strongest at the line center and decreases along the pressure broadened wing of the absorption line. However, due to water vapor continuum absorption, BTs at window frequencies near 30 GHz and 90 GHz are still affected. In contrast, the influence of liquid water is more dominant in the higher frequency window channels than in absorption channels due to increasing emission with frequency. This can be best seen under low humidity conditions by the increasing BT with increasing frequency. The near surface wind speed slightly alters the BTs through modification of surface reflectivity and emissivity as also shown in Fig. 2. This influence will act as a random source of error on the LWP and IWV retrievals as no independent information to correct for wind influence is available.

Figure 2 illustrates the difficulty of retrieving LWP and IWV as in certain channels (e.g., 90 GHz) it is indistinguishable whether BT changes result from changes in IWV or LWP. Therefore, a combination of at least two channels is needed for retrieving IWV or LWP. Note that measurement errors in any of the channels affect both I-IWV and LWP retrievals (Crewell and Löhnert, 2003). This means that a good retrieval of either IWV or LWP indicates a good retrieval capability of the other. Thus, an accurate IWV retrieval is a prerequisite of a good LWP retrieval. Note that in most LWP retrievals (e.g., Wentz and Meissner (2000)) the liquid is assumed to consist of cloud droplets only and therefore bulk approaches to calculate the liquid water absorption coefficients are used. However, for rain drops the Rayleigh approximation is not valid anymore and Mie effects need to be considered, though the discrimination of the cloud and rain signal using MWRs is difficult.




Figure 2. Sensitivity of brightness temperatures in the K-band and around 90 GHz to integrated water vapor (IWV), cloud liquid (CLWP), and rain (RWP) water path and 10 m surface wind (u). Dashed red, dotted blue and solid yellow lines show variations in IWV and LWP (CLWP or RWP) and u , respectively. Bandpasses of the HALO Microwave Package (HAMP) channels are indicated by gray bars. Calculations are based on a thermodynamic dropsonde profile and a synthetic cloud in nadir geometry above the ocean.

The HAMP MWR measures BT with one second integration time and a noise level of less than 0.5 K in the considered channels. Despite ground calibration using hot/cold targets on the air field, BT offsets were identified by comparison with forward simulated dropsondes. Flight dependent corrections were developed (Konow et al., 2018a) and corrected BTs are available in the Climate and Environmental Retrieval and Archive (CERA) (Konow et al., 2018b, c).

- 5 HAMP also includes a 35.5 GHz cloud radar with a sensitivity of -30 dBZ at 13 km distance in the NARVAL setup. Profiles of the radar reflectivity factor (Z) and the linear depolarization ratio are recorded with 30 m vertical and 1 s temporal resolution. To supplement HAMP, Vaisala RD94 dropsondes were launched from HALO to provide the thermodynamic conditions of the environment. In total 76 and 215 sondes were released during NARVAL1 and NARVAL2, respectively.

To distinguish between clear sky and cloudy conditions as good well as possible, Schnitt et al. (2017) derive a cloud mask for NARVAL1 based on the nadir spectral solar radiance measurements by HALO-SR (HALO Solar Radiation; Fricke et al. (2014)). Unfortunately, sun glint in August deteriorated the cloud mask retrieval during NARVAL2. Therefore, G6dde (2018) developed a cloud mask product using the imaging spectrometer specMACS (spectrometer of the Munich Aerosol and Cloud Scanner; Ewald et al. (2016)) which overcomes the sun glint problem. However, specMACS was not installed during NARVAL1. In order to have similar cloud mask performance during both campaigns the aerosol backscatter profile measured by the WATER vapor Lidar Experiment in Space (WALES) airborne demonstrator (Wirth et al., 2009) is used instead to provide an along track cloud mask with 1 s resolution.

WALES also provides profiles of water vapor molecular number density based on the differential absorption lidar (DIAL) principle. These profiles are converted to volume mixing ratio profiles using temperature and pressure data from ECMWF analyses. A resolution of about 200 m vertical and 12 s temporal was chosen as a compromise between accuracy and resolution.

- 20 The water vapor data is given on the vertical grid of the raw backscatter data which is 15 m, but smoothed with an averaging

kernel of 200 m width (full width at half maximum, FWHM). Water vapor profiles are provided down to about 250 m in cases with no or optically thin clouds, which can be penetrated by the lidar beam. Water vapor information is available below thin clouds, but the cloud itself is masked out in the profile.

The requirement of simultaneous measurements by all sensors reduces the dataset. While all research flights during NARVAL1 can be used, ~~hardware issues during no data is available for some~~ NARVAL2 ~~prevent having data during some flights~~ flight days due to hardware issues as summarized in Tab. 1. The spatial sampling differs even with the same temporal sampling due to footprints differences. The HAMP MWR has the largest beamwidth in its lowest frequency channel of 5.0° (FWHM). The corresponding surface footprint at 10 km altitude is about 870 m across and 1090 m along track. The HAMP radar beamwidth is 0.6° whereas WALES has a field of view of 1.6 mrad. The respective footprints are about $105\text{ m} \times 335\text{ m}$ and $16\text{ m} \times 216\text{ m}$. We reduce the along track sampling differences by averaging temporally, but the cross track sampling issues remain. This means, a cloud covering a lateral part of the MWR footprint might be missed by the lidar or even the radar. Cross track imagers such as specMACS could be used to assess these issues. However, specMACS was only installed on HALO for NARVAL2 and the detailed analysis of HAMP beam filling is beyond the scope of this study. The problem of different footprints and sensitivities of different NARVAL instruments for cloud masking is illustrated by Stevens et al. (2019, in press).

Table 1. Dataset availability. Days of research flights from which the datasets are used for the study of NARVAL1 and NARVAL2, respectively.

Dataset	NARVAL1	NARVAL2
	day in December 2013	day in August 2016
HAMP Radiometer	10, 11, 12, 14, 15, 16, 19, 20	8, 10, 12, 15, 17, 19, 22, 24, 26, 30
HAMP Radar	10, 11, 12, 14, 15, 16, 19, 20	8, 10, 12, 15, 17, 19, 22
Drosondes	10, 11, 12, 14, 15, 16, 19, 20	8, 10, 12, 15, 17, 19, 22, 24, 26, 30
WALES water vapor	10, 11, 12, 14, 15, 16, 19, 20	10, 12, 15, 17, 19, 22, 24, 26, 30
WALES cloud mask	10, 11, 12, 14, 15, 16, 19, 20	8, 12, 15, 19, 22, 24, 26, 30

2.3 Retrieval database

Recently, high spatial resolution simulations with the storm resolving ICON model were able to show how resolved convection and its associated circulation interact with and form the larger scale circulation within the Atlantic inter-tropical convergence zone (Klocke et al., 2017). These simulations serve for training and testing retrieval algorithms. The simulations were performed on a triangular grid with a horizontal spacing of about 1.25 km and 75 vertical levels. The simulations cover the area of 4°S to 18°N and 64°W to 42°W . The data was spatially sub-sampled to reduce the computational effort while still covering the variability of atmospheric profiles. To eliminate atmospheric columns with a high degree of correlation, columns are selected on a $0.5^\circ \times 0.5^\circ$ longitude-latitude grid, so that each time step includes 849 cases over the ocean as indicated in Fig. 1. Data from 24 days with hourly outputs each, spanning the period of each campaign are alternately separated into test and training

data. In general, the training and test data excludes cases with ~~ice or~~ LWP greater than 1000 g m^{-2} and cases with ice, i.e., 86 % of all profiles over the ocean ~~is~~ are used. This limitation is done as our focus is on liquid clouds and their transition to rain. Note that classical satellite algorithms (e.g., Wentz and Meissner (2000)) are trained with an upper LWP limitation of 300 g m^{-2} .

Synthetic HAMP measurements, i.e., BTs and radar reflectivity profiles in nadir view, are simulated for each model column based on its thermodynamic profile and hydrometeors (cloud liquid water, rain, cloud ice, snow, and graupel). The Passive and Active Microwave TRAnsfer code (PAMTRA; Maahn et al. (2015); Cadeddu et al. (2017)) is used. It is configured with 27 output levels to mimic different flight altitudes (6 km to 15 km). The ICON model was set up with a one-moment microphysics scheme (Baldauf et al., 2011). In PAMTRA, cloud and rain particles and their size distributions are described according to the microphysical scheme of ICON and the single scattering properties for each particle are approximated by the Lorentz-Mie theory. Cloud and rain particles are simulated with a $20 \mu\text{m}$ diameter mono-disperse and exponential distribution of water spheres, respectively. The exponential distribution has its intersect N_0 classically fixed to 0.08 cm^{-4} (Marshall and Palmer, 1948). Absorption coefficients of atmospheric gases (i.e., oxygen, water vapor, nitrogen) are calculated after Rosenkranz (1998) with corrections of the water vapor continuum absorption according to Turner et al. (2009) and the line width modification of the 22.235 GHz water vapor line as proposed by Liljegren et al. (2005). The emissivity and reflectivity of the sea ocean surface is calculated by the FAST microwave Emissivity Model version 5 (FASTEM5; Liu et al. (2011)), which is a modification of the Fresnel coefficients including corrections for ocean surface roughness and foam building as a function of wind speed.

To test the realism of the retrieval database, histograms of BTs were compared with their observed counterparts. Joint histograms of an absorption (22 GHz) and a window channel (31 GHz or 90 GHz) show that the relations between channel pairs are depicted in the model and observations in the same way (Fig. 3). In clear sky conditions absorption and window channels are highly correlated with both increasing with increasing moisture albeit the increase is less in the window channels. Clear sky cases with low BT_{31} and BT_{90} are ~~visibly~~ visible as a line of high occurrence and reveal the linear relation between absorption and window channel BTs as a function of IWV. The simulations and measurements show the same relations but differ slightly in terms of the BT combination distribution within this line as the underlying IWV sampling is slightly different. If liquid water clouds occur, the window channel BTs increase compared to clear sky cases (solid lines in Fig. 3). The window channel at 90 GHz has a higher sensitivity towards LWP compared to BT_{31} as it can be seen by the increased LWP line spread. Rainy cases show higher emissions in all channels (dotted lines in Fig. 3). For thick clouds and rain the most liquid sensitive channel (90 GHz) experiences saturation effects with BT_{90} approaching cloud temperatures. The joint histograms reveal the major signals by liquid and water vapor which are exploited within retrieval algorithms. However, multiple influence factors like the exact vertical structure lead to the variability illustrated in Fig. 3. Minor deviations between observations and simulations are visible in the frequency of combinations of BT_{31} and BT_{90} with high BT_{22} . Those combinations are associated with heavy precipitation and were observed less frequently than present in the model as flight patterns avoided heaviest precipitation.

2.4 Retrieval

The atmosphere emits radiation depending on the atmospheric state as illustrated in Fig. 2. In general, the retrieval of the atmospheric state from MWR measurements is under-determined as multiple atmospheric states can lead to the same set of




Figure 3. Relation between brightness temperatures (BT) in an absorption channel (22.24 GHz) and two window channels, i.e., 31.40 GHz (a, b) and 90.00 GHz (c, d). Two-dimensional histograms of occurrences in simulations (a, c) and HAMP measurements (b, d). Solid contours highlight BT combinations in simulations that mostly occur with LWP higher than 50 g m^{-2} , 250 g m^{-2} , and 500 g m^{-2} . Dotted lines highlight combinations where RWP mostly exceeds the respective threshold. LWP contours in b and d are taken from a and c for guidance. HAMP data from all NARVAL2 flights and ICON/PAMTRA data of the corresponding dates are used. Profiles and measurements with ice are excluded.

BTs. Statistical relations have to be established to link the measurement to the most common atmospheric state that can provide those measurements. To account for non-linearity in this inverse modeling problem, we use an artificial neural network (NN) model similar to Cadeddu et al. (2009) to relate BTs to IWV and LWP. The NN is configured with eight input neurons and 15 hidden neurons in one hidden layer.

- 5 For testing and training the retrievals, Gaussian noise of 0.5 K is added to the simulated BTs to account for uncertainties of the HAMP MWR, the radiative transfer, and absorption model. The database is separated by alternating days into test and training data. Retrieval parameters are derived for each PAMTRA output level to account for the altitude dependence of the microwave signal. The output levels are chosen such, that a HALO flight level never deviates more than 90 m from the next output level. The parameters at output levels closest to HALO's altitude are interpolated to HALO's altitude in the retrieval application.
- 10 Retrieval parameters are derived separately for both campaigns. For testing, each retrieval is applied to the test data of the campaign it is trained for.

In the retrievals, IWV and LWP, and later ~~cloud-liquid-water-path (CLWP)-and-rain-water-path (RWP)~~ CLWP and RWP are the integrals of the water vapor and liquid water over the whole column as seen from space. Despite the fact that HALO flies lower, we chose the total integrals as they prevent artificial flight level depended biases in statistics and allow a comparison with

satellite and model data. According to ICON model data, typically less than 0.1 kg m^{-2} water vapor is above a flight altitude of 10 km. About 1 kg m^{-2} of IWV is not seen by the MWR at the lowest NARVAL2 flight altitude of 6.4 km but is included in the retrieval. The LWP retrieval is trained with the integral of all liquid water, that is given by the model either as cloud water or rain water. The sum of both is used due to the difficulty of MWRs to distinguish clouds and rain (Fig. 2).

- 5 Neural network LWP retrievals are compared with linear regression (REG) models as used by Schnitt et al. (2017). The regression relates measured brightness temperatures BT_i to LWP including the quadratic terms of BT_i

$$LWP = c + \sum_i (b_i BT_i + a_i BT_i^2), \quad (1)$$

- where a_i , b_i and c are regression coefficients. Such REGs are less susceptible than NNs to extrapolation towards unforeseen input data, i.e., data values or combinations that are not covered by the training data. However, NNs are better in representing
 10 non-linear effects that are apparent in microwave radiative transfer and thus can better adjust to the extremes of the LWP target space. The application of the retrievals to test data reveals overall uncertainties between 0.5 kg m^{-2} and 0.6 kg m^{-2} for IWV for both approaches, i.e. NN and REG, and 22 g m^{-2} and 26 g m^{-2} for LWP using the NN and REG, respectively. For LWP the uncertainty strongly depends on atmospheric conditions as it will be investigated in Sec. 4.

- When retrieval algorithms are applied to HALO measurements, slight biases ~~are observed under clear sky which show slow~~
 15 ~~changes with time of LWP from 0 with slow changes over time are observed during clear sky scenes~~. To reduce these biases and to improve the retrieval of low LWP values, we follow the synergistic approach by van Meijgaard and Crewell (2005). Herein, we use the WALES cloud mask for clear sky identification. HAMP measurements are considered clear sky if no cloud is detected by WALES within ± 2 seconds flight time. The distance weighted average clear sky LWP within ± 30 minutes is then subtracted from each a priori retrieved LWP value.

- 20 In thick clouds we occasionally observed, that while the REG retrieval gave $LWP > 1000 \text{ g m}^{-2}$, the NN LWP time series showed a sudden decline. This is likely caused by the clipping of the NN retrieval at 1000 g m^{-2} , which is expected as the retrieval database is limited to $LWP < 1000 \text{ g m}^{-2}$ and thus BTs associated with higher amounts of liquid are unknown to the retrieval. To avoid this behavior, we use a second NN retrieval trained with an extended database up to 4000 g m^{-2} to flag scenes that are potentially above 1000 g m^{-2} . Overall, 0.76 % of the measurements were masked in this way. Note that these
 25 measurements often coincide with ice scattering depressions in channels at higher frequencies.

To retrieve the contribution of rain drops (RWP) to the total LWP, the vertically integrated radar reflectivity is used in addition to the MWR channels in another NN retrieval. The aim is separating the LWP into CLWP and RWP, i.e., splitting the contributions from small cloud droplets and larger rain drops by estimating the fraction

$$f = \frac{RWP}{LWP} = \frac{RWP}{RWP + CLWP}. \quad (2)$$

- 30 This retrieval is based on the hydrometeor classes of rain and cloud liquid water in the ICON model. The RWP is calculated by multiplying f and the retrieved total LWP.

3 Assessment of integrated water vapor

Three independent methods to derive IWV are available from HALO: the MWR retrieval, vertically integrated humidity from dropsondes, and vertically integrated humidity from WALES. Each of the three methods has its advantages and shortcomings. The microwave radiometry can not provide profile information but gives continuous IWV under nearly all sky conditions. The dropsondes provide in situ measurements, but no valid data up to about the first half kilometer below the aircraft because of the sensor's adjustment from the aircraft cabin conditions to the outside. Furthermore, wind drifts sondes out of the aircraft nadir with a typical horizontal drift during the decent of 4 km. The dropsonde relative humidity sensor has a repeatability of 2 % according to the manufacturer (Vaisala, 2017). This relates to an IWV accuracy of about 1.4 kg m^{-2} . WALES provides water vapor profiles, but they are only available when no cloud extinguishes the laser beam. This limits the application of WALES for the IWV retrieval to clear sky scenes.

To compute the numerical derivative in the DIAL equation, the first data point is at about 250 m above the sea surface and centered at the retrieval interval. Therefore, in the vertical integration, the missing near surface information is filled with the median mixing ratio in the lowest five range bins. The median is chosen to reduce any surface artifacts which can occur, when the first raw data signal point used in the retrieval contains the surface reflex. We estimate that the error of this assumption is about 0.3 kg m^{-2} by analyzing dropsonde humidity profiles. The IWV estimation is discarded if information of more than 400 m above sea-level is missing or there is a gap due to a thin cloud. Also, stability of the estimated WALES IWV is required, which means that the differences to the preceding and succeeding IWV estimations have to be smaller than 2 kg m^{-2} .

An example of water vapor retrievals on August 19, 2016 is shown in Fig. 4. An elevated moisture layer between 3 km to 4 km altitude is visible in the first half of the scene. Around 14:53 a plume of moist air reaching up into even higher levels causes an IWV gradient of nearly 10 kg m^{-2} (26 kg m^{-2} to 35 kg m^{-2}) over a distance of about 110 km. This gradient is captured well by WALES and HAMP. The two dropsondes that were released between 14:45 and 14:55 reconstruct this gradient, but both have a dry offset. This offset might be due to drifting of the sonde towards the drier air mass. After a short outage of WALES at around 15:00, shallow clouds below 2 km prevent the determination of lidar IWV frequently. Most of the IWV measurements from dropsondes agree with the coincident remote sensing estimates within the sondes' uncertainty.

A more quantitative comparison is achieved by considering all measurements from both campaigns which cover a wide variety of water vapor conditions (Fig. 5). Overall, the sondes agree well with HAMP over the whole observed range from very low (20 kg m^{-2}) to very high (60 kg m^{-2}) values of IWV (Fig. 5a). The root-mean-square deviation (RMSD) is 1.39 kg m^{-2} (1.28 kg m^{-2}) with a mean bias of 0.28 kg m^{-2} (0.47 kg m^{-2}) during NARVAL1 (NARVAL2) [as summarized in Tab. 2](#). The positive biases of HAMP is most likely caused by the retrieval, which is trained with the whole column IWV, whereas the sonde IWV is only integrated along its measurement path. Most sondes were released above 9 km which would miss an IWV of about 0.2 kg m^{-2} according to ICON data. Note that dropsondes released from below 6.5 km are discarded in the comparison to avoid an artificial bias. The random error between HAMP and sondes (1.2 kg m^{-2}) is smaller than the estimated uncertainties of the dropsonde (1.4 kg m^{-2}) and the MWR retrieval (0.6 kg m^{-2}) which indicates the high quality of the measurements as uncertainties due to spatio-temporal mismatch are included in the RMSD as well. Note that uncertainties due to MWR calibra-




Figure 4. Water vapor time series from NARVAL2 research flight 6 on August 19, 2016. Top: I WV time series by HAMP, WALES and dropsondes with their uncertainties. Bottom: WALES water vapor profiles. White areas denotes masked lidar data. The scene represents a circle around 14.8°N and 51.0°W over a distance of 520 km.

tion are largely compensated as offsets between measured BT from those derived by radiative transfer calculations based on dropsondes have been corrected by Konow et al. (2018a).




Figure 5. IWV comparison of HAMP MWR and dropsondes (a), HAMP MWR and WALES (b) and WALES and dropsondes (c) during NARVAL2. The colors indicate the flight days in August 2016. Scores are given in Tab. 2.

WALES IWV can be used for continuous comparison to HAMP IWV along the flight track in clear sky scenes. A comparison of all coincident measurements during NARVAL2 is depicted in Fig. 5b. The average bias between HAMP and WALES IWV is -0.59 kg m^{-2} . The bias is cut in half when considering only the 40 simultaneous measurements during which a dropsonde was launched (Tab. 2). The random error is smaller in contrast to the HAMP – dropsonde comparison. This is likely due to

Table 2. Comparison of IWV retrieved from HAMP, WALES and dropsondes. Pairwise observations of two instruments and the subsets of the observations for that all instruments were available. Bias, root-mean-square deviation (RMSD), and bias corrected RMSD (STD) in kg m^{-2} . “Observed by all” implies only small or no clouds.

		HAMP - sondes	HAMP - WALES	WALES - sondes
NARVAL1 observed pairwise	bias	0.28	0.92	−1.21
	RMSD	1.39	1.36	1.60
	STD	1.38	1.01	1.07
	count	43	2482	24
NARVAL1 observed by all	bias	0.32	1.70	−1.37
	RMSD	1.21	2.20	1.70
	STD	1.20	1.41	1.03
	count	21	21	21
NARVAL2 observed pairwise (Fig. 5)	bias	0.47	−0.59	0.73
	RMSD	1.28	1.21	1.38
	STD	1.19	1.06	1.19
	count	146	1632	47
NARVAL2 observed by all	bias	0.32	−0.25	0.57
	RMSD	1.16	0.82	1.23
	STD	1.12	0.79	1.11
	count	40	40	40

the better spatial match between the two nadir measurements compared to a drifting sonde. However, higher RMSD between HAMP and WALES IWV can be found during NARVAL1, which is mostly related to a higher bias. The bias increases to 1.70 kg m^{-2} in the HAMP – WALES comparison when only considering measurements during which a sonde was released. A bias of similar magnitude is apparent between WALES and the dropsondes. Most likely the dry bias of WALES is due to the method of how the 12 s water vapor profiles are derived. The profiles only contain raw profiles (within the 12 s), that are not blocked by a cloud. For small scale boundary layer convection, this means preferred sampling of downdraft regions. In these downdraft regions dry air is entrained from the rather dry free troposphere into the convection layer during NARVAL1 (Stevens et al., 2017). This results in biased sampling of rather dry profiles. During NARVAL2 humidity was reaching higher altitudes, which resulted in less entrainment of dry air in cloud gaps.

A small confounding effect from liquid water in cloudy scenes can be derived from the separation of the HAMP – dropsonde comparison into all (“observed pairwise”) and clear sky (“observed by all”, i.e., when also WALES is available) in Tab. 2. In the NARVAL1 dataset, the bias for cloudy sky sondes (0.24 kg m^{-2}) is somewhat smaller than that for clear sky (0.32 kg m^{-2}).

However, RMSD and STD in cloudy scenes are about 0.3 kg m^{-2} larger than in clear sky. NARVAL2 also shows a larger bias in cloudy sky of about 0.53 kg m^{-2} in comparison to clear sky (0.28 kg m^{-2}). The cloudy sky RMSD and STD of 1.32 kg m^{-2} and 1.21 kg m^{-2} , respectively, are only slightly larger than their clear sky counterparts. An increase of the random error for cloudy scenes is expected as also higher water vapor variations are expected in heterogeneous cloud fields.

- 5 With the exception of the HAMP – WALES comparison during NARVAL1, the RMSD between the different instrument pairs is found between 0.8 kg m^{-2} and 1.4 kg m^{-2} (Tab. 2). This corresponds to an error of 2 % to 7 % over the observed range of 20 kg m^{-2} to 60 kg m^{-2} . For comparison, Mears et al. (2015) found random IWV deviations between different spaceborne MWR and ground-based GPS (Global Positioning System) instruments of 1.7 kg m^{-2} to 2.0 kg m^{-2} over a similar IWV range using 26 small island stations located mainly in the tropics.
- 10 The HAMP IWV retrieval has a theoretical uncertainty of about 0.6 kg m^{-2} which is derived by applying the IWV retrieval to simulated measurements from the test database (Sec. 2.4) and is constant over a wide IWV range (not shown). This is well in line with the RMSD derived in the pairwise comparisons taking into account the estimated uncertainties of WALES and dropsondes as well as uncertainties due spatio-temporal mismatch. In summary, the pairwise comparisons in relation to the individual uncertainties indicate high HAMP IWV performance and the suitability of our retrieval approach.

15 4 Assessment of liquid water path

There are no independent measurements of sufficient quality to assess the quality of the LWP product. However, the large retrieval database (173 339 ice free cases in the test dataset) allows a theoretical in-depth analysis of the retrieval performance. This approach is supported by the good consistency between the BTs in the database and the HAMP measurements in terms of relation resemblance (Fig. 3) and performance of IWV retrieval (Sec. 3). We analyze the retrieval error as a function of the true LWP as well as of the retrieved LWP using the database.

- First, we analyze the difference of retrievals developed with all ice free cases of the training database (all sky) and with cloudy cases only, which reduces the dataset size to about one quarter. A model profile is regarded as cloudy if $\text{LWP} > 1 \text{ g m}^{-2}$. REG and NN retrievals are trained with the all sky and the cloudy sky datasets separately. The errors of retrieved LWP from the test database are calculated for bins of the true LWP. Both REG and NN show similar behavior of the RMSD between the retrieved LWP and the model truth with increasing LWP (Fig. 6a). The RMSD is constant for LWP below about 30 g m^{-2} and increases with LWP, e.g. 50 g m^{-2} at 500 g m^{-2} . For LWP values $> \sim 800 \text{ g m}^{-2}$ the number of test cases reduces strongly leading to less robust results. For $\text{LWP} < 30 \text{ g m}^{-2}$, the errors are smaller for REG and NN retrieval types if the clear sky cases are included in the training (compare Fig. 6a and 6b). Including clear sky in the training, the retrieval errors decrease slightly for a REG model and are almost cut to half for an NN. This shows the ability of an NN to non-linearly relate a variety of BT combinations to zero LWP. However, retrievals that are especially trained for all sky scenes have a considerably larger RMSD for $\text{LWP} > 20 \text{ g m}^{-2}$ than those trained with cloudy cases only as clear sky cases make up 77 % of the data. Since we are targeting clouds and not clear sky, we ~~decided for~~ chose a retrieval trained with data excluding the clear sky model profiles.
- 30

Instead [to include clear sky directly in the retrieval](#), we make use of lidar measurements, which are better suited than MWR for cloud masking.

Regarding cloudy sky retrievals, the RMSD for a given (true) LWP less than 40 g m^{-2} is smaller when using the NN retrieval instead of a REG model (Fig. 6a). This is related to a suppression of unphysical negative LWP values by the NN. Thus, in contrast to a REG which has a nearly Gaussian error characteristic, the NN tends to overestimate LWP. This results in a more negative mean LWP error (true - retrieved) of clouds with less than 10 g m^{-2} but also in a smaller interquartile range of errors when using the NN instead of the REG. However, the retrieval error for true LWP $< 10 \text{ g m}^{-2}$ remains in the order of 10 g m^{-2} to 18 g m^{-2} even when using the NN.

The bias errors visible in Fig. 6a can not be used to adjust the retrieved LWP as the true LWP value is not known in practice. For the application of the error analysis on measurements, it is important to analyze the LWP error as a function of the retrieved LWP. The RMSDs of the NN and REG retrievals are larger than 100 % for a retrieved LWP below 12 g m^{-2} which can be regarded as a detection limit (Fig. 7). Therefore, ancillary measurements with higher sensitivity are needed to detect these thin liquid water clouds. The RMSD is below 20 g m^{-2} for REG LWP $< 50 \text{ g m}^{-2}$ and NN LWP $< 100 \text{ g m}^{-2}$, and moderately increases with increasing LWP. Therefore, the relative RMSD decreases from 50 % for a retrieved LWP of about 40 g m^{-2} to 20 % for LWP $> 100 \text{ g m}^{-2}$ for both retrieval types. While the RMSD is rather similar for REG and NN, the NN succeeds in capturing the nonlinear retrieval providing a nearly zero bias across the full LWP range and is therefore preferred over REG.




Figure 6. Expected retrieval error as function of true LWP for neural network and linear regression LWP retrievals. Retrievals (a) trained for $1 \text{ g m}^{-2} < \text{LWP} < 1000 \text{ g m}^{-2}$. Retrievals (b) trained including clear sky cases ($\text{LWP} < 1000 \text{ g m}^{-2}$). Error measures (colored lines) for logarithmically distributed bins with ten bins per LWP power of ten. Gray dashed lines denote the corresponding relative LWP error.

Analyzing the retrieved LWP distribution for clear sky scenes is a widely used method to assess an LWP retrieval (e.g., Liu et al. (2001), Greenwald et al. (2018)) because this characterization can be made from measurements using ancillary observations that define clear sky scenes. We use WALES measurements for cloud and clear sky indication. The distributions of LWP from HAMP MWR are depicted in Fig. 8 for observed clear sky scenes (blue lines) along the track for both campaigns.




Figure 7. As Fig. 6, but with errors shown as function of retrieved LWP. Retrievals are trained and tested with $1 \text{ g m}^{-2} < \text{LWP} < 1000 \text{ g m}^{-2}$. First bin contains all data with retrieved LWP $< 2.5 \text{ g m}^{-2}$ (including negative).

The distributions are compared to the theoretical ones of retrieved LWP from all clear sky (true LWP $< 1 \text{ g m}^{-2}$) cases of the respective campaign in the ICON/PAMTRA database (orange lines in Fig. 8). The latter distributions are closely related to the retrieval uncertainty of the lowest LWP bin in Fig. 6a as this represents the retrieval uncertainty for true LWP $< 2.5 \text{ g m}^{-2}$. The distributions roughly resemble Gaussian behavior with mean values of about 10 g m^{-2} and widths of about 9 g m^{-2} . Some differences between NARVAL1 and -2 exist which are even stronger for the measured distributions. During NARVAL1, the measured distribution is skewed towards higher values. This might be caused by cloud patches that were only present in a lateral part of the MWR footprint such that the scene was falsely identified as clear sky by the lidar, which only slices through the center of the MWR footprint. As this effect is not visible for NARVAL2 measurements, it might be that clouds were generally smaller and more frequent during NARVAL1 (see Sec 6).

For both campaigns the similar widths and standard deviations of the retrieved LWP indicate a good agreement between simulations and measurements for clear sky (Fig.8). The apparent second mode at 20 g m^{-2} in the observed clear sky LWP distribution during NARVAL2 is caused by different mean deviations during different flights, probably influenced by the calibration. Overall, the narrow Gaussian widths (11.4 g m^{-2} and 8.3 g m^{-2} for NARVAL1 and -2) of the retrieved clear sky LWP distributions demonstrate the good performance of HAMP compared to evaluation studies by Liu et al. (2001) (28 g m^{-2} , airborne) and Greenwald et al. (2018) ($\sim 30 \text{ g m}^{-2}$, satellite). The better HAMP performance is likely due to its smaller footprint, additional frequency channels, and more recent technology. The sensor synergy of using the lidar cloud mask for clear sky bias correction (Sec. 2.4) reduces the bias in clear sky conditions to values barely above zero as small cloud patches can still be in

the outer area of the MWR footprint which is not transected by the lidar beam. The bias correction further narrows the clear sky LWP distributions. Note that a good agreement (small bias) is expected as the lidar cloud mask is also used to define clear sky. The deviations of the observed clear sky LWP distributions from delta distributions are due to the moving window in the bias correction.




Figure 8. LWP distributions retrieved during clear sky scenes only. Shown are the LWP retrieved with the neural network based on radiometer observations (blue lines) during (a) NARVAL1 and (b) NARVAL2, the retrieved LWP from the test database (orange lines) for profiles with $\text{LWP} < 1 \text{ g m}^{-2}$, and the observed LWP after applying the lidar clear sky correction (green lines). Mean (μ) and standard deviation (σ) are given for each distribution in g m^{-2} .

5 In summary, the ICON/PAMTRA database allows estimating the expected uncertainty of the LWP retrieval. This reveals a lower retrieval limit of about 12 g m^{-2} and an RMSD below 20 g m^{-2} for LWP below 100 g m^{-2} and below 20 % above for the NN retrieval. A narrow clear sky distribution of HAMP measurements ($\text{STD} \sim 10 \text{ g m}^{-2}$) is found that is in good agreement with the theoretical assessment, but a small bias in the order of 12 g m^{-2} remains which is eliminated by the clear sky correction. The synergy of MWR and lidar removes the bias and reduces the clear sky LWP noise to 5 g m^{-2} to 7 g m^{-2} .

10 5 Assessment of rain

To investigate the formation of rain with HAMP measurements, this section extends the applicability of the LWP retrieval to drizzle and light precipitation by combining cloud radar with MWR. As described in Sec. 2.4, RWP is retrieved as the fraction $f = \frac{\text{RWP}}{\text{LWP}}$ by a NN using eight BTs and integrated radar reflectivity as input. Two physical effects are considered in the retrieval: hydrometeor scattering, which becomes more important with increasing droplet size and microwave frequency, and radar backscatter being sensitive to D^6 , where D is the droplet diameter. The first effect is considered by including the 90 GHz channel as proposed by Cadeddu et al. (2017). For the latter effect, the vertically integrated (linear) radar reflectivity is used as retrieval input in addition to the MWR channels also used in the LWP retrieval. This integrated reflectivity as a columnar quantity is more comparable to a BT and less noisy than the reflectivity of a single range gate and is thus used as retrieval input.

The Gilbert skill score (GSS) (Hogan et al., 2010), also known as equitable threat score (ETS), is used to rate how well retrieval "yes" events correspond to true "yes" events while accounting for hits due to chance. "Yes" events mark RWP above a given threshold. The GSS is defined as

$$GSS = \frac{hits - hits_by_chance}{hits + misses + false_alarms - hits_by_chance} \quad (3)$$

5 using the common entries of the contingency table and the hits due to chance

$$hits_by_chance = \frac{(hits + misses) \times (hits + false_alarms)}{hits + misses + false_alarms + correct_negatives}. \quad (4)$$

GSS ranges from $-\frac{1}{3}$ to 1 with 1 being the perfect score.

The retrieval of RWP is evaluated for different RWP thresholds (Fig. 9). GSS shows good performance being higher than 0.75 for RWP thresholds from 10 g m^{-2} to about 50 g m^{-2} and higher than 0.5 for RWP up to 250 g m^{-2} . Note that 762, 295, and 62 of the test cases have RWP greater than 10 g m^{-2} , 50 g m^{-2} , and 250 g m^{-2} , respectively and only few samples with higher RWP are available. The hit rate is higher than 80 % for thresholds between 10 g m^{-2} and 250 g m^{-2} , but the 250 g m^{-2} threshold also generates 37 % false alarms. Especially, the high GSS for low RWP thresholds makes the f retrieval a useful tool combining cloud radar and MWR for detecting measurements, that contain warm precipitation.

A case study of two showering shallow cumuli is shown in Fig. 10 to illustrate the capabilities of retrieving ~~LWP and RWP~~ CLWP and RWP separately. The figure shows how HAMP is able to resolve spatial features of showering cells, which were observed with a cross section of several HAMP footprints. The precipitating core of both cells had maximum RWPs of probably more than 200 g m^{-2} . The ~~RWP shows~~ stronger relative gradients ~~than LWP (=CLWP + RWP), which indicates a narrow~~ of RWP compared to CLWP indicate the narrowness of the precipitating core. Note ~~that how the~~ higher horizontally resolved information by radar (MWR footprints 3.3° to 5.0° vs. radar footprint 0.6°) ~~is integrated in~~ contributes relatively stronger to the RWP retrieval ~~compared to the LWP than to the CLWP~~ retrieval. The RWP retrieval consistently indicates no rain except for the time when the radar signal touches the surface or when there is a clearly visible fallstreak (17:42:30). The two showering clouds reveal maximum ~~LWP of 716 g m^{-2} and more than 1000 g m^{-2} , respectively. The latter is likely a lower limit as the LWP~~ total LWP of more than 700 g m^{-2} and 1000 g m^{-2} . The second shower core likely contains more water than indicated, as the retrieval sets the clipping flag. This case study also demonstrates the higher sensitivity of the lidar and the (C)LWP retrieval which ~~deteet clouds shows cloud signals~~ between 17:38:30 and 17:39:10 of clouds which are too thin to be ~~visible in the radar data~~ detected by the radar.




Figure 9. Scores for detecting an atmospheric profile with RWP higher than the respective threshold. (a) Gilbert skill score (GSS), hit rate and false alarm ratio. (b) Median fraction of rainwater as a function of RWP threshold. (c) Number of alarms and events for retrieved and true RWP above the threshold, respectively.




Figure 10. Example scene of rain retrieval from NARVAL1 research flight 8 on December 20, 2013. Retrieved ~~LWP~~CLWP (top), retrieved RWP (middle) and radar reflectivity profile (bottom). Note that the scale in the top panel is piecewise linear with the scale change at 20 g m^{-2} . The IWV varies around $31.5 \pm 1.5 \text{ kg m}^{-2}$ in this scene ~~has a length of 117 km.~~

6 Comparison of dry and wet season

The synergy of lidar, radar and MWR is necessary to understand the difference of clouds in the dry and wet season as all instruments have their specific limitations. The lidar cloud mask indicates the more frequent occurrence of clear sky during the wet season (70.0 %) compared to the dry season (53.3 %, Tab. 3) even though the environment is characterized by less humid air in the dry season (Fig. 11a, b). The IWV distribution is clearly confined to moderate values with a mean of 28 kg m^{-2} in the dry season which is mainly due to a rather dry middle troposphere seen in the lidar water vapor profiles. During the wet season, IWV values up to 60 kg m^{-2} were observed distributed into two modes around 35 kg m^{-2} and 52 kg m^{-2} . These two modes are clearly distinct from the single mode observed in the dry season and reveal the expected humid character of the wet season. The most humid air during NARVAL1 was sampled in a deep convective system on the southernmost leg of research flight 2 on December 11, 2013. This was the NARVAL1 flight during which HALO was closest to the inter-tropical convergence zone (ITCZ). The NARVAL2 IWV distribution seems to be driven by the vicinity of the flight track to deeper convective systems and the ITCZ as it can be analyzed from satellite images and thus also by the selection of flight patterns.

Interestingly, clouds show a higher mean LWP of about 63 g m^{-2} in the dry season compared to a mean LWP of 45 g m^{-2} 40 g m^{-2} in the wet season. Likewise, thicker clouds ($\text{LWP} > 50 \text{ g m}^{-2}$) were more frequent in the dry season (Fig. 11c, d), i.e., 27.1 % of the time when a cloud was seen in the dry season it contained $\text{LWP} > 50 \text{ g m}^{-2}$, whereas only 18.6 % of the time in the wet season clouds exceeded this value. ~~However, the variability between flights (Fig. 11) is high and illustrates sampling issues due to flight planing considerations. An analysis of ground-based LWP measurements at the Barbados Cloud Observatory (Stevens et al., 2016) over the years 2013-2018 confirms the generally higher LWP values during December than August (not shown).~~

The dry season clouds tend to produce light precipitation more frequently than the wet season clouds as indicated by the more frequent exceedance of RWP thresholds (Tab. 3). The cumulative distributions of RWP occurrences of all cloudy measurements with $\text{LWP} > 50 \text{ g m}^{-2}$ are depicted in Fig. 11e and f for each flight in the two seasons, when radar measurements are available. The vast majority (NARVAL1: 91 %; NARVAL2: 96 %) of all these measurements show $\text{RWP} < 10 \text{ g m}^{-2}$. Higher amounts of light rain seem to be more frequent in the dry season dataset, although the small number of heavy RWP observations inhibits a statistical sound statement as $\text{RWP} > 100 \text{ g m}^{-2}$ was only observed for 162 s and 49 s in the radar-radiometer datasets of dry and wet season, respectively. These time spans exclude measurements flagged as clipping ($\text{LWP} > 1000 \text{ g m}^{-2}$) or frozen precipitation (ice scattering). While warm precipitation seems to occur less often, clouds associated with frozen precipitation were more often observed in the wet season (1.6 % of the time) than in the dry season (0.5 %). Therefore, the lower LWP of the wet season clouds might be due to a higher precipitation efficiency compared to the dry season.

The higher LWP in the dry season might partly be explained by the choice of flight patterns. However, an analysis of ground-based LWP measurements at the Barbados Cloud Observatory (Stevens et al., 2016) over the years 2013-2018 confirms the generally higher LWP values during December than August (not shown). Thus, also changes in the organization of clouds could cause the differences in cloud fraction and LWP. The fact that the medium LWP range from 100 g m^{-2} to 400 g m^{-2} is less frequent in the wet season could be due to the higher degree of organization causing more clear sky areas and more intense

clusters with higher amounts of precipitation. In that sense the latter would be missed by our flight patterns as we avoided strongly convective scenes with formation of large ice particles.




Figure 11. Frequency distribution functions of IWV during the dry season (NARVAL1, a) and wet season (NARVAL2, b), LWP during NARVAL1 (c) and NARVAL2 (d), and cumulative distribution functions of RWP during NARVAL1 (e) and NARVAL2 (f). Colors denote the day in the month of the respective study. Colors in (a) and (b) denote the contribution of each flight to the total distribution. The bin edges are represented as minor ticks in (c) to (f). LWP distributions only include measurements where the lidar cloud flag reports a cloud within ± 2 seconds. RWP distribution is based on non-clear-sky-corrected LWP dataset (see note ^a Tab. 3), where $\text{LWP} > 50 \text{ g m}^{-2}$.

Table 3. Comparing NARVAL1 and NARVAL2 cloud properties observe south of 20°N and with HALO altitude above 6 km. Percentages of flight time with available corresponding datasets during each study.

Dataset	NARVAL1	NARVAL2	
	December 2013	August 2016	
	radiometer, lidar and radar	radiometer and lidar	radiometer and radar
Clear sky	53.31 %	69.95 %	–
LWP > 20 g m ^{−2}	21.62 %	10.60 %	–
LWP > 50 g m ^{−2}	12.63 %	5.26 %	–
LWP > 500 g m ^{−2}	1.18 %	0.33 %	–
$\overline{\text{LWP}}$ of clouds	63 g m ^{−2}	40 g m ^{−2}	–
RWP ^a > 10 g m ^{−2}	1.85 %	–	0.30 %
RWP ^a > 50 g m ^{−2}	0.43 %	–	0.78 <u>0.07</u> %
Ice-flag	0.51 %	0.94 %	1.76 %
LWP clipping	0.97 %	0.45 %	0.53 %
Total hours	25:26:18	39:43:28	41:22:48

^a Based on non-clear-sky-corrected LWP as radar and lidar cloud mask were only available during 5 of 10 flights during NARVAL2.

7 Summary and conclusions

Clouds play a critical role in the development of the future climate and especially marine low level clouds have been identified as source of uncertainty. An important cloud macrophysical quantity is LWP. Global observations are limited by satellite resolution or accuracy and ground-based observations over the oceans are few. To fill this observational gap, the NARVAL studies were initiated to assess North Atlantic trade wind clouds using the HALO research aircraft. We use a multichannel microwave radiometer, a cloud radar, a lidar and a dropsonde system deployed to HALO to provide insights into clouds on the kilometer scale. For NARVAL1 (December 2013) and NARVAL2 (August 2016) a unique retrieval training and test database was developed based on ICON simulations with 1.25 km grid spacing. The database contains more than 350 000 physically consistent profiles that characterize the thermodynamic state of the atmosphere and the hydrometeor distributions during each of the two campaigns. Synthetic HAMP measurements in terms of BTs and radar reflectivity profiles in nadir view were simulated for each profile using PAMTRA. The synthetic BT measurements show bivariate relations that are consistent ~~to~~with those observed and therefore give trust that the database can be used to develop retrievals and assess LWP quality.

To estimate IWV, LWP, and RWP from HAMP measurements, artificial neural networks are trained with the retrieval database. BTs of seven K-band and the 90 GHz channel are used for IWV and LWP; vertically integrated radar reflectivity is used in addition for RWP.

Similar to LWP, a IWV retrieval is based on the spectral of BT characteristics between the same water vapor absorption and window channels. A good retrieval of either IWV or LWP is a prerequisite for the other. The IWV comparison to dropsonde measurements and the continuous along-track comparison to the water vapor lidar WALES shows good agreement with an RMSD smaller than 1.4 kg m^{-2} and no distinct error dependence of IWV itself. Overall, the IWV assessment shows the good practical performance of HAMP and the suitability of the ICON/PAMTRA database for developing microwave retrievals for NARVAL1 and NARVAL2.

LWP retrievals are theoretically assessed as functions of retrieved LWP and true LWP. A slight advantage of the neural network compared to a linear regression retrieval is evident, especially at the limits of the LWP range (1 g m^{-2} to 1000 g m^{-2}). Both approaches show relative errors greater than 100 % for a retrieved LWP $< 12 \text{ g m}^{-2}$, which can be regarded as detection limits. If more liquid water is contained in the column, the random error decreases to 20 % at LWP $\approx 100 \text{ g m}^{-2}$ and 10 % at LWP $\approx 800 \text{ g m}^{-2}$. Both retrievals show an offset error smaller than the random component for LWP $< 10 \text{ g m}^{-2}$ with different signs depending on whether it is analyzed as function of true or retrieved LWP. Because of the ambiguity of the error sign, we conclude that this bias can not be accounted for with the MWR retrieval alone and we developed a synergistic clear sky offset correction using the WALES lidar cloud mask. The cloud mask reduces the noise of clear sky LWP to 7.1 g m^{-2} and 5.0 g m^{-2} for NARVAL1 and NARVAL2, respectively.

To allow investigating the onset of precipitation, a neural network retrieval is trained to estimate the fraction between RWP and LWP from a combination of integrated radar reflectivity factors and BTs. Using the test database, a Gilbert skill score above 0.75 is found for RWP thresholds between 10 g m^{-2} and 50 g m^{-2} which shows good applicability for detection of rain or drizzle onset.

We used data from 36 flight hours in December 2013 (dry season, NARVAL1) and 64 flight hours in August 2016 (wet season, NARVAL2) to investigate differences between the seasons. The analysis shows that although clouds were more frequent and their LWP and RWP were higher during the flights in the dry season, more microwave scattering of ice was observed in the wet season indicating strong precipitation events. The difference between $\overline{\text{LWP}}_{\text{dry season}} \approx 63 \text{ g m}^{-2}$ and $\overline{\text{LWP}}_{\text{wet season}} \approx 40 \text{ g m}^{-2}$ is clearly larger than the LWP retrieval uncertainty. As expected, the IWV histograms reveal the dry season as dryer and more uniform, and the wet season as more humid. However, the IWV distributions also reveal sampling biases due to flight track choices especially for the wet season. Therefore, the airborne measurements need to be combined with long-term ground-based and spaceborne measurements to draw statistically sound conclusions. The fine scale airborne microwave observations such as the measurements obtained with HAMP can be used to investigate the sub-satellite-footprint inhomogeneity of LWP and rain for a better error characterization of satellite measurements. [Sound conclusions on the diurnal cycle can not be drawn from the data presented here, as the spatial variability of the clouds on the observed mesoscale was higher than an expected effect of the diurnal cycle.](#)

The synergy of active and passive microwave observations could further benefit from using an optimal estimation approach including the full radar profile and all MWR channels to improve the partition of rain and cloud droplets and frozen particles (e.g., Battaglia et al. (2016)). [With respect to trade wind cumuli, the products of the present study in combination with cloud boundary estimations from the radar and backscatter lidar will be used to evaluate the condensate loads of different shallow](#)

trade wind cumulus types in large eddy simulations. For example, radar and lidar both detect shallow convection or shallow outflow anvils as depicted in Fig. 10. But in addition, the lidar also allows detecting boundary layer driven clouds, which have tops around 1 km and are below the radar sensitivity.

An extension of the NARVAL observations is planned by the EUREC⁴A field study in early 2020 (“elucidate the cou-
5 plings between clouds, convection and circulation”; Bony et al. (2017)) ~~and the~~ which among other objectives will investigate
convective aggregation. The algorithms presented here will be applied ~~.-This and together with additional measurements a~~
better understanding of the governing processes that cause differences between dry and wet season will be analyzed. For that
the campaign will provide additional observations of large scale dynamics ~~and horizontal~~, horizontally resolved remote sensing
observations by a second and in situ observations by additional aircraft in the cumulus layer. Also, more locally targeted flights,
10 distributed over the daytime are planed to study the diurnal cycle. Together with ship, drone and buoy measurements a unique
dataset for a better understanding of precipitation onset will be generated.

Data availability. The time series of IWV, LWP, and RWP are available in the CERA (Jacob et al., 2019a, b). The HAMP MWR, HAMP
radar and dropsonde data are published and described by Konow et al. (2018a, b, c).

Author contributions. FA and SC were initiators of the DFG HAMP project. MG and MW derived the lidar cloud mask and water vapor
15 profiles. HK provided quality controlled HAMP and dropsonde data in a unified file format. MJ and SC conceptualized this study. MM
designed the PAMTRA simulations. MJ developed the HAMP retrievals, conducted the analysis, and wrote the paper with support and input
from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The work has been supported by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG) within
20 the DFG Priority Program (SPP 1294) “Atmospheric and Earth System Research with the Research Aircraft HALO (High Altitude and Long
Range Research Aircraft)” under grant CR111/10-11. We would like to thank Daniel Klocke for running the ICON simulations and the
German Climate Computing Centre (DKRZ) for storing and supplying the data. D. Klocke is supported by the Hans Ertel Centre for Weather
Research (HErZ) by the DWD. Sabrina Schnitt and Mareike Burba are thanked for comments improving the manuscript. We appreciate the
dedication of Bjorn Stevens as the driving force behind the tropical NARVAL activities. The authors gratefully acknowledge the constructive
25 comments by two anonymous referees who helped improving the paper.

References

- Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational Convective-Scale Numerical Weather Prediction with the COSMO Model: Description and Sensitivities, *Monthly Weather Review*, 139, 3887–3905, <https://doi.org/10.1175/MWR-D-10-05013.1>, 2011.
- 5 Battaglia, A., Mroz, K., Lang, T., Tridon, F., Tanelli, S., Tian, L., and Heymsfield, G. M.: Using a multiwavelength suite of microwave instruments to investigate the microphysical structure of deep convective cores, *Journal of Geophysical Research: Atmospheres*, 121, 9356–9381, <https://doi.org/10.1002/2016JD025269>, 2016.
- Bony, S. and Stevens, B.: Measuring area-averaged vertical motions with dropsondes, *Journal of the Atmospheric Sciences*, 76, 767–783, <https://doi.org/10.1175/JAS-D-18-0141.1>, 2019.
- 10 Bony, S., Stevens, B., Frierson, D., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T., Sherwood, S., Siebesma, A., Sobel, A., Watanabe, M., and Webb, M.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261–268, <https://doi.org/10.1038/ngeo2398>, 2015.
- Bony, S., Stevens, B., Ament, F., Bigorre, S., Chazette, P., Crewell, S., Delanoë, J., Emanuel, K., Farrell, D., Flamant, C., Gross, S., Hirsch, L., Karstensen, J., Mayer, B., Nuijens, L., Ruppert, J. H., Sandu, I., Siebesma, P., Speich, S., Szczap, F., Totems, J., Vogel, R., Wendisch, M., and Wirth, M.: EUREC4A: A Field Campaign to Elucidate the Couplings Between Clouds, Convection and Circulation, *Surveys in*
- 15 *Geophysics*, 38, 1529–1568, <https://doi.org/10.1007/s10712-017-9428-0>, 2017.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Clouds and Aerosols, book section 7, p. 571–658, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.016>, www.climatechange2013.org, 2013.
- 20 Byers, H. R. and Hall, R. K.: A census of cumulus-cloud height versus precipitation in the vicinity of puerto rico during the winter and spring of 1953–1954, *Journal of Meteorology*, 12, 176–178, [https://doi.org/10.1175/1520-0469\(1955\)012<0176:ACOCCH>2.0.CO;2](https://doi.org/10.1175/1520-0469(1955)012<0176:ACOCCH>2.0.CO;2), 1955.
- Cadeddu, M., Turner, D., and Liljegren, J.: A Neural Network for Real-Time Retrievals of PWV and LWP From Arctic Millimeter-Wave Ground-Based Observations, *IEEE Transactions on Geoscience and Remote Sensing*, 47, 1887–1900, <https://doi.org/10.1109/tgrs.2009.2013205>, 2009.
- 25 Cadeddu, M. P., Marchand, R., Orlandi, E., Turner, D. D., and Mech, M.: Microwave Passive Ground-Based Retrievals of Cloud and Rain Liquid Water Path in Drizzling Clouds: Challenges and Possibilities, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 6468–6481, <https://doi.org/10.1109/tgrs.2017.2728699>, 2017.
- Crewell, S. and Löhnert, U.: Accuracy of cloud liquid water path from ground-based microwave radiometry 2. Sensor accuracy and synergy, *Radio Science*, 38, 7–1–7–3, 2003.
- 30 Crewell, S., Ebell, K., Löhnert, U., and Turner, D. D.: Can liquid water profiles be retrieved from passive microwave zenith observations?, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036934>, 2009.
- Elsaesser, G. S., O’Dell, C. W., Lebsock, M. D., Bennartz, R., Greenwald, T. J., and Wentz, F. J.: The Multisensor Advanced Climatology of Liquid Water Path (MAC-LWP), *Journal of Climate*, 30, 10 193–10 210, <https://doi.org/10.1175/JCLI-D-16-0902.1>, 2017.
- Ewald, F., Kölling, T., Baumgartner, A., Zinner, T., and Mayer, B.: Design and characterization of specMACS, a multipurpose hyperspectral
- 35 cloud and sky imager, *Atmospheric Measurement Techniques*, 9, 2015–2042, <https://doi.org/10.5194/amt-9-2015-2016>, 2016.

- Fricke, C., Ehrlich, A., Jäkel, E., Bohn, B., Wirth, M., and Wendisch, M.: Influence of local surface albedo variability and ice crystal shape on passive remote sensing of thin cirrus, *Atmospheric Chemistry and Physics*, 14, 1943–1958, <https://doi.org/10.5194/acp-14-1943-2014>, 2014.
- Gödde, F.: Detecting Clouds in the Presence of Sunlint: An Approach Using Spectral Water Vapor Absorption, Master's thesis, Ludwig-Maximilians-University Munich, https://macsserver.physik.uni-muenchen.de/campaigns/documents/theses/master_thesis_felix_goedde_2018.pdf, 2018.
- Greenwald, T. J., Bennartz, R., Lebsock, M., and Teixeira, J.: An Uncertainty Data Set for Passive Microwave Satellite Observations of Warm Cloud Liquid Water Path, *Journal of Geophysical Research: Atmospheres*, 123, 3668–3687, <https://doi.org/10.1002/2017JD027638>, 2018.
- Hogan, R. J., Ferro, C. A. T., Jolliffe, I. T., and Stephenson, D. B.: Equitability Revisited: Why the "Equitable Threat Score" Is Not Equitable, *Weather and Forecasting*, 25, 710–726, <https://doi.org/10.1175/2009WAF2222350.1>, 2010.
- Jacob, M., Ament, F., Gutleben, M., Konow, H., Mech, M., Wirth, M., and Crewell, S.: Liquid water path and integrated water vapor over the tropical Atlantic during NARVAL-South, https://doi.org/10.26050/WDCC/HALO_measurements_5, 2019a.
- Jacob, M., Ament, F., Gutleben, M., Konow, H., Mech, M., Wirth, M., and Crewell, S.: Liquid water path and integrated water vapor over the tropical Atlantic during NARVAL2, https://doi.org/10.26050/WDCC/HALO_measurements_6, 2019b.
- Klepp, C., Ament, F., Bakan, S., Hirsch, L., and Stevens, B.: NARVAL Campaign Report, Reports on Earth System Science 164, Max-Planck-Institut für Meteorologie, http://www.mpimet.mpg.de/fileadmin/publikationen/Reports/WEB_BzE_164_last.pdf, 2014.
- Klocke, D., Brück, M., Hohenegger, C., and Stevens, B.: Rediscovery of the doldrums in storm-resolving simulations over the tropical Atlantic, *Nature Geoscience*, 10, 891–896, <https://doi.org/10.1038/s41561-017-0005-4>, 2017.
- Konow, H., Jacob, M., Crewell, S., Ewald, F., Hagen, M., Hirsch, L., Jansen, F., Mech, M., and Stevens, B.: A unified data set of airborne cloud remote sensing using the HALO Microwave Package (HAMP), *Earth System Science Data Discussions*, <https://doi.org/10.5194/essd-2018-116>, 2018a.
- Konow, H., Jacob, M., Ament, F., Crewell, S., Ewald, F., Hagen, M., Hirsch, L., Jansen, F., Mech, M., and Stevens, B.: HALO Microwave Package measurements during Next-generation Remote sensing for VALidation Studies - South (NARVAL-South), https://doi.org/10.1594/WDCC/HALO_measurements_2, 2018b.
- Konow, H., Jacob, M., Ament, F., Crewell, S., Ewald, F., Hagen, M., Hirsch, L., Jansen, F., Mech, M., and Stevens, B.: HALO Microwave Package measurements during Next-generation Remote sensing for VALidation Studies 2 (NARVAL2), https://doi.org/10.1594/WDCC/HALO_measurements_3, 2018c.
- Krautstrunk, M. and Giez, A.: The Transition From FALCON to HALO Era Airborne Atmospheric Research, in: *Atmospheric Physics*, pp. 609–624, Springer Berlin Heidelberg, https://doi.org/10.1007/978-3-642-30183-4_37, https://doi.org/10.1007/978-3-642-30183-4_37, 2012.
- Lebsock, M. D., L'Ecuyer, T. S., and Stephens, G. L.: Detecting the ratio of rain and cloud water in low-latitude shallow marine clouds, *Journal of Applied Meteorology and Climatology*, 50, 419–432, <https://doi.org/10.1175/2010JAMC2494.1>, 2011.
- Liljegren, J. C., Boukabara, S.-A., Cady-Pereira, K., and Clough, S. A.: The effect of the half-width of the 22-GHz water vapor line on retrievals of temperature and water vapor profiles with a 12-channel microwave radiometer, *Geoscience and Remote Sensing, IEEE Transactions on*, 43, 1102–1108, 2005.
- Liu, G., Curry, J. A., Haggerty, J. A., and Fu, Y.: Retrieval and characterization of cloud liquid water path using airborne passive microwave data during INDOEX, *Journal of Geophysical Research: Atmospheres*, 106, 28 719–28 730, <https://doi.org/10.1029/2000JD900782>, 2001.

- Liu, Q., Weng, F., and English, S. J.: An Improved Fast Microwave Water Emissivity Model, *IEEE T. Geosci. Remote*, 49, 1238–1250, <https://doi.org/10.1109/TGRS.2010.2064779>, 2011.
- Lohmann, U. and Neubauer, D.: The importance of mixed-phase and ice clouds for climate sensitivity in the global aerosol–climate model ECHAM6-HAM2, *Atmospheric Chemistry and Physics*, 18, 8807–8828, 2018.
- 5 Maahn, M., Löhnert, U., Kollias, P., Jackson, R. C., and McFarquhar, G. M.: Developing and Evaluating Ice Cloud Parameterizations for Forward Modeling of Radar Moments Using In Situ Aircraft Observations, *Journal of Atmospheric & Oceanic Technology*, 32, 880–903, <https://doi.org/10.1175/JTECH-D-14-00112.1>, 2015.
- Marshall, J. S. and Palmer, W. M. K.: The Distribution of Raindrops with Size, *Journal of Meteorology*, 5, 165–166, <https://doi.org/10/fdzhp6>, 1948.
- 10 Maschwitz, G., Löhnert, U., Crewell, S., Rose, T., and Turner, D. D.: Investigation of ground-based microwave radiometer calibration techniques at 530 hPa, *Atmospheric Measurement Techniques*, 6, 2641–2658, <https://doi.org/10.5194/amt-6-2641-2013>, 2013.
- Masunaga, H., Nakajima, T. Y., Nakajima, T., Kachi, M., and Suzuki, K.: Physical properties of maritime low clouds as retrieved by combined use of Tropical Rainfall Measuring Mission (TRMM) Microwave Imager and Visible/Infrared Scanner 2. Climatology of warm clouds and rain, *Journal of Geophysical Research: Atmospheres*, 107, 2002.
- 15 Mears, C. A., Wang, J., Smith, D., and Wentz, F. J.: Intercomparison of total precipitable water measurements made by satellite-borne microwave radiometers and ground-based GPS instruments, *Journal of Geophysical Research: Atmospheres*, 120, 2492–2504, <https://doi.org/10.1002/2014JD022694>, 2015.
- Mech, M., Orlandi, E., Crewell, S., Ament, F., Hirsch, L., Hagen, M., Peters, G., and Stevens, B.: HAMP – the microwave package on the High Altitude and Long range research aircraft (HALO), *Atmospheric Measurement Techniques*, 7, 4539–4553, [https://doi.org/10.5194/amt-](https://doi.org/10.5194/amt-7-4539-2014)
- 20 7-4539-2014, 2014.
- Painemal, D., Greenwald, T., Cadetdu, M., and Minnis, P.: First extended validation of satellite microwave liquid water path with ship-based observations of marine low clouds, *Geophysical Research Letters*, 43, 6563–6570, 2016.
- Rosenkranz, P. W.: Water vapor microwave continuum absorption: A comparison of measurements and models, *Radio Science*, 33, 919–928, <https://doi.org/10.1029/98RS01182>, 1998.
- 25 Schnitt, S., Orlandi, E., Mech, M., Ehrlich, A., and Crewell, S.: Characterization of Water Vapor and Clouds During the Next-Generation Aircraft Remote Sensing for Validation (NARVAL)-South Studies, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10, 3114–3124, <https://doi.org/10.1109/JSTARS.2017.2687943>, 2017.
- Sherwood, S. C., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing, *Nature*, 505, 37–42, <https://doi.org/10.1038/nature12829>, 2014.
- 30 Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z., Illingworth, A. J., O’connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., and Mitrescu, C. a.: THE CLOUDSAT MISSION AND THE A-TRAIN, *Bulletin of the American Meteorological Society*, 83, 1771–1790, <https://doi.org/10.1175/BAMS-83-12-1771>, 2002.
- Stevens, B., Farrell, D., Hirsch, L., Jansen, F., Nuijens, L., Serikov, I., Brüggmann, B., Forde, M., Linne, H., Lonitz, K., and Prospero, J. M.: The Barbados Cloud Observatory: Anchoring Investigations of Clouds and Circulation on the Edge of the ITCZ, *Bulletin of the American Meteorological Society*, 97, 787–801, <https://doi.org/10.1175/BAMS-D-14-00247.1>, 2016.
- 35 Stevens, B., Brogniez, H., Kiemle, C., Lacour, J.-L., Crevoisier, C., and Kiliani, J.: Structure and Dynamical Influence of Water Vapor in the Lower Tropical Troposphere, *Surveys in Geophysics*, 38, 1371–1397, <https://doi.org/10.1007/s10712-017-9420-8>, 2017.

- Stevens, B., Ament, F., Bony, S., Crewell, S., Ewald, F., Gross, S., Hansen, A., Hirsch, L., Jacob, M., Kölling, T., Konow, H., Mayer, B., Wendisch, M., Wirth, M., Wolf, K., Bakan, S., Bauer-Pfundstein, M., Brueck, M., Delanoë, J., Ehrlich, A., Farrell, D., Forde, M., Göttsche, F., Grob, H., Hagen, M., Jäkel, E., Jansen, F., Klepp, C., Klingebiel, M., Mech, M., Peters, G., Rapp, M., Wing, A. A., and Zinner, T.: A high-altitude long-range aircraft configured as a cloud observatory – the NARVAL expeditions, *Bulletin of the American Meteorological Society*, ahead of print, available online, <https://doi.org/10.1175/BAMS-D-18-0198.1>, 2019, in press.
- 5 Turner, D. D., Vogelmann, A. M., Austin, R. T., Barnard, J. C., Cady-Pereira, K., Chiu, J. C., Clough, S. A., Flynn, C., Khaiyer, M. M., Liljegren, J., Johnson, K., Lin, B., Long, C., Marshak, A., Matrosov, S. Y., McFarlane, S. A., Miller, M., Min, Q., Minimis, P., O’Hirok, W., Wang, Z., and Wiscombe, W.: Thin Liquid Water Clouds: Their Importance and Our Challenge, *Bulletin of the American Meteorological Society*, 88, 177–190, <https://doi.org/10.1175/BAMS-88-2-177>, 2007.
- 10 Turner, D. D., Cadetdu, M. P., Löhnert, U., Crewell, S., and Vogelmann, A. M.: Modifications to the water vapor continuum in the microwave suggested by ground-based 150-GHz observations, *Geoscience and Remote Sensing, IEEE Transactions on*, 47, 3326–3337, <https://doi.org/10.1109/TGRS.2009.202262>, 2009.
- Vaisala: Vaisala RD94 Dropsonde Datasheet, Technical data, Vaisala, <https://www.vaisala.com/sites/default/files/documents/RD94-Datasheet-B210936EN-B.pdf>, 2017.
- 15 van Meijgaard, E. and Crewell, S.: Comparison of model predicted liquid water path with ground-based measurements during CLIWA-NET, *Atmospheric Research*, 75, 201 – 226, <https://doi.org/10.1016/j.atmosres.2004.12.006>, 2005.
- Wentz, F. J. and Meissner, T.: Algorithm Theoretical Basis Document (ATBD), Version 2, AMSR Ocean Algorithm, Tech. rep., Remote Sensing Systems, http://images.remss.com/papers/rsstech/2000_121599A-1_Wentz_AMSR_Ocean_Algorithm_ATBD_Version2.pdf, 2000.
- Winker, D. M., Hunt, W. H., and McGill, M. J.: Initial performance assessment of CALIOP, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL030135>, 2007.
- 20 Wirth, M., Fix, A., Mahnke, P., Schwarzer, H., Schrandt, F., and Ehret, G.: The airborne multi-wavelength water vapor differential absorption lidar WALES: system design and performance, *Applied Physics B*, 96, 201–213, <https://doi.org/10.1007/s00340-009-3365-7>, 2009.
- Zhou, L., Liu, Q., Liu, D., Xie, L., Qi, L., and Liu, X.: Validation of MODIS liquid water path for oceanic nonraining warm clouds: Implications on the vertical profile of cloud water content, *Journal of Geophysical Research: Atmospheres*, 121, 4855–4876, 2016.
- 25 Zuidema, P., Westwater, E. R., Fairall, C., and Hazen, D.: Ship-based liquid water path estimates in marine stratocumulus, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2005JD005833>, 2005.