

<i>Reviewer #1 Comments, advice and requests</i>	<i>Authors' full compliance, response, and thanks</i>
<p><i>Since submitting responses to the reviewers, the authors have made some further changes in the spirit of the requests. A couple of Figure reference numbers have been corrected. They are recurrent features of Microsoft Word idiosyncrasies (sorry). Format should be more similar to Copernicus standards.</i></p>	
<p>There is no clear outline presented in the manuscript, so I had to create my own (copied below) in order to grasp the manuscript completely. The manuscript includes two separate methodology sections (Section 2 and 6) and two separate theory sections (Section 4 and 5), and their arrangement and transitions left me frequently confused.</p>	<p><i>We have included "An Outline of This Paper" immediately after the first paragraph, following the style suggested by R1. We will reformat this as AMT will ultimately decide. We believe that leads us to omit paragraphs Section 1.1, L120–L140, which are over-detailed superfluous.</i></p>
<p>Additionally, there are many instances of parenthetical asides, notes, and comments (e.g., L362–365, L399–409, L422–428, all of Section 6.4) that interrupt the flow of the manuscript and greatly impede its overall understandability.</p>	<p><i>These instances and one other have been replaced by a named section of the Supplementary Material, e.g. "See also SM for a Note on Initial Point." at L362. I have attempted to make all such references minimally disruptive to the flow of the paper. Following AMT guidelines, they are not fundamental to advancing the arguments of the paper.</i></p>
<p>The conversational tone of this manuscript additionally introduces confusion. For instance, L203 states "We now move to..." and it's unclear if this means in the following paragraphs or in the next section. In L312 the phrase "Recall that..." is unclear.</p>	<p><i>These are restated.:</i>  <i>L203: The next section provides motivation for and understanding of an alternate approach ...</i>  <i>L312: With this section, we illustrate tracer relationships that define our approach to EnRs and EFs in more detail and also in more difficult circumstances, e.g., where the MCE is difficult to estimate, for example because its range of applicability during continued flight sampling is not clear.</i></p>
<p>Also, the included figures are very difficult to understand, in part because their text, captions, and legends are frequently too small to read (esp. Figures 4, 8, and 9) and because full explanations of what are in the figures are found both within the figure captions themselves and within various portions of the manuscript body.</p>	<p><i>The figures have been largely redrafted to have larger text. Figures 4 and 9 have been redrafted to show labels more clearly. (An remaining error on some time markings will be corrected.) Explanations of Figure 4 are expanded:</i></p> <p><b>Figure 4.</b> (a) Timeline of sampling, for the period shown in Figure 3(a), Montana, of CO<sub>2</sub>+CO (blue, left axis) and the fire</p>

Overall, these makes the manuscript difficult to follow and the presented scientific concepts and results difficult to understand.

tracers CO and  $b_{\text{scat}}$  (red and green points, right axis). Orange-filled points were identified as clear plume points. Unfilled points were not, but might have some fire influence, especially near plume points. (b) scatter diagram of CO vs CO<sub>2</sub>+CO with arrows showing the time progression of aircraft sampling of identified plume points. Colors provide a key to times shown in (a). Light gray numerals give observation times in minutes. (c) a similar diagram of  $b_{\text{scat}}$  vs CO<sub>2</sub>+CO. Similar shapes of figures are noted in the text. (d) Timeline of sampling for the period shown in Figure 3(b), Coastal Transect. (e) scatter diagram of CO vs CO<sub>2</sub>+CO during the transect, like (b). (f) a similar diagram of  $b_{\text{scat}}$  vs CO<sub>2</sub>+CO for the Coastal Transect. The black bars graphed in (a) and (d) are estimates of non-fire influenced  $C_{\text{bkgd}}$ , see text. They and the non-plume points suggest air-mass changes in CO<sub>2</sub>+CO.

*Figure 9 has similar changes and changes to the caption*

**Figure 9.** Analyzed relation of tracers to carbon burned using the MERET technique for portions of SEAC4RS Flight 10 and ARCTAS Flight 14. Compare (a) through (d) with Figure 4, (b), (c), (e), and (f). Colors key the observations to times shown in the timelines, Figures 4(a) and 4(c). Light gray numerals give observation times in minutes.

*Figure 8 has been made larger, and a large display in the published paper is recommended. When points representing different tracers overlap in the figure, this truly signals something about the excellent precision of the individual measurements, and we do not attempt to distinguish them. The figure caption has been expanded:*

**Figure 8.** (Lower panel). Estimates of the 422 background  $\hat{x}_i^0 = \text{CO}_2 + \text{CO}$  concentrations implied based on the 8 fire tracers indicated in the legend. Contributing individual estimates  $\hat{x}_{ij}^0$  are shown by overlapping colored points, with the median estimate

	<p><math>\hat{x}_i^0</math> indicated by a black bar. Usually the colored points overlap closely, this indicates strong agreement (Upper panel) Estimates of <math>C_{\text{burn}} = x_i - \hat{x}_i^0</math> indicators of fuel carbon burned, in the heavy green line. The preliminary estimate of <math>C_{\text{burn}}</math> based on the consensus of tracer deviations (without variable EnR estimates) is also shown in a light line. A scale factor, maximizing overlap with the heavy line, was necessarily estimated by regression. Flight days are indicated by the days marked on the top axes, and individual plumes, separated by non-plume concentrations of longer than 10 minutes, are shown as vertical separator lines. A set of horizontal lines at ~400 ppm indicates selected intervals for optimizing numerics (see text, Section 6.3, item 7).</p>
<p>I feel that there are two different manuscripts here, or at least one manuscript with a large appendix or supplement that includes the majority of the theory (Sections 4, 5, pages 12 – 19). The forthcoming paper (Chatfield and Andreae (2019) appears to be a useful companion to this manuscript, and it is referenced several times (e.g., L669-672), but it is unclear if the two papers are meant to be considered together or if they are stand-alone manuscripts.</p> <p>While I believe that this manuscript has significant scientific value and falls within the scope of AMT, and that the work described and methodology proposed (the MERET method) has substantial value, the current structure and length imposes a significant impediment on its understandability and impact. There were many times in which I was confused or lost, and so while I feel like I understand much of what was presented, I am not confident that the manuscript has successfully communicated all that the authors intended. As such, I feel that</p>	<p><i>This is well-considered, but the authors find few other options. We have put much more into the Supplementary Material.</i></p> <p><i>Consequently,</i></p> <p><i>(a) Material not strictly necessary has been moved to the material.</i></p> <p><i>(b) A table of contents has been included, following the reviewer's first comment and suggestion above.</i></p> <p><i>(c) The fact that the paper contains a development of plume theory is more prominent in the abstract: A new theoretical development of plume theory for multiple tracers is developed after examining the aircraft samples</i></p> <p><i>If the editors of AMT allow, we could change the title to: Theory and Estimation of Emissions Relationships in Forest Fire Plumes: 1: Reducing Effect of Mixing Errors on Emission Factors</i></p>

<p>significant reorganization and clarification is needed before this can be recommended for publication.</p>	<p><i>(d) The authors do not think that the theory could stand alone without showing that it leads to apparently good statistical estimates, and are unwilling to begin the whole AMT review process again if we suggest a division.</i></p>
<p>The scientific value of understanding forest-fire plume properties, and in particular of quantifying the enhancement ratios (EnRs) for properties of interest via the MERET method, is very high and this manuscript is a significant contribution to the field. The descriptions of the relationships between EnRs, ERs, and EFs in Section 1 is informative, although it would be particularly valuable if additional descriptions of how EnRs “approximate emission ratios (ERs)” (L77) if they are sampled before atmospheric transformations can occur. What is the relation after transformations? This needs to be made clear in the introduction.</p>	<p><i>Besides rewriting the paragraph, we have added a note to the Supplementary Material which clarifies this: ‘More on the relationships of EnRs, ERs, and EFs is found in the Supplementary Material (SM), “Note on EnRs and ERs”. ‘ The reviewer appears to want more information about when ERs can be larger or smaller than EnRs. This seemed appropriate for a note. A helpful suggestion!</i></p>
<p>The interpretation of Figures 4b,c,e,f in Section 3 is extremely valuable, but I largely struggled with understanding what was being represented until the description of the different examples later in the manuscript (esp. Sections 4.2 and 5). Only on a second read-through was I able to follow the text and more completely understand what is presented in Figure 4.</p>	<p><i>We thank the reviewer for this observation. We have rewritten the introductory paragraph:</i></p> <p>This section provides some examples of <math>C_{\text{tot}}</math> and fire tracers. It illustrates the limitations of changes in <math>C_{\text{tot}}</math> along a sampling path as an indicator of fire influence, <math>C_{\text{burn}}</math>, for emissions estimation and the much greater similarities of the such changes of tracers that possess shorter transformation time-scales. These define our approach to EnRs and EFs. The relation of fire emissions to observed <math>C_{\text{tot}}</math> to <math>C_{\text{burn}}</math>, can be apparently simple or complex, depending on how the history of non-fire CO and CO<sub>2</sub> entrained into fire plume air parcels affects <math>C_{\text{tot}}</math>. We show this commonality of relationships will to motivate the theory of expanding plumes in Section 4. That theory will suggest a method worked out in Sections 5 and 6 to find the key variable, <math>C_{\text{bkgd}}</math>, that then provides <math>C_{\text{burn}}</math> and thus EnRs.</p>



	<p><i>We have also edited several places succeeding paragraph, not described here..</i></p>
<ul style="list-style-type: none"> <li>• L54: “Chatfield and Andreae (2017)” should be “Chatfield and Andreae (2019, in preparation)”</li> <li>• L66: “DCO<sub>tot</sub>” should be “DC<sub>tot</sub>”.</li> <li>• Table 1: The line labeled “Proportional to carbon burned: define” is confusing. What does define mean here? Is this a typo?</li> <li>• Figure 2 refers to a slope of 32.60458 while the text (L299) refers to a slope of <math>33 \times 10^{-3}</math>. This inconsistency is confusing.</li> <li>• The variable <math>C_j</math> used in L417-418 and other lines does not appear in the Table of Symbols (Table 2) and is only described on L418</li> <li>• L425: “...the same plume. <i>provided we...</i>” is confusing</li> <li>•</li> <li>•</li> <li>• Figure 6 has an x-axis label of <math>C_{tot}</math> while the text (L469) refers to <math>C_{burn}</math></li> <li>• L659: “However, we let the define the types...” seems to be missing a word.</li> <li>• I believe “Figure 9” on Line 733 should be “Figure 8”</li> </ul>	<p>✓</p> <p><math>\Delta C_{tot} = \Delta CO_2 + \Delta CO</math></p> <p><i>Yes, a typo. Now</i>  <i>Proportional to total burned material, as measured by <math>C_{burn}</math></i></p> <p><i>Chose ppb/ppm rather than ppm/ppm</i></p> <p><i>Included.</i></p> <p><i>Changed. Remarks placed in the Supplement. After the equation (12) we now have: For periods of expansion in which the entrained concentrations are constant. See also SM: Note on Varying Entrainment</i></p> <p><i>Changed.</i></p> <p>However, we let the statistical technique define these types,</p> <p><i>Changed</i></p>
<ul style="list-style-type: none"> <li>• The variable <math>C_j</math> used in L417-418 and other lines does not appear in the Table of Symbols</li> </ul> <p>(Table 2) and is only described on L418</p> <ul style="list-style-type: none"> <li>• L425: “...the same plume. <i>provided we...</i>” is confusing</li> </ul>	<p><i>This has been added. The variable is the constant of integration and is generally replaced by <math>\alpha_j = \exp(C_j)</math></i></p> <p><i>Now in Supplementary Material, This now uses alpha and beta for different possible positions, values of i, and re-worded “<math>\alpha</math> and <math>\beta</math>, in the same plume.</i></p>

<ul style="list-style-type: none"> <li>•</li> <li>•</li> <li>• Figure 6 has an x-axis label of <math>C_{\text{tot}}</math> while the text (L469) refers to <math>C_{\text{burn}}</math></li> <li>•</li> <li>• L659: “However, we let the define the types...” seems to be missing a word.</li> <li>•</li> <li>• I believe “Figure 9” on Line 733 should be “Figure 8”</li> <li>•</li> </ul>	<p><i>These are supposed chosen so that we know that <math>x^E</math> and all the <math>y_j^E</math> remain coanstant.</i></p> <p><i>Both <math>C_{\text{tot}}</math> and <math>C_{\text{burn}}</math> are used. The x axis has <math>C_{\text{tot}}</math> , units, while the increment beyond the vertical axis at 380, shows <math>C_{\text{burn}}</math>. This is now indicated.</i></p> <p>“However, we let the statistical technique define these types, and so apply basic clustering techniques.” We also added a sentence soon afterward: “NMF and <i>k-means</i> clustering are shown to be equivalent in cases appropriate to our work (Ding et al., 2005).”</p> <p><i>Yes, Figure 8, thank you!</i></p>
<ul style="list-style-type: none"> <li>• The phrase “affine dependence” is used several times L522-523: The suggestion that the reader should make their own calculations in order to understand the linear responses is unhelpful.</li> <li>•</li> </ul>	<p><i>True. “Linear” has several meaning in English. See Wikipedia. So we have now “An affine dependence (linear polynomial relationship including an intercept). Linear Transformations in linear algebra must omit the intercept, hence the unusual phrasing “linear polynomial.” Chatfield has considered the ramifications of this dependence NMF linear transformations considerably.</i></p> <p><i>The reader is relieved of calculations now:</i></p> <p>“Some similar calculations make it clear that the estimates respond in an appropriate averaging manner under varied assumptions.” We simply emphasize the linearity of the analysis.</p>
<ul style="list-style-type: none"> <li>• (e.g., L145) and is unfamiliar to me.</li> <li>• In Section 1.2, there are many places where I get lost. For instance, the equation on L168 lacks a sufficient description and I’m unsure what the “<math>a_{j \leftarrow \text{CO}}</math>” and</li> </ul>	<p><i>We have added an explanatory phrase: “the slope <math>a_{j \leftarrow \text{CO}}</math> of the regression estimates of an EnR of the species with respect to CO, multiplied by an attempted very careful estimate of the slope <math>a_{\text{CO} \leftarrow (\text{fire-added } C_{\text{burn}})}</math> EnR of CO with respect to fire-</i></p>

<p>“<math>a_{CO} \leftarrow (\text{fire} - \text{added CO}_2 + \text{CO})</math>” terms mean. I feel a more complete explanation is needed.</p> <ul style="list-style-type: none"> <li>• The use of the variable <math>x</math> for <math>C_{tot}</math> in Section 1.2 and other places is confusing, especially when <math>C_{tot}</math> and <math>x</math> are used together (e.g., L153-158).</li> <li>•</li> </ul>	<p>produced <math>C_{burn}</math>. The <math>a_{CO} \leftarrow (\text{fire} - \text{added CO}_2 + \text{CO})</math> was described using the Modified Combustion Efficiency,...</p>
<p>L528-529: I do not understand what is meant by “provides safety against a variable and incompletely described background” or “The median is not affected by undetected changes in background...”</p> <ul style="list-style-type: none"> <li>•</li> </ul>	<p><i>A good observation. We also needed to explain why we were concerned about this. I have changed this to “This graph also suggests that if there are more than three tracers (we use 8), then the median of all the estimates, median (<math>\hat{x}_{ij}^0</math>), is robust against errors resulting if a tracer <math>j</math> has a variable or poorly described background resulting in <math>\hat{x}_{ij}^0</math> at falling distinctly higher or lower than the others. We must be concerned about this since tracers can have occasionally important non-fire sources.”</i></p>
<p>Figure 4 is extremely difficult to understand as there is almost no description in the caption itself; the descriptions and explanations are found within the text body. Specifically:</p> <ul style="list-style-type: none"> <li>• The text and images are very small <ul style="list-style-type: none"> <li>○ The label “<math>b_{scat}</math>” in Figure 4a,c is too small</li> <li>○ The number labels in Figure 4b,c,e,f are too small</li> </ul> </li> <li>• There are many individual components that are confusing</li> </ul>	<p><i>We have put a lot of time to address this remark.</i></p> <p><i>All figures have been redrawn with larger lettering. The sequence of times of observations should be much clearer for Figure 4 and Figure 9. See above for the wording of the section introduction and the expanded figure caption.</i></p>
<p><i>Since submitting responses to the reviewers, the authors have made some further changes in the spirit of the requests.</i></p>	



# Emissions Relationships in Western Forest Fire Plumes:

## I. Reducing the Effect of Mixing Errors on Emission Factors

Robert B Chatfield<sup>1</sup>, Meinrat O Andreae<sup>2,3</sup>, ARCTAS Science Team\* and SEAC4RS Science Team\*

<sup>1</sup> NASA Ames Research Center, Moffett Field, CA 94035, USA

<sup>2</sup> Max Planck Institute for Chemistry P.O. Box 3060, D-55020 Mainz, Germany

<sup>3</sup> Scripps Institution of Oceanography, UCSD, La Jolla, CA 92093, USA

\* A full list of authors and their affiliations appears at the end of the paper.

**Abstract:** Studies of emission factors from biomass burning using aircraft data complement the results of lab studies and extend them to conditions of immense hot conflagrations. A new theoretical development of plume theory for multiple tracers is developed after examining aircraft samples. We illustrate and discuss emission relationships for 422 individual samples from many forest-fire plumes in the Western US. samples are from two NASA investigations: ARCTAS (Arctic Research of the Composition of the Troposphere from Aircraft and Satellites) and SEAC4RS (Studies of Emissions and Atmospheric Composition, Clouds, and Climate Coupling by Regional Surveys). This work provides sample-by-sample enhancement ratios (EnRs) for 23 gases and particulate properties. Many EnRs provide candidates for emission ratios (ERs, corresponding to the EnR at the source) when the origin and degree of transformation is understood and appropriate. From these, emission factors (EFs) can be estimated, provided the fuel dry mass consumed is known or can be estimated using the carbon mass budget approach. This analysis requires understanding the interplay of mixing of the plume with surrounding air. Some initial examples emphasize that measured  $C_{\text{tot}} = \text{CO}_2 + \text{CO}$  in a fire plume does not necessarily describe the emissions of the total carbon liberated in the flames,  $C_{\text{burn}}$ . Rather, it represents  $C_{\text{tot}} = C_{\text{burn}} + C_{\text{bgd}}$ , which includes possibly varying background concentrations for entrained air. Consequently, we present a simple theoretical description for plume entrainment for multiple tracers from flame to hundreds of kilometers downwind and illustrate some intrinsic linear behaviors. The analysis suggests a Mixed Effects Regression Emission Technique (MERET), which can eliminate occasional strong biases associated with the commonly used normalized excess mixing ratio (NEMR) method. MERET splits  $C_{\text{tot}}$  to reveal  $C_{\text{burn}}$  by exploiting the fact that  $C_{\text{burn}}$  and all tracers respond linearly to dilution, while each tracer has consistent EnR behavior (slope of tracer concentration with respect to  $C_{\text{burn}}$ ). The two effects are separable. Two or three or preferably more emission indicators are required as a minimum; here we used eight. In summary, MERET allows fine spatial resolution (EnRs for individual observations) and comparison of similar plumes distant in time and space. Alkene ratios provide us with an approximate photochemical timescale. This allows discrimination and definition, by fire situation, of ERs, allowing us to estimate emission factors.

Deleted: The

Deleted: when

Deleted: ten

Deleted: Limited variations in the EnRs for each tracer can be incorporated and the variations and co-variations analyzed. The percentage CO yield (or the modified combustion efficiency) plays some role. Other co-relationships involving nitrogen and organic classes are more prominent; these have strong relationships to the  $C_{\text{burn}}$  to  $\text{O}_3$  emission relationship.

## 45 1. Introduction

### 1.1. Importance and previous work

Biomass burning has a large influence on the atmospheric burden of ozone and aerosols, and consequently also affects climate (Crutzen et al., 1979; Crutzen and Andreae; 1990; Jaffe and Wigder, 2012; Andreae, 2019). Biomass burning emission factors that are useful to drive photo-chemical models are most often estimated by one of two sampling techniques (Akagi et al., 2011). The first approach, measurements on the ground close to an open fire or on laboratory fires that are controlled to approximate natural conditions, can provide the most detailed information on sources. The burning conditions can be readily assessed and fit into parameterizations of the emissions process, provided the correct mix of burn types typical of large fires can be estimated. It can, however, be difficult to mimic and safely sample truly intense flaming conflagrations. The second, measurements made from aircraft, provides a much wider sample of different fires and emissions from different regions of a single fire. However, the estimates can be difficult to classify as simply “flaming” or “smoldering” or even as defined mixtures of just two types. Adjoining areas with fires in various stages of combustion can merge into the same plume, or remain relatively distinct. These questions of classifications related to the originating fires are addressed statistically in a succeeding paper, Chatfield and Andreae (2020).

This work presents a rationale for careful attention in estimation of emission relationships and emission factors in contrasting case studies, and uses such studies to develop an entraining plume theory for emissions relationships. Illustrations show that the theory is robust. This theory implies a statistical regression technique; a second methodology section then gives details of implementation given the complexities of atmospheric sampling. The result, a key “equivalent background” estimate, is then applied to quantify the atmospheric signal of fuel burned,  $\text{CO}_2 + \text{CO}$ ; this allows quantification of emission factors.

Since this work takes a complex path winding through observations, simple theory, examples regression methodology, basic results, and finally emission factors, we offer this table of contents:

#### 1. Introduction

##### 1.1. Importance of previous work

##### 1.2. Development of EF estimation to date

#### 2. Methodology: defining an indicator dataset

#### 3. Observed behavior of $C_{\text{tot}}$ in fire plumes – Properties of tracers

#### 4. Theory: expanding plume for several species

##### 4.1. A general relationship

##### 4.2. Examples showing robustness of computations of idealized $C_{\text{tot}}$

#### 5. Theory: A regression relationship for EnRs

#### 6. Methodology:

##### 6.1. Finding the $\text{CO}_2 + \text{CO}$ background

##### 6.2. Practicalities: variable EnRs

##### 6.3. Summary of the MERET method

##### 6.4. Number of independent samples

#### 7. Results: Estimation of $x_{\text{f0}}$ and $C_{\text{burn}}$

#### 8. Estimates of emissions ratios: Two MERET examples

Deleted: P

Deleted: W

Deleted: 2017

Formatted: Font color: Background 1

Formatted: Subscript

Formatted: Paragraph

Formatted: Space Before: 0 pt, After: 2 pt, Don't add space between paragraphs of the same style

Formatted: Font: Italic

Formatted: Subscript

Formatted: Font: Italic

Formatted: Subscript

Formatted: Subscript

### 8.1. MERET results for our two examples

### 8.2. Table of several significant emissions

## 9. Conclusions

### 9.1. Questions for future research

## End material and References.

A further guide is given at the end of Section 1.

Let us introduce our view of enhancement ratios, emission ratios, and emission factors. Under appropriately defined circumstances, the amount of fuel carbon burned that is liberated to the atmosphere is the sum of carbon added to the ambient air in the form of all fire-originated gases and particles as a result of combustion:

$$\Delta x = C_{\text{burn}} = \Delta \text{CO}_2 + \Delta \text{CO} + \Delta \text{CH}_4 + \Delta (\text{particulate carbon species}) + \Delta (\text{O}) \text{VOCs} \quad (1)$$

where the  $\Delta$ 's refer to the enhancement relative to pre-burn air.

In deriving emissions factors, i.e., how much of a species is emitted per kg of biomass burned, it is usual to obtain the amount of carbon burned from the sum of excess mixing ratios,  $C_{\text{burn}} = \text{CO}_2 + \text{CO} + \text{other carbon-containing emissions, including aerosol particles}$ . To an accuracy within  $\geq 1.5\%$  (totals from the datasets we analyzed) to  $3\%$  (Andreae and Merlet, 2001), carbon burned or  $C_{\text{burn}}$  is approximated by the excess ( $\text{CO}_2 + \text{CO}$ ), as measured above a background concentration,  $C_{\text{bgd}}$  (Andreae et al., 1988). With this understanding, we will assume  $\Delta C_{\text{tot}} = \Delta \text{CO}_2 + \Delta \text{CO}$  in the following. In some measurement situations, frequent and accurate measurements of  $\text{CH}_4$  and particulate  $C$  are also available, and their inclusion could perceptibly add to the precision of estimates. Analysis proceeds similarly with these terms.

We will thus define the quantity  $C_{\text{burn}}$  as  $C_{\text{burn}} = \Delta x = \Delta \text{CO}_2 + \Delta \text{CO}$ . Hence,  $C_{\text{bgd}} + C_{\text{burn}} = C_{\text{tot}}$ . A complication arises from the fact that pre-burn/non-burn air may have various compositions, especially when we consider various sources for low-level inflow air, and especially air that is entrained in the smoke plume by the time of sampling. This is an important topic, which has been discussed in detail by Guyon et al. (2005) and Yokelson et al. (2013) and that we will focus on below. An Enhancement Ratio (EnR) for a species or property  $j$  with mixing ratio  $y_j$  is then  $\text{EnR}_j = \Delta y_j / \Delta x$ .

We will use the term "enhancement ratio," EnR, in this paper. For many species, EnRs sampled prior to substantial atmospheric transformation (e.g., chemistry or particulate processes) approximate ERs. More on the relationships of EnRs, ERs, and EFs is found in the Supplementary Material (SM), "Note on EnRs and ERs". ER estimation constitutes the analysis of atmospheric samples that contribute to EF. Emission factors are defined relative to the amount of fuel burned and are derived from emission ratios by accounting for the concentration of carbon in the biomass burned and adjustment of units (Andreae and Merlet, 2001). Separate methods of land analysis are employed. EFs can be derived from ERs by

$$\text{EF}_j = \text{ER}_j \times \frac{MW_j}{MW_c} \times C_{\text{BM}} \quad (2)$$

where  $\text{ER}_j$  is the emission ratio of species  $j$ ,  $MW_j$  and  $MW_c$  are the molecular weight of species  $j$  and the atomic weight of carbon, respectively, and  $C_{\text{BM}}$  is the carbon content of the dry biomass. We focus on clarifying appropriate methods of finding EnRs and ERs, which enable EF estimation.

One part of EF estimation concerns the amount of fuel consumed in fires, its carbon content, and the fraction liberated to the atmosphere (i.e., excluding char remaining on the ground); here

Formatted: Indent: Left: 0", Space Before: 0 pt, After: 2 pt, Don't add space between paragraphs of the same style

Formatted: Font: Italic

Deleted: O

Deleted: A

Deleted: emission ratios

Deleted: (

Deleted: Rs) and thus allow estimates of emission factors,

Deleted: They

Formatted: Font: Italic

Deleted: For use of emission factors in modeling, the circumstances of emission need to be described clearly, and the ER must be a true result of emissions. This work is focused on


Deleted: ,

Deleted: and

145 we will focus on the other part of the question, which concerns the relationship of emitted com-  
pounds to the *C* liberated to the atmosphere. Many of the EnRs we calculate appear good candi-  
dates for EF estimates. One remaining task, making specific links of particular EFs to appropri-  
150 ate fire conditions for which they apply, requires individualized trajectories and fuel characteri-  
zations. This important task is beyond reasonable treatment in this publication, which focuses on  
the improving the understanding of airborne samples. It seems likely to us that uncertainties on  
the relation of area and fuel burned contribute more error to emissions estimates than those con-  
tributions of minor *C*-containing species in the plume that were described above.

155 There are other uses for EnRs that arise in understanding fire plumes, which revolve around  
the evolution of relatively fresh smoke plumes, e.g., the enhancement of ozone, peroxy acetyl ni-  
trate, or other bound (not NO or NO<sub>2</sub>) nitrogen species (Alvarado and Prinn, 2009, Alvarado et  
al., 2009; 2010, Jaffe and Widger, 2012). These also should have a direct relation the fuel ~~carbon~~  
~~burned and other properties such as burning conditions, fuel moisture, and fuel N~~ content.

160 Two special sampling intensives utilizing NASA's fully instrumented DC-8 aircraft allowed  
us to investigate forest-burning emissions. In June, 2008, the aircraft sampled a variety of fire  
plumes around California (Jacob et al., 2010; Singh et al., 2010; 2012; Hornbrook et al. 2011)  
during the California ARCTAS (Arctic Research of the Composition of the Troposphere from  
Aircraft and Satellites) intensive period. In a later part of the campaign, the DC-8 sampled in  
Northern Canada (Simpson et al., 2011); we excluded these plumes as representing different for-  
est burn conditions. In 2013, the DC-8 made several samplings of forest fires in California and  
165 the Rocky Mountain West during SEAC4RS (Studies of Emissions and Atmospheric Composi-  
tion, Clouds and Climate Coupling by Regional Surveys; Toon et al., 2015). We analyzed all of  
these fire plumes, but excluded samples east of 102 °W, which were mostly from agricultural  
fires. Our aim was to understand a variety of plumes, but limit variation to a single general cate-  
gory (temperate forest fires) as used for three-dimensional simulation models and geographical  
170 summaries.



Deleted: C

Deleted:

Deleted: perhaps

Formatted: Font: Italic

Deleted: ¶

**Moved down [5]:** Flight tracks for the period and locations of major fires during these periods are shown in Figure 1. Analysis of the vertical variation of fire tracers suggested that plumes below 5 km ASL included recent and informative fires in our study. We saw no unequivocal variation of composition with height, possibly due to limitations on aircraft maneuvers low and near the fires. Consequently, the aircraft samples likely cannot adequately represent ground-hugging smoke flows.



**Figure 1.** ARCTAS and SEAC4RS flights analyzed in this work. Each flight is identified by the flight number of that series. Flights 5 and 6 were in the Western US but not included.

Flight tracks for the period and locations of major fires during these periods are shown in Figure 1. Analysis of the vertical variation of fire tracers suggested that plumes below 5 km ASL included recent and informative fires in our study. We saw no unequivocal variation of composition with height, possibly due to limitations on aircraft maneuvers low and near the fires. Consequently, the aircraft samples likely cannot adequately represent ground-hugging smoke flows.

## 1.2. Development of EF estimation to date

EnRs and EFs for biomass burning plumes have largely been based on measurements of the CO<sub>2</sub> or CO concentrations in the plumes. Typical analyses begin with measurements of  $C_{\text{tot}}$  and the concentrations of several tracers we may call  $y_j$ ,  $j = 1, \dots, N_{\text{Tracers}}$ . Multiple instances  $i = 1, \dots, N_{\text{instances}}$  are observed, e.g., every few seconds or few minutes within a plume. An affine dependence (linear polynomial relationship including an intercept) is observed between each of the tracers and  $C_{\text{tot}}$  with a y-intercept that depends most significantly on the local out-of-plume background values of CO<sub>2</sub>, CO, and each tracer individually.

$$C_{\text{tot}} = C_{\text{bkgd}} + C_{\text{burn}} \quad (3)$$

The following analysis suggests several complexities that must be addressed in order to understand these affine relationships. Several aspects of slopes, intercepts, and deviations from linearity of the relationship of tracer  $y_j$  to  $C_{\text{tot}}$  plots must be examined, and so we transition to graphic terminology with  $x$  representing  $C_{\text{tot}}$ . Later we will describe measurements of  $C_{\text{tot}}$  and tracers  $j$  at a given instance  $i$ ,  $x_i$  and  $y_{ij}$ . For a simple plume within a homogeneous mixed layer characterized by an  $x$  concentration  $x^E$  and  $y$  concentration  $y^E$ , we write

$$x = x^E + (x - x^E) \quad (4)$$

and

$$(y_j - y_j^E) = a_j(x - x^E) \quad (5)$$

with enhancement ratios,  $a_j$ , that can yield EFs directly. Early estimations (e.g., Greenberg et al., 1984; Andreae et al. 1988) used plots and regressions against CO<sub>2</sub> to estimate EnRs and EFs. These earliest techniques assumed fire was the main origin of CO<sub>2</sub>. Very early it was recognized that other effects, e.g., variation of photosynthesis, respiration, and mixing required a more careful approach (e.g., Guyon et al., 2005). Alternatively (e.g., Andreae et al., 1988; Hobbs et al., 2003; Lefer et al., 1994), EnRs were derived with respect to CO. Symbolically,

$$\text{EnR estimate} = a_{j \leftarrow \text{CO}} \cdot a_{\text{CO} \leftarrow (\text{fire-added } C_{\text{burn}})}$$

the slope  $a_{j \leftarrow \text{CO}}$  of the regression estimates of an EnR of the species with respect to CO, multiplied by an attempted very careful estimate of the slope  $a_{\text{CO} \leftarrow (\text{fire-added } C_{\text{burn}})}$ . EnR of CO with respect to fire-produced  $C_{\text{burn}}$ . The  $a_{\text{CO} \leftarrow (\text{fire-added } \text{CO}_2 + \text{CO})}$  was described using the Modified Combustion Efficiency

$$\text{MCE} = \Delta \text{CO}_2 / (\Delta \text{CO}_2 + \Delta \text{CO}) = 1 - \Delta \text{CO} / (\Delta \text{CO}_2 + \Delta \text{CO}) = 1 - \text{EnR}_{\text{CO}}. \quad (6)$$

with a best estimate of the domain of points for which a constant MCE could be assumed.

The method has become known as the normalized excess mixing ratios method (NEMR; Akagi et al, 2011). Yokelson and others (2013) described the care required to make sure that the MCE was well defined; otherwise, severe difficulties ensue. They describe a situation in which

Formatted: Font: Bold

Deleted: 1

Formatted: Indent: First line: 0"

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Normal

Moved (insertion) [5]

Deleted: Flight tracks for the period and locations of major fires during these periods are shown in Figure 1. Analysis of the vertical variation of fire tracers suggested that plumes below 5 km ASL included recent and informative fires in our study. We saw no unequivocal variation of composition with height, possibly due to limitations on aircraft maneuvers low and near the fires. Consequently, the aircraft samples likely cannot adequately represent ground-hugging smoke flows. We begin this work with a general overview of data analysis using conventional methods. We provide motivation for a new method, using  $C_{\text{tot}}$  and combustion tracers in situations observed during the SEAC4RS mission. We then use these values to describe  $C_{\text{burn}}$  as the difference between  $C_{\text{tot}}$  and  $C_{\text{bkgd}}$ . This involves finding an "equivalent background value" of  $C_{\text{tot}}$ , a value which can be described as summarizing the amount of mixing-in of background  $C_{\text{tot}}$  diluting a sample. This is indicated by the agreement of all fire tracers employed. We describe the tracers we employed and the removal of urban influences. Two examples where naïve EnRs appear biased are described, one in Montana and one in Northern California. A theoretical section then gives a synthesis of tracer dilution in fire plumes and leads to a suggested empirical regression equation. Graphical examples of dilution histories emphasize the point that the regression is a useful summary of more complex dilution processes. The "Mixed Effects Regression Emission Technique" or MERET algorithm is then described, a method that allows for spec

Formatted ... [2]

Deleted: <#>With  $C_{\text{burn}}$  established, we give EnRs for the tracers in the many plume samples. Esti- [3]

Formatted ... [4]

Deleted: <#>e...timation to dd ... [5]

Deleted: , ...M ... [6]

Formatted ... [7]

Deleted: ,

Formatted: Paragraph

Deleted: , and a... multiplied by an attempted very mor [8]

Deleted:

Formatted: Font: 10 pt

Formatted: Font: 11 pt

Formatted: Normal

Formatted: Font: 12 pt

Deleted:

$x^E$  and  $y_{CO}^E$  in a diluting plume took on two distinct values, a mixed-layer value and a free-troposphere value, during plume rise and transport. More than two values may be relevant, emphasizing their call for careful sampling of pre-fire air and its dilution environment. We describe below new methods to resolve many of the difficulties with  $x^E$  and to indicate unwanted effects of  $y_{CO}^E$  variability. These methods could provide EnRs for many species with reasonable precision under more conditions.

This need for caution was very evident in the ARCTAS and SEAC4RS observational situations. Some Western USA data we analyzed showed variations in background  $C_{tot} = CO_2 + CO$  (away from direct recent effects of respiration and photosynthesis) of 15 ppm (interquartile Range of 4 ppm), while other Western USA regions showed variations of ~8 ppm (according to the analyses in this paper that we present later in Figure 4). The contributions from fires were often comparable to this variation, ~2 – 40 ppm, mean ~6 ppm. Air flowed from the west into forest fires at low altitudes, or later diluted the smoke plume at intermediate levels. We could expect background air with a variety of histories of influence by photosynthesis (lower resultant  $CO_2$ ) or respiration (higher  $CO_2$ ), or urban-influenced air (higher  $CO_2$ ). Low-level inflow air could have been mostly affected by local forests, farming, etc. Some of the most problematic situations tend to be associated with plumes sampled early in the day, when air from a nocturnal boundary layer — strongly enriched with respiration  $CO_2$  — is mixed into the smoke plumes (Guyon et al., 2005). There could also have been substantial variations in  $C_{tot}$  due to intercontinental transport, the composition reflecting long-previous modification due to these same processes and to latitudinal gradients. Yates et al. (2011) reported and more fully referenced atmospheric sampling of Western air showing variations in  $CO_2$  and also  $CH_4$  and  $O_3$ . On the east side of the Pacific Anticyclone, the common pattern was for descent and horizontal shearing displacements, producing substantial  $C_{tot}$  variations in both horizontal and vertical directions (Barry and Chorley, 1998).

Previous analyses have been made for the ARCTAS data, by Simpson et al. (2011) for the large Canadian fires sampled and by Hornbuckle et al. (2011) for all fires. The Hornbuckle article usefully complements this paper by describing features and origins of the plumes sampled. Both groups described novel methods, but followed the traditional CO-emissions-ratio or NEMR methodology (Andreae et al., 1988; Hobbs et al., 2003; Akagi et al., 2011). Pfister et al. (2011) considered the emissions and transport of CO in the California ARCTAS samples. Analyses for the SEAC4RS fires have also been reported (Liu, 2018).

~~The following sections provide motivation for and understanding of~~ an alternate approach to the description of EnRs and EFs, the Mixed Effects Regression Estimation Technique, MERET; in some cases, MERET and NEMR ~~may~~ form complimentary supporting views of plume emissions. Whereas NEMR depends on multiple measurements in the same plume in an understood environment, MERET is typically applicable to individual measurements of similar EnR-determining fire conditions across many different plumes. It does instead require several informative fire tracer species be measured simultaneously, not simply  $CO_2$  or CO, as well as the tracer whose EnR is desired. It can also be used for good candidate EFs when the environmental history of the plume is not well characterized. It is applicable to any plumes encountered, without need for extensive measurement of that plume history.

The MERET technique attempts to use the simultaneous variability, sample by sample, of a large set fire tracer compounds and aerosol descriptors to find a single quantification,  $C_{burn}$ , of fire emissions, which it splits from  $C_{bgd}$  such that the sum is  $C_{tot}$ . To do so, it must also ascribe a set of EnRs to the fire tracers, and recognize that these EnRs may vary from sample to sample in

Deleted: We now move to motivate

a limited way. The interplay of these estimates contributing to  $C_{\text{burn}}$  and EnRs for each observation appears daunting. Section 3 will graphically illustrate how strongly effects beyond fire emissions describe variations in  $C_{\text{tot}}$ , also how similar and informative are various tracers as graphed against  $C_{\text{tot}}$ . Section 4 will describe a theory of multiple fire emissions co-emitted from a fire based on familiar plume concepts, and give examples. The examples show the linearity of the theory that such simple approaches with a limited number of parameter estimates yield a reasonable approximation to more complex behavior. Sections 5 and 6 describe a mixed-effects regression algorithm based on plume theory. Section 7 provides a limited number of EnR estimates and describes graphically how flight segments describing similar emissions conditions can be identified.

## 2. Methodology: defining an indicator dataset

An initial task is the identification of appropriate variables and sampling rates. The technique we describe requires the measurement of  $C_{\text{tot}} \approx \text{CO}_2 + \text{CO}$  and several concentrations of emitted species or similar, extensive, properties of emissions (e.g., dried-airstream scattering coefficients,  $b_{\text{scat}}$ ), which we will call *emission indicator species* or tracers. A set of indicator species was chosen for this publication to enable deriving relevant EnRs and to support our initial classification (e.g., flaming, smoldering, high- $N$  fuel, etc.). It is important to have as many differently behaving emission indicator species as possible, as different indicators may respond differently to different fuels and fire intensities (“fire chemistries”), and such variations are usually not known before analysis. We favored indicator species with rapid sampling rates, so as to define  $C_{\text{burn}}$  for the maximum number of instances, but certain variables like CO, CH<sub>4</sub>, and  $b_{\text{scat}}$  had special claims, as they can be maximally expressed in important types of fires. For our samples, methane and methanol showed significant idiosyncracies. Their cumulative probability distribution differed from all other tracers: prominently very high concentrations and much higher positive skewness. We surmise that this behavior resulted from other prominent sources, e.g. cattle raising, or that very long distance transport and long lifetimes caused very great non-fire sources, like CO<sub>2</sub>. It was convenient to use these same frequently measured indicator variables to define  $C_{\text{burn}}$  and also for classification of fire chemistries. For classification, we added intensive variables, essentially ratios that should be physically independent of  $C_{\text{burn}}$ .

The emissions indicator species that satisfied these requirements for both missions are shown in Table 1, along with references to the measurement techniques and observers. Only extensive quantities (proportional to  $C_{\text{burn}}$ ) are used in this paper. CO<sub>2</sub> was measured by Stephanie Vay (ARCTAS) and Andreas Beyersdorff (SEAC4RS) using the AVOCET instrument (Vay et al., 2011). In examining EnRs for various species, we also use the organic aerosol (OA) measurements of Jose Jimenez and ionic composition information from the Jack Dibb group. The ARCTAS and SEAC4RS data sites give full information, as instrumentation characteristics naturally vary somewhat between missions. (<https://www-air.larc.nasa.gov/cgi-bin/ArcView/arctas>, <https://www-air.larc.nasa.gov/cgi-bin/ArcView/seac4rs?DC8=1>)

Our techniques use algorithms that currently allow few missing observations among the variables. The sampling rates for emission indicators measured by PTRMS (proton-transfer ionization mass spectrometry) differed between the two aircraft missions. The SEAC4RS mission acquired suitably complete PTRMS-derived datasets at a once-per-minute rate, and this defined the data interval used for both datasets. Additionally, in SEAC4RS CO was measured only by (less

Deleted: Consequently, section 3

Deleted: provide some examples of the special nature of this interplay

Deleted: .

Formatted: Indent: First line: 0"

Formatted: Subscript

Moved (insertion) [2]

frequent) can samples for the first flights prior to the Rim Fire of 26 August, and CH<sub>4</sub> was sampled only by cans for all flights. These are important species: CO is the most commonly used tracer for fire plumes because of its favorable plume-to-background concentration ratio and readily available measurement instrumentation. It is also used to define the MCE in much of the biomass-burning literature (Yokelson et al., 1996; Jaffe and Widger, 2011). Consequently, SEAC4RS imposed additional difficulties and processing. However, we judged it important to include SEAC4RS in a combined analysis to broaden the fire chemistries analyzed, as the Rim Fire was exceptionally large, hot, and well sampled.

Table 1. Indicator Variables

Concentration / property	Abbreviation	Technique	Group	Reference
<b>Extensive quantities</b> <i>Proportional to total burned material, as measured by C<sub>burn</sub></i>				
Toluene	C <sub>6</sub> H <sub>5</sub> CH <sub>3</sub>	PTRMS	Wisthaler	Wisthaler, et al. 2013.
Benzene	C <sub>6</sub> H <sub>6</sub>	PTRMS	Wisthaler	Wisthaler, et al. 2013.
Formaldehyde	HCHO	LAS	Fried	Fried et al., 2008.
Acetonitrile	CH <sub>3</sub> CN	PTRMS	Wisthaler	Wisthaler, et al. 2013.
Absorption Coefficient Dry, Total, 532 nm	<i>b<sub>abs</sub></i> , Abs_5	Nephelometry	Anderson	Wagner et al., 2015, Anderson Langley Aerosol Group, LARGE
Scattering Coefficient, Dry, Submicron 550 nm	<i>b<sub>scat</sub></i> , Scat_5	Nephelometry	Anderson	"
Carbon monoxide	CO	LAS, GC	Diskin, Blake	Pfister et al., 2011.
Acetaldehyde	CH <sub>3</sub> CHO	PTRMS	Wisthaler	Wisthaler, et al. 2013.
<b>Intensive quantities</b> <i>Not proportional to carbon burned</i>				
Single Scattering Albedo	SSA	Nephelometry	Anderson	Wagner et al., 2015, Anderson
Ångström Exponent, scattering	ÅE	Nephelometry	Anderson	"
<b>Other variables used</b>				
	O <sub>3</sub> , NO <sub>x</sub> = NO + NO <sub>2</sub> , NO <sub>y</sub>	Chemiluminescence, UV	Weinheimer (ARCTAS) Ryerson (SEAC4RS)	Weinheimer, et al. et al. 1994. Ryerson et al., 2000
Methane	CH <sub>4</sub>	LAS, GC	Diskin, Blake	Pfister et al., 2011.
Methanol	CH <sub>3</sub> OH	PTRMS	Wisthaler	Wisthaler, et al. 2013.

Notes: PTRMS: proton transfer mass spectrometry, LAS: laser absorption spectrometry.  $1 - \omega$  is the single-scatter co-albedo; likewise, CO is linked to  $1 - \omega$  (Modified Combustion Efficiency), so that all values extend upwards from 0. CO<sub>2</sub> measurements: see text.

The selection of fire plumes required some care. While CH<sub>3</sub>CN is a highly specific tracer of fires (Singh et al. 2012), detailed analysis suggests that it is not the best quantitative tracer. (Further analysis suggested that CH<sub>3</sub>CN has variable EFs, so it signals fires well, but does not quantify C<sub>burn</sub> adequately.) Plumes were characterized by levels of CH<sub>3</sub>CN above 0.225 ppb,

#### Moved (insertion) [3]

**Deleted:** In order to provide a suitably complete dataset for SEAC4RS, we used the can samples to infer likely concentrations at one-minute intervals of key species, i.e., CH<sub>4</sub> for all flights and CO for the first few flights, using available can samples at slightly lower frequency. The **R** package for multiple imputations by chained equations (*mice()*) was employed, using the whole data period, but filling in observations with missing data. (Our assessment of the effect of imputation was informal and is reviewed again below.) It was highly desirable to include the imputed concentrations of methane, since it is commonly measured and appears to be a prominent signal of different types of "fire chemistry", i.e., enhanced emission of reduced species; methanol and acetone are often correlated with CH<sub>4</sub> and give support to this idea.

**Deleted:** carbon burned

**Deleted:** define

**Deleted:** Methane ... [11]

**Deleted:** Methanol ... [12]

#### Formatted Table

**Moved up [2]:** The emissions indicator species that satisfied these requirements for both missions are shown in Table 1, along with references to the measurement techniques and observers. Only extensive quantities (proportional to C<sub>burn</sub>) are used in this paper. CO<sub>2</sub> was measured by Stephanie Vay (ARCTAS) and Andreas Beyersdorff (SEAC4RS) using the AVOCET instrument (Vay et al., 2011). In examining EnRs for various species, we also use the organic aerosol (OA) measurements of Jose Jimenez and ionic composition information from the Jack Dibb group. The ARCTAS and SEAC4RS data sites give full information, as instrumentation characteristics naturally vary somewhat between missions. (<https://www-air.larc.nasa.gov/cgi-bin/ArcView/arctas>, <https://www-air.larc.nasa.gov/cgi-bin/ArcView/seac4rs?DC8=1>) Our techniques use algorithms that currently allow few missing observations among the variables. The sampling rates for emission indicators measured by PTRMS (proton-transfer ionization mass spectrometry) differed between the two aircraft missions. The SEAC4RS mission acquired suitably complete PTRMS-derived datasets at a once-per-minute rate, and this defined the data interval used for both datasets. Ad-


**Deleted:** Our techniques use algorithms that currently allow few missing observations among the variables. The sampling rates for emission indicators measured by PTRMS (proton-transfer ionization mass spectrometry) differed between the two aircraft missions.

**Moved up [3]:** In order to provide a suitably complete dataset for SEAC4RS, we used the can samples to infer likely concentrations at one-minute intervals of key species, i.e., CH<sub>4</sub> for all flights and CO for the first few flights, using available can samples at slightly lower frequency. The **R**

**Deleted:** The use of imputation seemed justified by three observations: (1) Checks made when both LAS and GC data were available suggested agreement. In an early period, missing tunable-laser absorption spectrometry data for

over four times the assumed background of 0.054 ppb. Since some plumes are known to be quite low in gas-phase emissions, a few samples with lower  $\text{CH}_3\text{CN}$  mixing ratios, but with  $b_{\text{scat}} > 2 \times 10^{-2}$  were allowed in. Plots of  $\text{CH}_3\text{CN}$  vs  $b_{\text{scat}}$  suggested that a linear combination of the two minimal conditions clearly separated a population of forest fire plumes from other high-particulate situations.

There were forest-fire plumes for which urban sources of CO and other fire tracers made attribution and quantification problematic, and so a further test based on CO was applied to exclude urban samples, using CO vs.  $\text{CO}_2$  plots for the years 2008 and 2013 separately (Figure 2). We used a  $\Delta\text{CO}/\Delta\text{CO}_2$  ratio of  $< 33 \text{ ppb/ppm}$  to exclude plumes with excessive urban contamination. The figure suggests that some plumes with modest levels of urban influence remained and a few genuinely uncertain situations were excluded where fire might still have been dominant. Species with sources other than biomass burning and with lifetimes sufficiently long to allow regional mixing can pose difficulties somewhat similar to  $\text{CO}_2$  variability, with solutions suggested in Section 8.2 and Chatfield and Andreae (2019). We noted some localized observations of perplexing, consistently negative  $\Delta\text{CH}_4/\Delta\text{C}_{\text{tot}}$  relationships in the ARCTAS data (but not other species) and removed these observation instances. Such relationships were found close to seaports or oil-producing regions.



**Figure 2.** Urban and forest fire plumes are separable by the ratio of CO to  $\text{CO}_2$ . Colors indicate a relative measure of  $\text{CH}_3\text{CN}$  above background, from blue (lowest,  $\sim 0.1 \text{ ppb}$ ) to red (highest,  $\sim 6 \text{ ppb}$ ) values.

### 3. Observed behavior of $\text{C}_{\text{tot}}$ in fire plumes – Properties of tracers

This section provides some examples of  $\text{C}_{\text{tot}}$  and fire tracers. It illustrates the limitations of changes in  $\text{C}_{\text{tot}}$  along a sampling path as an indicator of fire influence,  $\text{C}_{\text{burn}}$ , for emissions estimation and the much greater similarities of the such changes of tracers that possess shorter transformation time-scales. These define our approach to EnRs and EFs. The relation of fire emissions to observed  $\text{C}_{\text{tot}}$  to  $\text{C}_{\text{burn}}$ , can be apparently simple or complex, depending on how the history of non-fire CO and  $\text{CO}_2$  entrained into fire plume air parcels affects  $\text{C}_{\text{tot}}$ . We show this commonality of relationships will to motivate the theory of expanding plumes in Section 4. That theory will suggest a method worked out in Sections 5 and 6 to find the key variable,  $\text{C}_{\text{bgd}}$ , that then provides  $\text{C}_{\text{burn}}$  and thus EnRs.

Figure 3 describes two flights in which plume encounters show how the interpretation of  $\text{C}_{\text{tot}} = \text{C}_{\text{burn}} + \text{C}_{\text{bgd}}$  can be simple or complex. Figure 43(a) shows the locations of the plume samples

Deleted:  $\times 10^{-3}$

Formatted: Font: Not Bold

Formatted: Font: Italic

Deleted: Recall that in this work we wish to advance the use of multiple tracers to estimate EnRs and EFs in more detail and also in more circumstances, e.g., where MCE is difficult to estimate.

Deleted: t

Deleted: air

Formatted: Paragraph

Deleted: give either clear or misleading descriptions of EnRs

Deleted: Figure 3


Deleted: a



observed during SEAC4RS Flight 10 over Montana between 2 and 4.5 km MSL. Local topography ranged from 1 to 2 km elevation. This data set included samples from the very intense plume of the Rim Fire (discussed later), collected far downwind. Figure 43(b) shows ARCTAS Flight 14, which was over the Coastal Mountains of Northern California at 0.5 to 1.5 km altitude, with topography from 0 to 1 km.

Figure 4(a) (mapped in 3(a)) gives the time series of fire indicators from Flight 10. The fire tracers CO and  $b_{\text{scat}}$  appear generally well correlated with  $C_{\text{tot}}$ . This correlation is seen in b-c. Colors from blue to red give a key to sampling times. The large orange dots in Figure 4(a) and 4(d) distinguish the plume points selected (based on our plume tracers) from adjacent non-plume measurements made in the flight. The lines connecting the adjacent plume samples suggest two or perhaps three linear patterns pointing back to a no-fire background of  $C_{\text{tot}} \sim 392.5$  and  $394.5$  ppm. Separately, a few points near the horizontal axis seem to suggest a low EnR. These points occur in the middle of sampling, just after 11 LT. Patterns of variation related to  $b_{\text{scat}}$  (550 nm, c) are very similar to those of CO (b). Most plumes encountered suggest very similar slopes.

Whereas the SEAC4RS data (a-c) suggest mostly expected behavior, the ARCTAS Flight 14 measurements (d-f) show that  $C_{\text{tot}}$  variations, likely due to  $C_{\text{bkgd}}$  variability, can greatly complicate the attempt to estimate EnRs. The very first samples plotted and those after about 13:35 LT have very clean tracer levels. (Those between 13:03 and 13:15 LT did not quite qualify as plume points, but the tracers do indicate some fire influence.) In this case, the trace of  $C_{\text{tot}}$  does not reflect fire influence well at all. Both fire tracers shown in c-d show wildly varying relationships to  $C_{\text{tot}}$ , but are remarkably similar in that relationship.



**Figure 3.** Flight paths and locations of plumes for two fire samples. (a) Sampling over Montana during SEAC4RS Flight 10. (b) Sampling in a cross-mountain transect along the Northern California Coast. The size of the spheres indicates the relative amount of biomass burning contribution. Numerical values of the contribution are in a later figure, Figure 9. Some of the information was converted from GoogleMaps® using the R programming language.

Deleted: Figure 3... (b) ... shows ARCTAS Flight 14, which was over the Coastal Mountains of Northern California [13]

Deleted: a gives

Field Code Changed

Field Code Changed

Deleted: Simple differences would suggest similar EnRs (slopes). A ... eparately, a set of ... ew points near the hori... [14]

Field Code Changed ... [15]

Deleted: Separated ... ost plumes encountered at 10:45, 11:20, and after 12:00 LT ... [16]

Field Code Changed ... [17]

Deleted: can ... an greatly complicate the attempt to estimate EnRs. The large orange dots in Figure 4d distinguish... [18]

Field Code Changed

Deleted: ¶

Moved down [4]:

Deleted: ¶ ... [19]


Moved (insertion) [4]

Deleted:


Formatted: Font: Not Bold

Deleted: Figure 8.

Formatted ... [20]




(a)




(b)

(c)



(d)



(e)


(f)

Deleted: a ... [21]

Formatted: Left

Formatted: Font: Helvetica

Formatted Table




Deleted:

Formatted ... [22]

Deleted: b

Formatted ... [24]

Formatted ... [25]




Deleted:

Formatted ... [23]


Formatted ... [26]

Formatted ... [27]




Deleted:

Formatted ... [29]



Deleted:

Formatted ... [29]



**Figure 4.** (a) Timeline of sampling, for the period shown in Figure 3(a), Montana, of  $\text{CO}_2+\text{CO}$  (blue, left axis) and the fire tracers  $\text{CO}$  and  $b_{\text{scat}}$  (red and green points, right axis). Orange-filled points were identified as clear plume points. Unfilled points were not, but might have some fire influence, especially near plume points. (b) scatter diagram of  $\text{CO}$  vs  $\text{CO}_2+\text{CO}$  with arrows showing the time progression of aircraft sampling of identified plume points. Colors provide a key to times shown in (a). Light gray numerals give observation times in minutes. (c) a similar diagram of  $b_{\text{scat}}$  vs  $\text{CO}_2+\text{CO}$ . Similar shapes of figures are noted in the text. (d) Timeline of sampling for the period shown in Figure 3(b), Coastal Transect. (e) scatter diagram of  $\text{CO}$  vs  $\text{CO}_2+\text{CO}$  during the transect, like (b). (f) a similar diagram of  $b_{\text{scat}}$  vs  $\text{CO}_2+\text{CO}$  for the Coastal Transect. The black bars graphed in (a) and (d) are estimates of non-fire influenced  $C_{\text{bgd}}$ , see text. They and the non-plume points suggest air-mass changes in  $\text{CO}_2+\text{CO}$ .

The small black bars in both Figure 4(a) and 4(d) foreshadow our results of Section 7; they show our eventual estimates of  $C_{\text{bgd}}$  for the selected fire-plume cases. We suggest that they aid a plausible description, viz:  $C_{\text{tot}}$  sampled by the airplane increases due to higher non-fire  $C_{\text{bgd}}$  from just before 13:00 to 13:15, and then decreases gradually until about 13:28. At one sample around 13:21 and several at 13:20–13:30, the background is particularly low, 382–383 ppm. This is a plausible description of the mixing of air-masses with originally concentrations of  $C_{\text{tot}} \sim 382$  and  $\sim 388$  ppm. Wisps of less-mixed air occasionally interrupt a relatively continuing trend. A close examination of the  $\text{CO}$  and  $b_{\text{scat}}$  increments compared to  $C_{\text{tot}}$  increments in Figures 4d–4f agrees with this description suggested by the black dashes. The simpler case of SEAC4RS Flight 10 may one similar example. At 11:08 LST early in the flight, there is a brief excursion upwards of  $C_{\text{tot}}$  without any excursion in the tracers. The small black bars show this as a plausible excursion of  $C_{\text{bgd}}$ . Figures 4b and 4c show this as the 2–3 exceptional points near the horizontal axis.

The plausibility of these examples highlights ideas of fundamental similarities of the way plumes of different tracers behave with entrainment even as  $C_{\text{bgd}}$  varies in response to distant, non-fire processes. This leads us to a mathematical description of our observations in Section 4.

## 4. Theory: expanding plume for several species

### 4.1. A general relationship

Figure 5 gives a general description of the dilution process, showing by the size of cubes how a mole of near-flame air is diluted by non-fire material as entrainment occurs. (The boxes shown suggest volumes, but lofting adiabatically changes volume. Discussion in terms of moles simplifies the discussion of mixing ratios and EnRs.) The figure is based on observations of plume size and plume dilution during rise followed by largely horizontal dilution downwind (Lareau et al., 2017; Hanna et al., 1982), which are consistent with mixing ratios measured in this dataset and near-fire  $\text{CO}_2$  concentrations of  $1\text{--}2.5 \times 10^4$  ppm. The sizes are meant to be suggestive, but we found that they give a valuable frame of discussion of all lofted forest fire plumes. Some details are in Supplementary Material (SM): “Note on Volumes.”

Moved (insertion) [1]

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Italic

Formatted: Font: 12 pt, Not Bold, Subscript

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 13 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 13 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Italic

Formatted: Font: 12 pt, Not Bold, Subscript

Formatted: Normal

**Deleted:** (a) Variations of  $\text{CO}_2+\text{CO}$  and tracers showing complexities of fire plumes and background air. (b) Timeline of  $\text{CO}_2+\text{CO}$  and  $\text{CO}$  in SEAC4RS Flight 10 (Montana) samples. MERET analysis shown in a subsequent figure. (c) Variations of  $\text{CO}_2+\text{CO}$  and tracers in ARCTAS Flight 14 (Northern California). ¶

Figure 5 gives a general description of the dilution process, showing by the size of cubes how a mole of near-flame air is diluted by non-fire material as entrainment occurs. (The boxes shown suggest volumes, but lofting adiabatically changes volume. Discussion in terms of moles simplifies the discussion of mixing ratios and EnRs.) The figure is based on observations of plume size and plume dilution during rise followed by largely horizontal dilution downwind (Lareau et al., 2017; Hanna et al., 1982), which are consistent with mixing ratios measured in this dataset and near-fire  $\text{CO}_2$  concentrations of  $1\text{--}2.5 \times 10^4$  ppm. The sizes are meant to be suggestive, but we found that they give a valuable frame of discussion of all lofted forest fire plumes. For example, the sequence of increasingly large boxes emphasizes that the relative effect of entrained non-fire air is largest near the flames, but the absolute effect of entrained air on the composition of an observed parcel is often largest close to the parcel at its point of sampling. The following section gives a framework illustrating effects of emissions and entrainment on EnRs. (Parenthetically, the box volumes of parcels at different altitudes are similar to the mole amounts shown; to be consistent with adiabatic rise of parcels, volumes should be about  $\sim 11\%$ – $14\%$  larger in linear dimension for most plume tops.) ¶




Using Figure 5 as a guide, consider a parcel originating at a time  $t_1$  containing  $v = v_1$  moles, that expands with an exponential relative rate  $r_v = v^{-1}(dv/dt)$ . (For our illustrative examples and to rationalize the MERET method, we need not start at the flame. We suggest a reasonable starting point described below.) This rate of expansion  $r_v(t)$  of the molar volume varies considerably over time and fires are expected to have different magnitudes. Then molar mixing ratios will evolve with a law

$$\frac{dx}{dt} = -\frac{1}{v} \frac{dv}{dt} (x - x^E) \quad (7a)$$

$$\frac{dy_j}{dt} = -\frac{1}{v} \frac{dv}{dt} (y_j - y_j^E) \quad (7b)$$

where  $x^E$  is the mixing ratio of entraining  $C_{tot}$  and  $y_j^E$  is the entraining background mixing ratio of fire tracer species or property  $j$ . The effect of volume addition is captured by  $v(t)$  which varies with time and expansion. The use of the relative rate  $v(t)$  does not require that the dilution is exactly exponential, but does make the algebra somewhat simpler.



**Figure 5.** Inflow of air into an expanding fire plume; a likely near-fire aircraft sampling location would be near the cube on the upper right. Cubes are shown with three-d sizes proportional to the number of moles of entrained air. These may be considered volumes of air adjusted downward to compensate for the adiabatic expansion that rising plumes undergo. The smallest cube is taken to be near the flames, at roughly the point where fire emissions transition from mixing to entraining background air. Exact placement of this cube is not important to the analysis of entrainment, expansion, and tracer mixing ratio. Successively larger cubes have volumes roughly in the ratio of 1, 40 (partially raised), 140 (near neutral buoyancy level); sizes are consistent with a buoyant fire plume (Lareau et al., 2017). The rightmost cube has a ratio to the first of 400, consistent with horizontal Gaussian dispersion during travel downwind. See text for more details.

A general formulation of the expanding plume model consistent with the description appears to suggest formidable complexities. Using  $\tau$  to describe the integration through time of an expanding parcel,

Deleted: Figure 5

Formatted: Paragraph

Deleted: 5

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

$$y_j(t_{\text{sample}}) = \int_0^{t_{\text{sample}}} -v(\tau)(y_j(\tau) - y_j^E(\tau)) d\tau + y_j(t=0) \quad (8)$$

with a similar equation for  $C_{\text{tot}}$ , now generally referred to as  $x(t_{\text{sample}})$ , involving  $x(\tau)$  and  $x^E(\tau)$  where  $x(t=0)$  and  $y_j(t=0)$ . These are determined by the  $C_{\text{burn}}$  from the fuel consumed and the tracer compounds released at the same time, as well as background concentrations,  $C_{\text{bkgd}}$ , and pre-flame backgrounds of the tracer  $y_j^E$ . We leave aside as a separate problem of a fire-burning model the complexities of the actual flame and its incorporation of additional air up until the point that entrainment of non-burning air becomes dominant.

Our approach is to avoid describing the complete history of  $v(\tau)$  and any complex variation of  $x^E(\tau)$  and  $y_j^E(\tau)$ . These would require a complete description of air along the parcel trajectory and the turbulent physics of entrainment. We accomplish this by concentrating on the fact that the entrainment process affects both  $x(\tau)$  and all the  $y_j(\tau)$  in the same proportions. This is a single-parcel description ignoring complexities of the rest of the plume. For convenience of discussion, we stipulate that the environmental air entrained has  $x^E(\tau)$  and  $y_j^E(\tau)$  constant over long periods. For example, it is constant in the mixed layer surrounding the fire flames and initial plume and then again in the subsequent regions often in the free troposphere. Conceptually there may be several regions which contribute; the exact history is lost. Our idea is that regression analysis allows us to infer a characteristic sum of effects which is described by a single quantity. The analysis can only be as complex as the number of our measured quantities allows. [See also SM for a Note on Initial Point.](#)

Returning to the differential-equation view of the simple expanding plume model suggests a method for estimating the most important parameters. Solving each of the equations for the expansion rate and equating the expressions we obtain a form that eliminates the details of entrainment and emphasizes proportionality. We recommend the reader refer back to Table 2, Table of Symbols during the discussion of theory and then estimation details.

$$\frac{1}{(y_j - y_j^E)} \frac{dy_j}{dt} = -\frac{1}{v} \frac{dv}{dt} = \frac{1}{(x - x^E)} \frac{dx}{dt} \quad (9)$$

Since  $dy_j^E/dt = 0 = dx^E/dt$ , we get

$$\ln(y_j - y_j^E) = \ln(x - x^E) + C_j \quad (10)$$

$$(y_j - y_j^E) = a_j (x - x^E) \quad a_j = \exp(C_j) \quad (11)$$

Note that by our definitions, the reasonable interpretation of  $C_j$  is the EnR  $a_j$  for species  $j$ . Consider two observations of the same plume, each made at differing degrees of dilution  $v$ . For convenience, these are labeled  $\beta$  and  $\alpha$ , mnemonically “before” and “after.” Temporally they could be nearly coincident or  $\beta$  after  $\alpha$ .

$$(y_{\alpha j} - y_{\beta j}) - (y_j^E \alpha - y_j^E \beta) = a_j (x_{\beta} - x_{\alpha}) - a_j (x^E \alpha - x^E \beta) \quad (12)$$

[For periods of expansion in which the entrained concentrations are constant. See also SM: Note on Varying Entrainment.](#)

Deleted: ¶	... [31]
Formatted	... [32]
Formatted	... [33]
Formatted	... [34]
Formatted	... [35]
Formatted	... [36]
Formatted	... [37]
Formatted	... [38]
Formatted	... [39]
Formatted	... [40]
Formatted	... [41]
Formatted	... [42]
Formatted	... [43]
Formatted	... [44]
Formatted	... [45]
Formatted	... [46]
Formatted	... [47]
Formatted	... [48]
Formatted	... [49]
Formatted	... [50]
Formatted	... [51]
Formatted	... [52]
Formatted	... [53]
Formatted	... [54]
Formatted	... [55]
Formatted	... [56]
Deleted: b	
Deleted: a	
Deleted: b	
Deleted: a	
Deleted: .	
Deleted: For these observations,	
Deleted: a	
Deleted: b	
Deleted: a	
Deleted: Eb	
Deleted: b	
Deleted: a	
Deleted: a	
Deleted: b	
Deleted: ¶	
Deleted: This formula is the basis for the NEMR technique	

Table 2. Table of Symbols

Symbol	Signifies	Observed, Estimated, or Hypothetical
$C_{\text{tot}}$	CO <sub>2</sub> + CO (+ other carbon, ignored), ppm	<i>O</i>
$C_{\text{burn}}$	CO <sub>2</sub> + CO (+ other carbon, ignored) emitted from fire, present downwind, in plume sample, to be estimated as $(x_i - x_i^0)$	<i>E</i>
$C_{\text{bkgd}}$	CO <sub>2</sub> + CO <i>not</i> emitted from fire, present downwind in plume sample, thought of as a mixture of $C_{\text{tot}}$ entrained at various stages in plume expansion and rise, to be estimated as $\hat{x}_i^0$ or casually as $x_i^0$ . This is not necessarily air surrounding the plume sample!	<i>E H*</i>
$C_j$	A constant of integration, replaced by $a_j = \exp(C_j)$	<i>H</i>
$i$	Sample sequence number, organized for convenience by time of the sample	<i>O</i>
$j$	Tracer number for regression, here 1 to 10 or “CO” or “b <sub>Scat</sub> ”, ... . After regression estimation is completed, $j$ may be used similarly to specify any fire emission concentration or response, e.g. “propene” or “O <sub>3</sub> ”	<i>O</i>
$a$ or $b$	Location at beginning or end of a period of idealized plume development and entrainment	<i>H*</i>
$x_i$	$C_{\text{tot}}$ = CO <sub>2</sub> + CO at a plume sample location and time, used in algebraic development, shown on $x$ axis	<i>O</i>
$y_{ij}$	Tracer concentration, e.g. toluene, $b_{\text{Scat}}$ , at plume location and time.	<i>O</i>
$x^E$ or $x_A^E$	Environmental air “background” $C_{\text{tot}}$ concentrations existing at location A, e.g. beginning of our integration of the plume expansion equation. B signifies condition at the end of calculation	<i>H*</i>
$y^E$ or $y_A^E$	Environmental air “background” concentrations of fire tracers.	<i>H</i>
$x_i^0$	Equivalent background concentration characterizing composition at plume sample location and time	<i>E</i>
$y_j^0$	Background concentration of tracer $j$ . Typically estimated as a minimum value from observed probability density function for samples in a particular flight intensive, especially non-plume samples without signals of stratospheric air.	<i>E</i>
$a_j$	Slope relationship of $y_{ij}$ to $x_i$ for species $j$ , typically species $j$ under burning conditions for a “fire type” that is common for all species at instance (time) $i$ . These slopes then transform to EnRs and ERs.	<i>E</i>
$c_j$	Intercept relationship of $y_{ij}$ to $x_i$	<i>E</i>
$\hat{x}_{ij}^0$	One of several (10) estimates of $x_i^0$ based on tracer $j$ and fire type assigned by clustering for observation instance $i$ .	<i>E</i>
$\hat{x}_i^0$	The median of the $\hat{x}_{ij}^0$ over all the tracers $j$	<i>E</i>
$(x_i - x_i^0)$	The estimate of $C_{\text{burn}}$ for instance $i$ in theoretical development and then from regression. Regression results are properly $(x_i - \hat{x}_i^0)$	<i>E</i>

Note that symbols may transition from Hypothetical to Estimated as the discussion develops.

Formatted: Font: 10 pt

Formatted: Font: 10 pt


Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt



**Figure 6.** (a) Simulated dilution of three different fire tracers, EnRs as shown, and with environmental  $x^E$  of 380 ppm. These are nominally CO, benzene, and ethylene. Background concentrations of the tracers have been subtracted.  $C_{burn}$  is measured from the y-axis, where  $C_{tot}$  has the value 380 ppm. Larger dots highlight equivalent degrees of dilution. (b) Simulated dilution of one tracer, nominally CO above background, with different background  $x^E$ . Backgrounds are illustrated by the x intercepts. (c) Simulation of three tracers, varying EnRs and varying backgrounds (deducible from the x intercepts). Thin lines emphasize similar constant y values with different backgrounds, and constant  $x^E$  values with varying EnRs. (d) Simulations like (a), but with a change of the  $x^E$  entrained  $x^0 = C_{burn}$  background from 395 ppm to 375 ppm at the time of the 8th dilution step. A single applicable background value ~378 ppm is an appropriate linear interpolation between 395 and 375 ppm. (e) Simulation where background  $x^E$  remains constant, but background CO changes by 1.5 ppb during the time (CO drawn at twice height for visibility).

This formula is the basis for the NEMR technique mentioned above, with  $j = CO$  playing a particularly important role. The inequality restrictions should be evaluated for an EnR to be a candidate for an ER and then an EF. In some cases, the background values,  $x^{Ea}$ ,  $y_j^{Ea}$ , might be estimated from measurements made outside the plume. It can be somewhat more difficult to estimate  $x^{Eb}$ ,  $y_j^{Eb}$  upwind, especially for air entraining into the fire plume at its source. A plume may also entrain air from various backgrounds and at various times during lofting and spread. That is, the history of entrainment may well be more complex than two conditions, “a” and “b”, and the number of situations where we may estimate EnRs and then EFs is greatly limited. This important realization was described by Yokelson et al. (2013). The NEMR method can deal with most differences in  $x^{Eb}$  but not  $y_j^{Eb}$  for most tracers, and the quantity  $y_{CO}^{Eb}$  for carbon monoxide must be well sampled and well understood.

Deleted: Figure 6.

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Check spelling and grammar

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Not Italic

Formatted: Font: 12 pt, Not Bold, Not Superscript/ Subscript

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Subscript

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Not Italic

Formatted: Font: 12 pt, Not Bold, Not Superscript/ Subscript

Formatted: Font: 12 pt, Not Bold

## 4.2. Examples showing robustness of computations of idealized $C_{\text{tot}}$

We used this approach to produce the following concrete examples of increasing complexity. They illustrate the origin of the features seen in Figure 6 in terms of this simple plume dilution model. They helped motivate our solution techniques and indicate methods of analysis of individual plumes. These examples indicate possible limitations, but they also indicate a comforting averaging behavior of the linear differential equations as they describe our solutions. These uniformities and deviations also showed up in the analyses that we develop below. The examples also give some quantitative feel on the effects of deviations from the simplest hypothesis, e.g.,  $x^E$  and  $y_j^E$  remain constant through time. Figure 6 shows calculations describing behaviors of  $x$  and the  $y_j$  in several plausible situations. Each graph represents the development of plume mixing ratios for a period of plume doubling, similar to the analysis time chosen in Poppe et al. (1998), following their equations 7 and 8. The dots show equal increments of plume expansion. Most parameters defining the equations may be read from the graphs themselves. Each initial concentration is shown by the points to the upper right of the line, i.e., the points with maximum  $x = C_{\text{burn}}$  and  $y_j = \text{tracer concentration for each case considered}$ .

Figure 6(a) illustrates a plume history for  $x^E = 380$  ppm and EnRs with respect to airborne  $C_{\text{tot}}$  of  $95 \times 10^{-3}$  ppm/ppm,  $12 \times 10^{-6}$  ppm/ppm, and  $25 \times 10^{-6}$  ppm/ppm, which are reasonable values for carbon monoxide (in ppb), benzene, and ethylene (in ppt). In the figures, focus attention on the relative behavior of the tracers. It is assumed that there are no consequential production or destruction reactions and also that there is a constant background tracer concentration, which has been subtracted. The individual plots show situations of increasing complexity. Figure 6(a) shows the dilution behavior of the three species. A constant dilution rate is plotted; note that varying dilution rate changes the spacing of the dots but not the linear pattern. Larger dots highlight equivalent dilution of tracer and  $x = C_{\text{tot}}$  as would be observed in hypothesized discrete airplane samples. Figure 6(b) illustrates the dilution of CO in environments with differing entrained  $x^E$ . In Figure 6(a), the larger dots align vertically, in Figure 6(b), horizontally. Figure 6(c) illustrates the situation where both EnRs and backgrounds vary; the thin lines emphasize independent aspects of EnR and  $x^E$ . The points on the  $x$  axis where (excess) tracer is zero are important to our estimation technique, more important than  $x^E$ . Estimation of  $x^E$  utilizes data on the vertical lines, while EnRs utilize information from both the vertical and horizontal lines. Statistically speaking, the problem of estimation of both backgrounds and EnRs illustrates simultaneous effects that are “separable”. The reader may wish to extend the analysis to a large sequence of changes in entrained concentrations and note the essential linearity of this aspect of the formulation and that the solution expresses an appropriate averaging effect. We remark that the near uniqueness of the solutions obtained below (making small allowances for measurement error) will underline the robustness of the solutions.

However, the effect of uniform variations in background tracer concentrations  $y_{ij}^0$  is not completely solved in this work.  $y_{ij}^0$  can be estimated by examining the lowest values  $y_{ij}$  in non-plume air; it is best to exclude values that represent exotic air (e.g., stratospheric air) or possible measurement problems at very low mixing ratios (e.g. negative values). Restricting attention to larger values of  $x_i$  and  $y_{ij}$  greatly ameliorates problems arising from  $y_{ij}^0$ ; making comparisons from one variable  $j$  to another in very clean parcels may also help.

Formatted: Normal, Indent: First line: 0"

Deleted: in Figure 6

Deleted: . Figure 6

Deleted: Figure 6

Deleted: . Figure 6(

Deleted: . Figure 6

Deleted: Figure 6a

Deleted: Figure 6

Deleted: Figure 6c

## 1000 5. Theory: A regression relationship for EnRs

1005 Let us consider more broadly the equations that provide a basis for statistical estimation. For current purposes of explanation, we make the seemingly large assumption that points from different plumes have similar properties at the same degree of dilution and may be compared. That is, the  $a_j$  are consistent for all plumes. Effects of varying  $x^E$  and  $y_j^E$  between the plumes may be largely taken out by regression; that is our current concern. Later, we will describe our approach to address possible variations in the EnR relationships  $a_j$  for parcels in the same or different plumes.

1010 The basis of MERET considers the unmeasured extreme where  $y_{aj} = 0$ . To begin with, we consider the situation where (i) the emissions relationships  $a_j$  are constant for all observations and (ii) background values of the tracers are small enough in a relative sense, i.e.,  $|y_j^{Ea} - y_j^{Eb}| \ll |y_{aj} - y_{bj}|$ . That condition is common for many species that have loss timescales of less than a month and/or have small non-fire sources. Each of these restrictions can eventually be relaxed. In this case

$$y_a = a_j (x_a - \{x_b - x^{Eb} + x^{Ea}\}) + \{y_j^{Ea} - y_j^{Eb}\} \quad (14)$$

$$\backslash \quad y_a = a_j (x_a - \{x_b - x^{Eb} + x^{Ea}\}), \text{ for } |y_j^{Ea} - y_j^{Eb}| \ll |y_{aj} - y_{bj}| \quad (14a)$$

1015 notice that terms within braces can be estimated by regression as sums, varying by the situation  $b$ . What if the values of these terms change discretely in time, for example as a plume leaves a daytime mixed layer, or distinct upper-air plumes are encountered? Simple algebra with linear formulas suggests that estimates of the terms in braces change discretely. Gradual changes of entrained mixing ratios of course imply a continuity condition on these terms.

1020 We return to the illustrative dilution behaviors described in Figure 6. Figure 6(d) describes a sudden change of background  $x^E$  by 20 ppm,  $x^{Eb}$  to  $x^{Ea}$ , midway in the expansion/dilution; at this stage of plume evolution, 20 ppm is about four times larger than typical fire contributions to  $C_{\text{tot}}$ . Estimates of  $x_i^0$  from a few samples along these lines (without knowledge of the time of change) would be intermediate. Equation 14a suggests that the EnR estimate need not be affected. Some similar calculations make it clear that the estimates respond in an appropriate averaging manner under varied assumptions. Figure 6(e) shows a very contrasting behavior, when there is a sudden change in concentration of entraining tracer (CO) during plume dilution, a change of 1.5 ppm. In comparison, the addition of CO by burning at the start of the interval is ~3.5 ppm. We may distinguish this as  $\hat{x}_{ij}^0$ , where the  $j = \text{CO}$  and the hat indicates an estimate.

1030 This graph also suggests that if there are more than three tracers (we use 8), then the median of all the estimates, median ( $\hat{x}_{ij}^0$ ), is robust against errors resulting if a tracer  $j$  has a variable or poorly described background resulting in  $\hat{x}_{ij}^0$  at falling distinctly higher or lower than the others. We must be concerned about this since tracers can have occasionally important non-fire sources. Change in background of a tracer compared to observed change due to fire is critical in determining its usefulness in determining a useful estimate of the appropriate background  $x^0$  as the well as the quality of its EnR of a tracer. Methane in particular have long atmospheric lifetime and several sources; consequently, it can exhibit variations that are more than 10% of the fire emission contribution for well dispersed plumes.

1035 The preceding discussion suggests that we may use the specialized least-squares technique,

$$y_{ij} = a_j(x_i - x_i^0) + c_j + e_{ij} \quad (15)$$

Deleted: Figure 6. Figure 6d

Deleted: We reiterate the suggestion that if the reader makes sample calculations, it becomes

Deleted: linear

Deleted:

Deleted: Figure 6e

Deleted: 10

Deleted:  $\hat{x}_{ij}^0$

Deleted: provides safety against

Deleted: a

Deleted: and

Deleted: incompletely

Deleted: .


Deleted: The median is not affected by undetected changes in background for one tracer species.

Formatted: Font: 12 pt



where  $x_i^0$  expresses all the terms, and any other corrections not proportional to  $a_j$ . If  $-x^{Eb} + x^{Ea}$  remains constant in the samples considered,  $x_i^0$  can still be estimated. A term  $c_j$  is included; it should be zero, but a significant non-zero estimated value can alert us to inadequacies in our assumptions. We may call  $x_i^0$  an “effective background”. However, it is not a specific background, but actually summarizes the whole effect of changes in  $C_{bkgd}$  and also the degree of dilution. This means that the regression can synthesize information from not just one well-characterized plume but rather a variety of plumes with the same  $a_j$  behavior.

Figure 7 illustrates the use of regression using the ideas developed in Figure 6. Using the same formulas as above, we depict observations made of CO, C<sub>6</sub>H<sub>6</sub>, and C<sub>2</sub>H<sub>2</sub> made at three instances (times). The three tracers determine a value  $x^0$ , and given that information, the three tracer enhancements and therefor tracer EnRs are determinable. This simulation assumes no error in the measurements of CO<sub>2</sub>+CO or the tracers, and assumes no variation in the EnRs, so values are determined perfectly. It is useful for our following discussion to examine the uniqueness of the solutions for two tracers and two plume observation sample instances, and then two tracers and three instances. In the case of two samples, with  $m$  samples and  $n$  slopes (EnRs), we need to estimate  $x_1, x_2, a_1, a_2$ , and  $x_0$  using only  $y_{11}, y_{12}, y_{21},$  and  $y_{22}$ , there are not enough measured variables to determine a unique solution,  $N_{Tracer} + N_{Instance} + 1 > N_{Tracer} \cdot N_{Instance}$ , viz.,  $5 > 4$ . However, if there are three tracers,  $N_{Tracer} \cdot N_{Instance} > N_{Tracer} + N_{Instance} + 1$ , and we get a solution. Any measurement error would give conflicting solutions. Using linear regression with larger  $m$  and  $n$ , one obtains solutions and increasingly accurate error estimates.



**Figure 7.** Multiple tracers allow solution for an equivalent background  $x^0$ , illustrated by an idealized example largely replicating the conditions of fire plume sampling above an Amazonian mixed layer as described by Yokelson et al. (2013). The fact that the various colored lines associated with each  $x$  meet at the value with  $x$ -coordinate  $x^0$  and  $y$ -coordinate 0 represents the estimation that precisely solves equation (5) above. If we had included error in observations or variation in EnRs, there would be uncertainties in the positions of  $x^0$  and the slopes. See the text for details.

Deleted: Figure 7

Deleted: Figure 6.

Formatted: Font color: Auto

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Deleted: Figure 7.

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Not Italic

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Not Italic

Formatted: Font: 12 pt, Not Bold, Not Superscript/ Subscript

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Check spelling and grammar, Border: : (No border)

## 6. Methodology:

### 6.1. Finding the CO<sub>2</sub> + CO background

The use of tracers with backgrounds removed and then scaled to a common mean establishes a well-conditioned matrix problem and easier analysis of sensitivity effects. Let us proceed with the regression and begin to address some complications that arise. The mathematical problem we must solve is equation 15:

$$y_{ij} - y_j^0 = a_j(x_i - x_i^0) + c_j + e_{ij} \quad (15)$$

In the following development in this section, we will work with  $y_{ij}$  above background, i.e., set  $y_j^0 \leftarrow (y_{ij} - y_j^0)$ , at least until it becomes important to consider background values of the tracer variables again.

We must attempt to fit  $y_{ij}$  for each species  $j$  (1 to  $N_j$ ) and for each measurement instance  $i$  (1 to  $N_j$ ). The term  $e_{ij}$  describes the error. The mixed-effects formulations expresses the idea that all the fire-tracer variables  $y_{ij}$  must “point back” to a zero-point, where no  $C$  was added by the burn,  $C_{\text{burn}} \equiv x_i - x_i^0 = 0$ , but that each instance  $i$  may have a different zero-point. A regression formula should solve this in some way. However, it also shows why the emission factor problem is difficult: the equation is non-linear in the multiple regression sense: i.e., we must include  $a_j x_i^0$  as a regression term. Commonly, regression fits provide a  $y$ -intercept, i.e., the value at  $x = 0$ . Here we have an  $x$ -intercept to estimate, i.e., the concentration of the reference species at zero added fire emissions, and so the problem becomes non-linear in the regression sense. The problem is also a mixed-random-effects model: we must estimate the effects of two “random” variables affecting the  $a_j$  and the  $x_i^0$ . In summary, we have a non-linear random effects model (Pinheiro and Bates, 2000), which requires specialized techniques.

Why not simply reverse the problem and seek  $x$  as a function of  $y$ ?

$$x_i = \alpha_j y_{ij} - x_i^0 + e_{ij} \quad (16)$$

where the  $x_i^0$  are estimated with a regression model (specifically a fixed-effects model). The difficulty is that the trivial solution  $x_i = -x_i^0$  fits perfectly and was hard for us to avoid even when we attempted to restrict the solution with a non-linear solver. Is it not easier to convert  $y$  intercepts into  $x$ -intercepts? This appears more productive and should appeal to those not familiar with using a non-linear solver in this particular mode of non-linearity. We may write a regression equation with an intercept:

$$y_{ij} = a_j x_i + c_i^0 + e_{ij} \quad (17)$$

where the  $y$ -intercepts,  $c_i^0$ , are estimated for each instance and the  $e_{ij}$  are minimized by least squares. The  $c_i^0$  may be thought to represent an error in  $y_{ij}$  resulting from the use of  $x_i$  with an incorrect baseline (the unknown  $x_i^0$ , an error that is constant for all species  $j$ ). The **R** expression used for this was

```
main.lmer = lmer( y ~ x + (x - 1 | species.type) + ( 1 | id ) + 1
```

where `id` indicates the sequential observation number for the tracer species. The term `species.type` indicates the species number  $j$ . The word `type` signals a generalization described in the next section. The vertical lines indicate how factors are involved with variables, 1 indicates an intercept is to be described by a random effect, and `(x - 1 | species.type)` indicates that a slope that multiplies  $x$  is to be estimated, indexed by `species.type`. “No intercept estimated” is signaled by `-1`. The regression generates a set of fitted  $y$  values which we may

Formatted: Font: 11 pt



call  $\hat{y}_{ij}$ . to estimate  $x_i^0$ . A simple way is to use the slopes estimates  $\hat{a}_j$  of  $a_j$  by regression to find the several estimates  $\hat{x}_{ij}^0$  provided by

$$\hat{x}_{ij}^0 = (\hat{y}_{ij} - y_{ij}^0) / \hat{a}_j \quad (18)$$

(equivalently, use  $y_{ij} \leftarrow y_{ij} - y_j^0$ ), so that  $\hat{x}_{ij}^0 = \hat{y}_{ij} / \hat{a}_j$ , where we will (currently) assume that  $y_{ij}^0$  has been satisfactorily estimated as  $y_j^0$  as a single background value suitable for all samples, as described before. The use of a single  $\hat{a}_j$  for all observation instances of the same species and NMF-assigned cluster is a fairly large constraint on the resulting estimate. The use of a larger number of cluster classes will allow more ability to follow the EnR actually characteristic of the observation, but at the cost of parsimony and sensitivity to instrumental error for the species or property. We found that the conversion of y-intercepts to x-intercepts was more accurate when the values of nominal baseline  $C_{tot}$  were adjusted so that the median value was 391 ppm. Essentially, this use of a constant baseline makes the numerics better conditioned: the difference of small quantities, not the difference of large quantities. This adjustment made the individual  $\hat{x}_{ij}^0$  symmetric around  $\hat{x}_i^0$  and also more closely spaced. We then take

$$\hat{x}_i^0 = \text{median}_j \hat{x}_{ij}^0 \quad (19)$$

The estimation of  $\hat{x}_i^0$ , now allows estimates of the incremental carbon liberated to the atmosphere,  $C_{\text{burn}} = x_i - \hat{x}_i^0$ . We will drop the hat from  $\hat{x}_i^0$  below, writing  $x_i^0$  and  $C_{\text{burn}} = (x_i - x_i^0)$  except when we wish to emphasize their nature as estimates.

Emission factors for individual tracer species may be obtained directly by adding fixed and random effects on slopes for each species and each observation,  $\hat{a}_j$ . An enhancement ratio for any concentration or property  $y_j$  with a background, measured at time  $i$  in the aircraft sampling can be obtained using the carbon-burned estimate

$$\text{EnR for } y_{ij} = \frac{(y_{ij} - y_j^0)}{(x_i - x_i^0)} = (y_{ij} - y_j^0) / (C_{\text{burn}})_i \quad (20)$$

To repeat, the variable  $y_{ij}$  now stands for any property for which we seek an EnR, for example ozone, which is not one of the ten indicator variables.  $y_{ij}$  describes a non-fire-dependent background value. This ratio estimate is available for all tracers, and is preferred over a similar slope variable  $\hat{a}_j$  used to estimate  $x_i^0$  in the in equation 18 above.

Formulas for the statistics of ratio quantities with uncertainties in numerator and denominator can be theoretically complex, so we simply computed error estimates by simulation using computed Bernoulli trials. One thousand samples of normal distributions were calculated each for the numerator and the denominator, using their uncertainties as  $1\sigma$  values. Then the ratios of the first numerator and first denominator normal deviate sample, the second, ... to the 1000th normal deviate sample for numerator divided by the 1000th normal deviate sample were calculated, and the distribution of ratios summarized. For the numerator, the measured value and the suggested standard deviation (typically a percentage ratio) provided the parameters for the normal distribution. For the denominator, the mean was the  $C_{\text{burn}}$  estimate, and the standard deviation was a value of 0.25 ppm, documented as the measurement error (precision + bias) of  $\text{CO}_2$ . (See Table 1). Uncertainties in the calculation of  $\hat{x}_i^0$  were considered small and did not add to the dispersion of the denominator, especially since it is clear that any additive biases contributing to the quoted uncertainty of  $(\text{CO}_2 + \text{CO})$  cancel out. Sample calculations in the supplementary material (SM: "Note on Sensitivity to number of tracers used"), suggest errors typically of magnitude 0.03 ppm

Formatted: Font: Not Bold

due to variations in technique, and usually  $<0.1$  ppm. Additive errors should also cancel out for the numerator, since a background is subtracted. Indeed, some tracers like ethene appeared to have a negative background as determined from plots and simple regression calculations of  $y_{ij}$  on  $(C_{\text{burn}})_i$ . This is not unexpected, since these compounds are sampled into cans, where a small but self-limiting coating of the measured species on the can surfaces might cause such a negative offset, and yet the integrity of the can sample at larger values might be little affected.

## 6.2. Practicalities: variable EnRs

Equations 17–19 provide the basis of the MERET technique. There are however some details that increase its relevance and accuracy. First there is normalization. Common practice is to normalize all the tracer species  $j$  with respect to the mean of all observations of species  $j$ , after subtracting a baseline. This allows each tracer to influence  $y_{ij}$  equally. Assigning weights accomplishes the same purpose, but scaling allows better diagnostic graphs. In fact, the literature referenced above emphasizes how informative  $j = \text{CO}$  is, despite its relatively small variation in EnR or slope. Consequently, we give CO twice the weight of all the other species.

Secondly, we allow for a certain amount of true variation in the EnRs, expecting this to make equation 18 perform better. This is done by imagining that virtual species can be associated with “fire types” for example “flaming CO” or “smoldering CO” or “high-nitrogen-fuel  $\text{CH}_3\text{CN}$ ”. A “fire-type” is a value for each observation that applies to all tracer species at that instance. It expresses commonalities between different mixes of burning emissions, commonalities that may be more frequently or less frequently expressed in any given plume, e.g., “smoldering-CO fire-type”. We might speculate on the nature of the fire-type, e.g. “smoldering” or “derived from nitrogen-rich fuel”. However, we let the [statistical technique](#) define these types, and so apply basic clustering techniques. We used non-negative matrix factorization (NMF), but Mahalanobis clustering or other techniques seem to do as well. NMF is more fully described in a companion paper describing patterns linking EnRs for several compounds; [NMF and  \$k\$ -means clustering are shown to be equivalent in cases appropriate to our work \(Ding et al., 2005\)](#). We used the **R** routine `nmf()` with  $k = 6$  components and the [Lee estimation technique with singular value initialization \(Lee and Seung, 2001, et al, 2001, Boutsidis and Gallopoulos, 2008\)](#). Since all fire tracers are correlated, such clustering characterizations are much better defined if based on a rough normalization to the fuel burned. We used a consensus variable, essentially a first principal component of the tracers, to act as a normalizing agent, playing the role logically played by  $C_{\text{burn}} = (x_i - x_i^0)$ .

(This estimate cannot resolve situations in which the EnR for a tracer varies; also, we found it problematic to use regression to give it an accurate value of  $C_{\text{burn}}$  in ppm.) Exact quantitative calibration of  $C_{\text{burn}}$  in ppm is not required, just a relative scale. (Note: A certain degree of characterization of fire-types can be found in Chatfield and Andreae [\[2020\]](#), which explores striking relationships among the EnRs. In that work, we cluster the fire-types using a completed description of the estimated  $(C_{\text{burn}})_i$  values, the estimates described below in [Figure 8](#).)

In a separate analysis effort, we sought timescales that could be inferred from the data, which could distinguish the relative age of burning emissions. At greater distances from the fire, there is both aerosol transformation and photochemical loss/production of species. Photochemical processing appeared easier to diagnose. We followed the ideas of Roberts et al. (1984), McKeen and Liu (1984), Parrish et al. (2007), and Warneke et al. (2013). The Parrish et al. presentation was

Deleted: .

Deleted: 5

Deleted: Brunet

Deleted: approach

Deleted: Gaujoux, 2014

Deleted: 2019

Deleted: Figure 8)

most directly relevant. For considerations of these plume samples, a single origin strongly controlling mixing ratios made analysis simpler. Following Parrish's Equation 3, and using the symbols  $E$  and  $Y$  are the mixing ratios of ethEne, and ethYne, respectively,

$$\tau_{\text{age}}(\text{OH}) = -\frac{1}{k_E - k_Y} \left\{ \ln \frac{y_Y}{y_E} - \ln \frac{ER_Y}{ER_E} \right\} \quad (21)$$

In view of this we constructed estimates for each instance  $i$  of  $\log_{10}(y_i Y / y_i E) - \text{Constant}$ . The Constant can be estimated with similar results (a) so that the shortest times are about +15 minutes, or (b) from the highest observed values of  $\log_{10}(y_i Y / y_i E)$ . The values of longer times are determined by the assumed value of  $[\text{OH}]$ . The references cited describe the fact that most  $\tau_{\text{age}}(\text{OH})$  observations have a contribution from mixing as well as photochemistry, but this has little affect the relative ages. In view of the uncertainty of the history of  $[\text{OH}]$  during transport, we simply graph the log of the ratios. Data analysis suggested that the assumed background mixing ratios of the species of ethyne and ethene were small. The supplementary material provides some more details and one estimate of the associated times. [SM: "Note on Sensitivity to number of tracers used"](#)

### 6.3. Summary of the MERET method

Here is a summary of the MERET method as we currently propose it. It contains many steps, but we note that there are many processes that affect EnR estimation.

1. Select a dataset of plumes with sufficient mixing ratios of the tracer,  $\text{CH}_3\text{CN}$ , above 0.225 ppb and expectably minimum interfering signals from urban sources:  $(\text{CO} - \text{CO}^{\text{Backg}}) / (\text{CO}_2 - \text{CO}_2^{\text{Backg}})$  ratio of  $> 33 \times 10^{-3}$ . Exact background levels are not critical, but should correspond to the year, season, and location sampled (examine plume and non-plume dataset). Plumes from agricultural burning and with detectable urban influence are thus excluded.
2. Select  $N_{\text{Trace}}$  fire tracer datasets that can provide sufficient measurement instances (we suggest  $N_{\text{instance}} > 50$ ) for each of the tracers, preferably hundreds of points. Tracers must be measurable sufficiently frequently to populate the dataset.
3. Estimate background levels  $y_j^E$  typically entrained into fire plumes. Plume values of tracers should be at least four times higher. For numerical reasons, select an offset  $C_{\text{baseline}}$  for each plume (each day). This is a value fixed for each plume  $I$  that is 2 ppm  $\text{CO}_2 + \text{CO}$  smaller than the minimum plume value observed for that day (with enough samples, for that plume of  $> 4$  points). Subtract this from the  $x_i$  values but record it so as to add it to quantities like the estimated  $x_i^0$  later.
4. Normalize the tracers above estimated backgrounds. This simplifies the regression estimation and its interpretation. All tracers tend to contribute equally.
5. Roughly cluster plumes into  $N_{\text{Types}}$  clusters by estimated EnRs (this can be applied recursively, after other steps). The value of  $N_{\text{Types}}$  ~~was~~ observed to make little difference. Too small a value  $N_{\text{Types}}$  can however lead to negative  $\hat{y}_{ij}$ . Here, ~~six~~ types were used.  $N_{\text{Types}}$  should be at least one smaller than the number of fire tracers.

6. Make many estimates of  $\hat{x}_{ij}^0$  and  $\hat{a}_{ij}$  using a mixed effects regression like `main.lmer`, allowing random effects corresponding to species (or species and type of fire) and by instance. Calculate  $\hat{x}_{ij}^0 = (\hat{y}_{ij} - y_{ij}^0) / \hat{a}_{ij}$  from the fitted values and thus  $(x_i - \hat{x}_{ij}^0)$ . Although a single  $x_i^0$  is sought, each tracer will give an estimate  $\hat{x}_{ij}^0$  at each instance. Take [The results for  \$C\_{\text{burn}}\$  and tracer EnRs suggested to us that one likely source of uncertainty is](#)

~~Deleted:~~ Weighting inversely proportional to the quantities could be used instead.

~~Deleted:~~ is

~~Deleted:~~ five

that  $C_{\text{burn}}, \hat{x}_i^0$ , and the tracers may change very rapidly in comparison to our one-minute sampling intervals. Looking into this, we found that many of the  $C_{\text{burn}}$  estimates are of small magnitude, 12 of the 422 samples yielded  $C_{\text{burn}} < 1.5$  ppm. Even large jumps from sample to sample in estimated  $\hat{x}_i^0$  were not particularly associated with anomalous estimates of  $C_{\text{burn}}$ . The remaining, appealing possibility is occasional imprecise time alignment of all measurements, particularly of the  $\text{CO}_2$  measurements. Such imprecise alignment could happen at any stage, from sampling line delays to interpolation to one-minute time intervals. Such variations in  $\text{CO}_2$  would affect the  $\hat{x}_{ij}^0$  found for all tracers in a coordinated way, just as was observed. Note that estimates of  $C_{\text{burn}}$  were little affected, since significant  $\hat{x}_{ij}^0$  excursions were associated with large  $\text{CO}_2 + \text{CO}$  values. See SM: “Note on Examples of Enhancement Ratios.”

More technical observations are these: one implication of this tendency for homogeneity in the  $\hat{x}_{ij}^0$  estimates is that the MERET technique works best with a large set of alternative fire-type classes. MERET should work better with more tracers. Adding additional classes (clusters) also tends to add only minor variations in the slopes  $\hat{a}_j$ , and harmful effects seen in over-fitting of regression models are avoided by a requirement that the  $\hat{a}_j$ ’s be positive.

the robust estimate of  $x_i^0$  as the median of these  $\hat{x}_{ij}^0$  at each instance  $i$ .

7. Use of an offset in calculations: We subtracted a baseline,  $C_{\text{baseline}}$ , a value determined as a constant for contiguous intervals, shown later in Section 7, Figure 8, and yielding a  $\sim 1$  ppm offset. We found that this minimized skewness and variance in the  $\hat{x}_{ij}^0$  estimates for each observation instance  $i$ . It is comforting that the effect of differing offsets on the values of the median,  $x_i^0$  is small,  $< 1$  ppm. One must not forget to add the offset back in the  $C_{\text{baseline}}$  when reporting  $x_i^0$ !

8. Sharp positive and negative excursions of  $x_i^0$  are seen near dramatic spikes in  $x_i$ . However,  $(x_i - x_i^0)$  and consequently the EnRs are little affected. We can only speculate that small differences in the time averaging of  $\text{CO}_2$  and the tracers due to the instruments may explain these.

Note also that the number of parameters  $N_{\text{Trace}} \cdot N_{\text{Types}} + N_{\text{Instance}}$  for the mixed effects regression is  $\ll N_{\text{Trace}} \cdot N_{\text{Instance}}$  so that the mixed effects regression is very over-determined.

It is possible to use this method recursively, making presumably better classifications of fire types. In our experience, while it is possible to make convergent, recursive characterizations of the appropriate  $C_{\text{burn}}$  quantity, more appropriate clustering, and mixed-model *lmer()* estimates, the quantities  $x_i^0$  and  $(x_i - x_i^0)$  were almost unaffected by such care. If we had available fewer than 10 tracers, such recursion might be important. We will incrementally update documentation of the code (Chatfield, 2020). New applications of the code will suggest improvements.

#### 6.4. Number of independent samples

A natural broader question is: “How well do these mean EnRs for a species represent the EnRs that might be measured in a large suite of significant forest fires in the Western US?” Clearly, this question can only be asked in the context of the sample provided by the two campaigns. Instances when the aircraft continued to sample smoke for many minutes could contain several types of plumes, as we will see illustrated for the Rim Fire Plume of August 2013. The use of 10-sec averages (if available) would not provide six times as much information about fire plumes as 60-sec averages over the same measurement run. We tried a simple, approximate quantification of “independent instances” available to us using a frequently used formulation by Trenberth

Deleted: each flight

Deleted:

Deleted: on <https://github.com/Rob-ertBChatfield/FireEmissionsEstimate/>.

Deleted: Remark:

1250 1984). This can also be seen as providing one answer to the question “How many effectively independent samples of  $C_{\text{burn}}$  are there contributing to a mean, standard deviation, etc., of  $C_{\text{burn}}$ , ... or of CO tracer?” That could be useful if instruments appeared to give imprecise measurements that required averaging. Trenberth assessed the correlation of successive observations by estimating an autoregressive AR(1) model with parameter  $\phi$  and random error  $\varepsilon_i$ , i.e.,

1255 estimate of  $\xi_{i+1} = \phi \xi_i + \varepsilon_i$ . We applied this for  $x_i - x^0 (= C_{\text{burn}})$  and several fire tracers  $y_{ij}$ , like toluene. Most contiguous sampling periods suggested around 0.6; this suggested Trenberth’s “effective time between independent observations” as  $\approx (1 + \phi)/(1 - \phi)$ , about 4 minutes. Four minutes corresponds to about 15 km at lower-tropospheric airspeeds for the DC-8. The effect of this on the formal standard errors as described by a normal distribution was to increase

1260 them by a factor of  $\sim 2$ . Roughly similar effects are expected for the empirical descriptions of EnR variability described below. Undoubtedly, for plumes within minutes of the source, the number of degrees of freedom corresponds more closely to the number of 1-min observations, but the number of such samples is low.

1265 Not surprisingly, *residuals* in regressions of CO against  $C_{\text{burn}}$  are very little correlated. We surmise that such low correlation gives confidence in the mathematical determination of the mean regression slope. However, it does not provide help in answering the larger question, that of relevance in new situations. The sequential samples of plumes may have features like non-stationarity and selection bias; we hope that these ideas suggest more sophisticated analyses of relevance, left to future work.


1270

Formatted: Font: Italic

**Deleted: <#>Results: Estimation of and  $C_{\text{burn}}$**

<#>The important results of the mixed model are the background, and even more importantly the incremental carbon liberated to the atmosphere. The background estimates of for all samples and the contributing individual estimates are shown in Figure 9. The median is shown as a thin black line. The colored circles in the legend identify how the tracer species  $j$  contribute an individual, determining the median.

<#>What are the uncertainties in the estimates of and we have made? The uncertainty in estimated carbon burned plays an important role in the ultimate estimates, the emission factors. In this section, we will confine our exposition to this uncertainty for now. The graphs of and shown in Figure 8 provide a practical understanding of the uncertainty. Note the continuity in; this important observation is described below.



**Figure 8. (Lower panel).** Estimates of the 422 background  $\hat{x}_i^0 = \text{CO}_2 + \text{CO}$  concentrations implied based on the 8 fire tracers indicated in the legend. Contributing individual estimates  $\hat{x}_{ij}^0$  are shown by overlapping colored points, with the median estimate  $\hat{x}_i^0$  indicated by a black bar. Usually the colored points overlap closely, this indicates strong agreement (Upper panel) Estimates of  $C_{\text{burn}} = x_i - \hat{x}_i^0$  indicators of fuel carbon burned, in the heavy green line. The preliminary estimate of  $C_{\text{burn}}$  based on the consensus of tracer deviations (without variable EnR estimates) is also shown in a light line. A scale factor, maximizing overlap with the heavy line, was necessarily estimated by regression. Flight days are indicated by the days marked on the top axes, and individual plumes, separated by non-plume concentrations of longer than 10 minutes, are shown as vertical separator lines. A set of horizontal lines at ~400 ppm indicates selected intervals for optimizing numerics (see text, Section 6.3, item 7).

## 7. Results: Estimation of $x_i^0$ and $C_{\text{burn}}$

The important results of the mixed model are the background  $\hat{x}_i^0$ , and even more importantly the incremental carbon liberated to the atmosphere,  $C_{\text{burn}} = x_i - \hat{x}_i^0$ . The background estimates of  $\hat{x}_i^0$  for all samples and the contributing individual estimates  $\hat{x}_{ij}^0$  are shown in Figure 8. The median  $\hat{x}_i^0$  is shown as a thin black line. The colored circles in the legend identify how the tracer species  $j$  contribute an individual  $\hat{x}_{ij}^0$ , determining the median  $\hat{x}_i^0$ .

What are the uncertainties in the estimates of  $\hat{x}_i^0$  and  $C_{\text{burn}}$  we have made? The uncertainty in estimated carbon burned ( $x_i - \hat{x}_i^0$ ) plays an important role in the ultimate estimates, the emission factors. In this section, we will confine our exposition to this uncertainty for now. The graphs of  $\hat{x}_i^0$  and ( $x_i - \hat{x}_i^0$ ) shown in Figure 8 provide a practical understanding of the uncertainty. Note the continuity in  $\hat{x}_i^0$ ; this important observation is described below. Traditional estimation of uncertainties for ( $x_i - \hat{x}_i^0$ ) is complex due to the several steps involved and the use of median estimates. The advisability of using the median estimator and its statistical properties have long been recognized (Laplace, 1774; Lawrence, 2013). This variety of uncertainty estimation may be useful as the MERET technique is refined. However, we expect that the study of uncertainty depends more on evaluating sources of true variability in the EnRs and also on the conservation of tracer concentrations from the flames to the sampling point than on the mathematics of median estimation. Consequently, the following paragraphs explore these questions related to the number and choice of tracers. We suggest that the typical strong overlap of the individual-tracer values may high precision of the observer's techniques!


How does the number of tracers affect results? What are the effects of using alternate or simpler sets of tracers? How many tracers are required for stable estimates? We began to address these questions by examining estimates made with fewer tracers in the intercept-determining set: the selection of the set of  $j$ 's. The supplementary material gives two examples of subsets. (SM: "Note on Sensitivity to Number of Tracers Used") Here is a summary of that material. The two sets chosen are those that are the most unambiguous indicators of  $\hat{x}_{ij}^0$  based on their mutual agreement with  $\hat{x}_i^0$  from the full set of 10 tracers. They are Set 1 (CO, Scat\_5, and HCHO) and Set 2 (CO, Scat\_5, HCHO, acetaldehyde, and toluene). These are indicated by a careful examination of Figure S8 in the Supplementary Material. (Abs\_5 contributed  $\hat{x}_{ij}^0$  most varying from  $\hat{x}_{ij}^0$ .) Set 2 gave variations around 0.02 ppm, the smaller set, Set 1, gave very similar variation except for the flights of June 22 and 25, where many observation instances varied by around 0.1 ppm, but with 11 points out of 422 differing by 0.3 ppm. This level of agreement surprised us. More significantly to our aims, the relative error in  $C_{\text{burn}}$  was only about 2%. When sets containing the less correlated tracers were used, deviations ranged up to 0.2–0.4 ppm, which appeared still remarkably small.

Observation-to-observation consistency in  $\hat{x}_i^0$ -estimates, seen for most plumes observed in Figure 8, is the strongest argument for the precision of the  $C_{\text{burn}}$  estimates. Recall, our theory does not use sequential time information: thus, successive estimates are essentially independent of each other. There is of course the dependency due to each observation's contribution to the estimate as one component of the entire dataset. This continuity is maintained even though the magnitudes of  $\text{CO}_2 + \text{CO}$  and estimated  $C_{\text{burn}}$  can change dramatically as the sampling aircraft enters and leaves each plume. Smooth excursions seen early in the flight marked 8.27 are explicable in terms of large changes in sampling altitude and location around the Rim Fire on that day. There are variations in  $\hat{x}_i^0$  from plume to plume and day to day.

Formatted: Indent: First line: 0.25"

Deleted: ¶

Deleted: ¶



Deleted: ... [58]

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Deleted: A note: methane and

Deleted: , and each with separate patterns of disagreement among the samples

Deleted: .


Deleted: 9

Deleted: Note that the

Formatted: Font: Italic



In contrast to this typical continuity of  $\hat{x}_{ij}^0$ -estimates, there are 15 to 20 brief and large excursions which deserve some attention. Of course, these may be disregarded in getting a general picture of EnRs. All the tracers suggest these excursions of the median, although there is a larger variation between the individual tracer-based estimates  $\hat{x}_{ij}^0$ . These excursions are always associated with large changes in  $\text{CO}_2 + \text{CO}$  and  $C_{\text{burn}}$ , but often they occur one minute later. We examined these excursions in detail. They do not seem to relate to changes in the EnRs  $\hat{a}_j$  (as qualified by fire-type) estimated simultaneously. The observations  $y_{ij}$  and the fitted  $\hat{y}_{ij}$  agree well, as well as for non-excursion points. Note however, that we may only use a single set of fire types, independent of  $j$ , to construct set of  $\hat{a}_j$ 's and  $\hat{y}_{ij}$  to make the  $\hat{x}_{ij}^0$  estimates.




**Figure 9.** Analyzed relation of tracers to carbon burned using the MERET technique for portions of SEAC4RS Flight 10 and ARCTAS Flight 14. Compare (a) through (d) with Figure 4, (b), (c), (e), and (f). Colors key the observations to times shown in the timelines, Figures 4(a) and 4(c). Light gray numerals give observation times in minutes.


Deleted: consistent

Deleted: , including the effect of fire-type classes on  $\hat{a}_j$ 's, and that this enforces considerable homogeneity in the  $\hat{x}_{ij}^0$  estimates.


Deleted: The results for  $C_{\text{burn}}$  and tracer EnRs suggested to us that one likely source of uncertainty is that  $C_{\text{burn}}$ , and the tracers may change very rapidly in comparison to our one-minute sampling intervals. Looking into this, we found that many of the  $C_{\text{burn}}$  estimates are of small magnitude, 12... of 99




Deleted:



Deleted:



Deleted:



Deleted:

Formatted: Font: 12 pt, Not Bold

Deleted: ...Colors key the observations to times shown in the timelines, Figures 4(a) and 4(c). Light gray numerals give observation times in minutes. Lightly colored marks on the axes show intercepts associated with the analysis for each inst ... [60]



The results for  $C_{\text{burn}}$  and tracer EnRs suggested to us that one likely source of uncertainty is that  $C_{\text{burn}}$ ,  $\hat{x}_i^0$ , and the tracers may change very rapidly in comparison to our one-minute sampling intervals. This would seem to be a concern since some  $C_{\text{burn}}$  estimates are of small magnitude: 12 of the 422 samples yielded  $C_{\text{burn}} < 1.5$  ppm. However, large jumps from sample to sample in estimated  $\hat{x}_i^0$  were not particularly associated with anomalous estimates of  $C_{\text{burn}}$ . The remaining, appealing possibility is occasional imprecise time alignment of all measurements, particularly of the  $\text{CO}_2$  measurements. Such imprecise alignment could happen at any stage, from sampling line delays to interpolation to one-minute time intervals. Such variations in  $\text{CO}_2$  would affect the  $\hat{x}_{ij}^0$  found for all tracers in a coordinated way, just as was observed. Note that estimates of  $C_{\text{burn}}$  were little affected, since significant  $\hat{x}_{ij}^0$  excursions were associated with large  $\text{CO}_2 + \text{CO}$  values.

MERET should work better with more tracers, since more fire-types may be revealed. However, additional classes (clusters) tend to add only minor variations in the slopes  $\hat{a}_j$ . Furthermore, harmful effects often seen in over-fitting of regression models should be minimized by a requirement that the  $\hat{a}_j$ 's be positive. (Positive and negative regression coefficients cannot be added that allow fitting of just a few points.)

## 8. Estimates of emissions ratios: Two MERET examples

### 8.1. MERET results for our two examples

The usefulness of our estimates of  $\hat{x}_i^0$  and  $C_{\text{burn}} = x_i - \hat{x}_i^0$  is seen in the MERET analysis (Fig. 9) of the two case studies analyzed above using the NEMR approach, where portions of Flights 10 and 14 were shown in Figure 4. The tracers CO and  $b_{\text{scat}}$  appear much better correlated with the  $C_{\text{burn}}$  estimated from MERET, especially in Flight 14. The plots for both CO and scattering imply linear relationships with an implied intercept near 0, i.e., background values of  $y_i$  have been satisfactorily removed. Difficulties with a variable  $C_{\text{bgkd}}$  appear to be resolved. However, the slopes of all the lines do not all agree. The Montana scatterplots (a) and (b) appear to suggest two slightly different linear features. The California Transect scatterplots (c) and (d) show more separated linear features, though the slopes are parallel. We expect that these might correspond to varying fire types, perhaps varying MCE's, to be discussed in Chatfield and Andreae (2020), or to variations in background values  $y_j^0$ , which are much harder to detect with either MERET or NEMR. A combined approach, using MERET to locate regions of similar MCE, might be useful here. Also, note that the variation of slope is more evident for  $b_{\text{scat}}$  than CO emphasizing the special role of CO as a single best fire tracer, closely followed by  $b_{\text{scat}}$ .

### 8.1 Table of several significant emissions

Table 3 provides a summary of the EnR relationships for some of the most significant gaseous emissions and particulate properties. In many cases, these EnRs can be converted to ERs and emission factors when the relationship of airborne  $C_{\text{burn}}$  to surface fuel consumed can be established. For the most highly reactive species, these EnRs will tend to be underestimates. An interpretive paper (Chatfield and Andreae, 2020) will give additional information on the photochemical age of the observation in many cases. Ozone and peroxy acetyl nitrate (PAN) are not emis-

Formatted: Subscript

Formatted: Subscript

Formatted: Font color: Custom  
Color(147,0,0)

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Indent: First line: 0"

Deleted: More technical observations are these: one implication of this tendency for homogeneity in the  $\hat{x}_{ij}^0$  estimates is that the MERET technique works best with a large set of alternative fire-type classes.

Deleted: Adding

Deleted: also

Deleted: s

Deleted: , and

Deleted: are

Deleted: avoided

Deleted: .

Deleted: .

Formatted: Font: Italic

Deleted: For CO, the x and y intercepts for each segment joining observations are shown in a light red color near the origin.

Deleted: background not evident

Deleted:

Deleted: T

Deleted: ,

Deleted: however. The observation points are marked by colors following the fire-type class we obtained using all tracers in the rough categorization of correlated variations in slopes. (Refer back to Section 6.2 and Section 6.3, step 5 – "Roughly cluster".) The colors are correlated from one observation to the next, suggesting that the somewhat differing plumes characterized are resolved with our sampling frequency. Variation of slope is more evident for  $b_{\text{scat}}$  than CO. Generally, the slopes for CO tend to be more uniform in all flights analyzed.

Deleted: ¶ ... [61]

Deleted: 19

1480

sions, but produced in the plumes. The relationships to fuel burned, and their variations are nevertheless interesting. Descriptions of variation are given as the 16<sup>th</sup> and 84<sup>th</sup> percentiles of all the estimates. These are similar to error estimates if nothing more is known about the origin and age of the particular samples, a matter more fully discussed in Chatfield and Andreae (2020). EnR's as they varied in time, and in relationship to measures of photochemical processing, are shown in the SM: "Note on Examples of Enhancement Ratios Variation in Time."

Deleted: 2019

Formatted: Font: Not Bold

Table 3. EnR Estimates for Fire Emissions Considered

Fire Emission	EnR estimate	Percentile 16	Percentile 84	Unit	Conversion factor to EF
CO	74	62	85	ppb ppm <sup>-1</sup>	1.17
CH <sub>4</sub>	8.6	2.3	13.0	ppb ppm <sup>-1</sup>	0.67
Ethyne	0.26	0.205	0.31	ppb ppm <sup>-1</sup>	1.08
Ethene	0.88	0.65	1.07	ppb ppm <sup>-1</sup>	1.17
Ethane	0.70	0.57	0.80	ppb ppm <sup>-1</sup>	1.25
Propene	0.056	0.005	0.100	ppb ppm <sup>-1</sup>	1.75
Propane	0.16	0.12	0.19	ppb ppm <sup>-1</sup>	1.83
n-Butane	0.028	0.019	0.037	ppb ppm <sup>-1</sup>	2.17
Benzene	0.094	0.073	0.134	ppb ppm <sup>-1</sup>	3.25
Toluene	0.054	0.023	0.067	ppb ppm <sup>-1</sup>	3.88
Methanol	2.1	1.7	3.1	ppb ppm <sup>-1</sup>	1.33
HCHO	1.15	0.81	1.62	ppb ppm <sup>-1</sup>	1.25
Acetaldehyde	0.56	0.24	0.71	ppb ppm <sup>-1</sup>	1.83
Acetone	0.74	0.54	1.14	ppb ppm <sup>-1</sup>	2.42
CH <sub>3</sub> CN	0.13	0.11	0.16	ppb ppm <sup>-1</sup>	1.25
NO <sub>x</sub> (as N)	0.051	0.024	0.131	ppb ppm <sup>-1</sup>	0.63
O <sub>3</sub>	14.8	8.5	25.1	ppb ppm <sup>-1</sup>	(2.0)
PAN	0.26	0.17	0.38	ppb ppm <sup>-1</sup>	(3.17)
Scat <sub>5</sub> , b <sub>Scat</sub>	79	50	100	m <sup>-1</sup> ppm <sup>-1</sup>	0.042
Abs <sub>5</sub> , b <sub>Abs</sub>	3.2	2.2	4.4	m <sup>-1</sup> ppm <sup>-1</sup>	0.042
Ammonium	0.32	0.19	0.47	μg m <sup>-3</sup> ppm <sup>-1</sup>	0.032
Nitrate	0.28	0.11	0.60	μg m <sup>-3</sup> ppm <sup>-1</sup>	0.107
Sulfate	0.156	0.063	0.290	μg m <sup>-3</sup> ppm <sup>-1</sup>	0.164

Notes: Conversions assume a C to dry biomass ratio of 0.5. Conversions to μg m<sup>-3</sup> assume 25 °C and 1013 hPa. O<sub>3</sub> and PAN are not directly produced by fires. HCHO is produced but often decreases rapidly. Under appropriate conditions indicated in Chatfield and Andreae (2020), the EnR estimates can be used as ERs. For tracers that are rapidly removed or transformed, these tend to be the higher values.

Deleted: 19

Deleted: ¶

9. Conclusions

A major problem with the estimation of fire enhancement ratios and emission factors is inherent in their character: flames promote mixing in their plumes. Total carbon liberated to the atmosphere (approximately C<sub>burn</sub> = CO<sub>2</sub>+CO) is mixed with background air at different points in the plume's evolution, and removal of that mixing effect has been a difficulty. The NEMR technique

1485

often uses CO as unique tracer, but the EnR of CO is variable, adding uncertainty to the estimation of the EFs. Given the variability of CO due to combustion efficiency (MCE) and environmental variability, it has been emphasized that the NEMR technique can only be confidently applied in situations in which conditions affecting the ratio of CO to (CO<sub>2</sub>+CO) can be well determined, ideally from source to sampling (Yokelson et al., 2013). The method also tends to emphasize the use of samples of CO and tracer collected over many minutes, so that the regression method for EnRs of tracer relative to CO, defining  $a_{CO} \leftarrow (\text{fire-added } C_{\text{burn}})$  becomes stable, and a conversion to fuel carbon burned becomes possible.

We sought to decompose  $C_{\text{tot}}$  into  $C_{\text{bkgd}}$  and  $C_{\text{burn}}$ . However, meteorology and mixing allow significant variations in  $C_{\text{bkgd}}$  due to other powerful processes, e.g., CO<sub>2</sub> from respiration/photosynthesis in mixed-layer air. Once lofted,  $C_{\text{bkgd}}$  varies little unless the plume enters layers of free tropospheric air from long-range transport with different  $C_{\text{bkgd}}$ , which further dilute the plume (Yokelson et al., 2013). We noted such problems using NEMR in analyzing a significant number of plumes for enhancement ratios studied in the Western US during two campaigns, ARCTAS-California and SEAC4RS, with 422 one-minute samples in all.

The problem of deriving an accurate  $C_{\text{bkgd}}$  is solved by noticing that there are two different kinds of information provided by multiple observational instances of a tracer and multiple tracers at a single instance. Information about the various EnRs and  $C_{\text{bkgd}}$  are mixed, but not inextricably. There is a solution based on mixed-effects (also called random effects) regression modeling. We propose a Mixed Effects Regression Emissions Technique (MERET) to replace or at least to check on NEMR, for which we used the **R** routine *lmer()*.

The MERET technique is related to traditional entraining-plume models for parcels. We presented a synthesis describing multiple tracers from fire to sampling location. Sample calculations with the model suggest that it deals appropriately (linearly) with several varied histories for plume mixing. This motivates a regression equation for an “equivalent background”  $x_i^0$  for each observation that is related to entraining concentrations  $x^E(t)$  along the trajectory, and is appropriate for each tracer species (Figure 8). The theory then allows this  $x_i^0$  to be used to define  $C_{\text{burn}}$  and thus to define the EnR for any appropriate fire-derived variable. This technique makes EnRs appear appropriate even in difficult situations, and allows estimates of EnRs for individual samples.

EnRs are appropriate for the estimation of emission factors when the plume age is short compared to the transformation timescale of the measured fire tracer, and we provide an approximate diagnostic for this age for most samples. Formaldehyde, acetaldehyde, the alkenes, benzene, NO<sub>x</sub>,  $b_{\text{Scat}}$ , and  $b_{\text{Abs}}$  particularly require such attention.

Carbon monoxide is usually the best single tracer that correlates with fire emissions ( $C_{\text{burn}}$ ), supporting the use of the NEMR technique. Our analysis suggests other tracers had EnR variations that collectively helped to distinguish  $C_{\text{burn}}$  from CO in regression. The NEMR methodology depends on a full analysis of the history of CO influences on a sample to obtain a reliable MCE. MERET allows estimates of MCE as well as  $C_{\text{burn}}$  for each sample. Thus it demarcates sampling periods with nearly homogeneous MCE. However, possible large variations in the entraining background of CO should still be considered carefully in dilute plumes with  $C_{\text{burn}} < 2$  ppm.

Deleted:

Deleted: "

Deleted: "

Deleted: Figure 8).

Deleted: Analysis of the entire dataset from the Rim Fire gave relatively good comparisons of NEMR and MERET. However, NEMR had to be applied for a very long time period (> 30 min); shorter periods gave puzzling results. Variability of the MCE and the  $C_{\text{bkgd}}$  each seemed to have a role. Our MERET analysis suggested various fire conditions and ages for the airborne sample during this period. The time series of EnRs suggested appropriate sub-periods where the NEMR could be more skillfully used. ¶

Deleted: NMER

### 9.1. Questions for future research

We conclude with some questions for future research; these also review the suggested conclusions of this paper and acknowledge the limitations of a single publication.

- (1) How well can the use of one or a few tracers, e.g., CO,  $b_{\text{scat}}$ , HCHO, actually constrain EnRs and EFs when only a few instruments may be used? How many variables need to be measured or how fresh should the plumes be to allow CO to be used both as a fire tracer and to allow useful estimates of MCE?
- (2) Can MERET be used to identify time periods of relatively homogeneous MCE, and can that MCE value be used with NEMR to create suitable EnRs? Since NEMR uses differences sample by sample (in time), no minimum value of another tracer need be estimated. (Consider that MERET does allow some evaluation of the minimum value estimate to be assessed and a better minimum assigned.)
- (3) Can the MERET/NEMR and better near-fire non-plume sampling help us to prevent misattribution of fire emissions? These would include observations for fire intake air, air likely to be entrained in ascent, and air surrounding a plume and likely to be entrained as a plume spreads downwind. Can simulations of entraining plumes aid this effort?
- (4) What do “fire types” represent and which species or properties tend to correlate in their EnRs (Chatfield and Andreae, 2020).

#### *Science Team List*

##### **ARCTAS Science Team:**

James Crawford (NASA Langley Research Center), Henry Fuelberg (Florida State University), Chris Hostetler (NASA Langley Research Center), Daniel Jacob (Harvard University), Hal Maring (NASA Headquarters), Philip B. Russell (NASA Ames Research Center), Kent Shiffer (NASA Ames Research Center), Hanwant Singh (NASA Ames Research Center), Kathy Thompson (Science Systems and Applications, Inc.), Bruce E. Anderson (NASA Langley Research Center), Eric Apel (National Center for Atmospheric Research), Donald R. Blake (University of California Irvine), William H. Brune (Pennsylvania State University), Christopher A. Cantrell (National Center for Atmospheric Research), Ronald C. Cohen (University of California Berkeley), Jack Dibb (University of New Hampshire), Glenn S. Diskin (NASA Langley Research Center), Alan Fried (University of Colorado), Johnathan W. Hair (NASA Langley Research Center), L. Greg Huey (Georgia Tech), Jose-Luis Jimenez (CIRES, University of Colorado), Yutaka Kondo (University of Tokyo), Richard E. Shetter (University of North Dakota), Robert Talbot (University of New Hampshire), Stephanie A. Vay (NASA Langley Research Center), Rodney Weber (Georgia Tech), Andrew Weinheimer (National Center for Atmospheric Research), Paul Wennberg (Cal Tech), Armin Wisthaler (Universität Innsbruck and Universitet i Oslo), John Barrick (NASA Langley Research Center), Anthony Bucholtz (Naval Research Laboratory), Antony Clarke (University of Hawaii), Lawrence C. Freudinger (NASA Armstrong Flight Center), Charles Gatebe (NASA Goddard Space Flight Center), Athanasios Nenes (Georgia Tech), James Podolske (NASA Ames Research Center), Jens Redemann (University of Oklahoma), Sebastian Schmidt (CU Boulder), Anthony Strawa (NASA Ames Research Center), Brian Cairns (NASA Goddard Institute for Space Science), Richard Ferrare (NASA Langley Research Center), Samuel Oltmans (NOAA ESRL, CU Boulder), Glenn Shaw (University of Alaska Fairbanks), Anne M. Thompson (NASA Goddard Space Flight Center), Bob Curry (NASA Armstrong Flight Center), David Easmunt (NASA Goddard Space Flight Center), Michael S. Wusk (NASA Langley Research Center).

Deleted: 2019

Formatted: Normal, No bullets or numbering

Formatted: Font: Italic

**SEAC4RS Science Team:**

- 1595 Hal Maring (NASA Headquarters), Ken Jucks (NASA Headquarters), Jassim Al-Saadi (NASA Headquarters), Richard Eckman (NASA Headquarters), Alex Pszenny (NASA Headquarters), Kent Shiffer (NASA Ames Research Center), Jhony Zavaleta (NASA Ames Research Center), Michael Craig (NASA Ames Research Center), David Jordan (NASA Ames Research Center), Quincy Allison (NASA Ames Research Center), Dan Chirica (NASA Ames Research Center),
- 1600 Susan Tolley (NASA Ames Research Center), Erin Czech (NASA Ames Research Center), Erin Justice (NASA Ames Research Center), Katja Drdla (NASA Ames Research Center), Brian Toon (University of Colorado), Eric Jensen (NASA Ames Research Center), Rich Ferrare (NASA Langley Research Center), David Diner (NASA Jet Propulsion Laboratory), Anthony Bucholtz (Naval Research Laboratory), Lance Christensen (NASA Jet Propulsion Laboratory),
- 1605 Matt McGill (NASA Goddard Space Flight Center), Steve Platnick (NASA Goddard Space Flight Center), Brian Cairns (NASA Goddard Space Flight Center), Jim Anderson (Harvard University), Rushan Gao (NOAA ESRL), Bob Herman (NASA Jet Propulsion Laboratory), MJ Mahoney (NASA Jet Propulsion Laboratory), Rushan Gao (NOAA ESRL), Steve Wofsy (Harvard University), Sebastian Schmidt (University of Colorado), Elliot Atlas (University of Miami),
- 1610 Paul Bui (NASA Ames Research Center), Hanwant Singh (NASA Ames Research Center), Jack Dibb (University of New Hampshire), Phillip Russell (NASA Ames Research Center), Jose-Luis Jimenez (University of Colorado), Charles Brock (NOAA ESRL), Simone Tanelli (NASA Jet Propulsion Laboratory), Andreas Beyersdorf (NASA Langley Research Center), Anthony Bucholtz (Naval Research Laboratory), Samuel Hall (National Center for Atmospheric Research),
- 1615 Paul Wennberg (Cai Tech), Thomas Ryerson (NOAA ESRL), Glenn Diskin (NASA Langley Research Center), Armin Sorooshian (University of Arizona), Alan Fried (National Center for Atmospheric Research), John Hair (NASA Langley Research Center), Glenn Diskin (NASA Langley Research Center), Greg Huey (Georgia Tech), Rushan Gao (NOAA ESRL), Thomas Hanisco (NASA Goddard Space Flight Center), Bruce Anderson (NASA Langley Research Center), Paul Bui (NASA Ames Research Center), Karl Froyd (NOAA ESRL), Greg Huey (Georgia Tech),
- 1620 Armin Wisthaler (Universität Innsbruck), Jack Dibb (University of New Hampshire), Paul Lawson (SPEC Incorporated), Sebastian Schmidt (University of Colorado), Ronald Cohen (University of California Berkeley), Donald Blake (University of California Irvine), Joshua Schwarz (NOAA ESRL), J. Vanderlei Martins (University of Maryland Baltimore County), Lenny Pfister (NASA Ames Research Center), Henry Fuelberg (NASA Langley Research Center), Rennie Selkirk (Universities Space Research Association), Daniel Jacob (Harvard University), Arlindo Da Silva (NASA Goddard Space Flight Center), Paul Konopka (Georgia Tech), Ed Hyer (Naval Research Laboratory), Greg Charmichael (University of Iowa), Jennifer Olson (NASA Langley Research Center), Louisa Emmons (National Center for Atmospheric Research), Gao Chen (NASA Langley Research Center), Chip Trepte (NASA Langley Research Center), Jay Mace (University of Utah), Pat Minnis (NASA Langley Research Center), Eric Ray (NOAA ESRL), Karen Rosenlof (NOAA ESRL), Daniel Jacob (Harvard University), Robert Yokelson (University of Montana), Jeffery Reid (Naval Research Laboratory), Jens Redemann (NASA Ames Research Center), David Starr (NASA Goddard Space Flight Center), Johnny Luo (City College of New York), Jay Mace (University of Utah), Laura Pan (National Center for Atmospheric Research), David Starr (NASA Goddard Space Flight Center), Jay Mace (University of Utah), Johnny Luo (City College of New York), James Crawford (NASA Langley Research Center), Frank Culter (NASA Armstrong Flight Center), Patrick Lloyd (NASA Armstrong Flight Center), Rick Shetter

1640 (National Suborbital Education and Research Center), Adam Webster (National Suborbital Edu-  
cation and Research Center), Eric Buzay (National Suborbital Education and Research Center),  
David VanGilst (National Suborbital Education and Research Center), Jane Peterson (National  
Suborbital Education and Research Center), Tim Moes (NASA Armstrong Flight Center), Chris  
Miller (NASA Armstrong Flight Center), Mike Kapitzke (NASA Armstrong Flight Center),  
1645 Denis Steele (NASA Armstrong Flight Center), Stuart Broce (NASA Armstrong Flight Center),  
Jan Nystrom (NASA Armstrong Flight Center), Kevin Kraft (NASA Armstrong Flight Center).

1650 Some of the team have changed affiliations since their work was documented. Two members of  
NASA's Earth Science Projects Office (ESPO) may direct you on how to make enquiries regard-  
ing either team, marilyn.vasques@nasa.gov and bernadette.luna@nasa.gov. Also, a leader at  
NASA Headquarters, has some knowledge of both teams' members, barry.lefer@nasa.gov.

1655 *Data availability:* ARCTAS and SEAC4RS data are available for download from these perma-  
nent NASA data repositories: Arctic Research of the Composition of the Troposphere from  
Aircraft and Satellites (ARCTAS, <https://www-air.larc.nasa.gov/missions/arctas/arctas.html>, and  
SEAC4RS – Studies of Emissions and Atmospheric Composition, Clouds and Climate Coupling  
by Regional Surveys <https://www-air.larc.nasa.gov/missions/seac4rs/>.

1660 *Supplement:* The supplement related to this article is available on- line at: (TBD)

#### 1660 *Acknowledgements*

Particular acknowledgment goes to the instrumental groups in Table 1 and Section 4 and the  
many contributing efforts of the ARCTAS and SEAC4RS Science Teams and staff. Chatfield  
pursued this research under NASA grant 13-ACCDAM13-0110, and the effort grew out of re-  
search funded by California Air Resources Board and San Joaquin Valley air regional authority,  
1665 where difficulties with previous methodology became apparent. We appreciate reviews by S. Pa-  
lacios, M. Segal-Rozenhaimer, E. Yates, and especially R. Koppmann. M. O. Andreae acknowl-  
edges extensive support by the German Max Planck Society and facilitation of recent research by  
the Scripps Institute of Oceanography. Robert F. Esswein helped with certain data reformatting.  
1670 We acknowledge the great value of the R programming language and the lme4 and NMF pack-  
ages.

#### 1675 *Author contributions*

MOA contributed motivating ideas and approach following on Yokelson et al. (2013). RBC  
brought data into suitable format and evaluated suitability of tracers. He contributed the theory of  
parallel plume rise expressions for fire tracers and the mixed-effects modeling approach. RBC  
wrote the publication with considerable contributions of MOA.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Normal, No bullets or numbering

Formatted: Font: Italic

Formatted: Normal, No bullets or numbering

## 10. References

- 1680 Akagi, S. K., Yokelson, R. J., Burling, I. R., Meinardi, S., Simpson, I. J., Blake, D. R., McMeeking, G. R., Sullivan, A., Lee, T., Kreidenweis, S., Urbanski, S., Reardon, J., Griffith, D. W. T., Johnson, T. J. and Weise, D. R.: Measurements of reactive trace gases and variable O<sub>3</sub> formation rates in some South Carolina biomass burning plumes, *Atmospheric Chemistry and Physics*, 13, 1141–1165, doi:[10.5194/acpd-12-25255-2012](https://doi.org/10.5194/acpd-12-25255-2012), 2013.
- 1685 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D. and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, doi:[10.5194/acpd-10-27523-2010](https://doi.org/10.5194/acpd-10-27523-2010), 2011.
- Akagi, SK, et al. Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmos. Chem Phys.*, 12, 1397–1421, 2012.
- 1690 Alvarado, M. J. and Prinn, R. G.: Formation of ozone and growth of aerosols in young smoke plumes from biomass burning: 1. Lagrangian parcel studies, *J. Geophys. Res.-Atmos.*, 114, doi:[10.1029/2008jd011144](https://doi.org/10.1029/2008jd011144), 2009.
- Alvarado, M. J., Wang, C. and Prinn, R. G.: Formation of ozone and growth of aerosols in young smoke plumes from biomass burning: 2. Three-dimensional Eulerian studies, *J. Geophys. Res.-Atmos.*, 114, doi:[10.1029/2008jd011186](https://doi.org/10.1029/2008jd011186), 2009.
- 1695 Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K. E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., St Clair, J. M., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C. and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10(20), 9739–9760, doi:[10.5194/acp-10-9739-2010](https://doi.org/10.5194/acp-10-9739-2010), 2010.
- 1700 Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – An updated assessment: *Atmos. Chem. Phys. Discuss.*, 2019, 1–27, doi:[10.5194/acp-2019-303](https://doi.org/10.5194/acp-2019-303), 2019.
- 1705 Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochemical Cycles*, 15(4), 955–966, doi:[10.1029/2000GB001382](https://doi.org/10.1029/2000GB001382), 2001.
- Andreae, M. O., Browell, E. V., Garstang, M., Gregory, G. L., Harriss, R. C., Hill, G. F., Jacob, D. J., Pereira, M. C., Sachse, G. W., Setzer, A. W., Silva Dias, P. L., Talbot, R. W., Torres, A. L., and Wofsy, S. C.: Biomass burning emissions and associated haze layers over Amazonia, *J. Geophys. Res.*, 93, 1509–1527, 1988.

Barry, R.G., Chorley, R.J.: Atmosphere, Weather and Climate (7th edition), Routledge (London), 1998.

Bates D, Maechler M, Bolker B and Walker S: lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-7, <http://CRAN.R-project.org/package=lme4>, <http://arxiv.org/abs/1406.5823>, 2014.

Baylon, P., Jaffe, D. A., Wigder, N. L., Gao, H. and Hee, J.: Ozone enhancement in western US wildfire plumes at the Mt. Bachelor Observatory: The role of NO<sub>x</sub>, Atmospheric Environment, 109, 297–304, doi:10.1016/j.atmosenv.2014.09.013, 2015.

Boutsidis, C., and Gallopoulos, E.: SVD based initialization: A head start for nonnegative matrix factorization, In: Pattern Recognition 41., 1350–1362. issn: 00313203. doi: 10.1016/j.patcog.2007.09.010, 2008.

Burling, I. R., Yokelson, R. J., Griffith, D. W. T., Johnson, T. J., Veres, P., Roberts, J. M., Warnke, C., Urbanski, S. P., Reardon, J., Weise, D. R., Hao, W. M. and de Gouw, J.: Laboratory measurements of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States, Atmospheric Chemistry and Physics, 10(22), 11115–11130, doi:10.5194/acp-10-11115-2010, 2010.

Chatfield, R.B.: package Estim.ARC.SEAC.ERs, <https://github.com/RobertBChatfield/FireEmissionsEstimate/>, 2002.

Chatfield, R.B., and Andreae, M.O.: Emissions relationships among western forest fire plumes: II. Plume typing for sources of ozone and aerosol absorption, advanced draft in preparation for Atmos. Chem. Phys. 2020.

Crutzen, P.J., Heidt, L.E., Krasnec, J.P., Pollock, W.H., Seiler, W., Biomass burning as a source of atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl and COS. Nature 282, 253e256, 1979. P J. Crutzen, L E. Heidt, J P. Krasnec, W H. Pollock and W Seiler: Biomass burning as a source of atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl, and COS, Nature, 282, 253–256, 1979.

Crutzen, P.J., and Andreae, M.O.: Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles, Science, 250, no. 4988, 1669–1678, DOI:10.1126/science.250.4988.1669, 1990.

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, Megan J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath II, T., Docherty, K. S., Worsnop, and D. R., Jimenez, J.L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, Anal. Chem. 78, 8281-8289, 2006.

Ding, C., He, X., and Simon, H.D.: On the Equivalence of Nonnegative Matrix Factorization and Spectral Clustering, Proceedings of the 2005 SIAM International Conference on Data Mining, 10.1137/1.9781611972757.70, also permalink <https://doi.org/10.1137/1.9781611972757.70>, 2005.

Deleted: 19

Formatted: Font: Times New Roman, 12 pt, Font color: Auto

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto



- Forrister, H., Liu, J., Scheuer, E., Dibb, J., Ziemba, L., Thornhill, K. L., Anderson, B., Diskin, G., Perring, A. E., Schwarz, J. P., Campuzano-Jost, P., Day, D. A., Palm, B. B., Jimenez, J. L., Nenes, A., and Weber, R. J.: Evolution of brown carbon in wildfire plumes, *Geophys. Res. Lett.*, 42, 4623–4630, <https://doi.org/10.1002/2015GL063897>, 2015.
- Frenklach, M., Reaction mechanism of soot formation in flames: *Physical Chemistry Chemical Physics*, 4, 2028–2037, doi:10.1039/b110045a, 2002.
- Fried, A., Walega, J. G., Olson, J. R., Crawford, J. H., Chen, G., Weibring, P., Richter, D., Roller, C., Tittel, F. K., Heikes, B. G., Snow, J. A., Shen, H., O'Sullivan, D. W., Porter, M., Fuelberg, H., Halland, J., and Millet, D.: Formaldehyde Over North America and the North Atlantic During the Summer 2004 INTEX Campaign: Methods, Observed Distributions, and Measurement-Model Comparisons, *J. Geophys. Res.*, {113, }D10302, <https://doi.org/10.1029/2007JD009185>, 2008.
- Galanter, M., Levy II, H., and Carmichael, G. R.: Impacts of biomass burning on tropospheric CO, NO<sub>x</sub>, and O<sub>3</sub>, *J. Geophys. Res.*, 105, 6633–6653, doi:10.1029/1999JD901113, 2000.
- Gaujoux, R., Seoighe, C.: A flexible R package for nonnegative matrix factorization, *BMC Bioinformatics* 11.1 p. 367. issn: 1471-2105. doi: 10.1186/1471-2105-11-367, 2010.
- Gaujoux, R.: An Introduction to NMF package, <https://cran.r-project.org/web/packages/NMF/vignettes/NMF-vignette.pdf>, 2014.
- Greenberg J. P., Zimmerman P. R., Heidt L., and Pollock W.: Hydrocarbon and carbon monoxide emissions from biomass burning in Brazil, *J. Geophys. Res.* 89 (D1), 1350–1354. 1984.
- Gueniche, H. A., Biet, J., Glaude, P. A., Fournet, R., and Battin-Lederc, F.: A comparative study of the formation of aromatics in rich methane flames doped by unsaturated compounds: *Fuel*, 88, 1388–1393, doi:10.1016/j.fuel.2009.03.006, 2009.
- Guyon, P., Frank, G. P., Welling, M., Chand, D., Artaxo, P., Rizzo, L., Nishioka, G., Kolle, O., Fritsch, H., Silva Dias, M. A. F., Gatti, L. V., Cordova, A. M., and Andreae, M. O.: Airborne measurements of trace gases and aerosol particle emissions from biomass burning in Amazonia: *Atmos. Chem. Phys.*, 5, 2989–3002, 2005.
- Hanna, S.R., Briggs, G.A., Hosker, R.P.: Handbook on Atmospheric Dispersion. Prepared for the US Department of Energy, Technical Information Center, US Department of Energy(DOE/TIC-11223), 1982.
- Hobbs, P. V., Sinha, P., Yokelson, R. J., Christian, T. J., Blake, D. R., Gao, S., Kirchstetter, T. W., Novakov, T., and Pilewskie, P.: Evolution of gases and particles from a savanna fire in South Africa, *J. Geophys. Res.*, 108, 8485, doi:10.1029/2002JD002352, 2003.

1785 Hornbrook, R. S., Blake, D. R., Diskin, G. S., Fried, A., Fuelberg, H. E., Meinardi, S., Mikoviny, T., Richter, D., Sachse, G. W., Vay, S. A., Walega, J., Weibring, P., Weinheimer, A. J., Wiedinmyer, C., Wisthaler, A., Hills, A., Riemer, D. D., and Apel, E. C.: Observations of non-methane organic compounds during ARCTAS – Part 1: Biomass burning emissions and plume enhancements, *Atmos. Chem. Phys.*, 11, 11103-11130, <https://doi.org/10.5194/acp-11-11103-2011>, 2011.

Jaffe, D., Wigder, N.: Ozone production from wildfires: a critical review. *Atmos. Environ.* 51, 1–10, <http://dx.doi.org/10.1016/j.atmosenv.2011.11.063>, 2012.

1790 Janhäll, S., Andreae, M. O., and Pöschl, U.: Biomass burning aerosol emissions from vegetation fires: particle number and mass emission factors and size distributions, *Atmos. Chem. Phys.*, 2001, 10, 1427-1439, doi:10.5194/acp-10-1427-2010.

1795 Koppmann, R., A. Khedim, J. Rudolph, D. Poppe, M. O. Andreae, G. Helas, M. Welling, and T. Zenker: Emissions of organic trace gases from savanna fires in southern Africa during the 1992 Southern African Fire Atmosphere Research Initiative and their impact on the formation of tropospheric ozone, *J. Geophys. Res.*, 102, 18,879–18,888, 1997.

1800 Koppmann, R., von Czapiewski, K., and Reid, J. S.: A review of biomass burning emissions, part I: gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds, *Atmos. Chem. Phys.*, 5, 10455-10516, <https://doi.org/10.5194/acpd-5-10455-2005>, 2005

1805 Langridge, J. M., Richardson, M. S., Lack, D. A., Law, D. C., and Murphy, D. M.: Aircraft instrument for comprehensive characterization of aerosol optical properties, Part I: Wavelength-dependent optical extinction and its relative humidity dependence measured using cavity ring-down spectroscopy, *Aerosol Science and Technology*, 45, 1305-1318  
doi10.1080/02786826.2011.592745, 2011.

Laplace, P.S., *Mémoire sur la Probabilité des Causes par les Événemens*, *Mémoires de Mathématique et de Physique, Présentés à l'Académie Royale des Sciences, Par Divers Savans & Lûs Dans ses Assemblées*, 6, 621-656, 1774.

1810 Lawrence, J.: Distribution of the Median in Samples from the Laplace Distribution, *Open Journal of Statistics*, 3 (6), DOI:10.4236/ojs.2013.36050, 2013.

Lee, D.D., and Seung, H.S.: Algorithms for non-negative matrix factorization, *Advances in Neural Information Processing Systems*, url: <http://scholar.google.com/scholar?q=intitle:Algorithms+for+non-negative+matrix+factorization>, 2001.

1815 Leeuwen, T. T., and G. R. van der Werf: Spatial and temporal variability in the ratio of trace gases emitted from biomass burning. *Atmos. Chem. Phys.*, 11, 3611-3629, 2011.

Deleted: ¶

Lefer, B. L., Talbot, R. W., Harriss, R. C., Bradshaw, J. D., Sandholm, S. T., Olson, J. O., Sachse, G. W., Collins, J., Shipman, M. A., Blake, D. R., Klemm, K. I., Gorzelska, K., and Barrick, J.: Enhancement of acidic gases in biomass burning impacted air masses over Canada, *J. Geophys. Res.* 99, 1721–1737, doi: 10.1029/93JD02091, 1994. ¶  
Leighton, P.: *The Photochemistry of Air Pollution*, Academic Press, 1961. ¶

Moved (insertion) [6]

Deleted: ¶

- Lefer, B. L., Talbot, R. W., Harriss, R. C., Bradshaw, J. D., Sandholm, S. T., Olson, J. O., Sachse, G. W., Collins, J., Shipman, M. A., Blake, D. R., Klemm, K. I., Gorzelska, K., and Bar-  
rick, J.: Enhancement of acidic gases in biomass burning impacted air masses over Canada, J.  
Geophys. Res. 99, 1721–1737, doi: 10.1029/93JD02091, 1994.
- 1830 Leighton, P.: The Photochemistry of Air Pollution, Academic Press, 1961.
- Liu, X., Huey, L. G., Yokelson, R. J., Selimovic, V., Simpson, I. J., Müller, M., Jimenez, J. L.,  
Campuzano-Jost, P., Beyersdorf, A. J., Blake, D. R., Butterfield, Z., Choi, Y., Crounse, J. D.,  
Day, D. A., Diskin, G. S., Dubey, M. K., Fortner, E., Hanisco, T. F., Hu, W., King, L. E., Klein-  
man, L., Meinardi, S., Mikoviny, T., Onasch, T. B., Palm, B. B., Peischl, J., Pollack, I. B.,  
1835 Ryerson, T. B., Sachse, G. W., Sedlacek, A. J., Shilling, J. E., Springston, S., St. Clair, J. M.,  
Tanner, D. J., Teng, A. P., Wennberg, P. O., Wisthaler, A., and G. M. Wolfe.: Airborne measure-  
ments of western U.S. wildfire emissions: comparison with prescribed burning and air quality  
implications, J. Geophys. Res.-Atmos., 122, 6108–6129, <https://doi.org/10.1002/2016JD026315>,  
2017.
- 1840 McKeen, S. A., Liu, S. C., Hydrocarbon ratios and photochemical history of air masses, Ge-  
ophys. Res. Lett., <http://dx.doi.org/10.1029/93GL02527>, 1984.
- McKeen, S.A., Liu, S.C., Hsie, E.-Y., Lin, X., Bradshaw, J.D., Smyth, S., Gregory, G.L., and  
Blake, D.R.: Hydrocarbon ratios during PEM-WEST A: A model perspective, J. Geophys. Res.,  
101, 2087–2109, doi: 10.1029/95JD02733, 1996.
- 1845 Morton, B. R., G. I. Taylor, and J. S. Turner, 1956: Turbulent gravitational convection from  
maintained and instantaneous sources. Proc. Roy. Soc. London, A234,1-23.
- Poppe, D., Koppmann, R. and Rudolph, J.: Ozone formation in biomass burning plumes: Influ-  
ence of atmospheric dilution, Geophys. Res. Lett., 25, 3823–3826, 1998.
- National Research Council: Estimating Mortality Risk Reduction and Economic Benefits from  
1850 Controlling Ozone Air Pollution. National Academies Press, Washington D.C., 2008.
- Pfister, G. G., Avise, J., Wiedinmyer, C., Edwards, D. P., Emmons, L. K., Diskin, G. D., Po-  
dolske, J., and Wisthaler, A.: CO source contribution analysis for California during ARCTAS-  
CARB, Atmos. Chem. Phys., 11, 7515-7532, <https://doi.org/10.5194/acp-11-7515-2011>, 2011.
- Pinheiro, J.C., and Bates, D.M.: Mixed-Effects Models in S and S-PLUS, Springer, New York,  
1855 2000.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emis-  
sions part II: intensive physical properties of biomass burning particles, Atmos. Chem. Phys., 5,  
799-825, <https://doi.org/10.5194/acp-5-799-2005>, 2005.

Moved down [7]: Leighton, P.: The Photochemistry of  
Air Pollution, Academic Press, 1961.¶

Moved (insertion) [7]

Deleted: ¶

Moved up [6]: Leeuwen, T. T., and G. R. van der Werf:  
Spatial and temporal variability in the ratio of trace gases  
emitted from biomass burning. Atmos. Chem. Phys., 11,  
3611-3629, 2011.¶

- Roberts, J. M., F. C. Fehsenfeld, S. C. Liu, M. J. Bollinger, C. Hahn, D. L. Albritton, and R. E. Sievers: Measurements of aromatic hydrocarbon ratios and NO<sub>x</sub> concentrations in the rural troposphere: Estimates of air mass photochemical age and NO<sub>x</sub> removal rate, *Atmos. Environ.*, 18, 2421–2432, 1984.
- 1870 Ryerson, T. B., Williams, E. J., and Fehsenfeld, F.C.: An efficient photolysis system for fast-response NO<sub>2</sub> measurements, *J. Geophys. Res.-Atmos.*, doi:10.1029/2000JD900389, 2000.
- Seinfeld, J.H. and S.N. Pandis, *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 3rd Edition, Wiley, 2016.
- 1875 Simpson, I. J., Akagi, S. K., Barletta, B., Blake, N. J., Choi, Y., Diskin, G. S., Fried, A., Fuelberg, H. E., Meinardi, S., Rowland, F. S., Vay, S. A., Weinheimer, A. J., Wennberg, P. O., Wiering, P., Wisthaler, A., Yang, M., Yokelson, R. J., and Blake, D. R.: Boreal forest fire emissions in fresh Canadian smoke plumes: C1-C10 volatile organic compounds (VOCs), CO<sub>2</sub>, CO, NO<sub>2</sub>, NO, HCN and CH<sub>3</sub>CN, *Atmos. Chem. Phys.*, 11, 6445–6463, <https://doi.org/10.5194/acp-11-6445-2011>, 2011.
- 1880 Singh, H. B., and the ARCTAS Science Team: Pollution influences on atmospheric composition and chemistry at high northern latitudes: Boreal and California forest fire emissions, *Atmos. Environ.*, 44, 4553–4564, 2010.
- 1885 Singh, H. B., Cai, C., Kaduwela, A., Weinheimer, A., and Wisthaler, A.: Interactions of fire emissions and urban pollution over California: Ozone formation and air quality simulations, *Atmos. Environ.*, 56, 2, 45–51, 2012.
- Squires, P. and Turner, J.S.: An entraining jet model for cumulo-nimbus updraughts. *Tellus A*. 10.3402/tellus.v14i4.9569.1962.
- 1890 Toon, O. B., Maring, H., Dibb, J., Ferrare, R., Jacob, D. J., Jensen, E. J., Luo, Z. J., Mace, G. G., Pan, L. L., Pfister, L., Rosenlof, K. H., Redemann, J., Reid, J. S., Singh, H. B., Thompson, A. M., Yokelson, R., Minnis, P., Chen, G., Jucks, K. W., Pszenny A.: Planning, implementation and scientific goals of the Studies of Emissions and Atmospheric Composition, Clouds and Climate Coupling by Regional Surveys (SEAC4RS) field mission, *J. Geophys. Res. – Atmos.*, 121, 4967–5009, doi:10.1002/2015JD024297, 2016.
- 1895 Trenberth, K.: Some effects of finite sample size and persistence on meteorological statistics. Part I: Autocorrelations, *Monthly Weather Review*, 112, 2359–2368, 1984.
- Trentmann, J., Andreae, M. O., Graf, H.-F., Hobbs, P. V., Ottmar, R. D., and Trautmann, T.: Simulation of a biomass burning plume: Comparison of model results with observations: *J. Geophys. Res.*, 107, 4013, doi:10.1029/2001JD000410, 2002.
- 1900 U.S. EPA: U.S. Environmental Protection Agency 2002 National Emissions Inventory. Available: <http://www.epa.gov/ttn/chief/net/2002invent>, 2006.

- Wagner, N. L., Brock, C. A., Angevine, W. M., Beyersdorf, A., Campuzano-Jost, P., Day, D., de Gouw, J. A., Diskin, G. S., Gordon, T. D., Graus, M. G., Holloway, J. S., Huey, G., Jimenez, J. L., Lack, D. A., Liao, J., Liu, X., Markovic, M. Z., Middlebrook, A. M., Mikoviny, T., Peischl, J., Perring, A. E., Richardson, M. S., Ryerson, T. B., Schwarz, J. P., Warneke, C., Welti, A., Wisthaler, A., Ziemba, L. D., and Murphy, D. M.: In situ vertical profiles of aerosol extinction, mass, and composition over the southeast United States during SENEX and SEAC<sup>4</sup>RS: observations of a modest aerosol enhancement aloft, *Atmos. Chem. Phys.*, 15, 7085-7102, <https://doi.org/10.5194/acp-15-7085-2015>, 2015.
- Warneke, C., de Gouw, J. A., Edwards, P. M., Holloway, J. S., Gilman, J. B., Kuster, W. C., Graus, M., Atlas, E., Blake, D., Gentner, D. R., Goldstein, A. H., Harley, R. A., Alvarez, S., Rappenglueck, B., Trainer, M., and Parrish, D. D.: Photochemical aging of volatile organic compounds in the Los Angeles basin: Weekday-weekend effect, *J. Geophys. Res.-Atmos.*, 118, 5018–5028, <https://doi.org/10.1002/jgrd.50423>, 2013.
- Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO<sub>x</sub>, NO<sub>y</sub>, and other species in the lower stratosphere and upper troposphere during AASE II, *Geophys. Res. Lett.*, 21, 2583–2586, 1994.
- Wang, H.: Formation of nascent soot and other condensed-phase materials in flames: Proceedings of the Combustion Institute, 33, 41-67, doi:10.1016/j.proci.2010.09.009, 2011.
- Wilson, E. B., First and second laws of error. *Journal of the American Statistical Association* 18 841-851, 1923.
- Wisthaler, Armin; Crawford, J.H; Haidahcer, S.; Hanel, G.; Hartungen, E.; Jordan, A.; Märk, L.; Mikoviny, T.; Müller, M.; Mutschlechner, P.; Schottkowsky, R. & Sulzer: Development of a compact PTR-ToF-MS for suborbital research on the earth's atmospheric composition, In A Hansel & J. Dunkel (ed.), 6th International Conference on Proton Transfer Reaction Mass Spectrometry and its Applications. Contributions. Innsbruck University Press. ISBN 978-3902811912. pp 96 – 96, 2013.
- Yates, E.L., Schiro, K., Lowenstein, M., Sheffner, E.J., Iraci, L.T., Tadić, J.M.; Kuze, A: Carbon Dioxide and Methane at a Desert Site—A Case Study at Railroad Valley Playa, Nevada, USA. *Atmosphere* 2011, 2, 702-714.
- Yokelson, R.J., D.W.T. Griffith, and D.E. Ward: Open-path Fourier transform infrared studies of large-scale laboratory biomass fires, *J. Geophys. Res.*, 101, 21067021080,11996, 1996.
- Yokelson, R. J., Karl, T., Artaxo, P., Blake, D. R., Christian, T. J., Griffith, D. W. T., Guenther, A., and Hao, W. M.: The Tropical Forest and Fire Emissions Experiment: overview and airborne fire emission factor measurements, *Atmos. Chem. Phys.*, 7, 5175-5196, <https://doi.org/10.5194/acp-7-5175-2007>, 2007.
- Yokelson, R.J., M. O. Andreae, and S. K. Akagi, Pitfalls with the use of enhancement ratios or normalized excess mixing ratios measured in plumes to characterize pollutant sources and aging *Atmos. Meas. Tech.*, 6, 2155–2158, 2013.

1940

Vay, S. A., Choi, Y., Vadrevu, K. P., Blake, D. R., Tyler, S. C., Wisthaler, A., Hecobian, A., Kondo, Y., Diskin, G. S., Sachse, G. W., Woo, J.-H., Weinheimer, A. J., Burkhardt, J. F., Stohl, A., Wennberg, P. O.: Patterns of CO<sub>2</sub> and radiocarbon across high northern latitudes during IPY 2008, J. Geophys. Res., 116, doi:10.1029/2011JD015643, 2011.

$\alpha$

▲ **Page 5: [1] Deleted** RChatfield 4/19/20 4:31:00 PM

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲ **Page 5: [2] Formatted** RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

▲



Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [2] Formatted RChatfield 4/19/20 4:29:00 PM

Underline color: Auto

Page 5: [3] Deleted RChatfield 3/25/20 5:22:00 PM

Page 5: [4] Formatted RChatfield 4/19/20 4:29:00 PM

Font: Not Italic, Underline color: Auto

Page 5: [4] Formatted RChatfield 4/19/20 4:29:00 PM

Font: Not Italic, Underline color: Auto

Page 5: [4] Formatted RChatfield 4/19/20 4:29:00 PM

Font: Not Italic, Underline color: Auto

Page 5: [5] Deleted RChatfield 4/19/20 4:35:00 PM

▼

1.1.

Page 5: [5] Deleted RChatfield 4/19/20 4:35:00 PM

Page 5: [6] Deleted RChatfield 4/19/20 7:10:00 PM

Page 5: [6] Deleted RChatfield 4/19/20 7:10:00 PM

Page 5: [7] Formatted RChatfield 4/19/20 4:39:00 PM

Font: Not Italic

Page 5: [7] Formatted RChatfield 4/19/20 4:39:00 PM

Font: Not Italic

Page 5: [8] Deleted RChatfield 4/6/20 2:18:00 PM

Page 5: [8] Deleted RChatfield 4/6/20 2:18:00 PM

Page 8: [9] Deleted RChatfield 4/13/20 10:07:00 PM

Page 8: [10] Deleted RChatfield 4/13/20 11:59:00 PM

Page 8: [11] Deleted RChatfield 4/2/20 3:31:00 PM

Page 8: [12] Deleted RChatfield 4/2/20 3:31:00 PM

Page 10: [13] Deleted RChatfield 4/2/20 3:37:00 PM

Page 10: [13] Deleted RChatfield 4/2/20 3:37:00 PM

Page 10: [13] Deleted RChatfield 4/2/20 3:37:00 PM

Page 10: [14] Deleted RChatfield 4/12/20 10:38:00 PM

Page 10: [14] Deleted RChatfield 4/12/20 10:38:00 PM

Page 10: [14] Deleted RChatfield 4/12/20 10:38:00 PM

Page 10: [14] Deleted RChatfield 4/12/20 10:38:00 PM

▼ ▲  
**Page 10: [14] Deleted** RChatfield 4/12/20 10:38:00 PM

▼ ▲  
**Page 10: [14] Deleted** RChatfield 4/12/20 10:38:00 PM

▼ ▲  
**Page 10: [15] Change** Unknown  
Field Code Changed

▲  
**Page 10: [15] Change** Unknown  
Field Code Changed

▲  
**Page 10: [16] Deleted** RChatfield 4/12/20 10:45:00 PM

▼ ▲  
**Page 10: [16] Deleted** RChatfield 4/12/20 10:45:00 PM

▼ ▲  
**Page 10: [17] Change** Unknown  
Field Code Changed

▲  
**Page 10: [17] Change** Unknown  
Field Code Changed

▲  
**Page 10: [18] Deleted** RChatfield 4/12/20 11:26:00 PM

▼ ▲  
**Page 10: [18] Deleted** RChatfield 4/12/20 11:26:00 PM

▼ ▲  
**Page 10: [18] Deleted** RChatfield 4/12/20 11:26:00 PM

▼ ▲  
**Page 10: [18] Deleted** RChatfield 4/12/20 11:26:00 PM

▼ ▲  
**Page 10: [18] Deleted** RChatfield 4/12/20 11:26:00 PM

▼ ▲  
**Page 10: [19] Deleted** RChatfield 4/14/20 12:18:00 AM

▼ ▲  
**Page 10: [20] Formatted** RChatfield 4/19/20 10:33:00 PM  
Font: Not Bold

▲  
**Page 10: [20] Formatted** RChatfield 4/19/20 10:33:00 PM  
Font: Not Bold

▲  
**Page 11: [21] Deleted** RChatfield 4/12/20 12:44:00 PM

▲ Page 11: [22] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [22] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [23] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [23] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [24] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [24] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [25] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [25] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [26] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [26] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [27] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [27] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [28] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [28] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica, Not Italic

▲ Page 11: [29] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [29] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [29] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲ Page 11: [29] Formatted RChatfield 4/12/20 12:52:00 PM

Font: Helvetica

▲

Page 11: [30] Formatted	RChatfield	4/12/20 12:52:00 PM
-------------------------	------------	---------------------

Font: Helvetica

Page 11: [30] Formatted	RChatfield	4/12/20 12:52:00 PM
-------------------------	------------	---------------------

Font: Helvetica

Page 14: [31] Deleted	RChatfield	3/30/20 3:45:00 PM
-----------------------	------------	--------------------

Page 14: [32] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [33] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [34] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [35] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [36] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [37] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [38] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [39] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [40] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [41] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [42] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [43] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [44] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [45] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [46] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

Page 14: [47] Formatted	RChatfield	3/28/20 3:23:00 PM
-------------------------	------------	--------------------

Font: 11 pt

▲ Page 14: [48] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [49] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [50] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [51] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [52] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [53] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [54] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [55] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [56] Formatted RChatfield 3/28/20 3:23:00 PM

Font: 11 pt

▲ Page 14: [57] Deleted RChatfield 4/19/20 10:34:00 PM

✖ Page 27: [58] Deleted RChatfield 4/19/20 8:59:00 PM

▲ Page 28: [59] Deleted RChatfield 4/19/20 10:37:00 PM

▼  
▲ Page 28: [60] Deleted RChatfield 4/13/20 12:48:00 AM

✖  
▲ Page 28: [60] Deleted RChatfield 4/13/20 12:48:00 AM

✖  
▲ Page 29: [61] Deleted RChatfield 4/14/20 12:21:00 AM