

Interactive comment on “Validation analysis of deriving acetonitrile (CH₃CN) profiles by observations of SMILES from the International Space Station, in the stratosphere and lower mesosphere” by T. Fujinawa et al.

T. Fujinawa et al.

fujinawa.tamaki@nies.go.jp

Received and published: 12 December 2019

Dear Dr. Hugh C. Pumphrey,

We deeply appreciate for valuable comments and suggestions. Please find the manuscript with several revisions. Answers to your comments/questions are given by point by point. We hope that current manuscript is significant for the publication in Atmospheric Measurement Techniques.

Sincerely yours, Tamaki Fujinawa.

C1

Specific comment 1-1

Page 2 line 25: “Acetonitrile [. . .] is one of the dominant gases emitted during wildfire events (90%-95%)”. This statement is either wrong or confusing as wildfires emit far more CO and CO₂ than they do CH₃CN. What I suspect that the authors mean is that 90%-95% of CH₃CN comes from wildfires. If that is what the authors mean, they should say so explicitly.

Answer to specific comment 1-1

We appreciate your comment. As you mentioned, the sentence is confusing. To avoid misunderstanding, we replace it as below.

Revisions to specific comment 1-1

Line 25: “[. . .] as it is one of the dominant gases emitted during wildfire events (90–95%)” was replaced by

→

Line 25: “as 90-95 % of CH₃CN comes from wildfires”.

Specific comment 1-2

Page 2 line 27: “...ocean uptake and the reaction with hydroxyl radicals (OH)([. . . refs . . .]).” This sentence appears to be missing some words at the end. Maybe it should read “...ocean uptake and the reaction with hydroxyl radicals (OH)([. . . refs . . .]) being the main loss mechanisms.”

Answer to specific comment 1-2

We appreciate your valuable comment. As you mentioned, some words which you recommended is added as below.

Revisions to specific comment 1-2

Line 27 : “[. . .] being the main loss mechanisms.” was added.

C2

Specific comment 1-3

Page 5 line 58 “maximum a posteriori solution” should be “maximum a posteriori probability solution”. When mentioning the MAP solution it is surely obligatory to cite Rodgers (2000).

Answer to specific comment 1-3

We appreciate you pointing this out. As you mentioned, we add the word ‘probability’ and cite Rodgers (2000).

Revisions to specific comment 1-3

Line 58 : “maximum a posteriori solution” was replaced by “maximum a posteriori probability solution (Rodgers, 2000)”

Specific comment 1-4

Page 16, Figure 10. I struggle to interpret this figure. This is partly because the individual panels are rather small; I do not know what might be done about this as the arrangement of the figure is useful. The MLS data appear to have a lot of gaps in; the number of gaps increases with altitude. Now, it is clear that some of the gaps are there because the authors have chosen to show MLS data only for times when there is also SMILES data available. I would suggest that for the middle column of data they might want to show MLS data for all days, so that the eye can more easily pick out the patterns in the data. The actual MLS data do not become more sparse with altitude as the figure suggests: I just plotted them up to check. However, they do become negative on average, as can be seen in Figure 9 (left). I would suggest that the authors use a colour scale which spans the range of both the MLS and SMILES data. Clearly, the MLS data are wrong in the sense that the atmosphere can not contain less than none of a constituent. But the time-latitude dependence of the MLS data is actually rather similar to that of the SMILES data – it would be nice if the plot could bring that out. I am pleased to see a sequential colour scale used for the actual quantities and a diverging

C3

one for the SMILES-MLS differences. Less satisfactory is the choice of sequential colour scale; the authors have used the notorious “jet” scale, or something very like it. (See <https://hughpumphrey.wordpress.com/2017/06/29/colours-for-contours/> for some thoughts and some useful links.) They might want to consider whether a scale other than “jet” might be appropriate in this figure.

Answer to specific comment 1-4

We deeply appreciate your valuable comment and link. As you mentioned, we revised Fig. 10 about the following points: (1) in the middle column in Fig. 10, the MLS data for all day, which is including the not-coincident points, were considered, (2) the color scale was expanded to negative values, (3) the sequential color map of ‘jet’ was changed to ‘magma’. We hope these revisions help you to interpret this figure.

Revisions to specific comment 1-4

Please see the revised Figure 10 (Fig. 1 HERE). Revised points are as following. (1) In the middle column in Fig. 10, the MLS data for all day, which is including the not-coincident points, were considered. (2) the color scale was expanded to negative values. (3) the sequential color map of ‘jet’ was changed to ‘magma’.

Specific comment 1-5

Page 16 Lines 194-196: The time-latitude dependence of the MLS data is actually quite clear and easy to see as long as you plot the data up with a colour scale that goes into the negative. Clearly, the MLS data have a negative bias of over 100% in the upper stratosphere, but this does not, of itself, prevent the seasonal behaviour being observed.

Answer to specific comment 1-5

We are grateful for pointing this out. We expanded the color scale of the MLS data to negative values at upper pressure levels in Fig. 10. As you mentioned, results of the MLS also showed the time-latitude dependence clearly in the upper stratosphere

C4

despite of the large negative bias. We think this comment is partly included in the specific comment 2-4. Therefore, please see also the answer and revisions against specific comment 2-4 in detail.

Revisions to specific comment 1-5

This comment seems to be almost same as Specific comment 1-4. Please see the revised Figure 10.

Specific comment 1-6

Page 16 Lines 198-199 and page 17 lines 210-211: It is not at all clear to me why there should be a connection between the maximum observed at 1hPa – 5 hPa in February and the timing of the biomass-burning season. Tropospheric source gases such as CH₃CN enter the stratosphere via the tropical tropopause and take over a year to ascend from 100hPa (16km) to 10hPa (32km). This “tape recorder” effect was first observed in water vapour (Mote et al., 1996) and subsequently in HCN (Pumphrey et al., 2008, 2018) among other species. Figure 1 shows that CH₃CN is similar to HCN, although the tape recorder signal is only clear in the 2016-18 period. There was a large influx of both HCN and CH₃CN at that time due to a very strong El Niño event.

Answer to specific comment 1-6

We deeply appreciate pointing this out and your valuable comment. The other anonymous referee also pointed out the same point. We thought there is seasonality of CH₃CN in lower stratosphere from 28 km to 36 km as can be seen in Figure 7, even if the error bars (one sigma standard deviation) were taken into account. Therefore, we thought that the enhancement which can be seen in Figure 7 would be caused by biomass burning event. However, as you mentioned, our description about the enhancement might cause a misunderstanding that biomass burning plume emitted during December to March directly caused the seasonal maximum measured in February.

Revisions to specific comment 1-6

C5

Lines 5-6: “We estimated the systematic and random errors to be ~5.8 ppt (7.8 %) and 25 ppt (60 %) for a single observation at 15.7hPa, respectively, in the Tropics, where the CH₃CN measurements are enhanced.” was replaced by

→

Lines 5-6: “We estimated the systematic and random errors to be ~5.8 ppt (7.8 %) and 25 ppt (60 %) for a single observation at 15.7hPa, respectively, in the Tropics.”

Lines 198-199: “, which is consistent with our understanding that most biomass burning occurs from December–March.” was removed.

Lines 198-199: “These seasonal maximum CH₃CN levels measured in February are consistent with the yearly high period of BB events from December–March.” was removed.

Technical correction 2-1~2-5

Page 1 Line 1: Maybe replace “one of the volatile organic compounds” with “a volatile organic compound”.

Page 1 line 7: Maybe replace “a pressure broadening” with “pressure broadening” or “the pressure broadening coefficient”.

Page 2 line 33: “lower stratosphere Kopp and Arnold et al. (1978); Schneider et al. (1997).” should be “lower stratosphere (Kopp and Arnold et al., 1978; Schneider et al., 1997)” (In LATEX this would be a \citep, not a \citet.)

Page 2 Lines 51-53: Figures should be called out in numerical order. Here, Figure 2 is mentioned in the running text before Figure 1.

Page 17 line 216: AMT prefers datasets to be referenced in the same way as papers, with the DOI included. The full reference for the MLS CH₃CN data is Santee and Read (2015).

C6

Answer to technical correction 2-1~2-5

We appreciate your valuable suggestions. As you mentioned, we revised several points as following. Detailed revisions are described in “Revision to technical correction 3-1~3-5”.

Revisions to technical correction 2-1~2-5

Page 1 Line 1: “[. . .] one of the volatile compounds [. . .]” was replaced by

→

“[. . .] a volatile organic compound [. . .]”

Page 1 line 7: “[. . .] a pressure broadening [. . .]” was replaced by

→

“[. . .] the pressure broadening coefficient [. . .]”

Page 2 line 33: “lower stratosphere Kopp and Arnold et al. (1978); Schneider et al. (1997).” was replaced by

→

“lower stratosphere (Kopp and Arnold et al., 1978; Schneider et al.,1997)”

Page 2 Lines 51-53: “[. . .] as shown in Fig. 2. SMILES employed [. . .] The date of observation made by AOS1 and AOS2 are shown in Fig. 1.” was replaced by

→

“[. . .] as shown in Fig. 1. SMILES employed [. . .] The date of observation made by AOS1 and AOS2 are shown in Fig. 2.”

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2019-261, 2019.

C7




Fig. 1. Revised figure from Fig. 10 in the manuscript.

C8