



# First eddy covariance flux measurements of gaseous elemental mercury (Hg<sup>0</sup>) over a grassland

Stefan Osterwalder<sup>1,2\*</sup>, Werner Eugster<sup>3</sup>, Iris Feigenwinter<sup>3</sup>, Martin Jiskra<sup>1</sup>

<sup>1</sup>Environmental Geosciences, University of Basel, 4056 Basel, Switzerland
 <sup>2</sup>CNRS, IRD, IGE, University Grenoble Alpes, 38058 Grenoble, France
 <sup>3</sup>Institute of Agricultural Sciences, ETH Zurich, 8092 Zurich, Switzerland

Correspondence to: Stefan Osterwalder (stefan.osterwalder@unibas.ch)

Abstract. Direct measurements of the net ecosystem exchange (NEE) of gaseous elemental mercury (Hg<sup>0</sup>) are crucial to improve the understanding of global Hg cycling und ultimately human and wildlife Hg exposure. The lack of long-term, ecosystem-scale measurements causes large uncertainties in Hg<sup>0</sup> flux estimates. Today it remains unclear whether terrestrial ecosystems are net sinks or sources of atmospheric Hg<sup>0</sup>. Here we show a detailed validation of the eddy covariance technique for direct Hg<sup>0</sup> flux measurements (*Eddy Mercury*) based on a Lumex mercury monitor RA-915AM. The flux detection limit derived from a zero-flux experiment in the laboratory was 0.22 ng m<sup>-2</sup> h<sup>-1</sup> (maximum) with a 50 % cut-off at 0.074 ng m<sup>-2</sup> h<sup>-1</sup>. The statistical estimate of the Hg<sup>0</sup> flux detection limit under real-world outdoor conditions at the site was 5.9 ng m<sup>-2</sup> h<sup>-1</sup> (50 % cut-off). We present the first successful eddy covariance NEE measurements of Hg<sup>0</sup> over a low-Hg level soil (41–75 ng Hg g<sup>-1</sup> topsoil [0-10 cm]) in summer 2018 at a managed grassland at the Swiss FluxNet site in Chamau, Switzerland

(CH-Cha). We measured a net summertime re-emission over a period of 34 days with a median Hg<sup>0</sup> flux of 2.5 ng m<sup>-2</sup> h<sup>-1</sup> (-0.6 to 7.4 ng m<sup>-2</sup> h<sup>-1</sup>, range between 25<sup>th</sup> and 75<sup>th</sup> percentiles). We observed a distinct diel cycle with higher median daytime fluxes (8.4 ng m<sup>-2</sup> h<sup>-1</sup>) than nighttime fluxes (1.0 ng m<sup>-2</sup> h<sup>-1</sup>). Drought stress during the

measurement campaign in summer 2018 induced partial stomata closure of vegetation which led to a midday depression in CO<sub>2</sub> uptake which did not recover during the afternoon. Thus, the cumulative net CO<sub>2</sub> uptake was only 8 % of the net CO<sub>2</sub> uptake during the same period in the previous year 2017. We suggest that partial stomata closure dampened Hg<sup>0</sup> uptake by vegetation, resulting in a NEE of Hg<sup>0</sup> dominated by soil re-emission.

Finally, we give suggestions to further improve the precision and handling of the *Eddy Mercury* system in order to assure its suitability for long-term NEE measurements of  $Hg^0$  over natural background surfaces with low soil Hg concentrations (< 100 ng g<sup>-1</sup>). The system, improved in the suggested way, has the potential to be integrated in global networks of micrometeorological tower sites (FluxNet) and provide the long-term observations on terrestrial atmosphere  $Hg^0$  exchange necessary to validate regional and global mercury models.



40



#### 30 1 Introduction

Mercury (Hg) is a top priority environmental pollutant that is transported through the atmosphere as gaseous elemental  $Hg^0$  (> 95 % of total atmospheric Hg). Anthropogenic Hg emissions into the atmosphere exceed natural emissions by a factor of about five (Outridge et al., 2018). Atmospheric Hg has a lifetime of 8–13 months allowing for long-range transport before being deposited back onto the Earth surface also at remote locations far

35 from pollution sources (Saiz-Lopez et al., 2018). Once deposited, Hg can be transformed into methylmercury which can bioaccumulate in the freshwater and marine food webs, thereby posing a threat for human and ecosystem health (Watras et al., 1998; Fitzgerald et al., 2007; Mason et al., 2012; Braune et al., 2015).

Atmospheric Hg deposition to terrestrial surfaces occurs predominantly as Hg<sup>0</sup> dry deposition through stomatal uptake by vegetation or as wet deposition after oxidation in the atmosphere to more soluble reactive mercury (Hg(II)) (Lindberg et al., 2007; Driscoll et al., 2013; Jiskra et al., 2018). Wet deposition of Hg(II) via rain and snowfall is relatively well quantified by Hg deposition networks such as the National Atmospheric Deposition

Program (NADP), the European Monitoring and Evaluation Programme (EMEP) and the Asia Pacific Mercury Monitoring Network (APMMN).

Dry deposition of  $\mathrm{Hg}^0$  has been identified as the dominating Hg deposition pathway over vegetated surfaces

- 45 based on Hg stable isotope fingerprinting. Dry deposition of Hg<sup>0</sup> though stomatal uptake contributes 65–90 % of the total Hg deposition to soils (Demers et al. 2007; Jiskra et al., 2015; Enrico et al., 2016; Zhang et al., 2016; Zheng et al., 2016; Obrist et al., 2017). However, Hg<sup>0</sup> dry deposition remains poorly constrained due to the lack of long-term monitoring networks (Obrist et al., 2018). Reduction of Hg(II) in terrestrial surface pools and subsequent re-emission of Hg<sup>0</sup> back to the atmosphere prolongs the cycling of anthropogenic Hg emissions in
- 50 the environment and can thereby delay the effects of curbing primary anthropogenic emissions on human Hg exposure (Zhu et al., 2016; Wang et al., 2016; Obrist et al., 2018). The net ecosystem exchange (NEE) of Hg<sup>0</sup>, the sum of Hg<sup>0</sup> dry deposition and the re-emission from foliage and soils, represents a major factor in how fast the environment will recover from anthropogenic Hg pollution. On a global scale, estimates of the terrestrial NEE of Hg<sup>0</sup> remain uncertain. In the most recent global mercury assessment soil re-emission estimates were
- 55 lowered to 1000 Mg a<sup>-1</sup> (UNEP, 2019) relative to 2200 Mg a<sup>-1</sup> in the 2013 assessment (UNEP, 2013), however the associated uncertainties remain large. A recent review of 132 direct flux measurement studies revealed a NEE Hg<sup>0</sup> flux between -513 and 1653 Mg a<sup>-1</sup> (range of 37.5<sup>th</sup> and 62.5<sup>th</sup> percentiles, the central 25 % of the distribution) (Agnan et al. 2016). The database predominantly contains Hg<sup>0</sup> flux measurements performed with dynamic flux chambers (85 % of all studies) that are ideal for short-term, mechanistic studies but less suitable for
- 60 quantitative flux estimations, especially over vegetated surfaces (Gustin et al., 1999; Eckley et al., 2016; Osterwalder et al., 2018). Year-round NEE measurements of Hg<sup>0</sup> at the landscape scale are compelling to reduce measurement uncertainties. However, there are only four year-round whole-ecosystem Hg<sup>0</sup> flux studies published, all using micrometeorological (MM) techniques, including the modified Bowen-ratio and aerodynamic gradient methods (Fritsche et al., 2008a; Castro and Moore, 2016; Obrist et al., 2017), and the
- 65 relaxed eddy accumulation (REA) technique (Osterwalder et al. 2017). These approaches use instruments that do not fulfill the criterion of fast response of the Hg sensor as it is required for eddy covariance (EC) flux measurements. Hence, these are not direct flux measurements and are thus dependent on a number of assumptions. The main difficulty using the modified Bowen-ratio and aerodynamic gradient method is to resolve a significant concentration gradient during turbulent conditions. During calm conditions, in contrast, it is challenging to determine a significant eddy diffusivity. Further drawbacks are (1) the potentially different





sink/source characteristics of the footprint due to the two measurement heights, (2) temporally intermittent sampling between the two sampling inlets, and (3) the fact that transport characteristics are based on reference scalars like heat, water or  $CO_2$  (Businger et al., 1986; Stannard et al., 1997; Edwards et al., 2005, Sommar et al., 2013a). The REA technique (Businger and Oncley, 1990) circumvents most of these difficulties. However,

- 75 uncertainties in Hg<sup>0</sup> flux calculations are introduced by the determination of the proportionality coefficient (β-value) and system dependent shortcomings such as a biased offset between the updraft and downdraft sampling lines or difficulties in controlling the air flow from the air inlets to the analyzer. Thus, it remains challenging to accurately measure very small concentration differences with REA (typically < 0.1 ng m<sup>-3</sup>) between updrafts and downdrafts over natural surfaces with low substrate Hg concentrations (Cobos et al., 2002; Bash and Miller, 2008; Sommar et al., 2013b; Osterwalder et al., 2016, Kamp et al., 2018).
  - The EC technique has been under development since the late 1940s to measure the surface–atmosphere exchange of heat, mass, and momentum in the surface boundary layer in the lowest 20–50 m of the atmosphere (Montgomery, 1948; Obukhov 1951; Swinbank 1951). In order to estimate a vertical turbulent flux, the covariance of two concurrently measured variables is calculated, (1) the scalar quantity of interest (in our case
- Hg<sup>0</sup>) and (2) the turbulent fluctuations of the vertical wind velocity, both measured at high temporal resolution. Since the 1990s a new generation of digital three-axis ultrasonic anemometers, infrared gas analyzers and comprehensive software packages have facilitated land–atmosphere exchange measurements of CO<sub>2</sub> and H<sub>2</sub>O (McMillen 1988). Today, the EC technique is considered the standard method to determine evapotranspiration and the NEE of energy and trace gases such as CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, O<sub>2</sub>, O<sub>3</sub> and volatile organic compounds using
- 90 high resolution (10–20 Hz), sometimes portable, and generally very reliable equipment (Aubinet et al., 2012). The first successful application of the EC technique to measure NEE of Hg<sup>0</sup> reported a re-emission flux of 849 ng m<sup>-2</sup> h<sup>-1</sup> over contaminated soils (85 mg Hg kg<sup>-1</sup> dry soil) during a pilot campaign in Nevada, USA (Pierce et al., 2015). The EC system was based on a fast response (25 Hz), field deployable pulsed cavity ring-down spectrometer (CRDS) (Faïn et al., 2010; Pierce et al., 2013). The minimum detection limit of 32 ng m<sup>-2</sup> h<sup>-1</sup>
  95 however did not allow Hg<sup>0</sup> flux measurements over soils exhibiting background Hg concentrations (typically

100 ng Hg g<sup>-1</sup>; Grigal et al., 2003) (Pierce et al., 2015).

110

Here we present the first EC measurements to determine the NEE of  $Hg^0$  over a grassland with background soil Hg levels. Our novel EC system is based on a Lumex mercury monitor RA-915AM (Lumex Ltd., St. Petersburg, Russia) atomic absorption spectrometer with Zeeman background correction allowing to measure  $Hg^0$  in ambient

- 100 air at a sampling frequency of 1 Hz (Sholupov et al., 1995, 2004). Ambient air Hg<sup>0</sup> measurement comparison studies between the more frequently used Tekran® 2537 analyzer (Tekran Inc., Toronto, Canada) and the mercury monitor's precursor, the Lumex RA-915+ mercury analyzer, were performed by the European Committee for Standardization's (CEN) Technical Committee 264 "Air Quality" EN 15852 and showed good agreement between the two instruments (Brown et al., 2010). Among other applications, the Lumex RA-915AM
- 105 mercury analyzer was successfully deployed in the Global Mercury Observation (GMOS) project at two sites in Russia and Suriname (Sprovieri et al., 2016).

The objective of the study was to test the performance of the RA-915AM as fast response analyzer and its suitability for EC flux measurements with the goal to reliably measure the NEE of gaseous elemental mercury or  $Hg^0$  (GEM-NEE) over terrestrial ecosystems. Hereinafter the new system is referred to as *Eddy Mercury*. We provide a description of the *Eddy Mercury* sampling system and present the data analysis procedure to calculate

the EC Hg<sup>0</sup> flux in detail. We discuss patterns in the GEM-NEE that was measured over a grassland during a 34-





day pilot campaign and give suggestions to improve the reliability and precision of the *Eddy Mercury* system for future long-term applications.

#### 2 Material and methods

#### 115 **2.1 Site description and instrumentation**

The *Eddy Mercury* system was tested between 20 July and 6 September 2018 at the Swiss FluxNet site Chamau (CH-Cha), located in central Switzerland, about 30 km southwest of Zurich (47° 12′ 36.8″ N, 8° 24′ 37.6″ E; 393 m a. s. l.). In this study NEE of Hg<sup>0</sup> and CO<sub>2</sub> was measured concurrently with two independent EC systems over the intensively managed grassland which is used for forage production. Details on grassland species composition, harvest and fertilization practices are described in Zeeman et al. (2010), Merbold et al. (2014) and Fuchs et al. (2018). The tower for long-term EC measurements was located between two adjacent grassland parcels (Fig. 1). The northern parcel (measured when up-valley winds prevail) was over-sown with clover in March 2015 and April 2016 to investigate the N<sub>2</sub>O emission reduction potential in comparison to the

conventionally fertilized grassland of the southern parcel (measured primarily when down-valley winds prevail).

- 125 The soil type is a gleysol–cambisol, with a bulk density of about 1 g cm<sup>-3</sup>, 30.6 % sand, 47.7 % silt and 21.7 % clay in the top 10 cm (Roth, 2006). The topsoil pH was 5.3 and was determined by adding 25 ml of 0.01 M CaCl<sub>2</sub>-solution to 10 g dry soil (Labor Ins AG, Kerzers, Switzerland, in 2014). The 24 year (1994-2017) average annual temperature measured at the nearby SwissMetNet surface weather station in Cham (CHZ, 444.5 m a. s. l.) was 10.1 °C and the average annual precipitation was 997 mm.
- 130 The *Eddy Mercury* system was mounted about 3 m west of a fully equipped long-term EC tower measuring greenhouse gas exchange (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>, H<sub>2</sub>O) and meteorological variables at 2 m height. The CO<sub>2</sub> flux system consisted of a 3D ultrasonic anemometer (Solent R3-50, Gill Instruments, Lymington, UK) and an open-path infrared gas analyzer for CO<sub>2</sub> and H<sub>2</sub>O concentrations running at 20 Hz resolution (IRGA, LI-7500, LI-COR Biosciences, Lincoln, NE, USA). From the 20 Hz IRGA measurements, 30 min flux averages were
- calculated using the LI-COR EddyPro<sup>®</sup> software. The 30 min CO<sub>2</sub> flux are recorded continuously since 2005 (Eugster and Zeeman, 2006; Zeeman et al., 2010). The measured meteorological variables included temperature and relative humidity (Hydroclip S3 sensor, Rotronic AG, Switzerland), net all-wave radiation (CNR1, Kipp &Zonen B.V., Delft, Netherlands), incoming and reflected photosynthetic active radiation (PARlite, Kipp and Zonen, Delft, Netherlands), and precipitation (0.5 m height; tipping bucket rain gauge from LAMBRECHT
- 140 meteo GmbH, Göttingen, Germany). In addition, soil temperatures were recorded at 0.05, 0.1, 0.15, 0.25, 0.4 m depth (T107, Campbell Scientific Inc., Logan, UT, USA).

#### 2.2 Soil sampling and total mercury analysis

145

Topsoil samples (0–10 cm) were taken in a circular arrangement around the EC tower (Fig. 1) using a core drill. The soil samples were transported to the laboratory in sealed plastic bags, and stored in a fridge at 4 °C. The samples were filled into aluminum shells, weighed and dried at 40 °C, until their weight remained constant. The samples were pestled and sieved through a 2 mm mesh to separate the fine earth and the skeleton. The fine earth was ground to powder using a laboratory scale ball mill. To get rid of all potential humidity, the ground samples were stored in small paper bags in a desiccator and dried again at 40 °C. The 22 topsoil samples were analyzed for total Hg using a DMA-80 Direct Mercury Analyzer (MLS Mikrowellen GmbH, Leutkirch im Allgäu,





150 Germany). Certified Hg standard solution (NIST 3133) was gravimetrically diluted to concentrations of 10 ng g<sup>-1</sup> to 1000 ng g<sup>-1</sup> and used for the calibration of the instrument. Repeated measurements of standard reference material (ERM-CC141 loam soil) 90.3  $\pm$  7.8 ng g<sup>-1</sup> (mean  $\pm$  SD, n=3) were in agreement with the certified value (83  $\pm$  17 ng g<sup>-1</sup>).

#### 2.3 Description of the Eddy Mercury system

- 155 The core of the *Eddy Mercury* apparatus to measure GEM-NEE is the RA-915AM mercury monitor (Lumex Analytics GmbH, Germany). The RA-915AM uses atomic absorption spectrometry (AAS) with Zeeman background correction to continuously measure Hg<sup>0</sup> in ambient air (Sholupov et al., 2004). The multi-path sample cell of the RA-915AM has an optical path length of 9.6 m and a cell volume of 0.7 L. Baseline corrections (zero drift) are performed automatically by the instrument using Hg-free air at user defined intervals. Span corrections are done using an inbuilt calibration cell that contains Hg<sup>0</sup> vapor. The measurement range can
- be set between 0 and 2000 ng m<sup>-3</sup> and its detection limit is 0.5 ng m<sup>-3</sup> according to the analytical specifications by the manufacturer. The air flow rate was increased to 14.3 L min<sup>-1</sup> by bypassing the instrument pump in order to reduce the residence time in the measurement cell (normal flow: 7 L min<sup>-1</sup>). For this, a stronger external pump was connected (model MAA-V109-MD, GAST Manufacturing, MI, USA). The instrument was placed in a
- 165 weatherproof, air-conditioned box (Elcase, Marthalen, Switzerland) to protect the sensitive RA-915AM from rain and reduce temperature fluctuations. A USB-to-RS232 serial data interface was used to establish a one-way communication link from the RA-915AM to the data acquisition computer. The air inlet was mounted 24 cm below the center of the head of the three dimensional (3D) ultrasonic anemometer (Gill R2A, Solent, UK) used for wind vector measurements that was installed 2 m above ground. A micro-quartz fiber filter (Grade MK 360,
- 47 mm diameter, Ahlstrom–Munksjö, Sweden) was installed in a 47 mm Perfluoralkyl-polymere (PFA) single stage filter assembly (Savillex, Eden Prairie, USA) at the air inlet. The air inlet was connected to the RA-915AM by a 2.8 m intake hose with 11 mm inner diameter (ID) attached to a 0.35 m, 4 mm ID sample intake hose. Both hose segments were unheated, insulated PFA tubing. The median lag time of the turbulent airflow (Reynolds number of > 5000) from the tube inlet to the analyzer was in the order of 1.15 s.

# 175 2.4 Eddy covariance flux measurements

The RA-915AM analyzer was configured to measure  $Hg^0$  concentrations at 1 Hz. The  $Hg^0$  concentrations and the 3D wind vectors were measured from 20 July to 6 September 2018 using four different settings of the RA-915AM analyzer with respect to the length of the measurement interval between two auto-calibration cycles (zero and span): (1) 24 hour intervals from 20–26 July 2018; (2) 4 hour intervals from 1–26 August 2018; (3) 1

180 hour intervals from 27–31 August 2018; (4) 4 minute intervals from 31 August until 6 September 2018. The ultrasonic anemometer had an internal sampling frequency of 1000 Hz, which was averaged (8 records of each acoustic sensor pair for each direction) to 20.83 Hz. The 1 Hz RA-915AM data was merged with the ultrasonic anemometer's data stream by oversampling as described in Eugster and Plüss (2010). Data were collected on a Linux-based Raspberry Pi computer equipped with a real-time clock chip and internet access. Because data transfer via the USB port from the embedded Windows 7 system of the RA-915AM was highly unreliable, only the system time stamps were synchronized with the Linux data acquisition system every second via a Windows PowerShell script. In cases when also this communication failed, an approximate time synchronization was done by polling the RA-915AM timestamp via the Samba file sharing protocol. Thus, in extension of the



195

215



synchronization method described by Eugster and Plüss (2010) the merging of  $Hg^0$  measurements with wind vector data had to be done offline in a separate data workup step. Fluxes were calculated over 60 minute intervals to account for the low sampling frequency of  $Hg^0$  signals. Thus, under modes (1) and (2) 3600  $Hg^0$ measurements were used for each 1 hour flux average.

### 2.5 Eddy covariance Hg<sup>0</sup> flux calculations

Calculation of the GEM-NEE required some modifications of the standard procedure that is established for  $CO_2$  fluxes (e.g. Aubinet et al., 2012). The modifications were done according to the five steps described in detail below.

#### 2.5.1 Preparation of raw Hg<sup>0</sup> measurements

The RA-915AM raw data files provide the following information at 1 Hz resolution: Date and time of measurement, photomultiplier current (arb. unit), air flow rate (L min<sup>-1</sup>), temperature of analyzed air (°C), 200 temperature of RA-915AM (°C), sample cell pressure (kPa), Hg<sup>0</sup> raw concentration (ng m<sup>-3</sup>, including all online corrections), status code and status description. The status code (a numerical value) and status description (a text variable) are redundant and provide the necessary information to distinguish ambient air concentration measurements from zero and span calibration measurements. The Hg<sup>0</sup> flux was calculated based on the Hg<sup>0</sup> raw concentration. To account for drift and baseline drift, which both are unavoidable when longer measurement 205 periods are used between calibration events, we proceeded as follows. After a calibration event, the Hg<sup>0</sup> raw concentration was considered to be the best empirical estimate of the true Hg<sup>0</sup> concentration. Until the end of a measurement period (begin of next calibration cycle), in a first step a linear drift correction was applied to bring the Hg<sup>0</sup> raw concentration before the next calibration event to the level of the next calibration result (offset correction). Since visual inspection of the data clearly indicated that there is more drift than a simple linear trend 210 in the data (see examples in Fig. 2), a high-pass filter approach was used to minimize drift and optimize the determination of Hg<sup>0</sup> fluctuations for EC flux measurements (Sect. 2.5.4).

#### 2.5.2 Preparation of the ultrasonic anemometer data

The ultrasonic anemometer data contained the three wind speed components of the wind vector (all in m s<sup>-1</sup>), the speed of sound (m s<sup>-1</sup>), and the information sent from the RA-915AM to the data acquisition system via the serial data link. Speed of sound *c* was converted to virtual sonic temperature  $T_v \approx c^2/403$  in Kelvin (Kaimal and Gaynor 1991). The vertical wind speed *w* was despiked using an iterative  $7\sigma$  filter that discards *w* outside the range of the 6 hour mean  $\pm$  7 standard deviations.

# 2.5.3 Merging of ultrasonic anemometer data with Hg<sup>0</sup> time series

After preparation of the two datasets they were merged by accounting for the time difference between the RA-915AM and the Linux data acquisition using the information that could be transferred via the serial link from the RA-915AM to the Linux system (accurate to within 1 second). If no such information was received from the RA-915AM, the time difference between the two systems was determined using a network time drift fallback option specifically added to the Linux system to overcome the problems with serial output from the RA-915AM: during the field experiment we polled the most recent data record acquired by the RA-915AM every five minutes using the Samba filesharing protocol, and associated that timestamp with the one of the EC system. This





(somewhat less accurate) information was then adjusted during periods where both approaches overlapped to determine the time difference required to shift the  $Hg^0$  raw data relative to the ultrasonic anemometer data before merging the two datasets. To ascertain that  $Hg^0$  are lagging the sonic data we added a  $\approx 1.5$  s safety margin in the interpretation of the available time synchronization information received either via serial link or Samba filesharing.

230

235

260

# 2.5.4 Determination of time lag between vertical wind speed and $\mathrm{Hg}^{0}$ fluctuations

The merged dataset was then divided into 1 hour segments for  $Hg^0$  flux calculations. Within each 1 hour segment the time lag between the two time series was fine-tuned using a cross-correlation procedure to find the best positive or negative correlation within a reasonable time window (0–4 s) around the physically expected time difference (1.15 s physical delay plus 1.5 s safety margin used in step 3). Because considerable non-turbulent drift of the  $Hg^0$  signal was still present after correcting for online calibration (Sect. 2.5.1), we detrended each 1

- hour segment using a third-order polynomial fit (Eq. 5) before computing the cross-covariance between the detrended  $Hg^0$  signal and w (Sect. 3.2.1). To account for the different sampling rates of w (20.83 Hz) and  $Hg^0$  (1 Hz), we used simple linear interpolation between individual  $Hg^0$  measurements and to bridge across calibration
- 240 gaps. After a first automatic run each best estimate for time lag was visually inspected and updated by a narrower search window for each 1 hour segment that narrowed in the search procedure to the most realistic cross-correlation peak (positive or negative). Note that calibration gaps are relevant data gaps with setting 4 (Sect. 2.4) but less problematic with settings 1–3. In all cases, the lack of variance in Hg<sup>0</sup> data during the gaps reduces the computed Hg<sup>0</sup> flux. Thus, our flux estimates are conservative estimates with respect to flux magnitudes.

#### 2.5.5 Computation of Hg<sup>0</sup> EC fluxes

After all data preparations according to Sect. 2.5.1 and Sect. 2.5.4 the Hg<sup>0</sup> flux  $F_{Hg0}$  was calculated as the covariance

$$F_{Hg0} = \overline{w'\chi'},\tag{1}$$

250 with  $\chi$  being the calibrated, detrended and linearly gap-filled Hg<sup>0</sup> concentration in ng m<sup>-3</sup> and *w* the vertical wind speed. For improved readability  $F_{Hg0}$  was converted from ng m<sup>-2</sup> s<sup>-1</sup> to ng m<sup>-2</sup> h<sup>-1</sup> before reporting. In the notation used here, primes denote short-term deviations from the mean (after detrending according to Sect. 2.5.4) over an averaging period (1 hour) and overbars denote the mean of a variable. Hg<sup>0</sup> flux computations were done using R version 3.5.2 (R Core Team, 2018).

# 255 **2.6 Determine the Hg<sup>0</sup> flux detection limit**

To determine whether a calculated  $Hg^0$  flux is significantly different from a zero-flux we used two approaches: (1) an indoor zero-flux experiment, and (2) a statistical estimate of the flux detection limit following the concept by Eugster and Merbold (2015) which is a further improvement of the concept presented by Eugster et al. (2007). The indoor zero-flux experiment was set up in the laboratory on the two days before installing all equipment in the field. The low turbulence conditions in combination with absence of local  $Hg^0$  sources in the laboratory allowed us to see what fluxes are resulting with the procedure described above when there is no real  $Hg^0$  flux. Such zero-flux experiments however tend to underestimate the flux detection limit under real-world outdoor conditions, where the second approach quantifies the statistical uncertainty of a calculated flux. The flux



265

270



(covariance) is the product of the correlation coefficient  $r_{w,\chi}$  between w and  $\chi$  and the square-root of the variances of the two variables (e.g. Eugster and Merbold 2015),

$$\overline{w'\chi'} = r_{w,\chi} \cdot \sqrt{w'^2} \cdot \sqrt{\chi'^2} = r_{w,\chi} \cdot \sigma_w \cdot \sigma_\chi \tag{2}$$

The significance of  $r_{w,\chi}$  can be estimated using Student's *t* test (see Eugster and Merbold 2015) for details. For each 1 hour period we thus computed the value of  $r_{w,\chi}$  that is significant at p = 0.05, and multiplied this value with measured  $\sigma_w$  and  $\sigma_{\chi}$  to obtain a more realistic estimate for the flux detection limit. It should be noted that this concept has been brought forward long ago by Wienhold et al. (1996) using a visual empirical approach, whereas Eugster and Merbold (2015) further developed the visual approach to a more objective time series statistical approach to perform the quantification of the flux detection limit. The threshold of significance of  $r_{w,\chi}$  can be estimated as

$$r_{w,\chi_p} = \frac{t_p}{\sqrt{n-2+t_p^2}} , \qquad (3)$$

275

where  $t_p$  is Student's t value for the significance level p (e.g., 0.05), and n is the auto-correlation corrected number of independent samples in the time series,

 $n \simeq N \frac{1-\rho_1}{1+\rho_1},\tag{4}$ 

where *N* is the number of samples in a time series, and  $\rho_1$  is the lag 1 auto-correlation coefficient of the scalar product time series  $w \gamma$ .

# 280 2.7 Eddy covariance CO<sub>2</sub> flux calculations and quality control flags

The 30 min CO<sub>2</sub> flux was quantified in the conventional way established in ecosystem studies (see Aubinet et al., 2012) using the Eddy Pro (LI-COR Inc., Lincoln, NE, USA) software (see Fuchs et al., 2018 for specific information related to the Chamau field site). For each 30-minute CO<sub>2</sub> flux interval a flux quality control (QC) flag was determined: 0 (best data quality for detailed investigations), 1 (good data for longer-term studies), and 2 (poor quality), after Mauder and Foken (2004). Since there are no established quality control procedures for Hg<sup>0</sup> fluxes yet, we used the QC information from the CO<sub>2</sub> flux measurement to retain or reject concurrent Hg<sup>0</sup> flux measurements. Thus, we solely present Hg<sup>0</sup> flux measurements with CO<sub>2</sub> flux quality flags < 2. During CO<sub>2</sub> flux processing using the EddyPro software, coordinate rotation for tilt correction, angle of attack correction for wind components, Webb–Pearman–Leuning terms for compensation of density fluctuations (Webb et al., 1980) and analytical corrections for high-pass (Eugster and Senn, 1995; Moncrieff et al., 2004) and low-pass filtering effects (Horst, 1997) were applied. Furthermore, a self-heating correction for the open-path gas analyzer was conducted (Burba et al., 2008) and CO<sub>2</sub> fluxes > 50 µmol m<sup>-2</sup> s<sup>-1</sup> and < -50 µmol m<sup>-2</sup> s<sup>-1</sup> were discarded.

#### **3 Results and Discussion**

#### **3.1 Environmental conditions**

In 2018, the annual mean air temperatures in Switzerland reached 6.9 °C, the largest value ever recorded since the onset of meteorological measurements in 1864. This nationwide average temperature was 1.5 °C warmer





compared to the average of the normal period of 1981-2010. Total precipitation measured from April to November 2018, was only 69 % of the long-term average (1981-2010). Thus, the period from April to November 2018 was the third driest period ever recorded in Switzerland (MeteoSchweiz, 2019). From the 300 beginning of the growing season until the end of our measurement campaign (April to September 2018), air temperatures at the Cham (CHZ) SwissMetNet surface weather station were elevated by 2.2 °C compared to the long-term average from 1994 to 2017 (15.8 °C) during the same period. Total precipitation from April to September 2018, was only 72 % (467 mm) of the long-term average (648 mm) calculated for the period between 1994 and 2017. These specific conditions reduced CO<sub>2</sub> uptake compared to the same period in 2017 (Sect. 3.3) 305 and led to lower grassland productivity and yields of only 6.8 t dry matter (DM) ha<sup>-1</sup> a<sup>-1</sup> in 2018 compared to an average yield of 12.7 t DM ha<sup>-1</sup> a<sup>-1</sup> quantified from 2015 to 2017 (start of the clover experiment). Over the course of the 34 day-campaign (20 July 2018, 02:00-24 July 2018, 08:00 and 09 August 2018, 12:00-06 September 2018, 17:00; all times are in Central European Time, CET= UTC+1) sunny conditions prevailed with a mean solar irradiation (Rg) of 310 W m<sup>-2</sup> during daytime and a mean irradiation of 609 W m<sup>-2</sup> at 13:00. The hourly 310 mean air and soil surface temperature ranged from 13.6 °C (06:00)-24.1 °C (15:00) and from 18.1 °C (08:00)-21.5 °C (18:00), respectively. The median daytime (Rg  $\geq$  5 W m<sup>-2</sup>) and nighttime (Rg < 5 W m<sup>-2</sup>) wind speed was 0.96 m s<sup>-1</sup> (range 0.04–5.77 m s<sup>-1</sup>) and 0.37 m s<sup>-1</sup> (range 0.05–2.49 m s<sup>-1</sup>), respectively. The prevailing wind direction during the day was N-NW (44 %) and E-SE (54 %) at night.

#### 3.2 Performance of the Eddy Mercury system

#### 315 3.2.1 High-frequency signal analysis

Two examples of the raw data used to compute fluxes (Eq. 1) are shown in Fig. 2, one from period 1 with 24 hour calibration intervals (Fig. 2a,c,e) and one with frequent calibrations every four minutes (Fig. 2b,d,f) which strongly reduces the instrument drift (Fig. 2d) as compared to the long calibration intervals (Fig. 2c). In principle, block-averaging raw data within a sampling interval is the best approach to compute EC fluxes (Aubinet et al. 2012). In case of substantial instrument drift as it is seen with the RA-915AM (Fig. 2c) it is necessary to remove the drift by some adequate procedure. Because of the curvature of the drift of the analyzer a simple linear detrending did not lead to satisfactory results, hence we used a third-order polynomial regression fit,

$$\chi' = \chi + \alpha_0 + \alpha_1 \cdot t + \alpha_2 \cdot t^2 + \alpha_3 \cdot t^3 , \qquad (5)$$

- 325 with *t* elapsed time within the averaging interval of 1 h. The turbulent  $Hg^0$  fluctuations after this additional detrending led to the time series shown in Fig. 2e and f. Lengthy discussions on possible shortcomings of such a detrending can be found in Lee et al. (2005) and Aubinet et al. (2012) and thus are not repeated here. It is however clear that in order to obtain higher quality EC fluxes than what we can present here it is required to improve the long-term stability of the instrument (Sect. 3.4).
- 330 Drift of the current version of the *Eddy Mercury* system is substantial (Fig. 3a), an effect that is common with experimental sensor set-ups, but is no longer prevalent with present-day CO<sub>2</sub> sensors. The Allan variance plot (Fig. 3b, see Allan (1966) and Werle et al. (1993)) indicates that the optimum averaging time is ca. 54s. For comparison, a CH<sub>4</sub> analyzer tested by one of the authors (Eugster and Plüss, 2010) shows an optimum average time which is roughly three times as long (ca. 180 s) before the instrument drift starts to dominate the Allan variance. Figure 3b shows that the Allan variance caused by drift at integration times beyond 550 s exceeds the





variance associated with turbulence at the 1 second integration time (see blue arrow in Fig. 3b), whereas in a more ideal instrument (see e.g. Eugster and Plüss, 2010) the long-term drift is smaller than the short-term variance of interest for EC measurements. Despite these findings, Fig. 3 clearly shows the potential and quality of the instrument for  $Hg^0$  flux measurements.

- 340 This interpretation is also supported by spectral and cospectral analyses (Fig. 4). Figure 4a shows an example spectrum of  $Hg^0$  measurements obtained over a 1 hour interval. The difference between the red and black lines in Fig. 4a visualizes the effect of polynomial detrending on the power spectrum of  $Hg^0$ , which is relatively small and of no real concern. Since the RA-915AM only delivers 1 Hz raw data, we had to oversample this digital  $Hg^0$  signal to match the 20.83 Hz resolution of the ultrasonic anemometer. Spectral densities at high frequencies >
- 0.5 Hz (the Nyquist frequency of the RA-915AM, which is ½ of the sampling frequency) are reflecting the effect of oversampling which in the case of the RA-915AM leads to local minima in spectral densities at 1 Hz and all its harmonic multiples (2, 3, 4, ... Hz), which is the result of linear interpolation between measurements. Between these local minima the spectral density obey the f<sup>-1</sup> power law (line "r" in Fig. 4a), which is very close to the inertial subrange slope f<sup>-2/3</sup> (line "i" in Fig. 4a). A damped signal (first order damping; see Eugster and
- Senn, 1995) would follow a  $f^{83}$  power law (line "d" in Fig. 4a), thus it is obvious that our set-up had an adequate flow rate through the RA-915AM that did not lead to substantial damping of the turbulent Hg<sup>0</sup> fluctuations. With the oversampling used here, the white noise level (blue band "w" in Fig. 4a) is artificially reduced below the level that we would obtain without oversampling.
- After having applied an adequate time lag correction to synchronize the detrended Hg<sup>0</sup> signal with vertical wind
   speed fluctuations w', the cospectra of fluxes that are significantly different from a random pattern are closely agreeing with the theoretical idealized cospectrum for neutral atmospheric stability derived from Kaimal et al. (1972) (see Eugster and Senn, 1995) shown by the solid blue line in Fig. 4b. Some minor signs of damping are seen at higher frequencies where the green spline deviates from the solid blue line (Fig. 4b). The comparison of cospectral densities with theoretical damped cospectra (dashed blue lines in Fig. 4b) clearly confirm the finding from the spectral analysis that the flow rate was high enough in the RA-915AM sample cell to prevent
- significant damping effects that tend to be a problem with closed-path EC flux measurements. On occasion of a clear Hg<sup>0</sup> flux which was statistically different from a zero-flux the cross-correlation peak was well defined (Fig. 5a,b). In some occasions with low fluxes relative to the flux detection limit (Sect. 3.2.2) the automatic detection of the cross-correlation peak was not successful. The peak often does not extend very
- 365 strongly beyond the (expected) noise level, as shown in Fig. 5c. However, when zooming in (Fig. 5d) the peak becomes rather clear, although only marginally above the range of insignificant correlations shown with blue background in Fig. 5. To minimize erroneous peak detections and thus wrong flux estimates, we fine-tuned the search window (red band in Fig. 5) for each 1-hour data segment by visually inspecting and selecting the search window within which the local maximum of the absolute correlation coefficient between w and  $\chi$  was found.

#### **370 3.2.2 Flux detection limit**

375

The flux detection limit was calculated for each 1 hour flux period (Sect. 2.6). The significance threshold for  $r_{w,\chi}$  was calculated for an error probability p = 0.05 and the product of this threshold  $r_{w,\chi}$  and measured  $\sigma_w$  and  $\sigma_\chi$  was determined as the flux detection limit for that specific 1 hour period. Figure 6 shows the probability density function of the flux detection limits from all 1 hour data segments. For comparison, the results from the 14 hour zero-flux experiments in the laboratory are added as a blue boxplot to Fig. 6. This comparison clearly shows that



380

385



a zero-flux experiment in the laboratory highly overestimate the quality of  $Hg^0$  flux measurements with a median (maximum) flux detection limit of 0.074 (0.22) ng m<sup>-2</sup> h<sup>-1</sup>. The more realistic flux detection limits based on statistically significant (p < 0.05) correlations are rather in the order of 5.9 (50 % cutoff) to 24 ng m<sup>-2</sup> h<sup>-1</sup> (99 % cutoff) with a 95 % cutoff at 13.7 ng m<sup>-2</sup> h<sup>-1</sup>. Using the same approach but in a qualitative way, Pierce et al. (2015) estimated the flux detection limit of their system to be around 32 ng m<sup>-2</sup> h<sup>-1</sup>.

# 3.2.3 Comparison of detection limits for Eddy Mercury, gradient-based and REA systems

The *Eddy Mercury* system circumvents major sources of uncertainty compared to gradient-based and REA systems, which are related to assumptions on similarity or equivalence of the eddy diffusivities of the scalar transfer coefficients (sensible heat flux, latent heat flux and trace gases). Testing the performance of the *Eddy Mercury* system revealed that 49.7 % of the  $Hg^0$  fluxes (363 out of 731 hours) were significantly different from zero.

Generally, land–atmosphere  $Hg^0$  flux measurements using MM methods are scarce and information on detection limits even rarer. For gradient-based systems a minimum resolvable  $Hg^0$  concentrations gradient (MRG) is determined by mounting the sampling lines at the same height for several days (same-air test) and compute the

- 390 concentration differences between the lines that are used for flux calculations. The MRG threshold is usually defined as the average plus one standard deviation of the concentration difference obtained by the same-air test. Fluxes are considered significant when the Hg<sup>0</sup> concentration difference is above the MRG. Exemplarily, Edwards et al., (2005) derived a flux gradient system-specific MRG of 0.01 ng m<sup>-3</sup> and a flux detection limit of 1.5 ng m<sup>-2</sup> h<sup>-1</sup>. To calculate the flux detection limit the gradient sampling system, site characteristics and
- 395 atmospheric conditions have to be considered (see Eq. 8 in Edwards et al., 2005). Fritsche et al. (2008a) derived a MRG of 0.02 ng m<sup>-3</sup> for their setup. The minimum determinable gradient-based Hg<sup>0</sup> deposition flux was between -0.5 and -4.6 ng m<sup>-2</sup> h<sup>-1</sup> (Fritsche et al., 2008b). Converse et al. (2010) and Zhu et al. (2015b) reported a similar MRG for their gradient-based MM systems of 0.07 and 0.06 ng m<sup>-3</sup>. During Hg<sup>0</sup> flux studies over agricultural land in China, 57 and 62 % of the aerodynamic and modified Bowen-ratio measurements were
- 400 significant (Zhu et al., 2015b). For the only two  $Hg^0$  REA systems, currently applied over terrestrial ecosystems, Zhu et al. (2015b) reported that the absolute precision in the updraft and downdraft  $Hg^0$  concentration difference was concentration (C) dependent at 0.069  $\pm$  0.022 C [ng m<sup>-3</sup>], while Osterwalder et al. (2017) determined a detection limit of 0.05 and 0.04 ng m<sup>-3</sup> for both gold cartridge pairs in their system. Over wheat canopy, 55 % of the fluxes were significant (Zhu et al., 2015a) while 52 % of the fluxes were significant over a boreal peatland
- 405 (Osterwalder et al., 2017). The share of significant  $Hg^0$  fluxes for gradient-based, REA and EC methods is in a similar range (~50 %), when applying the statistical significance test for *Eddy Mercury*. However, the same-air tests applied to determine the detection limit of gradient-based and REA fluxes is more appropriate to compare with our approach to determine the zero-flux in the laboratory. With a median zero-flux of 0.074 ng m<sup>-2</sup> h<sup>-1</sup> as measured in the laboratory, the share of significant fluxes measured with *Eddy Mercury* would increase to 99.7
- 410 %, which, however, is not realistic for measurements outside the laboratory environment. Generally, the reported mean fluxes derived from gradient-based, REA and *Eddy Mercury* should include data below the detection limit because otherwise the magnitudes of the average exchange rates would be overestimated (see Fritsche et al., 2008a; Osterwalder et al., 2016).





#### 3.3 Net ecosystem exchange of Hg<sup>0</sup> over grassland

The median (interquartile range, IQR)  $Hg^0$  flux measured at the Chamau (CH-Cha) research site using the *Eddy* 415 Mercury system was 2.5 (-0.6 to 7.4) ng m<sup>-2</sup> h<sup>-1</sup>. The Hg<sup>0</sup> flux revealed a distinct diel pattern with median (IQR) daytime and nighttime fluxes of 8.4 (1.9 to 15) ng m<sup>-2</sup> h<sup>-1</sup> and 1.0 (-0.9 to 3.3) ng m<sup>-2</sup> h<sup>-1</sup>, respectively. The minimum hourly median  $Hg^0$  flux (0.5 ng m<sup>-2</sup> h<sup>-1</sup>) was detected at 21:00 (Fig 7a). Re-emission of  $Hg^0$  reached a maximum between 11:00 and 14:00 (hourly median 10.8 ng m<sup>-2</sup> h<sup>-1</sup>). The diel Hg<sup>0</sup> variation corresponded with solar radiation with the highest mean level of irradiance at 13:00 (609 W m<sup>-2</sup>). The flux of CO<sub>2</sub> changed from net 420 soil respiration to net uptake by vegetation with sunrise (Fig. 7b). At noon, CO<sub>2</sub> fluxes were 26 % lower compared to the most negative flux occurring between 10:00 and 11:00 (-0.1 mg C m<sup>-2</sup> s<sup>-1</sup>). The absence of a midday maximum CO<sub>2</sub> uptake indicates a midday depression due to plant stress by exceptionally hot and dry conditions. The partial closure of their stomata during the warmest period of the day minimizes water loss 425 through transpiration with the consequence of lower CO<sub>2</sub> uptake. Overall the cumulative CO<sub>2</sub> NEE during our measurement campaign in 2018 was only 8 % compared to the same period in 2017 which exhibited average climatic conditions (Fig. 7b). The median CO<sub>2</sub> uptake in 2018 was 0.008 mg C m<sup>-2</sup> s<sup>-1</sup> compared to 0.08 mg C m<sup>-2</sup> <sup>2</sup> s<sup>-1</sup> measured in 2017. We suggest that the increased stomatal resistance of vegetation during the campaign due to high drought stress not only led to the above discussed minimized uptake of CO<sub>2</sub> but damped stomatal gas exchange in general, thus also the uptake of Hg<sup>0</sup>. Subsequently, soil re-emission was the dominating factor 430

driving GEM-NEE during summer 2018.

The Hg<sup>0</sup> flux measured at the CH-Cha site is comparable to fluxes (n=38) reported for other grassland sites globally (Zhu et al., 2016). A median Hg<sup>0</sup> flux of 0.4 ng m<sup>-2</sup> h<sup>-1</sup> and a flux range between -18.7 and 41.5 ng m<sup>-2</sup> h<sup>-1</sup> (site-based average fluxes) was reported for nine studies (Poissant and Casimir, 1998; Schroeder et al, 2005;

- Ericksen et al. 2006; Obrist et al. 2006; Fu et al. 2008a,b; Fritsche et al. 2008a,b; Converse et al. (2010). Several studies reported a net re-emission of Hg<sup>0</sup> during summer. Converse et al. (2010) reported net average Hg<sup>0</sup> re-emission of 2.5 ng m<sup>-2</sup> h<sup>-1</sup> from a high-elevation wetland meadow in Virginia, USA. Zhang et al. (2001) measured a Hg<sup>0</sup> flux of 7.6 ± 1.7 ng m<sup>-2</sup> h<sup>-1</sup> from an open background site in Michigan, USA. The average Hg<sup>0</sup> flux from a grassland in Québec, CA, was 2.95 ± 2.15 ng m<sup>-2</sup> h<sup>-1</sup> and a correlation of the dial flux cycle with solar radiation was reported (Poissant and Casimir 1998). Average net Hg<sup>0</sup> emission of 1.1 ng m<sup>-2</sup> h<sup>-1</sup> was
- recorded from a pasture in Ontario (Schroeder et al, 2005). The mean  $Hg^0$  flux from four grassland sites in the USA ranged from 0.3–2.5 ng m<sup>-2</sup> h<sup>-1</sup> between May 2003 and 2004 (Ericksen et al. 2006). Fu et al. (2008a) reported average  $Hg^0$  fluxes ranging from -1.7 to 13.4 ng m<sup>-2</sup> h<sup>-1</sup> from three grasslands in China in August 2006. Mechanism driving  $Hg^0$  re-emission from grasslands is not fully understood yet. Photoreduction has been
- reported to enhance Hg<sup>0</sup> emission from the soil and foliage surface and from Hg within foliar tissue (Gustin et al., 2002; Moore and Carpi, 2005; Choi and Holsen, 2009; Yuan et al., 2019). Soil warming has been suggested to promote Hg<sup>0</sup> re-emission (Poissant et al., 1999; Zhang et al., 2001; Gustin et al., 2002; Almeida et al., 2009), likely due to increased decomposition of organic material (Fritsche et al., 2008c) and facilitated mass transfer of Hg<sup>0</sup> through the topsoil to the atmosphere (Lin and Pehkonen, 1999). Zhang et al. (2001) reported a strong correlation of Hg<sup>0</sup> re-emission fluxes with solar radiation and soil temperature. A solar shielding experiment resulted in a 65 % decrease of soil Hg<sup>0</sup> re-emission suggesting that photoreduction is a major factor but also soil

temperature cannot be neglected. Few grassland studies have shown net Hg<sup>0</sup> dry deposition. Fritsche et al. (2008a) reported an average Hg<sup>0</sup> flux of -1.7 ng m<sup>-2</sup> h<sup>-1</sup> (modified Bowen-ratio) and -4.3 ng m<sup>-2</sup> h<sup>-1</sup> (aerodynamic gradient) during the vegetation period



475

485



over a sub-alpine grassland at Fruebuel in central Switzerland, 15 km SW of our study site. More summertime Hg<sup>0</sup> fluxes from three Central European grasslands were measured on a campaign basis and average grassland– atmosphere Hg<sup>0</sup> fluxes ranged from -4.3 to 0.3 ng m<sup>-2</sup> h<sup>-1</sup>. The highest variability of the fluxes was recorded for the Neustift site in Austria with a range of -76 to 37 ng m<sup>-2</sup> h<sup>-1</sup> (Fritsche et al., 2008b). A second full year Hg<sup>0</sup> flux study was performed at an upland meadow in Maryland, USA (Castro and Moore 2016). The hourly mean summertime Hg<sup>0</sup> flux was -1.2 ng m<sup>-2</sup> h<sup>-1</sup> and ranged between -224 and 354 ng m<sup>-2</sup> h<sup>-1</sup>.

We found that the southern Hg<sup>0</sup> source area of our grassland site has a 28 % higher Hg substrate concentration (mean = 59.4  $\pm$  8.4 ng Hg g<sup>-1</sup>) compared to the northern source area (mean = 46.4  $\pm$  5.1 ng Hg g<sup>-1</sup>) (Wilcoxon two sample t-test, p < 0.05, Fig. 8a). The *Eddy Mercury* system was able to resolve a marginally significant greater daytime (+44 %, p = 0.0515) Hg<sup>0</sup> flux (Fig. 8c) and insignificantly greater nighttime Hg<sup>0</sup> flux (+68 %, p

465 = 0.296) (Fig. 8b) originating from the Hg enriched southern source area. The proportionality of Hg<sup>0</sup> re-emission to soil Hg concentration has been shown across Hg-enriched soils (Eckley et al., 2015; Zhu et al., 2018; Osterwalder et al., 2019) but no significant correlation has been observed for low-Hg level background soils (Agnan et al., 2016). There are two possible explanations for the lack of a significant relationship between Hg<sup>0</sup> flux and soil Hg concentration: (i) analytical uncertainty of Hg<sup>0</sup> flux measurements or (ii), at vegetated surfaces, a masking of Hg<sup>0</sup> re-emission by stomatal uptake of Hg<sup>0</sup> which is independent on the soil Hg concentration.

3.4 Suggestions to improve the Eddy Mercury system

During the pilot campaign and the data analysis we found room for optimization of the *Eddy Mercury* system. Here we propose a number of adjustments that are expected to improve the system's performance in particular by 1) facilitating data transfer and processing, 2) increasing the measurement frequency and sample air flow through the RA-915AM and 3) achieving more stable temperature conditions in the field.

<u>Improve data transfer:</u> The determination of the time lag between the wind speed measurement and the  $Hg^0$  concentration measurement bearded a considerable source of uncertainty and cross-correlation peaks had to be visually verified (Sect. 2.5.4.). In the future, we aim for a real time transfer of raw data to the serial port instead of data transfer via the USB port on the embedded Windows 7 system of the RA-915AM. This will allow a better synchronization between the  $Hg^0$  measurements and the ultrasonic anemometer (Sect. 2.4) and

better synchronization between the Hg<sup>0</sup> measurements and the ultrasonic anemometer (Sect. 2.4) and significantly facilitate post-acquisition data treatment.
 <u>Increase measurement frequency:</u> The pilot campaign was performed with a measurement frequency of 1 Hz. In

the future, we wish to increase the measurement frequency up to 8-15 Hz. Such an increase in measurement frequency is possible through software adaptations of the RA-915AM and will make the oversampling of the Hg<sup>0</sup> signal performed here (Sect. 3.2) redundant and result in better counting statistics.

<u>Increase sample flow rate:</u> During this pilot study we connected a more powerful pump to the RA-915AM and managed to increase the flow rate from standard operation of 7–10 L min<sup>-1</sup> to 14.3 L min<sup>-1</sup> resulting in a two times lower residence time in the measurement cell. The lower residence time in the cell reduced the dampening of the signal (Sect. 3.2). However, this high flow led to a reduction in the cell pressure (approx. 700 mbar) which affects the detection limit for Hg<sup>0</sup> concentration measurements. In the future, we propose to further reduce the residence time of the air in the measurement cell by increasing the sample air flow by another 30 % to 20 L min<sup>-1</sup> using an external pump. To account for pressure drop we propose to minimize the constrictions present in the

RA-915AM by increasing the internal diameter of the valves and the inlet tubing.





Improve the long-term stability of the instrument: The stability of RA-915AM Hg<sup>0</sup> concentration measurements is temperature dependent (Sect. 3.2). We encountered strong diurnal temperature fluctuations of the instrument during the pilot campaign. We took several measures already during the campaign to increase the temperature stability (e.g. placing the pump outside the temperature controlled analyzer box, isolation of the analyzer box and shading it from direct sunlight). To improve the temperature stability in the future, we suggest to place the RA-915AM in an instrument box that has a better isolation and more powerful temperature control or ideally to place it in a climate controlled instrumental hut. For long-term deployments of the *Eddy Mercury* the sampling hose can be extended to bridge the distance between the air inlet, located close to the sonic anemometer and the instrumental hut where the system is placed. In that case it is important to guarantee a turbulent flow in the tube (Reynolds number of > 3000–3500; Lenschow and Raupach, 1991; Leuning and King, 1992), an adequate refresh rate in the sampling cell and to ensure that the pressure drop in the sampling cell is within the requirements of the instruments (> 600 mbar; pers. communication with Lumex Ltd.).

# 4 Conclusion

This study demonstrates the first successful application of the EC method for Hg<sup>0</sup> flux measurements over terrestrial surfaces with background soil Hg concentrations (< 100 ng  $g^{-1}$ ). We tested and validated the system at a Central European grassland site. The maximum flux detection limit derived from a zero-flux experiment in the lab was 0.22 ng m<sup>-2</sup> h<sup>-1</sup>. The statistical estimate of the flux detection limit under real-world conditions was 5.9 510 (50 % cutoff) to 13.7 ng m<sup>-2</sup> h<sup>-1</sup> (95 % cutoff). The *Eddy Mercury* system overcomes major uncertainties of other micrometeorological methods previously used for Hg<sup>0</sup> flux measurements associated with the intermittent sampling at two different levels (aerodynamic methods) and the stringent sampling and analytical requirements (relaxed eddy accumulation). The system will considerably facilitate ecosystem-scale Hg<sup>0</sup> flux measurement 515 because it features a fully automated operation, cutting down operation costs for technical maintenance by experienced staff, argon supply and consumables. Eddy Mercury bears the potential to be established as a standard micrometeorological method for long-term Hg<sup>0</sup> measurements over grassland and other terrestrial ecosystems. Such a standardization of measurements is strongly required to obtain comparable data and properly evaluate controlling factors on the net ecosystem exchange of Hg<sup>0</sup> on larger spatial- and temporal scales (Obrist 520 et al., 2018). Ultimately, the Eddy Mercury system could complement air pollution and greenhouse gas measurements within the global network of micrometeorological tower sites (FluxNet) assessing the impacts of controlling Hg emissions on deposition and re-emission (Baldocchi et al., 2001). The Eddy Mercury system also comes at an opportune time to include net ecosystem exchange measurements of Hg<sup>0</sup> in the joint WHO and UN Environment project to "develop a plan for global monitoring of human exposure to and environmental

525 concentration of mercury".





# Data availability

The data will be submitted to a data repository after final publication of the manuscript.

#### Author contributions

530

All authors contributed to designing the study, testing the RA-915AM in the laboratory and performing fieldwork. WE analyzed the data. Soil samples were taken by IF and analyzed for total mercury by MJ. IF analyzed the  $CO_2$  flux and meteorological data. MJ and SO coordinated the study. SO, WE and MJ wrote the paper with contributions of IF.

#### **Competing interests**

The authors declare that they have no conflict of interest.

# 535 Acknowledgements

This research was funded by the Institute of Agricultural Sciences, ETH Zurich; the Department of Environmental Geosciences, University of Basel; and the Freiwillige Akademische Gesellschaft (FAG) Basel. MJ received funding from the Swiss National Science Foundation, Ambizione grant (PZ00P2\_174101). We want to acknowledge Prof. Dr. Nina Buchmann for scientific and Paul Linwood for on-site technical support.

- 540 During fieldwork we appreciated the technical and logistical assistance by Rudolf Osterwalder from Mühlau (AG). We thank Dr. Ingvar Wängberg of the Swedish Environmental Research Institute (IVL) in Gothenburg for providing preliminary 1 Hz data of Hg<sup>0</sup> in ambient air to encourage the authors to carry out the present study. We thank the Federal Office of Meteorology and Climatology MeteoSwiss for providing data from the Cham (CHZ) weather station. We gratefully acknowledge the Lumex Instruments staff members, namely Vladimir
- 545 Ryzhov, Dr. Sergey Sholupov and Dr. Georg Debus for their enthusiasm and invaluable technical support on the RA-915AM instrument and fruitful discussions during the field campaign and data analysis.





### References

| 550 | Agnan, Y., Le Dantec, T., Moore, C. W., Edwards, G. C. and Obrist, D.: New constraints on terrestrial surface–<br>atmosphere fluxes of gaseous elemental mercury using a global database, Environ. Sci. Technol., 50,<br>507–524, https://doi.org/10.1021/acs.est.5b04013, 2016.  |
|-----|---|
|     | Allan, D. W.: Statistics of atomic frequency standards, Proceedings of the IEEE, 54, 221–231,<br>https://doi.org/10.1109/PROC.1966.4634, 1966.  |
| 555 | Almeida, M. D., Marins, R. V., Paraquetti, H. H. M., Bastos, W. R., and Lacerda, L. D.: Mercury degassing from<br>forested and open field soils in Rondônia, Western Amazon, Brazil, Chemosphere, 77, 60–66,<br>https://doi.org/10.1016/j.chemosphere.2009.05.018, 2009.  |
|     | Aubinet M, Vesala T and Papale, D.: Eddy Covariance: A Practical Guide to Measurement and Data Analysis,<br>Springer, Dordrecht, Heidelberg, London, New York, 438 pp., 2012.   |
| 560 | <ul> <li>Baldocchi, D., Falge, E., Gu, L. H., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K.,</li> <li>Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X. H., Malhi, Y., Meyers, T., Munger,</li> <li>W., Oechel, W., U, K., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K.</li> <li>and Wofsy, S.: FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale</li> <li>carbon dioxide, water vapor, and energy flux densities, Bull. Amer. Meteorol. Soc., 82, 2415–2434,</li> <li>https://doi.org/10.1175/1520-0477(2001)082&lt;2415:FANTTS&gt;2.3.CO;2, 2001.</li> </ul> |
| 565 | Bash, J. O. and Miller, D. R.:: A relaxed eddy accumulation system for measuring surface fluxes of total gaseous mercury, J. Atmos. Ocean. Technol. 25, 244–257, https://doi.org/10.1175/2007JTECHA908.1, 2008.   |
| 570 | <ul> <li>Braune, B., Chetelat, J., Amyot, M., Brown, T., Clayden, M., Evans, M., Fisk, A., Gaden, A., Girard, C., Hare, A., Kirk, J., Lehnherr, I., Letcher, R., Loseto, L., Macdonald, R., Mann, E., McMeans, B., Muir, D., O'Driscoll, N., Poulain, A., Reimer, K. and Stern, G.: Mercury in the marine environment of the Canadian Arctic: Review of recent findings, Sci. Total Environ., 509, 67–90, https://doi.org/10.1016/j.scitotenv.2014.05.133, 2015.</li> </ul>   |
| 575 | Brown, R., Pirrone, N., van Hoek, C., Horvat, M., Kotnik, J., Wängberg, I., Corns, W., Bieber, E., and Sprovieri,<br>F.: Standardization of a European measurement method for the determination of total gaseous mercury:<br>results of the field trial campaign and determination of a measurement uncertainty and working range,<br>Accredit. Qual. Assur., 15, 359–366, https://doi.org/doi:10.1007/s00769-010-0636-2, 2010.   |
|     | Burba, G. G., McDermitt, D. K., Grelle, A., Anderson, D. J. and Xu, L.: Addressing the influence of instrument<br>surface heat exchange on the measurements of CO <sub>2</sub> flux from open-path gas analyzers, Glob. Change<br>Biol., 14, 1854–1876, https://doi.org/10.1111/j.1365-2486.2008.01606.x, 2008.   |
| 580 | Businger, J. A.: Evaluation of the Accuracy with Which Dry Deposition Can Be Measured with Current<br>Micrometeorological Techniques, J. Climate Appl. Meteor., 25, 1100–1124,<br>https://doi.org/10.1175/1520-0450(1986)025<1100:EOTAWW>2.0.CO;2, 1986.  |
|     | Businger, J. and Oncley, S.: Flux Measurement with Conditional Sampling, J. Atmos. Ocean. Technol., 7, 349–352, https://doi.org/10.1175/1520-0426(1990)007<0349:FMWCS>2.0.CO;2, 1990.   |





| Castro, M. S. and Moore, C. W.: Importance of Gaseous Elemental Mercury Fluxes in West<br>585 Atmosphere 7, 110, https://doi.org/10.3390/atmos7090110, 2016.  |  |  |  |  |  |  |  |  |  |  |  |
|---|--|--|--|--|--|--|--|--|--|--|--|
|   | Choi, H. D. and Holsen, T. M.: Gaseous mercury emissions from unsterilized and sterilized soils: The effect of<br>temperature and UV radiation, Environ. Poll., 157, 1673–1678,<br>https://doi.org/10.1016/j.envpol.2008.12.014,2009.  |  |  |  |  |  |  |  |  |  |  |
| 590   | Cobos, D. R., Baker, J. M., and Nater, E. A.: Conditional sampling for measuring mercury vapor fluxes, Atmos.<br>Environ., 36, 4309–4321, https://doi.org/10.1016/S1352-2310(02)00400-4, 2002.   |  |  |  |  |  |  |  |  |  |  |
|   | Converse, A. D., Riscassi, A. L. and Scanlon, T. M.: Seasonal variability in gaseous mercury fluxes measured in<br>a high-elevation meadow, Atmos. Environ. 44, 2176–2185,<br>https://doi.org/10.1016/j.atmosenv.2010.03.024, 2010.  |  |  |  |  |  |  |  |  |  |  |
| 595   | Demers, J. D., Driscoll, C. T., Fahey, T. J. and Yavitt, J. B.: Mercury cycling in litter and soil in different forest types in the Adirondack region, New York, USA, Ecol. Appl., 17, 1341–1351, 2007.  |  |  |  |  |  |  |  |  |  |  |
|   | Driscoll, C. T.; Mason, R. P.; Chan, H. M.; Jacob, D. J. and Pirrone, N.: Mercury as a global pollutant: sources, pathways, and effects, Environ. Sci. Technol., 47, 4967–4983, https://doi.org/10.1021/es305071v, 2013.   |  |  |  |  |  |  |  |  |  |  |
| 600   | Eckley, C. S., Blanchard, P., McLennan, D., Mintz, R. and Sekela, M.: Soil–Air Mercury Flux near a Large<br>Industrial Emission Source before and after Closure (Flin Flon, Manitoba, Canada), Environ. Sci<br>Technol., 49, 9750–9757, https://doi.org/10.1021/acs.est.5b01995, 2015.   |  |  |  |  |  |  |  |  |  |  |
|   | Eckley, C. S., Tate, M. T., Lin, CJ., Gustin, M., Dent, S., Eagles-Smith, C., Lutz, M. A., Wickland, K. P.,<br>Wang, B., Gray, J. E., Edwards, G. C., Krabbenhoft, D. P. and Smith, D. B.: Surface-air mercury fluxes<br>across Western North America: A synthesis of spatial trends and controlling variables, Sci. Total<br>Environ., 568, 651–665, https://doi.org/10.1016/j.scitotenv.2016.02.121, 2016. |  |  |  |  |  |  |  |  |  |  |
| 605 Edwards, G. C., Rasmussen, P. E., Schroeder, W. H., Wallace, D. M., Halfpenny-Mitchell, L., Dia<br>Kemp, R. J. and Ausma, S.: Development and evaluation of a sampling system to determin<br>mercury fluxes using an aerodynamic micrometeorological gradient method, J. Geophys. Res<br>110, https://doi.org/10.1029/2004JD005187, 2015. |  |  |  |  |  |  |  |  |  |  |  |
| 610   | Enrico, M., Roux, G. L., Marusczak, N., Heimbürger, LE., Claustres, A., Fu, X., Sun, R. and Sonke, J. E.<br>Atmospheric Mercury Transfer to Peat Bogs Dominated by Gaseous Elemental Mercury Dry<br>Deposition, Environ. Sci. Technol., 50, 2405–2412, http://doi.org/10.1021/acs.est.5b06058, 2016.   |  |  |  |  |  |  |  |  |  |  |
|   | Ericksen, J. A., Gustin, M. S., Xin, M., Weisberg, P. J. and Fernandez, G. C. J.: Air-soil exchange of mercury<br>from background soils in the United States, Sci. Total Environ., 366, 851–863,<br>https://doi.org/10.1016/j.scitotenv.2005.08.019, 2006.   |  |  |  |  |  |  |  |  |  |  |
| 615   | Eugster, W. and Merbold, L.: Eddy covariance for quantifying trace gas fluxes from soils, SOIL 1, 187–205, https://doi.org/10.5194/soil-1-187-2015, 2015.  |  |  |  |  |  |  |  |  |  |  |
|   | Eugster, W. and Plüss, P.: A fault-tolerant eddy covariance system for measuring CH <sub>4</sub> fluxes. Agricultural and Forest Meteorology, Special Issue on Eddy Covariance (EC) flux measurements of CH <sub>4</sub> and N <sub>2</sub> O, 150, 841–851, https://doi.org/10.1016/j.agrformet.2009.12.008, 2010.  |  |  |  |  |  |  |  |  |  |  |

620 Eugster, W. and Senn, W.: A cospectral correction model for measurement of turbulent NO<sub>2</sub> flux, Bound.-Lay.



630

635

640

645

650

655



Meteorol., 74, 321-340, https://doi.org/10.1007/BF00712375, 1995.

- Eugster, W. and Zeeman, M. J.: Micrometeorological techniques to measure ecosystem-scale greenhouse gas fluxes for model validation and improvement, Int. Congr. Ser., 1293, 66–75, https://doi.org/10.1016/j.ics.2006.05.001, 2006.
- 625 Eugster, W., Zeyer, K., Zeeman, M., Michna, P., Zingg, A., Buchmann, N. and Emmenegger, L.: Methodical study of nitrous oxide eddy covariance measurements using quantum cascade laser spectrometry over a Swiss forest, Biogeosciences, 4, 927–939, https://doi.org/10.5194/bg-4-927-2007, 2007.
  - Faïn, X., Moosmueller, H. and Obrist, D.: Toward real-time measurement of atmospheric mercury concentrations using cavity ring-down spectroscopy, Atmos. Chem. Phys., 10, 2879–2892, https://doi.org/10.5194/acp-10-2879-2010, 2010.
  - Fitzgerald, W. F., Lamborg, C. H. and Hammerschmidt, C. R.: Marine biogeochemical cycling of mercury, Chem. Rev., 107, 641–662, https://doi.org/10.1021/cr050353m, 2007.
  - Fritsche, J., Obrist, D., Zeeman, M. J., Conen, F., Eugster, W. and Alewell, C.: Elemental mercury fluxes over a sub-alpine grassland determined with two micrometeorological methods, Atmos. Environ., 42, 2922– 2933, https://doi.org/10.1016/j.atmosenv.2007.12.055, 2008a.
  - Fritsche, J., Wohlfahrt, G., Ammann, C., Zeeman, M., Hämmerle, A., Obrist, D. and Alewell, C.: Summertime elemental mercury exchange of temperate grasslands on an ecosystem-scale, Atmos. Chem. Phys. 8, 7709–7722, https://doi.org/10.5194/acp-8-7709-2008, 2008b.
  - Fritsche, J., Obrist, D. and Alewell, C.: Evidence of microbial control of Hg<sup>0</sup> emissions from uncontaminated terrestrial soils, J. Plant Nutr. Soil Sci., 171, 200–209, https://doi.org/10.1002/jpln.200625211, 2008c
    - Fu, X. W., Feng, X. B., and Wang, S. F.: Exchange fluxes of Hg between surfaces and atmosphere in the eastern flank of Mount Gongga, Sichuan province, southwestern China, J. Geophys. Res.-Atmos., 113, D20306, doi:10.1029/2008JD009814, 2008a.
  - Fu, X. W., Feng, X. B., Wang, S. F., Qiu, G. L., and Li, P.: Mercury flux rate of two types of grasslands in Guiyang, Huanjing Kexue Yanjiu, 20, 33–37, 2008b (in Chinese with English Abstract).
  - Fuchs, K., Hörtnagl, L., Buchmann, N., Eugster, W., Snow, V. and Merbold, L.: Management matters: testing a mitigation strategy for nitrous oxide emissions using legumes on intensively managed grassland, Biogeosciences, 15, 5519–5543, https://doi.org/10.5194/bg-15-5519-2018, 2018.
  - Grigal, D. F.: Mercury sequestration in forests and peatlands: A review, J. Environ. Qual. 2003, 32, 393–405, 2003.
  - Gustin, M. S., Lindberg, S., Marsik, F., Casimir, A., Ebinghaus, R., Edwards, G., Hubble-Fitzgerald, C., Kemp, R., Kock, H., Leonard, T., London, J., Majewski, M., Montecinos, C., Owens, J., Pilote, M., Poissant, L., Rasmussen, P., Schaedlich, F., Schneeberger, D., Schroeder, W., Sommar, J., Turner, R., Vette, A., Wallschlaeger, D., Xiao, Z. and Zhang, H.: Nevada STORMS project: Measurement of mercury emissions from naturally enriched surfaces, J. Geophys. Res.-Atmos., 104, 21831–21844, https://doi.org/10.1029/1999JD900351, 1999.





| Gustin, | M. S., Bi | ester, H.,  | and Kim, C  | . S.: Inv | vestigation | of the | light-enhand | ed emission    | of mercury  | from  |
|---------|-----------|-------------|-------------|-----------|-------------|--------|--------------|----------------|-------------|-------|
|         | naturally | enriched    | substrates, | Atmos.    | Environ.,   | 36,    | 3241–3254,   | https://doi.or | g/10.1016/S | 1352- |
|         | 2310(02)  | 00329-1, 20 | 002.        |           |             |        |              |                |             |       |

- Horst, T. W.: A simple formula for attenuation of eddy fluxes measured with first-order-response scalar sensors, Bound.-Lay. Meteorol., 82, 219–233, https://doi.org/10.1023/A:1000229130034, 1997.
- Jiskra, M., Wiederhold, J. G., Skyllberg, U., Kronberg, R. M., Hajdas, I. and Kretzschmar, R.: Mercury deposition and re-emission pathways in boreal forest soils investigated with Hg isotope signatures, Environ. Sci. Technol., 49, 7188–7196, https://doi.org/10.1021/acs.est.5b00742, 2015.
- Jiskra, M., Sonke, J. E., Obrist, D., Bieser, J., Ebinghaus, R., Myhre, C. L., Pfaffhuber, K. A., Wängberg, I., Kyllönen, K., Worthy, D., Martin, L. G., Labuschagne, C., Mkololo, T., Ramonet, M., Magand, O. and Dommergue, A.: A vegetation control on seasonal variations in global atmospheric mercury concentrations, Nat. Geosci., 11, 244–250, https://doi.org/10.1038/s41561-018-0078-8, 2018.
- 670 Kaimal, J. C. and Gaynor, J. E.: Another look at sonic thermometry, Bound.-Lay. Meteorol., 56, 401–410, https://doi.org/10.1007/BF00119215, 1991.
  - Kaimal, J. C., Wyngaard, J. C., Izumi, Y. and Coté, O. R.: Spectral characteristics of surface-layer turbulence, Q. J. Roy. Meteor. Soc., 98, 563–589, https://doi.org/10.1002/qj.49709841707, 1972.
- Kamp, J., Skov, H., Jensen, B. and Sørensen, L. L.: Fluxes of gaseous elemental mercury (GEM) in the High
   Arctic during atmospheric mercury depletion events (AMDEs). Atmos. Chem. Phys., 18, 6923–6938. https://doi.org/10.5194/acp-18-6923-2018, 2018.
  - Kljun, N., Calanca, P., Rotach, M. W., and Schmid, H. P.: A simple two-dimensional parameterisation for Flux Foot Footprint Prediction (FFP), Geosci. Model Dev., 8, 3695–3713, https://doi.org/10.5194/gmd-8-3695-2015, 2015.
- Lee, X., Finnigan, J. and Paw U, K.T.: Coordinate Systems and Flux Bias Error, in: Handbook of Micrometeorology: A Guide for Surface Flux Measurement and Analysis, edited by: Lee, X., Massman, W. and Law, B., Atmospheric and Oceanographic Sciences Library, Springer Netherlands, Dordrecht, 33–66, https://doi.org/10.1007/1-4020-2265-4\_3, 2005.
- Lenschow, D. H. and Raupach, M. R.: The attenuation of fluctuations in scalar concentrations through sampling
   tubes, J. Geophys. Res., 96, 15259–15268, https://doi.org/10.1029/91JD01437, 1991.
  - Leuning, R. and King, K. M.: Comparison of eddy-covariance measurements of CO<sub>2</sub> fluxes by open- and closedpath CO<sub>2</sub> analysers, Bound.-Lay. Meteorol., 59, 297–311, https://doi.org/10.1007/BF00119818, 1992.
  - Lin, C. J. and Pehkonen, S. O.: The chemistry of atmospheric mercury: a review, Atmos. Environ., 33, 2067–2079, https://doi.org/10.1016/S1352-2310(98)00387-2, 1999.
- 690 Lindberg, S.; Bullock, R.; Ebinghaus, R.; Engstrom, D.; Feng, X.; Fitzgerald, W.; Pirrone, N.; Prestbo, E. and Seigneur, C.: A synthesis of progress and uncertainties in attributing the sources of mercury in deposition, Ambio, 36, 19–32, https://doi.org/10.1579/0044-7447(2007)36[19:asopau]2.0.co;2, 2007
  - Mason, R. P., Choi, A. L., Fitzgerald, W. F., Hammerschmidt, C. R., Lamborg, C. H., Soerensen, A. L. and Sunderland, E. M.: Mercury biogeochemical cycling in the ocean and policy implications, Environ.





| 695 | Res., 119, 101–117, https://doi.org/10.1016/j.envres.2012.03.013, 2012   |
|-----|--|
|     | Mauder, M. and Foken, T.: Documentation and instruction manual of the eddy covariance software package TK2, Work Report, University of Bayreuth, Bayreuth, 2004.   |
|     | McMillen, R. T.: An Eddy-Correlation Technique with Extended Applicability to Non-Simple Terrain, Bound<br>Lay. Meteorol., 43, 231–245, https://doi.org/10.1007/BF00128405, 1988.  |
| 700 | Merbold, L., Eugster, W., Stieger, J., Zahniser, M., Nelson, D., and Buchmann, N.: Greenhouse gas budget<br>(CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O) of intensively managed grassland following restoration, Glob. Change Biol., 20,<br>1913–1928, https://doi.org/10.1111/gcb.12518, 2014.                                |
|     | MeteoSchweiz 2019: Klimareport 2018. Bundesamt für Meteorologie und Klimatologie MeteoSchweiz, Zürich, 94 S.   |
| 705 | Moncrieff, J., Clement, R., Finnigan, J., and Meyers, T.: Averaging, detrending, and filtering of eddy covariance<br>time series, in: Handbook of Micrometeorology: a guide for surface flux measurement and analysis, 29,<br>edited by: Law, B. E., Lee, X., and Massmann, W. J., Kluwer Academic, Dordrecht, 7–31, 2004.                 |
|     | Montgomery, R. B.: Vertical eddy flux of heat in the atmosphere, J. Meteor., 5, 265–274,<br>https://doi.org/10.1175/1520-0469(1948)005<0265:VEFOHI>2.0.CO;2, 1948.   |
| 710 | Moore, C. and Carpi, A.: Mechanisms of the emission of mercury from soil: Role of UV radiation, J. Geophys. ResAtmos., 110, D24302, https://doi.org/10.1029/2004JD005567, 2005.  |
|     | Obrist, D., Conen, F., Vogt, R., Siegwolf, R., and Alewell, C.: Estimation of Hg <sup>0</sup> exchange between ecosystems and the atmosphere using <sup>222</sup> Rn and Hg <sup>0</sup> concentration changes in the stable nocturnal boundary layer, Atmos. Environ., 40, 856–866, https://doi.org/10.1016/j.atmosenv.2005.10.012, 2006. |
| 715 | Obrist, D., Agnan, Y., Jiskra, M., Olson, C. L., Colegrove, D. P., Hueber, J., Moore, C. W., Sonke, J. E. and<br>Helmig, D.: Tundra uptake of atmospheric elemental mercury drives Arctic mercury pollution, Nature<br>547, 201-204, https://doi.org/10.1038/nature22997, 2017.  |
| 720 | Obrist, D., Kirk, J. L., Zhang, L., Sunderland, E. M., Jiskra, M. and Selin, N. E.: A review of global environmental mercury processes in response to human and natural perturbations: Changes of emissions, climate, and land use, Ambio, 47, 116–140, https://doi.org/10.1007/s13280-017-1004-9, 2018.                                   |
|     | Obukhov A. M.: Characteristics of the Micro-structure of the Wind in the Surface Layer of the Atmosphere, Izv. AN SSSR, ser. Geofiz. 3:49–68, 1951.  |
| 725 | Osterwalder, S., Fritsche, J., Alewell, C., Schmutz, M., Nilsson, M. B., Jocher, G., Sommar, J., Rinne, J. and<br>Bishop, K.: A dual-inlet, single detector relaxed eddy accumulation system for long-term measurement<br>of mercury flux, Atmos. Meas. Tech., 9, 509–524, https://doi.org/10.5194/amt-9-509-2016, 2016.                   |
|     | Osterwalder, S., Bishop, K., Alewell, C., Fritsche, J., Laudon, H., Åkerblom, S. and Nilsson, M. B.: Mercury evasion from a boreal peatland shortens the timeline for recovery from legacy pollution, Sci. Rep., 7, 16022, https://doi.org/10.1038/s41598-017-16141-7, 2017.   |
| 730 | Osterwalder, S., Sommar, J., Åkerblom, S., Jocher, G., Fritsche, J., Nilsson, M. B., Bishop, K. and Alewell, C.:   |



735

745

760



Comparative study of elemental mercury flux measurement techniques over a Fennoscandian boreal peatland, Atmos. Environ., 172, 16–25, https://doi.org/10.1016/j.atmosenv.2017.10.025, 2018.

- Osterwalder, S., Huang, J.-H., Shetaya, W. H., Agnan, Y., Frossard, A., Frey, B., Alewell, C., Kretzschmar, R., Biester, H. and Obrist, D.: Mercury emission from industrially contaminated soils in relation to chemical, microbial, and meteorological factors, Environ. Pollut., 250, 944–952, https://doi.org/ 10.1016/j.envpol.2019.03.093, 2019.
  - Outridge, P. M., Mason, R. P., Wang, F., Guerrero, S. and Heimbürger-Boavida, L. E.: Updated Global and Oceanic Mercury Budgets for the United Nations Global Mercury Assessment 2018., Environ. Sci. Technol., 52, 11466–11477, https://doi.org/10.1021/acs.est.8b01246, 2018
- 740 Pierce, A., Obrist, D., Moosmüller, H., Faïn, X. and Moore, C.: Cavity ring-down spectroscopy sensor development for high-time-resolution measurements of gaseous elemental mercury in ambient air. Atmos. Meas. Tech., 6, 1477–1489, https://doi.org/10.5194/amt-6-1477-2013, 2013.
  - Pierce, A. M., Moore, C.W., Wohlfahrt, G., Hörtnagl, L., Kljun, N. and Obrist, D.: Eddy covariance flux measurements of gaseous elemental mercury using cavity ring-down spectroscopy, Environ. Sci. Technol., 49, 1559–1568, https://doi.org/10.1021/es505080z, 2015.
  - Poissant, L. and Casimir, A.: Water-air and soil-air exchange rate of total gaseous mercury measured at background sites, Atmos. Environ., 32, 883–893, https://doi.org/10.1016/S1352-2310(97)00132-5, 1998.
- Poissant, L., Pilote, M., and Casimir, A.: Mercury flux measurements in a naturally enriched area: Correlation 750 with environmental conditions during the Nevada Study and tests of the release of mercury from soils (STORMS), J. Geophys. Res.-Atmos., 104, 21845–21857, https://doi.org/10.1029/1999JD900092, 1999.
  - R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. URL , 2018.
- 755 Roth, K.: Bodenkartierung und GIS-basierte Kohlenstoffinventur von Graslandböden, Diploma Thesis, University of Zurich, 2006.
  - Saiz-Lopez, A., Sitkiewicz, S. P., Roca-Sanjuán, D., Oliva-Enrich, J. M., Dávalos, J. Z., Notario, R., Jiskra, M., Xu, Y., Wang, F., Thackray, C. P., Sunderland, E. M., Jacob, D. J., Travnikov, O., Cuevas, C. A., Acuña, A. U., Rivero, D., Plane, J. M. C., Kinnison, D. E. and Sonke, J. E.: Photoreduction of gaseous oxidized mercury changes global atmospheric mercury speciation, transport and deposition, Nat. Commun., 9, 4796, https://doi.org/10.1038/s41467-018-07075-3, 2018.
    - Schroeder, W. H., Beauchamp, S., Edwards, G., Poissant, L., Rasmussen, P., Tordon, R., Dias, G., Kemp, J., Van Heyst, B. and Banic, C. M.: Gaseous mercury emissions from natural sources in Canadian landscapes, J. Geophys. Res.-Atmos., 110, https://doi.org/10.1029/2004JD005699, 2005.
- 765 Sholupov, S. and Ganeyev, A.: Zeeman atomic absorption spectrometry using high frequency modulated light polarization, Spectrochim. Acta B, 50, 1227–1236, https://doi.org/doi:10.1016/0584-8547(95)01316-7, 1995.



770

790



| Sholupov, | S., Pogarev,  | S., R      | yzhov, V   | /., N | Mashyanov,     | N.,   | and   | Stroganov    | , A.:  | Zeeman    | atomic   | absor   | ption |
|-----------|---------------|------------|------------|-------|----------------|-------|-------|--------------|--------|-----------|----------|---------|-------|
| sj        | ectrometer R. | A-915+     | ⊦ for dire | ct d  | etermination   | of 1  | nercu | ury in air a | and co | mplex m   | atrix sa | imples, | Fuel  |
| Р         | ocess. Techno | ol., 85, 4 | 473–485,   | http  | os://doi.org/d | oi:10 | 0.101 | 6/j.fuproc.2 | 2003.1 | 1.003, 20 | 004.     |         |       |

- Sommar, J., Zhu, W., Lin, C.-J. and Feng, X.: Field Approaches to Measure Hg Exchange Between Natural Surfaces and the Atmosphere A Review, Crit. Rev. Environ. Sci. Technol. 43, 1657–1739, https://doi.org/10.1080/10643389.2012.671733, 2013a.
- Sommar, J., Zhu, W., Shang, L., Feng, X. and Lin, C.-J.: A whole-air relaxed eddy accumulation measurement
   system for sampling vertical vapour exchange of elemental mercury, Tellus Ser. B-Chem. Phys.
   Meteorol., 65, 19940, https://doi.org/10.3402/tellusb.v65i0.19940, 2013b.
- Sprovieri, F., Pirrone, N., Bencardino, M., D'Amore, F., Carbone, F., Cinnirella, S., Mannarino, V., Landis, M., Ebinghaus, R., Weigelt, A., Brunke, E.-G., Labuschagne, C., Martin, L., Munthe, J., Wängberg, I., Artaxo, P., Morais, F., Barbosa, H. D. M. J., Brito, J., Cairns, W., Barbante, C., Diéguez, M. D. C., Garcia, P.E., Dommergue, A., Angot, H., Magand, O., Skov, H., Horvat, M., Kotnik, J., Read, K. A., Neves, L. M., Gawlik, B. M., Sena, F., Mashyanov, N., Obolkin, V., Wip, D., Feng, X. B., Zhang, H., Fu, X., Ramachandran, R., Cossa, D., Knoery, J., Marusczak, N., Nerentorp, M. and Norstrom, C.: Atmospheric mercury concentrations observed at ground-based monitoring sites globally distributed in the framework of the GMOS network, Atmos. Chem. Phys., 16, 11915–11935, https://doi.org/10.5194/acp-16-11915-2016, 2016.
  - Stannard, D. I.: A theoretically based determination of bowen-ratio fetch requirements, Bound.-Lay. Meteorol., 83, 375–406, https://doi.org/10.1023/A:1000286829849, 1997.
  - Swinbank, W.C.: The measurement of vertical transfer of heat and water vapor by eddies in the lower atmosphere, J. Meteor., 8, 135–145, https://doi.org/10.1175/1520-0469(1951)008<0135:TMOVTO>2.0.CO;2, 1951.
    - UN Environment, 2013. Global Mercury Assessment 2013: Sources, Emissions, Releases and Environmental Transport. UNEP Chemicals Branch, Geneva, Switzerland
    - UN Environment, 2019. Global Mercury Assessment 2018. UN Environment Programme, Chemicals and Health Branch Geneva, Switzerland.
- 795 Wang, X., Lin, C.-J., Yuan, W., Sommar, J., Zhu, W. and Feng, X.: Emission-dominated gas exchange of elemental mercury vapor over natural surfaces in China, Atmos. Chem. Phys., 16, 11125–11143, https://doi.org/10.5194/acp-16-11125-2016, 2016.
  - Watras, C. J., Back, R. C., Halvorsen, S., Hudson, R. J. M., Morrison, K. A. and Wente, S. P.: Bioaccumulation of mercury in pelagic freshwater food webs, Sci. Total. Environ., 219, 183–208, https://doi.org/10.1016/S0048-9697(98)00228-9,1998.
    - Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour transfer, Q. J. Roy. Meteor. Soc., 106, 85–100, https://doi.org/10.1002/qj.49710644707, 1980.
    - Werle, P., Mücke, R. and Slemr, F.: The limits of signal averaging in atmospheric trace-gas monitoring by



805

810

815



- tunable diode-laser absorption spectroscopy (TDLAS), Appl. Phys. B, 57, 131–139, https://doi.org/10.1007/BF00425997, 1993.
  - Wienhold, F. G., Fischer, H. and Harris, G. W.: Fast response tunable diode laser spectroscopy for trace gas flux measurements, Infrared Phys. Techn., 37, 67–74, https://doi.org/10.1016/1350-4495(95)00114-X, 1996.
- Yuan, W., Sommar, J., Lin, C.-J., Wang, X., Li, K., Liu, Y., Zhang, H., Lu, Z., Wu, C. and Feng, X.: Stable Isotope Evidence Shows Re-emission of Elemental Mercury Vapor Occurring after Reductive Loss from Foliage. Environ. Sci. Technol., 53, 651–660, https://doi.org/10.1021/acs.est.8b04865, 2019.
  - Zeeman, M. J., Hiller, R., Gilgen, A. K., Michna, P., Plüss, P., Buchmann, N., and Eugster, W.: Management and climate impacts on net CO<sub>2</sub> fluxes and carbon budgets of three grasslands along an elevational gradient in Switzerland, Agr. Forest Meteorol., 150, 519–530, https://doi.org/10.1016/j.agrformet.2010.01.011, 2010.
  - Zhang, H., Lindberg, S. E., Marsik, F. J. and Keeler, G. J.: Mercury Air/Surface Exchange Kinetics of Background Soils of the Tahquamenon River Watershed in the Michigan Upper Peninsula, Water Air Soil Poll., 126, 151–169, https://doi.org/10.1023/A:1005227802306, 2001.
- Zhang, L., Wu, Z., Cheng, I., Wright, L. P., Olson, M. L., Gay, D. A., Risch, M. R., Brooks, S., Castro, M. S.,
   Conley, G. D., Edgerton, E. S., Holsen, T. M., Luke, W., Tordon, R. and Weiss-Penzias, P.: The estimated six-year mercury dry deposition across North America, Environ. Sci. Technol., 50, 12864–12873, https://doi.org/10.1021/acs.est.6b04276, 2016.
  - Zheng, W., Obrist, D., Weis, D. and Bergquist, B. A.: Mercury isotope compositions across North American forests, Global Biogeochem. Cy., 30, 1475-1492, https://doi.org/10.1002/2015GB005323, 2016.
- 825 Zhu, W., Sommar, J., Lin, C.-J. and Feng, X.: Mercury vapor air-surface exchange measured by collocated micrometeorological and enclosure methods – Part I: Data comparability and method characteristics, Atmos. Chem. Phys., 15, 685–702, https://doi.org/10.5194/acp-15-685-2015, 2015a
  - Zhu, W., Sommar, J., Lin, C.-J. and Feng, X.: Mercury vapor air–surface exchange measured by collocated micrometeorological and enclosure methods – Part II: Bias and uncertainty analysis, Atmos. Chem. Phys., 15, 5359–5376, https://doi.org/10.5194/acp-15-5359-2015, 2015b.
  - Zhu, W., Lin, C.-J., Wang, X., Sommar, J., Fu, X. and Feng, X.: Global observations and modeling of atmosphere–surface exchange of elemental mercury: a critical review, Atmos. Chem. Phys., 16, 4451– 4480, https://doi.org/10.5194/acp-16-4451-2016, 2016.
- Zhu, W., Li, Z., Li, P., Yu, B., Lin, C.-J., Sommar, J. and Feng, X.: Re-emission of legacy mercury from soil
   adjacent to closed point sources of Hg emission, Environ. Poll., 242, 718–727, https://doi.org/10.1016/j.envpol.2018.07.002, 2018.







Figures and captions

**Figure 1:** (a) Schematic of the experimental setup at Chamau (CH-Cha) research site with exact location of topsoil samples for total Hg analysis (n=22) and Eddy covariance (EC) flux measurements of  $Hg^0$ ,  $CO_2$  and  $H_2O$  conducted between 20 July and 6 September 2018. (b) Footprint contour lines of 10 % to 90 % in 10 % steps representing the flux source area during our measurement period. Numbers indicate the distance in meter from the EC station (black cross). The footprint was calculated applying the footprint model presented in Kljun et al. (2015). The figure is a direct output from the online tool: http://footprint.kljun.net/.



**Figure 2:** Examples of raw data time series over a 1 hour data segment, (a, c, e) during period 1 with 24 hour calibrations only (21 July 2018, 10:00–11:00), and (b, d, f) during period 4 with 4 min instrument calibration intervals (6 September 2018, 01:00–02:00). Vertical wind speed (a, b) was not detrended. Hg<sup>0</sup> concentrations are shown before (c, d) and after detrending (e, f). While the 4 min calibration intervals clearly reduce the longer-term drift (d) compared to daily calibrations (c), the gaps during calibrations had to be filled by linear interpolation before calculating fluxes.







**Figure 3:** Allan variance plot using 14 h of continuous measurements in the laboratory (zero-flux experiment), starting 18 July 2018, 18:00. (a) Raw time series, (b) Allan variance as a function of integration time.



**Figure 4:** Example spectrum (a) of  $Hg^0$  fluctuation measurements and (b) cospectrum of the 1 hour averaged  $Hg^0$  flux, 31 August 2018 14:00–15:00. The power spectrum (panel a) before (red line) and after detrending (black line) is shown, and the theoretical slopes in the inertial subrange are shown for ideal conditions (i, solid line,  $f^{2/3}$  slope), for a rectangular oversampling at frequencies > 1 Hz (r, broken line,  $f^{-1}$  slope), and for a first-order damped spectrum (d, dashed line,  $f^{8/3}$  slope). The approximated white noise level is shown with a color band (w,  $f^{+1}$  slope). The flux cospectrum (b) shows absolute values of cospectral densities with black symbols denoting positive contributions to  $w'\chi'$ , and red symbols denoting negative contributions. The light green bold line is a local polynomial regression fit to the data points, whereas the blue line denotes an idealized cospectrum. The two dashed blue lines show damped cospectra with a damping constant of 0.1 and 0.3 s.







**Figure 5:** Cross-correlation analysis to determine the time lag between vertical wind speed (w) and Hg<sup>0</sup> time series ( $\chi$ ). (a,b) Example with a clearly positive Hg<sup>0</sup> flux (21 July 2018, 10:00–11:00), and (c,d) with a marginally positive flux (6 September 2018, 01:00–02:00). Panels (a) and (c) show the cross-correlation within a time lag window of ±10 s, and (b,d) zoom in to the search window used in this study (vertical red band). The blue horizontal band shows the range of zero-fluxes (cross-correlation  $r_{w,\chi} \neq 0$  with  $p \ge 0.05$ ).



Figure 6: Flux detection limit empirical probability distribution of the magnitude of flux measurements under outdoor conditions (black line). The boxplot insert shows the range of the magnitude of measured fluxes during the zero-flux experiment in the laboratory without  $Hg^0$  sources under very low turbulence conditions. The black line shows the theoretical detection limit based on the statistical significance (p < 0.05) of the correlation coefficient between vertical wind speed and  $Hg^0$  fluctuations.







**Figure 7:** Hourly aggregated diel cycle of (a)  $Hg^0$  fluxes and (b) simultaneously recorded CO<sub>2</sub> fluxes and fluxes measured in 2017 for the same period. Each hour of day represents the quantiles obtained from a three-hour window centered at the respective hour of all technically valid observations. The bold black line represents median flux values. The interquartile range (IQR) is the range of the middle 50 % of the data. The 70 % and 80 % confidence intervals (CI) and the number of measurements per hour (n) are given. The median CO<sub>2</sub> flux in 2017 is displayed (red bold dashed line). The IQR (red vertical lines) and 70 % CI (lightered vertical lines) are indicated.



**Figure 8:** Boxplots display (a) the total topsoil Hg concentration (0-10 cm) in the northern and the southern parcels as well as the Hg<sup>0</sup> flux over the respective parcels (b) during the night (21:00–05:00) and (c) during the day (10:00–17:00).