Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2019-287-SC1, 2019 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.





Interactive comment

Interactive comment on "Aerosol retrievals from the ACEPOL Campaign" *by* Guangliang Fu et al.

Sergey Korkin

sergey.v.korkin@nasa.gov

Received and published: 13 August 2019

The 'alpha * Rpol' part in Eq.(1), p.4, seems unclear to me. Please provide explicit form for the Rpol matrix like it is done for the RPV-part. Reference to the Maignan et al, 2009 paper gives little help:

1) Sections 3.2, 3.3, and 3.5 in Maignan et al. 2009 discuss different models - which particular one was used in the paper under review?

2) As far as i know, Maignan et al. 2009 does not define the Mueller matrix of the surface - only BPDF, which is based on the F12=F21=Fp element of the Fresnel matrix. To get the Mueller matrix, shall one compute matrix exponential of the Fresnel matrix, or, vice versa, create a matrix of "scalar" exponents of elements of the Fresnel matrix?

3) The complete 4x4 (or reduced 3x3 if V is ignored) Mueller matrix of the surface is



Discussion paper



required only to simulate the surface reflection of diffuse radiation (including multiple bouncing of light between the atmosphere and surface). How strong and important is that effect for polarization components?

4) Is 'alpha' in Eq.(1) band-dependent?

Thank you!

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2019-287, 2019.



Interactive comment

Printer-friendly version

Discussion paper

