

Supplementary Information for

**Retrieval of aerosol optical properties over Beijing: a comparison
between SKYNET and AERONET**

Xianyi Yang^{1,2}, Huizheng Che², Hitoshi Irie³, Quanliang Chen¹, Ke Gui², Ying Cai³,
Yu Zheng², Linchang An⁴, Hujia Zhao², Lei Li², Yuanxin Liang², Yaqiang Wang²,
Hong Wang², Xiaoye Zhang²

¹Plateau Atmospheric and Environment Laboratory of Sichuan Province, College of Atmospheric Science, Chengdu University of Information Technology, Chengdu 610225, China

²State Key Laboratory of Severe Weather (LASW), Institute of Atmospheric Composition, Chinese Academy of Meteorological Sciences, Beijing 100081, China

³Center for Environmental Remote Sensing, Chiba University, Chiba, Japan

⁴National Meteorological Center, CMA, Beijing 100081, China

Correspondence to: Huizheng Che (chehz@cma.gov.cn)

This file includes:

Figure S1 to S4

Supplementary Figure captions:

Figure S1: Comparison of SSA values retrieved from SKYNET (SR-CEReS) with SSA values retrieved after adjusting the SVA by a 1% step size at the 400-, 500- and 670-nm channels. Panels (a1–a3) and (b1–b3) show the decreases and increases in the SVA, respectively.

Figure S2: Comparison of SSA values retrieved from SKYNET (SR-CEReS) with SSA values retrieved after adjusting the SA by a 0.01 step size at the 400-, 500- and 670-nm channels. Panels (a1–a3) and (b1–b3) show the decreases and increases in the SA, respectively.

Figure S3: Daily averaged wind fields at (a) 850 hpa and (b) the surface in Beijing from 27 December 2016 to 9 January 2017 based on the ERA-Interim data.

Figure S4: Vertical variation of aerosol subtypes from the CALIPSO satellite products on 30 December 2016.




Figure S1: Comparison of SSA values retrieved from SKYNET (SR-CEReS) with SSA values retrieved after adjusting the SVA by a 1% step size at the 400-, 500- and 670-nm channels. Panels (a1–a3) and (b1–b3) show the decreases and increases in the SVA, respectively.





Figure S2: Comparison of SSA values retrieved from SKYNET (SR-CEReS) with SSA values retrieved after adjusting the SA by a 0.01 step size at the 400-, 500- and 670-nm channels. Panels (a1–a3) and (b1–b3) show the decreases and increases in the SA, respectively.






Figure S3: Daily averaged wind fields at (a) 850 hpa and (b) the surface in Beijing from 27 December 2016 to 9 January 2017 based on the ERA-Interim data.




Figure S4: Vertical variation of aerosol subtypes from the CALIPSO satellite products on 30 December 2016.